

BIOLOGY**BOOKS - OBJECTIVE NCERT AT YOUR FINGERTIPS BIOLOGY****CELL CYCLE AND CELL DIVISION****CELL CYCLE**

1. Amitosis usually occurs in

- A. eukaryotic cells
- B. prokaryotic cells
- C. meristems
- D. spore mother cells.

Answer: B

 **Watch Video Solution**

2. Which one is the correct sequence of a cell cycle?

- A. $G_2 \rightarrow M \rightarrow G_1 \rightarrow S$
- B. $S \rightarrow G_2 \rightarrow M \rightarrow G_1$


C. $G_1 \rightarrow S \rightarrow G_2 \rightarrow M$

D. $M \rightarrow G_1 \rightarrow S \rightarrow G_2$

Answer: C

[▶ Watch Video Solution](#)

3. Identify A,B,C and D in the given diagram depicting cell cycle and select the correct option.



A. $\{(A,B,C,D),(G_0),G_1,S,G_2)\}$

B. $\{(A,B,C,D),(G_1),S,G_2,G_0)\}$

C. $\{(A,B,C,D),(G_1),G_0,S,G_2)\}$

D. $\{(A,B,C,D),(S,G_0),G_1,G_2)\}$

Answer: A

 [Watch Video Solution](#)

4. Yeast cell divide once in approximately every

A. 90 minutes

B. 9 minutes

C. 24 hours

D. 24 secodns.

Answer: A

 [Watch Video Solution](#)

5. Human cells in culture show a cell cycle to be completed in approximately.

A. 42 hours

B. 24 hours

C. 24 minutes

D. 24 seconds.

Answer: B

 [Watch Video Solution](#)

6. Which phase occupies the maximum part of cell cycle?

- A. Mitrotic phase
- B. Meiotic phase
- C. Interphase
- D. Cytokinesis

Answer: C

 [Watch Video Solution](#)

7. This phase of cell cycle is a period of intense synthesis and growth. It constitutes 95% of the duration of cell cycle. It is

- A. interphase
- B. telophase
- C. prophase
- D. anaphase.

Answer: A

8. Read the following statements about cell division and select the correct ones.

- (i) M phase represents the phase when actual cell division occurs and I phase represents the phase between two successive M phases.
- (ii) In the 24 hours average duration of cell cycle of a human cell, cell division proper lasts for only about an hour.
- (iii) M phase constitutes more than 95% of the duration of cell cycle.

- A. i and ii
- B. ii and iii
- C. i and iii
- D. i,ii and iii

Answer: A

9. Which of the following phases of the cell cycle is not a part of interphase?

- A. S
- B. G_1
- C. G_0
- D. M

Answer: D

 [Watch Video Solution](#)

10. A cell cycle includes

- A. interphase and M phase
- B. prophase, metaphase, anaphase and telophase
- C. `G\_(1),S` and `G\_(2)` phases
- D. Karyokinesis and cytokinesis.

Answer: A

 [Watch Video Solution](#)

11. In which stage DNA replication takes place ?

- A. Metaphase
- B. `G\_(1)` phase
- C. S-phase
- D. `G\_(2)` phase

Answer: C

 [Watch Video Solution](#)

12. Synthesis of histone proteins occurs in

- A. `G\_(1)` phase
- B. interphase
- C. anaphase
- D. `G\_(0)` phase.

Answer: B

 [Watch Video Solution](#)

13. Select the incorrect statement regarding S phase of interphase.

- A. It occurs between `G\_(1)` and `G\_(2)`
- B. DNA replicates in the nucleus in this phase.
- C. Centrioles duplicate in the cytoplasm.
- D. As DNA doubled, number of chromosomes also doubles.

Answer: D

 [Watch Video Solution](#)

14. The DNA content of individual cells and the number of cells in each phase of a cell cycle" can be determined using flow cytometry. Which of the following combinations of "phase of a cell cycle and its corresponding DNA content" can be considered normal?

- (i) Diploid cells found in the `G\_(0)` or `G\_(1)` phase.

(ii) Cells with twice the normal DNA content in the early M phase.

(iii) Cells with twice the normal DNA content in the `G\_(2)` phase.

- A. i and ii
- B. ii and iii
- C. iii and iv
- D. i,ii,iii and iv

Answer: D

 [Watch Video Solution](#)

15. Cell would normally proceed to mitosis without interruption

- A. once it has entered the S phase
- B. once it has entered the `G\_(2)` phase
- C. at any time during cell division activity
- D. none of these

Answer: A

 [Watch Video Solution](#)

16. The cells that do not divide further, exit `G\_(1)` phase to enter an inactive stage called__ of the cell cycle.

- A. M stage
- B. `G\_(2)` stage
- C. S stage
- D. `G\_(0)` stage

Answer: D

 [Watch Video Solution](#)

17. Which of the following statements is correct ?

- A. Animals can show mitotic divisions in both haploid and diploid cells.
- B. After S phase the number of chromosomes becomes double i.e., $2n$ to $4n$.
- C. During the `G\_(2)` phase, proteins are synthesised in preparation for mitosis while cell growth continues.
- D. S or synthesis phase marks the period during which RNA synthesis takes place.

Answer: C

 [Watch Video Solution](#)

18. In which of the following stages, a chromosome is minimum coiled?

- A. interphase
- B. Metaphase

C. Prophase

D. anaphase.

Answer: A

 [Watch Video Solution](#)

19. Which of the following statements is correct regarding `G\_(0)` phase?

A. Mitogens are present in `G\_(0)` phase.

B. Mitogens are present but energy rich compounds are absent.

C. Both mitogens and energy rich compounds are present.

D. Neither mitogens nor energy rich compounds are present.

Answer: D

 [Watch Video Solution](#)

MITOSIS

1. Best material for studying mitosis in laboratory is

A. (a) leaf tip

B. (b) shoot tip

C. (c) root tip

D. (d) gamete

Answer: C

 [Watch Video Solution](#)

2. Mitotic spindle is mainly composed of ___ proteins.

A. tublin

B. myosin

C. actin

D. actomyosin

Answer: A

 [View Text Solution](#)

3. Microtubules are absent in

A. mitochondria

B. flagella

C. spindle fibres

D. centriole.

Answer: A

 [View Text Solution](#)

4. Centrosome undergo duplication during (i) of (ii) and begin to move towards opposite poles of the cell during (iii) state of (iv).

- A. `{"(i)","(ii)","(iii)","(iv)","S phase","Interphase","Prophase","Mitosis"}:`
- B. `{"(i)","(ii)","(iii)","(iv)","S phase","Interphase","Anaphase","Mitosis"}:`
- C. `{"(i)","(ii)","(iii)","(iv)","Prophase","Mitosis","Metaphase","Mitosis"}:`
- D. `{"(i)","(ii)","(iii)","(iv)","Prophase","Mitosis","Anaphase","Mitosis"}:`

Answer: A

 [View Text Solution](#)

5. ____ is the best stage to count the number and study the morphology of chromosomes.

- A. Prophase
- B. metaphase
- C. Anaphase
- D. Telophase

Answer: B

 [Watch Video Solution](#)

6. _____ is characterised by all the chromosomes coming to lie at the equator, with one chromatid connected by its kinetochore to spindle fibres from one pole and its sister chromatid connected by its kinetochore to spindle fibres from the opposite pole.

- A. Prophase
- B. Metaphase
- C. Anaphase
- D. Telophase

Answer: B

 [Watch Video Solution](#)

7. Read the following statements.

- (i) In mitotic cell division chromosome number is halved.
- (ii) Centromere is the point where two sister chromatids are held together.
- (iii) The period between two successive mitotic divisions is known as telophase.
- (iv) In `G<sub>1</sub>` phase of cell cycle protein and RNA are synthesised.

Which of the above given statements are correct?

- A. a) i and iii only
- B. b) ii and iii only
- C. c) i and iv only
- D. d) ii and iv only

Answer: D

 [Watch Video Solution](#)

8. You are provided with floral buds of Chrysanthemum in your class and are asked to count the chromosomes, then which of the following stages would you prefer to look into?

- A. Prophase
- B. Metaphase
- C. anaphase
- D. Interphase

Answer: B

 [Watch Video Solution](#)


9. During karyokinesis, the spindle fibres get attached to condensing chromosome at a highly differentiated region. This region is called as

- A. chromomere
- B. chromocentre
- C. centriole
- D. kinetochore.

Answer: D

 [Watch Video Solution](#)

10. Identify the structure indicated by labels (i),(ii),(iii) and (iv) and select the correct option.



A. a) i-Chromatid, ii-Centriole,
iii-Centromere, iv-Chromosome

B. b) i-Chromosome, ii-Centriole,
iii-Centromere, iv-Chromatid

C. c) i-Chromatid, ii-Centromere,
iii-Centriole, iv-Chromosome

D. d) i-Chromosome, ii-Centromere,
iii-Centriole, iv-Chromatid

Answer: A

11. Which of the following statements is not correct regarding colchicine?

- A. It prevents assembly of microtubules.
- B. It inhibits chromosome replication.
- C. It is an alkaloid.
- D. It is called as mitotic poison.

Answer: B

 [Watch Video Solution](#)

12. Colchicine is a cell poison which arrests cell division at __ and can induce ____

- A. metaphase, parthenocarphy
- B. anaphae, parthenocarpy
- C. metaphase, polyploidy
- D. anaphase, polyploidy

Answer: C

 [Watch Video Solution](#)

13. Spindle formation can be distrupted by exposing cell to the microtubule poison such as

- A. high concentration of oxygen
- B. vitamin A
- C. cholesterol
- D. colchicine.

Answer: D

 [Watch Video Solution](#)

14. At which stage of mitosis, the two daughter chromatids separate from each other, migrate towards the opposite poles and are now referred to as chromosomes of the future daughter nuclei?

- A. Prophase
- B. Metaphase
- C. Anaphase
- D. Telophase

Answer: C

 [Watch Video Solution](#)

15. During anaphasic movements of chromosomes, ____ of each chromosome is/are towards the pole and ____ of the chromosome trail(s) behind.

- A. centromere, arms

- B. arms, centromere
- C. chromatids, centromere
- D. none of these

Answer: A

 [Watch Video Solution](#)

16. Select the correct match

- A. Quiescent phase-`G\_(2)` phase
- B. Synthesis phase-`G\_(1)` phase
- C. Centromere splitting -Anaphase
- D. Chromosomal condensation-Telophase

Answer: C

 [Watch Video Solution](#)

17. The separation of two chromatids of each chromosome at early anaphase is initiated by

- A. (a) the interaction of centromere with the chromosomal fibres
- B. (b) the elongation of metaphasic spindle
- C. (c) the force of repulsion between the divided kinetochores
- D. (d) all of these

Answer: C

 [Watch Video Solution](#)

18. Match column I with column II and select the correct option from the given codes.

`{("Column I","Column II"),("V-shaped at anaphase",(ii)"Acrocentric chromosome"),("L-shaped at anaphase",(ii)"Metacentric chromosome"),("J-shaped at anaphase",(iii)"Telocentric chromosome"),("I-shaped at anaphase",(iv)"Sub-metacentric chromosome");}`

A. iv,ii,i,iii

B. ii,iv,i,iii

C. ii,iv,iii,i

D. iv,iii,ii,i

Answer: B

 [Watch Video Solution](#)

19. Which of the following is key event of anaphase of mitotic division?

A. Chromosomes are moved to spindle equator and get aligned through spindle fibres to both poles.

B. Centromeres split and chromatids separate.

C. Chromosomes cluster at opposite spindle poles and their identity is lost as discrete elements.

D. Both b and c

Answer: B

 [View Text Solution](#)

20. Select the incorrect statement regarding mitotic cell division.

{:(i)"Prophase",-,"Chromosomes begin to uncoil"),((ii)"Metaphase",-,"Chromatids move apart"),
((iii)"Telophase",-,"The nuclear membrane reappears"),((iv)"Late",-,"Each chromosome consists of two"),
((v)"Interphase",-,"Chromosomes are not distinct")}:`

- A. ii and iv only
- B. i and iii only
- C. ii, iv and v only
- D. i and v only

Answer: A

 [View Text Solution](#)


21. What is true about telophase stage of mitosis?

- A. Chromosomes lose their identity as discrete elements.
- B. Chromosomes cluster at opposite spindle poles.
- C. Nuclear envelope, nucleolus, Golgi complex and ER reform.
- D. All of these


Answer: D

 [View Text Solution](#)

22. Refer to the given figures and answer




A



B



C



D

Identify the given stages of mitosis and select the correct option

A. `{: (A,B,C,D), ("Prophase", "Metaphase", "Telophase", "Anaphase") :}`

B. `{: (A,B,C,D), ("Metaphase", "Anaphase", "Prophase", "Telophase") :}`


C. `{: (A,B,C,D), ("Anaphase", "Metaphase", "Prophase", "Telophase") :}`

D. `{: (A,B,C,D), ("Prophase", "Metaphase", "Anaphase", "Telophase") :}`


Answer: C

 [View Text Solution](#)

23. Refer to the given figures and answer



A



B



C



D

Which of the following shown the correct sequence of the given mitotic stages?

A. `DrarrCrarrBrarrA`

B. `CrarrBrarrDrarrA`

C. `BrarrArarrCrarrD`


D. `CrarrBrarrArarrD`

Answer: D




Watch Video Solution

24. Refer to the given figures and answer




A



B



C



D

At which of the given stages of mitosis, chromosomes appear in V,L,J and I shapes.

A. A

B. B


C. C

D. D


Answer: A

 Watch Video Solution

25. Refer to the given figures and answer



A



B



C



D

Read the given statements which represent the features of the figures A,B,C and D Match them correctly and select the correct option.

- (i) Chromosomes appear like a ball of wool (spireme stage)
- (ii) Reformation of nuclear envelope, nucleolus, Golgi complex and ER
- (iii) Formation of equatorial plate
- (iv) Splitting of centromeres

A. iv,iii,i,ii

B. iii,iv,i,ii

C. ii,iii,i,iv

D. iv,ii,iii,i

Answer: A

 [View Text Solution](#)

26. Match the column I with column II and select the correct option from the given codes.

{:("Column I","Column II"),("Disintegration of nuclear membrane",(i)"Anaphase"),("Appearance of

nucleolus", (ii) "Prophase"), ("Division of centromere", (iii) "Telophase"), ("Replication of DNA", (iv) "S-phase")):

A. ii, iii, i, iv

B. ii, iii, iv, i

C. iii, ii, i, iv

D. iii, ii, iv, i

Answer: A

 [Watch Video Solution](#)

27. Which phase of mitosis is essentially the reverse of prophase in terms of nuclear changes?

A. S-phase

B. Anaphase


C. Telophase

D. Interphase

Answer: C

 [Watch Video Solution](#)

28. The given graph shows the change in DNA content during various phases (A to D) in a typical mitotic cell cycle. Identify the phases and select the correct option.



A. $\{(A,B,C,D),(G_2,G_1,S,M):\}$

B. $\{(A,B,C,D),(G_2,S,G_1,M):\}$

C. $\{(A,B,C,D),(G_1,S,G_2,M):\}$

D. $\{(A,B,C,D),(M,G_1,S,G_2):\}$

Answer: C

[Watch Video Solution](#)

29. A cell's division time is 1 minute. In 20 minutes, a culture tube (culture medium) is $\frac{1}{8}^{th}$ filled with cells. When the tube will be fully filled?

A. (a) 21 minutes

B. (b) 23 minutes

C. (c) 60 minutes

D. (d) 160 minutes

Answer: B

 [Watch Video Solution](#)

30. Cytokinesis in an animal cell takes place by ____ method in ____ direction; while in a plant cell it occurs by _____method in ____ direction.

A. (a) furrowing, centrifugal, cell plate, centripetal

B. (b) furrowing, centripetal, cell plate, centrifugal

C. (c) cell plate, centrifugal, furrowing, centripetal

D. (d) cell plate, centripetal, furrowing, centrifugal

Answer: B

 [Watch Video Solution](#)

31. If a tissue has at a given time 1024 cells, how many cycles of mitosis had the original paraental single cell undergone?

A. 512

B. 10

C. 1024

D. 256

Answer: B

 [Watch Video Solution](#)

32. Spindle usually persists in the form of ____ during ____ method of cytokinesis.

- A. phragmoplast, cleavage
- B. phragmoplast, cell plate
- C. cell plate, cell plate
- D. cell plate, cleavage

Answer: B

 [Watch Video Solution](#)

33. Match column I with column II and select the correct option from the given codes.


`{:(("Column I","Column II"),("Division of nucleus",(i)"Interphase"),("Division of cytoplasm",(ii)"Cytokinesis"),("DNA replication",(iii)"Syncytium"),("Karyokinesis not followed by cytokinesis",(iv)"Karyokinesis"):}`

- A. a.ii,b.iv,c.i,d.iii
- B. a.iv,b.ii,c.i,d.iii
- C. a.iv,b.ii,c.iii,d.i
- D. a.iii,b.ii,c.iv,d.i

Answer: B

 [Watch Video Solution](#)

34. The given diagram depicts cell plate method of cytokinesis in plant cells. Identify A,B and C



A. `:(A,B,C),("Daughter nucleus","Phramgmoplast","Vesicles").:`

B. `:(A,B,C),("Daughter nucleus","Vesicles","Phragmoplast").:`

C. `{: (A,B,C), ("Parent nucleus", "Vesicles", "Phragmoplast") :}`

D. `{: (A,B,C), ("Parent nucleus", "Phragmoplast", "Vesicles") :}`

Answer: A

 [Watch Video Solution](#)

35. Phragmoplast is related to

- A. division of nucleolus
- B. cell elongation
- C. Cytokinesis
- D. assemblage of chromonomes at metaphase.

Answer: C

 [Watch Video Solution](#)

36. The role of mitosis is not merely to divide a cell into two daughter cells but to ensure genetic continuity from one cell generation to another cell generation. The mechanism ensuring genetic continuity is

- A. formation of cells with new chromosomes
- B. formation of two daughter cells
- C. formation of two cells with identical DNA
- D. halving the chromosome number between the two new cells.

Answer: C

 [Watch Video Solution](#)

37. Chromosome duplication without nuclear division refers

- A. meiosis
- B. mitosis
- C. androgenesis
- D. endomitosis.

Answer: D

 [Watch Video Solution](#)

MEIOSIS

1. Meiosis consists of

- A. two cell divisions without any DNA replication
- B. two cell divisions in which chromosome number is reduced to half
- C. two cell divisions with only two rounds of chromosome replication
- D. a single cell division with chromosome replication.

Answer: B

 [Watch Video Solution](#)

2. Which of the following is not the feature of meiosis?

- A. Meiosis involves two sequential cycles of nuclear and cell division, meiosis I and meiosis II but only a single cycle of DNA replication.
- B. Meiosis I is initiated after the parental chromosomes have replicated to produce identical sister chromatids at the S-phase.
- C. Meiosis involves pairing of non-homologous chromosomes and recombination between them.
- D. Four haploid cells are formed at the end of meiosis II.

Answer: C

 [Watch Video Solution](#)

3. Best material to study meiosis is

- A. root tip
- B. ovary
- C. young anther
- D. pollen grain.

Answer: C

 [Watch Video Solution](#)

4. The term "meiosis" was given by

- A. Johanssen
- B. Knoll and Ruska
- C. A. Flemming
- D. Farmer and Moore.

Answer: D

 [Watch Video Solution](#)

5. At which of the following stages, the chromosomes appear single, thin and thread like?

- A. Leptotene
- B. Zygotene
- C. Pachytene
- D. Diplotene

Answer: A

 [Watch Video Solution](#)

6. Zygotene of prophase-I is characterised by

- A. chromomeres

- B. synaptonemal complex
- C. crossing over
- D. terminalisation of chiasmata.

Answer: B

 [Watch Video Solution](#)


7. Crossing over in diploid organisms is responsible for

- A. dominance of genes
- B. linkage between genes
- C. segregation of alleles
- D. recombination of alleles.

Answer: D

 [Watch Video Solution](#)

8. Which of the following is correct regarding the given figure?



A. 3 6 12

B. 3 12 6

C. 6 6 12

D. 6 12 6

Answer: B

 [Watch Video Solution](#)

9. The members of a homologous pair of chromosomes

A. are identical in size and appearance

B. contain identical genetic information

C. separation and move to opposite poles of the cell during mitosis

D. are found only in haploid cells.

Answer: A


[Watch Video Solution](#)

10. Which of the following correctly shows a pair of homologous chromosomes at the start of meiosis?




A.

(b)




B.

(c)



C.

(d)



D.

Answer: D

[▶ Watch Video Solution](#)

11. Which of the following is correct about bivalent?

- (i) Bivalents are tetrads.
- (ii) A bivalent means 4 chromatids and 2 centromeres.
- (iii) One bivalent consists of 2 homologous chromosomes.
- (iv) Bivalents form in zygotene.

A. i,ii,iii and iv

B. iii only

C. iii and iv

D. iv only

Answer: A

 [Watch Video Solution](#)

12. The process of crossing over is assisted by which of the following enzymes?

- A. Endonuclease
- B. Polymerase
- C. Ligase
- D. Both a and c

Answer: D

 [Watch Video Solution](#)

13. To build up food reserves in the cytoplasm, chromosomes become unfolded to start transcription of mRNA and rRNA, during which phase of meiosis I ?

- A. Diakinesis
- B. Zygotene
- C. Diplotene
- D. Leptotene

Answer: C

 [Watch Video Solution](#)

14. At which stage, the homologous chromosomes separate due to repulsion, but are yet held by chiasmata?

- A. Zygotene
- B. Pachytene
- C. Diplotene
- D. Diakinesis

Answer: C

 [Watch Video Solution](#)

15. Diplotene stage of prophase-I is characterised by

- A. dissolution of synaptonemal complex
- B. separation of synapsed homologous chromosomes except at the site of crossovers
- C. formation of X-shaped structures called chiasmata
- D. all of these

Answer: D

 [Watch Video Solution](#)

16. (i) Thin thread like chromosomes with a beaded appearance

(ii) Appearance of recombination nodules

(iii) Formation of bivalents/tetrads

(iv) Terminalisation of chiasmata

(v) Appearance of chiasmata

Identify the different stages with respect to the above given features and select the correct option.

- A. `{"(i)","(ii)","(iii)","(iv)","(v)","Leptotene","Zygotene","Pachytene","Diplotene","Diakinesis")}`
- B. `{"(i)","(ii)","(iii)","(iv)","(v)","Leptotene","Zygotene","Pachytene","Diakinesis","Diplotene")}`
- C. `{"(i)","(ii)","(iii)","(iv)","(v)","Leptotene","Pachytene","Zygotene","Diakinesis","Diplotene")}`
- D. `{"(i)","(ii)","(iii)","(iv)","(v)","Leptotene","Pachytene","Diplotene","Zygotene","Diakinesis")}`

Answer: C



Watch Video Solution

17. (i) Thin thread like chromosomes with a beaded appearance

(ii) Appearance of recombination nodules

(iii) Formation of bivalents/tetrads

(iv) Terminalisation of chiasmata

(v) Appearance of chiasmata

Arrange the given statements in the correct sequence of their occurrence during prophase I.

- A. `irarriiirarriirarrvarriv`
- B. `irarriirarriiirarrivarrv`
- C. `irarrivarrvvarriiirarrii`
- D. `irarriiirarriirarrivarrv`

Answer: A

 [Watch Video Solution](#)

18. Match the column I with column II and select the option from given codes.

`{("Column I","Column II"),("Synaptonemal complex",(i)"Pachytene"),("Recombination nodule",
(ii)"Zygotene"),("Terminalisation of chiasmata",(iii)"Telophase I"),("Formation of dyad cell",(iv)"Diakinesis")};`

A. ii,i,iv,iii

B. i,ii,iv,iii


C. iii,i,iv,ii

D. ii,i,iii,iv

Answer: A


 [Watch Video Solution](#)

19. Identify the given figures showing meiotic phases and select the correct option.




A

A



B

B



C

C

A. `:(A,B,C),("Metaphase","Anaphase","Telophase");)`

B. `:(A,B,C),("Methaphase I","Anaphase I","Telophase I");)`


C. `:(A,B,C),("Metaphase II","Anaphase II","Telophase II");)`

D. `:(A,B,C),("Anaphase I","Metaphase I","Telophase I");)`

Answer: B


[Watch Video Solution](#)

20. The figure given below shows a cell undergoing meiosis.




Which of the option below shows the next stage in the process?

(a)




A.

(b)




B.

(c)



C.

(d)



D.

Answer: B

[Watch Video Solution](#)

21. Disjunction refers to

- A. the separation of homologous chromosomes at anaphase I
- B. the type of chromosomal aberration in which there is loss of a part of a chromosome
- C. incompatibility in fungi and other thallophytes
- D. modification of gene action by a nonallelic gene.

Answer: A

 [Watch Video Solution](#)

22. Lampbrush chromosomes are seen in which typical stage?

- A. Mitotic anaphase
- B. Mitotic prophase
- C. Mitotic metaphase
- D. Meiotic prophase

Answer: D

 [Watch Video Solution](#)

23. Match column I with column II and select the correct option from the given codes.

`{:(("Column I","Column II"),("Chromosomes move to equator",(i)"Pachytene"),("Centromere splits and ",

(ii)"Zygotene"),("Pairing between homologous chromosomes",(iii)"Anaphase"),("Crossing over between homologous chromosomes",(iv)"Metaphase");}`

A. i,ii,iii,iv

B. ii,iii,iv,i

C. iv,iii,ii,i

D. iii,i,iv,ii

Answer: C



Watch Video Solution

24. During meiosis I,chromosome number

A. is reduced to half

B. doubles up

C. remains the same


D. either a or b

Answer: A



Watch Video Solution

25. Refer to the given stages A,B,C and D of meiosis I and select the incorrect statement regarding them.



A. The last stage of A is diakinesis which is marked by terminalisation of chiasmata.

B. In stage B, microtubules from the opposite poles of the spindle attach to the pair of homologous chromosomes.

C. In stage C, homologous chromosomes separate. While sister chromatids remain associated at their centromeres.

D. In stage D, nuclear membrane and nucleolus disappear, cytokinesis follows and this is called as dyad of cells.

Answer: D

 [Watch Video Solution](#)

26. During meiosis I in humans, one of the daughter cells receives

A. only maternal chromosomes

B. a mixture of maternal and paternal chromosomes

C. same number of chromosomes as present in parent cell

D. none of these.

Answer: B

 [Watch Video Solution](#)

27. The number of chromosomes is reduced to half during

A. mitosis

B. meiosis II

C. Meiosis I

D. fertilisation.

Answer: C

 [Watch Video Solution](#)

28. Four different steps that occur during meiosis are given in the following list.

(i) Complete separation of chromatids

(ii) Pairing of homologous chromosomes

(iii) Lining up of paired chromosomes on equator

(iv) Crossing over between chromatids

Select the correct sequential arrangement of the steps.

A. ii,iii,iv,i

B. iii,ii,iv,i

C. ii,iv,iii,i

D. iii,i,ii,iv

Answer: C

 [Watch Video Solution](#)

29. Select the incorrectly matched pair.

A. Phragmoplast - Persistent spindle

B. Reductional division - Meiosis I

C. Equational division - Meiosis II

D. Crossing over - Non-homologous chromosomes

Answer: D

 [Watch Video Solution](#)

30. Splitting of centromere and hence separation of chromatids occur during

A. anaphase of mitosis

B. anaphase of meiosis I

C. anaphase of meiosis II

D. Both a and c

Answer: D

 [Watch Video Solution](#)


31. If gametes are produced after reduction division, they are termed as

- A. coenogametes
- B. mitogametes
- C. pseudogametes
- D. meiogametes.


Answer: D

 [Watch Video Solution](#)


32. If $2n=4$, then identify the figures A,B and C, as per the following codes and select the correct option



A



B



C

Anaphase of meiosis I=(i)

Anaphase of mitosis =(ii)

Anaphase meiosis II =(iii)

A. ii,i,iii

B. iii,ii,i

C. i,ii,iii

D. iii,i,ii

Answer: A

 [Watch Video Solution](#)

33. What does (I) and (II) represent in the given flowchart? Parent cell $\xrightarrow{(M-I)}$ 2 Daughter cells
 $\xrightarrow{(M-II)}$ 4 Daughter cells

A. (I)=2n

(II)=n

B. (I)=n

(II)=n

C. (I)=n

(II) =2n


D. (i)=2n

(ii)=2n


Answer: B



Watch Video Solution

**(i)**


?

(ii)**(iii)**

In above sequence of figures showing different stages of cell division, the missing stage (ii) is


In above sequence of figures showing different stages of cell division, the missing stage (ii) is

(a)





A.

(b)




B.

(C)



C.

(d)




D.

Answer: A

[Watch Video Solution](#)


35. Consider the given cell at metaphase-I stage undergoing normal meiosis.



Which of the following gametes will not be formed from this cell?


this cell?

(a)




A.

(b)




B.

(c)



C.

(d)



D.

Answer: D

 [Watch Video Solution](#)

36. While in mitosis, the daughter cells resemble each other and also the parent cell, in meiosis they differ not only from parent cell in having half the number of chromosomes. But also differ among themselves qualitatively in genetic constitution due to


- A. segregation and crossing over only
- B. independent assortment and segregation only
- C. independent assortment and crossing over only

D. crossing over, independent assortment and segregation.

Answer: D

[Watch Video Solution](#)

37. Refer to the the given and select the correct statement



A. In stage B homologous chromosomes are interconnected and chromosomes occurs in pairs.

B. Stage A is divisible into five substages


C. In stage D, chormosomes are not enclosed by a nuclear envelope.

D. In stage C centromeres divide and chromosomes are single stranded.


Answer: D

[Watch Video Solution](#)


38. Refer to the given figure.



A



B



C


Identify A,B and C and select the correct option.

- A. `{: (A,B,C), ("Prophase I", "Metaphase I", "Anaphase I") :}`
- B. `{: (A,B,C), ("Metaphase I", "Prophase II", "Anaphase II") :}`
- C. `{: (A,B,C), ("Metaphase I", "Metaphase II", "Anaphase") :}`
- D. `{: (A,B,C), ("Prophase II", "Metaphase II", "Anaphase II") :}`

Answer: B


 [Watch Video Solution](#)

39. Refer to the given figure of cell division.




Which of the following options show previous stage of this process?

(a)




A.

(b)




B.

(c)



C.

(d)



D.

Answer: A

 [Watch Video Solution](#)

40. In which of the following ways are mitosis and meiosis similar?

- A. Both have pairing of homologous chromosomes.
- B. Both are preceded by DNA replication.
- C. Both occur in all kinds of cells.
- D. Both include separation of paired chromomes.

Answer: B

 [Watch Video Solution](#)

41. At what phase of meiosis there are two cells, each with separated sister chromatids that have been moved to opposite spindle poles?

- A. Anaphase II
- B. Anaphase I
- C. Telophase II
- D. Telophase I

Answer: A

 [Watch Video Solution](#)

42. An anther has 1200 pollen grains. How many PMCs must have been there to produce them?

- A. 1200
- B. 300
- C. 150
- D. 2400

Answer: B

 [Watch Video Solution](#)

43. Meiosis does not occur in

- A. bacteria
- B. cyanobacteria
- C. plant cell
- D. both a and b

Answer: D

 [Watch Video Solution](#)

44. In animals meiotic division occurs during gamete formation. This gametic meiosis results in

- A. haplontic life cycle
- B. diplontic life cycle
- C. diplohaplontic life cycle
- D. none of these

Answer: B

 [Watch Video Solution](#)

45. The durations of mitotic stages in two situations. (A and B) are tabulated below.

Phase	Duration of Mitotic Stages (in Minutes)	
	A	B
Interphase	1356(22.6 h)	870 (14.5 h)
Prophase	126	54
Metaphase	24	14
Anaphase	5	3
Telophase	22	11
Total	1533(25.6 h)	952(15.9 h)

Following are some interpretations:

1. 'A' and 'B' indicate the same plant tissue grown at higher and lower temperature respectively.
- II. 'A' indicates a slow growing plant species and 'B' indicates a fast growing plant species.
- III. Both 'A' and 'B' indicate dormant plant tissues with excessively long interphase.

The correct interpretations is/are

- A. i and iii only
- B. II and III
- C. III only
- D. II only

Answer: D





 [Watch Video Solution](#)

46. Given graphs P,Q,R and S show four stages of cell cycle i.e., `G\_(1)`,S,`G\_(2)` and M, but in random order. Identify the stages and match with the activities of the cell.

I. Taxol treatment, which prevent microtubule depolymerization, arrests the cell at this tage .

II. With a mitogen treatment , such as an epidermal growth factor, and arrested cell at this stage proceeds to the next stage of the cell cycle.

III. The cell cycle check point at this stage confirms. that DNA duplication is complete before the cell proceeds to the next stage



A. I-P,II-Q,III-R

B. I-Q,II-S,III-R


C. I-R,II-Q,III-S

D. I-P,II-S,III-Q

Answer: A


47. Which of the following graphs shows the relative change in the amount of mitochondrial DNA of a cell undergoing mitosis?

(a)




A.

(b)




B.

(c)



c.

(d)




D.

Answer: A

[Watch Video Solution](#)


48. The given figure is a schematic break-up of the phases/stages of cell cycle. Select the correct option regarding it.



- A. a' represents karyokinesis which is the division of cytoplasm.
- B. b' is telophase which is just reverse of prophase.
- C. c' is the best phase to count total number of chromosomes in any species.
- D. in 'd' stage, replication of DNA takes place on the template of the existing DNA.

Answer: D

49. Given diagram shows variations in the amount of DNA of a developing eukaryote. What the arrow denotes?



- A. First meiotic anaphase
- B. Second meiotic anaphase
- C. Mitotic anaphase
- D. Mitotic telophase

Answer: B

[Watch Video Solution](#)

1. Meiosis in diploid organisms results in

- A. production of gametes
- B. reduction in the number of chromosomes
- C. introduction of variation
- D. all of the above

Answer: D

 [Watch Video Solution](#)

2. At which stage of meiosis does the genetic constitution of gametes is finally decided?

- A. Metaphase I
- B. Anaphase II
- C. Metaphase II
- D. Anaphase I

Answer: D

 [Watch Video Solution](#)

3. Meiosis occurs in organisms during

- A. sexual reproduction
- B. vegetative reproduction
- C. both sexual and vegetative reproduction
- D. none of these

Answer: A

 [Watch Video Solution](#)

4. During anaphase I of meiosis

- A. homologous chromosomes separate
- B. non-homologous chromosomes separate
- C. sister chromatids chromosomes separate
- D. non-sister chromatids chromosomes separate.

Answer: A

 [Watch Video Solution](#)

5. Mitosis is characterised by

- A. reduction division
- B. equal division
- C. both reduction and equal division

D. paring of homologous chromosomes.

Answer: B

 [Watch Video Solution](#)

6. A bivalent of meiosis i consists of

- A. two chromatids and one centromere
- B. two chromatids and two centromeres
- C. four chromatids and two centromeres
- D. four chromatids and four centromeres.

Answer: C

 [Watch Video Solution](#)

7. Cells which are not dividing are likely to be at

- A. `G\_(1)` phase
- B. `G\_(2)` stage
- C. `G\_(0)`
- D. S phase.

Answer: C

 [Watch Video Solution](#)

8. Which of the events listed below is not observed during mitosis?

- A. chromatic condensation
- B. Movement of centrioles to opposite poles
- C. Appearance of chromosomes with two chromatids joined together at the centromere
- D. Crossing over

Answer: D

 [Watch Video Solution](#)

9. Identify the wrong statement about meiosis.

- A. Pairing of homologous chromosomes.
- B. Four haploid cells are formed.
- C. At the end of meiosis number of chromosomes are reduced to half.
- D. Two cycles of DNA replication occur

Answer: D

 [Watch Video Solution](#)

10. Select the correct statement about `G\_(1)` phase.

- A. Cell is metabolically inactive.
- B. DNA in the cell does not replicate.
- C. It is not a phase of synthesis of macromolecules.
- D. Cell stops growing.

Answer: B

 [Watch Video Solution](#)

CORNER

1. Assertion : Interphase occupies 75-95% of the total generation time.

Reason: Interphase (I-phase) is the long non-dividing phase.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: B

 [Watch Video Solution](#)

2. Assertion: Some cells enter G_0 phase leading to inactivation of cell cycle.

Reason: G_0 phase occurs due to non-availability of mitogen and energy rich compounds.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: A

 **Watch Video Solution**

3. Assertion: `G\_(1)` phase is the interval between mitosis and initiation of DNA replication.

Reason: The cell is metabolically inactive during `G\_(1)` phase.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C

 **Watch Video Solution**

4. Assertion: Prophase is the first stage of mitosis which follows S and `G\_(1)` phases of interphase.

Reason: Prophase is marked by the initiation of clusters of chromosomes.

- A. If both assertion and reason are true and reason is the correct explanation of assertion

- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: D

 [Watch Video Solution](#)

5. Assertion: Small disc-shaped structures at the surface of the centromeres are called kinetiochores.

Reason: Kinetochores serve as the sites of attachment of spindle fibres to the centreomeres.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explnantion of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: B

 [Watch Video Solution](#)

6. Assertion: During anaphase, centromere of each chromosome splites and chromatids separate.

Reason: During anaphase, chromatids move to opposte poles.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explnantion of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: B

 [Watch Video Solution](#)

7. Assertion: Karyokinesis follows cytokinesis.

Reason: Karyokinesis is the division of cytoplasm into two daughter cells.

A. If both assertion and reason are true and reason is the correct explanation of assertion

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: D

 [Watch Video Solution](#)

8. Assertion: Cell growth results in disturbing the ratio between the nucleus and cytoplasm.

Reason: Mitosis helps the cell to restore the nucleocytoplasmic ratio:

A. If both assertion and reason are true and reason is the correct explanation of assertion

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: B

 [Watch Video Solution](#)

9. Assertion: The process of pairing of the chromosomes is called synapsis.

Reason: Synapsis occurs during leptotene stage.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

10. Assertion: Crossing over leads to recombination of genetic material on the two chromosomes.

Reason: Crossing over is the exchange of genetic material between two homologous chromosomes.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: A

 [Watch Video Solution](#)

11. Assertion: The crossing over is an enzyme-mediated process.

Reason: The enzyme involved in crossing over is lyase.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

12. Assertion: The final stage of meiotic prophase I is diplotene.

Reason: Diplotene is marked by terminalisation of chiasmata.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: D

 [Watch Video Solution](#)

13. Assertion: The stage between the two meiotic divisions is called interkinesis.

Reason: Interkinesis is generally short lived.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: B

 [Watch Video Solution](#)

14. Assertion: Metaphase II begins with splitting of centromere of each chromosome into two.

Reason: In Anaphase II chromosomes align at the equator.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: D

 [Watch Video Solution](#)

15. Assertion: Variations are important for the process of evolution.

Reason: Meiosis increases the genetic variability in the population of organisms from one generation to the next.

- A. If both assertion and reason are true and reason is the correct explanation of assertion
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: B



Watch Video Solution