

CHEMISTRY**BOOKS - MHT CET ENGINEERING ENTRANCES****CHEMICAL KINETICS****Exercise 1**

1. Rate of a reaction can be defined as

- A. change in concentration of a reactant in unit time
- B. change in concentration of a product in unit time
- C. Both (a) and (b)
- D. None of the above

Answer: C

 [Watch Video Solution](#)

2. Identify the incorrect statements .

- A. rusting of iron in the presence of the air and moisture, is a slow reaction
- B. inversio of a cane sugar occurs at a moderate rate

C. hydrolysis of starch is a fast reaction

D. ionic reactions are the examples of fast reactions

Answer: C

 [View Text Solution](#)

3. For a general reaction $R \rightarrow P$, then r_{inst} will be?

A. $\frac{d[R]}{dt} = \frac{d[P]}{dt}$

B. $\frac{d[R] - d[P]}{dt}$

C. $-\frac{d[R]}{dt} = +\frac{d[P]}{dt}$

D. $\frac{d[R] + d[P]}{dt}$

Answer: C

 [Watch Video Solution](#)

4. For a gaseous reaction at constant temperature

A. concentration \propto (1 / "partial pressure of species")

B. concentration \propto "partial pressure of species"

C. concentration = partial pressure of species

D. None of the above

Answer: B

 [Watch Video Solution](#)

5. Unit of rate of a reaction is

- A. concentration time^{-1}
- B. $\text{concentration}^{-1}$ time
- C. concentration time
- D. $\text{concentration}^{-1} \text{time}^{-1}$

Answer: A

 [Watch Video Solution](#)

6. In the reaction alternative.

$2\text{HI}(\text{g}) \rightarrow \text{H}_2(\text{g}) + \text{I}_2(\text{g})$ Choose the correct alternative.

- A. Rate of reaction $= \frac{1}{2} \frac{\Delta[\text{HI}]}{\Delta t} = \frac{\Delta[\text{H}_2]}{\Delta t} = \frac{\Delta[\text{I}_2]}{\Delta t}$
- B. Stoichiometric coefficients of HI (reactant) and H_2 and I_2 (products) are not same.
- C. Rate of consumption of HI = 2 (rate of formation of H_2 Or I_2)
- D. All of the above

Answer: D

 [Watch Video Solution](#)

7. Rate of a reaction can be expressed by following rate expression, $\text{Rate} = k[A]^2[B]$, if concentration of A is increased by 3 times and concentration of B is increased by 2 times, how many times rate of reaction increases?

- A. 9 times
- B. 27 times
- C. 18 times
- D. 8 times

Answer: C

 [Watch Video Solution](#)

8. $A + 2B \rightarrow C$, the rate equation for this reaction is given as

$\text{Rate} = k[A][B]$.

If the concentration of A is kept the same but that of B is doubled what will happen to the rate itself?

- A. Halved
- B. Same
- C. Doubled
- D. Quadrupled

Answer: C

 [Watch Video Solution](#)

9. The rate constant for the reaction

$2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$, is $3.0 \times 10^{-5} \text{ s}^{-1}$. If the rate is

$2.40 \times 10^{-5} \text{ mol L}^{-1} \text{ s}^{-1}$ then the concentration of

N_2O_5 (in mol L^{-1}) is

- A. 1.4
- B. 1.2
- C. 0.04
- D. 0.8

Answer: D

 [Watch Video Solution](#)

10. In a reaction, $2\text{A} \rightarrow \text{products}$, the concentration of A decreases from 0.5 mol L^{-1} to 0.4 mol L^{-1} in 10 min.

The rate during this interval is

- A. $0.05 \text{ mol L}^{-1} \text{ min}^{-1}$
- B. $0.42 \text{ mol L}^{-1} \text{ min}^{-1}$
- C. $0.005 \text{ mol L}^{-1} \text{ min}^{-1}$
- D. $0.5 \text{ mol L}^{-1} \text{ min}^{-1}$

Answer: C

 [Watch Video Solution](#)

11. The reaction, $2\text{NO}(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{NO}_2(\text{g})$, is of first order. If the volume of reaction vessel is reduced to $\frac{1}{3}$, the rate of reaction would be

- A. $\frac{1}{3}$ "times"
- B. $\frac{2}{3}$ "times"
- C. 3 times
- D. 6 times

Answer: C

 [Watch Video Solution](#)

12. On increasing the pressure three fold, the rate of reaction of $2\text{H}_2\text{S} + \text{O}_2 \rightarrow \text{products}$ would increase"

- A. 3 times
- B. 39times
- C. 12 times
- D. 27 times

Answer: D

 [Watch Video Solution](#)

13. Which of these does not influence the rate of reaction?

- A. Nature of the reactants
- B. Concentration of the reactants
- C. Temperature of the reaction
- D. Molecularity of the reaction

Answer: D

 [Watch Video Solution](#)

14. Units of rate constant of first and zero order reactions in terms of molarity `M` are respectively:

- A. s^{-1}, Ms^{-1}
- B. s^{-1}, M
- C. Ms^{-1}
- D. M, s^{-1}

Answer: A

 [Watch Video Solution](#)


15. With increase in temperature , rate of reaction

- A. increases
- B. decreases
- C. remains same

D. may increase

Answer: A

 [View Text Solution](#)



What is the ratio of the rate of decomposition of N_2O_5 to rate of formation of NO_2

A. 1:2

B. 2:1

C. 1:4

D. 4:1

Answer: B

 [Watch Video Solution](#)

17. Order of a chemical reaction is defined as the

A. difference in powers of the concentration of the reactants in the rate law expression

B. sum of powers of the concentration of the reactants in the rate law expression

C. sum of powers of the concentration of the products in the rate law expression

D. difference in powers of the concentration of the products in the rate law expression

Answer: B

 [Watch Video Solution](#)

18. Which one of the following statements for the order of a reaction is incorrect ?

- A. Order of reaction is always whole number
- B. Order can be determined only experimentally
- C. Order is not influenced by stoichiometric coefficient of the reactants
- D. Order of reaction is sum of the power to the concentration terms of reactants to express the rate of reaction

Answer: A

 [Watch Video Solution](#)

19. The order of a reaction with rate equal to $kC_A^{3/2}C_B^{-1/2}$ is

- A. 1
- B. $-(1)/(2)$
- C. $-(3)/(2)$
- D. 2

Answer: A

 [Watch Video Solution](#)

20. The unit $\text{mol L}^{-1}\text{s}^{-1}$ is meant for the rate constant of the reaction having the order

- A. 0
- B. 2
- C. 1
- D. 3

Answer: A

 [Watch Video Solution](#)

21. The relative order of rate of esterification of acids is

- A. 0
- B. first
- C. second order
- D. pseudo first order

Answer: C

 [Watch Video Solution](#)

22. Decomposition of ammonium nitrite is an example of

- A. bimolecular reaction

- B. unimolecular reaction
- C. Both (a) and (b)
- D. None of the above

Answer: B

 [Watch Video Solution](#)

23. When one reactant is present in excess in a chemical reaction between two substances, then the reaction is known as

- A. first order reaction
- B. second order reaction
- C. zero order reaction
- D. pseudo first order reaction

Answer: D

 [Watch Video Solution](#)

24. For the rate law, $r = k[A]^{3/2}[B]^{-1}$ the overall order of a reaction is

- A. zero
- B. half
- C. one

D. two

Answer: B

 [Watch Video Solution](#)

25. For a zero order reaction a graph of conc. (along Y axis) and time (along X-axis) is linear with

- A. a zero intercept and a + ve slope
- B. a zero intercept and a - ve slope
- C. a non-zero intercept and a - ve slope
- D. a non-zero intercept and a + ve slope

Answer: C

 [Watch Video Solution](#)

26. For a chemical reaction Can never be a fraction

- A. half-life
- B. molecularity
- C. Order
- D. rate constant

Answer: B

 [Watch Video Solution](#)

27. The rate of law for the reaction $x\text{A} + y\text{B} = m\text{P} + n\text{Q}$ is $\text{Rate} = k[\text{A}]^c[\text{B}]^d$. What is the total order of reaction ?

- A. $(x+y)$
- B. $(m+n)$
- C. $(c+d)$
- D. $(x)/y$

Answer: C

 [Watch Video Solution](#)

28. For which type of the reactions, order and molecularity have the same value ?

- A. first order reaction
- B. Bimolecular reaction
- C. Termolecular reaction
- D. Elementary reaction

Answer: D

 [Watch Video Solution](#)

29. Zero order reaction means that

- A. rate of reaction $\propto 1/(\text{"zero power of concentration of reactants"})$
- B. rate of reaction \propto zero power of concentration of reactants
- C. rate of reaction = zero power of concentration of products
- D. None of the above

Answer: B

 [Watch Video Solution](#)

30. Value of $t_{1/2}$ for first order reaction is

- A. $(0.693)/k$
- B. $(0.2303)/k$
- C. $[R]/2$
- D. $(0.301)/k$

Answer: A

 [Watch Video Solution](#)

31. In a first order reaction, reactant concentration 'C' varies with time 't' as

- A. C decreases with $(1)/t$
- B. $\log C$ decreases with $(1)/t$
- C. $(1)/c$ increases linearly with t

D. $\log C$ decreases linearly with t

Answer: D

 [Watch Video Solution](#)

32. Which of the following statement (s) is/are true?

A. For zero order reaction, $t_{1/2} \propto [R]_0$

B. For first order reaction, $t_{1/2}$ "is independent of" $[R]_0$

C. Both (a) and (b)

D. None of the above

Answer: C

 [Watch Video Solution](#)

33. $\frac{d[\text{A}]}{dt}$ dependence of rate is called differential rate equation. Choose the suitable word to replace A.

A. Concentration

B. Volume


C. Order

D. Pressure

Answer: A

 Watch Video Solution

34. Consider the following,



(I) The above reaction is an example of first order kinetics. (II) Rate of the reaction will be given as $\text{Rate} = k[\text{C}_2\text{H}_2]$.

Which of the above statement(s) is/are correct? Choose the correct option.

- A. Only I
- B. Only II
- C. Both (a) and (b)
- D. None of the above

Answer: C

 Watch Video Solution

35. A first order reaction is found to have a rate constant, $k = 4.2 \times 10^{-12} \text{ s}^{-1}$. Find the half-life of the reaction.

- A. $1.26 \times 10^{13} \text{ s}$
- B. $1.65 \times 10^{11} \text{ s}$
- C. $1.65 \times 10^{11} \text{ s}$
- D. $1.26 \times 10^{13} \text{ s}$

Answer: B

 [Watch Video Solution](#)

36. The rate of the backward reaction in a reversible reaction

- A. positive
- B. negative
- C. Either (a) or (b)
- D. None of the above

Answer: A

 [View Text Solution](#)

37. Expression for the half-life of zero order reaction is given as

- A. $t_{1/2} = (2[R]_0)/(K)$
- B. $t_{1/2} = ([R]_0)/(2K)$
- C. $t_{1/2} = (0.693)/(K)$
- D. $t_{1/2} = (0.301)/(K)$

Answer: B

 [Watch Video Solution](#)

38. The rate constant of a zero order reaction is $0.2 \text{ mol L}^{-3} \text{ h}^{-1}$. If the concentration of the reactant after 30 min is 0.05 mol dm^{-3} , then its initial concentration would be

- A. 0.01 mol dm^{-3}
- B. 0.15 mol dm^{-3}
- C. 0.25 mol dm^{-3}
- D. 4.00 mol dm^{-3}

Answer: B

 [Watch Video Solution](#)

39. The rate constant for the first order reaction is 60 s^{-1} . How much time will it take to reduce the concentration of the reactant to $1/16$ th value ?

- A. 0.046 s
- B. 0.025 s
- C. 0.098 s
- D. 0.060 s

Answer: A

 [Watch Video Solution](#)

40. Inversion of cane sugar is a \dots order reaction

$\text{Cane sugar } (\text{C}_{12}\text{H}_{22}\text{O}_{11}) + \text{H}_2\text{O} \xrightarrow{\text{H}^+} \text{C}_6\text{H}_{12}\text{O}_6 + \text{C}_6\text{H}_{12}\text{O}_6$

underset("Fructose")C`

Rate =D

Here A,B,C and D respectively are

A. `A="first,B=fructose",C=C_(6)H_(12)O_(6),D=K[H_2O]`

B. `A="second,B=fructose",C=C_(7)H_(14)O_(7),D=K[H_2O]`

C. `A="pseudo first, B=glucose,"C=C_(6)H_(12)O_(6),`

`D=K[C_(12)H_(22)O_(11)]`

D. `A="pseudo first, B=fructose,"C=C_(6)H_(12)O_(6),`

`D=K[C_(12)H_(22)O_(11)][H_(2)O]`

Answer: C

 [Watch Video Solution](#)

41. The rate constant 'K' for pseudo first order reaction is

A. $\frac{2.303}{t} \log C_{(0)} - C$

B. $\frac{2.303}{t} \log \frac{C}{C_{(0)}}$

C. $\frac{2.303}{t} \log C_{(0)} + C$

D. $\frac{2.303}{t} \log C_{(0)} / C$

Answer: D

 [Watch Video Solution](#)

42. Half-life period of a first order reaction is 10 min. Starting with initial concentration 12 M, the rate after 20 min is

- A. 0.693 M min^{-1}
- B. $0.693 \times 3 \text{ M min}^{-1}$
- C. $0.0693 \times 3 \text{ M min}^{-1}$
- D. $0.0693 \times 4 \text{ M min}^{-1}$

Answer: C

 [Watch Video Solution](#)

43. The rate of first-order reaction is $1.5 \times 10^{-2} \text{ M min}^{-1}$ at 0.5 M concentration of reactant. The half-life of reaction is

- A. 0.383 min
- B. 23.1 min
- C. 8.73 min
- D. 7.53 min

Answer: B

 [Watch Video Solution](#)

44. For a first order reaction half life is 14 sec. The time required for the initial concentration to reduce $1/8$ of the value is

A. $(14)^3 \text{ s}$

B. 28s

C. 42

D. $(14)^2 \text{ s}$

Answer: C

 [Watch Video Solution](#)

45. The half life period of a substance is 50 minutes at a certain initial concentration. When the concentration is reduced to one half of the initial value, the half-life period is 25 minutes. Calculate the order of the reaction.

A. 0

B. $(1/2)$

C. $(3/2)$

D. 2

Answer: C

 [Watch Video Solution](#)

46. How much time is required for two - third completion of a first order reaction having,

$$k = 5.48 \times 10^{-14} \text{ s}^{-1}?$$

A. $2.01 \times 10^{11} \text{ s}$

B. $2.01 \times 10^{13} \text{ s}$

C. $8.08 \times 10^{13} \text{ s}$

D. $16.04 \times 10^{11} \text{ s}$

Answer: B

 [Watch Video Solution](#)

47. $t_{1/4}$ can be taken as the time taken for concentration of reactant to drop to $\frac{3}{4}$ of its initial value. If the rate constant for a first order reaction is k , then $t_{1/4}$ can be written as:

A. $0.75/k$

B. $0.69/k$

C. $0.29/k$

D. $0.10/k$

Answer: C

 [Watch Video Solution](#)

48. For a reaction, the rate constant is 2.34 s^{-1} . The half-life period for the reaction is

- A. 0.30s
- B. 0.60s
- C. 3.3s
- D. data is insufficient

Answer: A

 [Watch Video Solution](#)

49. 75% of a first order reaction was completed in 32 min. When was 50% of the reaction completed ?

- A. 16 min
- B. 8 min
- C. 4 min
- D. 32 min

Answer: A

 [Watch Video Solution](#)

50. Rate constant for a reaction is 10^{-3} s^{-1} . How much time is required to reduce the initial concentration of reactant to 25%

- A. 693 s
- B. 1386 s

C. 6930 s

D. 2029 s

Answer: B

 [Watch Video Solution](#)

51. A first order reaction is `10%` complete in 20 min. the time taken for `19%` completion is :

A. 30 min

B. 40 min

C. 50 min

D. 38 min

Answer: B

 [Watch Video Solution](#)

52. For a zero order reaction, the integrated rate equation is

A. $kt = ([A])/([A]_0)$

B. $kt = [A] - [A]_0$

C. $[A] = -kt + [A]_0$

D. $[A] = kt - [A]_0$

Answer: C

 [Watch Video Solution](#)

53. If the half-time for a particular reaction is found to be constant and independent of the initial concentration of the reactants, then the reaction is of

- A. first order
- B. zero order
- C. second order
- D. None of the above

Answer: A

 [Watch Video Solution](#)

54. The integrated rate equation is $kt = \log C_0 - \log C_t$.

The straight line graph is obtained by plotting

- A. $\log C_t$ vs time
- B. $\frac{1}{\text{time}}$ vs C_t
- C. time vs C_t
- D. $\frac{1}{\text{time}}$ vs $\frac{1}{C_t}$

Answer: A

 [Watch Video Solution](#)

55. On the basis of Arrhenius equation, consider the following statement (s)

I. E_a is activation energy.

II. Unit of E_a is J mol^{-1}

III. R is gas constant.

Which of these is/are true statement(s)?

A. I and II

B. II and III

C. I and III

D. I, II and III

Answer: D

 [Watch Video Solution](#)

56. What will happen, on increasing the temperature or decreasing the activation energy (answer on the basis of Arrhenius equation)?

A. Rate of reaction will increase

B. An exponential increase in rate constant

C. Both (a) and (b)

D. None of the above

Answer: C

 [Watch Video Solution](#)

57. Rate constant k of a reaction is dependent on temperature:

$$k = A e^{-E_a/RT}$$

k has the least value at

- A. high T and high E_a
- B. high T and small E_a
- C. low T and low E_a
- D. low T and high E_a

Answer: D

 [Watch Video Solution](#)

58. The rate constant of a reaction is given by $k = 2.1 \times 10^{10} \exp(-2700/RT)$. It means that

- A. $\log k$ vs $1/T$ "will be a curved line with slope" $-(2700)/(2.303R)$
- B. $\log k$ vs $1/T$ will be a straight line with intercept on $\log k$ axis $-\log 2.1 \times 10^{10}$
- C. the number of effective collisions are $2.1 \times 10^{10} \text{ cm}^{-3} \text{ s}^{-1}$
- D. half-life of the reaction increases with increase of temperature

Answer: B

 [Watch Video Solution](#)

59. Rate of a reaction can be expressed by Arrhenius equation as:

$$k = Ae^{(-E_a/RT)}$$

In this equation, E_a represents:

- A. the energy above which all the colliding molecules
- B. the energy below which colliding molecules will not react
- C. the total energy of the reaction molecules at a temperature, T
- D. the fraction of molecules with energy greater than the activation energy of the reaction

Answer: A

 [Watch Video Solution](#)

60. The decomposition of hydrocarbon follows the equation $k = (4.5 \times 10^{11} \text{ s}^{-1}) e^{(-28000 \text{ K}/T)}$

Calculate E_a .

- A. $232.79 \text{ kJ mol}^{-1}$
- B. $425.25 \text{ kJ mol}^{-1}$
- C. 300 kJ mol^{-1}
- D. $885.2 \text{ kJ mol}^{-1}$

Answer: A

 [Watch Video Solution](#)

61. The activation energy for a reaction at temperature T K was found to be or $2.303 RT \text{ J mol}^{-1}$. The ratio of the rate constant to Arrhenius factor is

- A. 0.01
- B. 0.1
- C. 0.02
- D. 0.001

Answer: B

 [Watch Video Solution](#)

Exercise 2

1. The activation energy for the reaction $2\text{HI}(g) \rightarrow \text{H}_2(g) + \text{I}_2(g)$


is $209.5 \text{ kJ mol}^{-1}$ at 581K . Calculate the fraction of molecules of reactants having energy equal to or greater than activation energy ?

- A. 1.82×10^{-18}
- B. 1.47×10^{-19}
- C. 2.67×10^{-16}
- D. 3.89×10^{-19}

Answer: B

 [Watch Video Solution](#)

2. The chemical reaction $2\text{O}_3 \rightarrow 3\text{O}_2$ proceeds as follows :



The rate law expression should be :

A. $r = k[\text{O}_3]^2$

B. $r = k[\text{O}_3]^2[\text{O}_2]^{-1}$

C. $r = k[\text{O}_3][\text{O}_2]^{-1}$

D. Unpredictable

Answer: B

 [Watch Video Solution](#)

3. The rate constant of a reaction increases by 5% when its temperature is raised from 27°C to 28°C .

The activation energy of the reaction is

A. 36.6kJ/mol

B. 16.6kJ/mol

C. 46.6kJ/mol

D. 26.6kJ/mol

Answer: A

 [Watch Video Solution](#)

4. The time required for 10% completion of a first order reaction at 298 K is equal to that required for its 25% completion at 308 K. If the value of A is $4 \times 10^{10} \text{ s}^{-1}$.

Calculate the rate constant, k at 318 K.

$2.89 \times 10^{-2} \text{ s}^{-1}$

$3.26 \times 10^{-2} \text{ s}^{-1}$

$1.03 \times 10^{-2} \text{ s}^{-1}$

$0.03 \times 10^{-2} \text{ s}^{-1}$

- A. the energy below which colliding molecules will not react
- B. the total energy of the reacting molecules at a temperature, T
- C. The fraction of molecules with energy greater than the activation energy of the reaction
- D. $0.03 \times 10^{-2} \text{ S}^{-1}$

Answer: C

 [View Text Solution](#)

5. When initial concentration of a reactant is doubled in a reaction, its half-life period is not affected. The order of the reaction is

- A. zero
- B. first
- C. second
- D. more than zero but less than first

Answer: B

 [Watch Video Solution](#)

6. 1 g of ${}^{198}\text{Au}$ ($t_{1/2} = 65 \text{ h}$) gives stable mercury by β -emission. What amount of mercury will left after 260 h?

- A. 0.9374 g
- B. 0.3758 g
- C. 0.7586 g
- D. 0.9000 g

Answer: A

 [View Text Solution](#)

7. An endothermic reaction, $A \rightarrow B$ have an activation energy 15 kcal/mol and the heat of the reaction is 5 kcal/mol . The activation energy of the reaction, $B \rightarrow A$ is:

- A. 20 kcal/mol
- B. 15 kcal/mol
- C. 10 kcal/mol
- D. zero

Answer: C

 [Watch Video Solution](#)

8. The rate of a reaction quadruples when the temperature changes from 293K to 313K . Calculate the energy of activation of the reaction assuming that it does not change with temperature.

A. 48.625kJ mol^{-1}

B. 654.35kJ mol^{-1}

C. 354.20kJ mol^{-1}

D. 52.854kJ mol^{-1}

Answer: D

 [Watch Video Solution](#)

9. The rate law for the reaction

$\text{RCI} + \text{NaOH}(\text{aq}) \rightarrow \text{ROH} + \text{NaCl}$ is given by

Rate $= k[\text{RCI}]$. The rate of the reaction will be

A. is doubled by doubling the concentration of NaOH

B. is halved by reducing the concentration of RCI by one half

C. is increased by increasing the temperature of the reaction

D. is unaffected by change in temperature

Answer: B

 [Watch Video Solution](#)

10. The decomposition of 'A' into product has value of 'k' as $4.5 \times 10^3 \text{ s}^{-1}$ at 10°C and energy of activation of 60 kJ mol^{-1} . At what temperature would 'k' be $1.5 \times 10^4 \text{ s}^{-1}$?

- A. 273.15 k
- B. 24.01°C
- C. 280.39 K
- D. 45.29°C

Answer: B

 [Watch Video Solution](#)

11. The rate constant ' k_1 ' of one reaction is double of the rate constant (k_2) of another reaction. Then the relationship between the corresponding activation energies of the two reactions ' E_{a1} ' and ' E_{a2} ' will be

- A. $E_{a1} > E_{a2}$
- B. $E_{a1} = 4E_{a2}$
- C. $E_{a1} = E_{a2}$
- D. $E_{a1} < E_{a2}$

Answer: D

 [Watch Video Solution](#)

12. The rate constant of the chemical reaction doubled for an increase of 10 k in absolute temperature from 295 k. Calculate the (activation energy), E_a .

- A. 51.8 kJ mol^{-1}
- B. 82.1 kJ mol^{-1}
- C. 23.8 kJ mol^{-1}
- D. 62.1 kJ mol^{-1}

Answer: A

 [Watch Video Solution](#)

13. Powdered magnesium element catches fire more rapidly than magnesium wire of the same mass because

- A. Surface area of magnesium wire is larger than their powdered form
- B. density of magnesium wire is greater than that of their powdered form
- C. powdered magnesium have larger surface area
- D. None of above

Answer: C

 [Watch Video Solution](#)


14. For the reaction $2\text{H}_2 + \text{O}_2 \rightarrow 2\text{H}_2\text{O}$, the rate law expression is $r = k[\text{H}_2]^n$. When the concentration of H_2 is doubled, the rate of reaction found to be quadrupled. The value of n is

- A. 0
- B. 1
- C. 2
- D. 3

Answer: C

 [Watch Video Solution](#)

15. consider the following reaction,



The rate of change of concentration for nitrogen is $-0.3 \times 10^{-4} \text{ Ms}^{-1}$. The rate of change of concentration of ammonia is

- A. $0.2 \times 10^{-4} \text{ Ms}^{-1}$
- B. $0.4 \times 10^{-4} \text{ Ms}^{-1}$
- C. $0.6 \times 10^{-4} \text{ Ms}^{-1}$
- D. $-0.6 \times 10^{-4} \text{ Ms}^{-1}$

Answer: C

 [Watch Video Solution](#)

16. For a reaction $\text{R} \rightarrow \text{P}$, the concentration of a reactant changes from 0.03 M to 0.02M in 25 minutes. Calculate the average rate of the reaction using the units of seconds.

A. 6.66×10^{-5}

B. 6.6×10^{-6}

C. 5.67×10^{-5}

D. 7.26×10^{-6}

Answer: B

 [Watch Video Solution](#)

17. Consider the following reaction,



If the concentration of PO_2 increases by $5.2 \times 10^{-3} \text{M}$ in 100 s, the rate of a reaction is

A. $0.5 \times 10^{-4} \text{Ms}^{-1}$

B. $2.5 \times 10^{-5} \text{Ms}^{-1}$

C. $1.3 \times 10^{-5} \text{Ms}^{-1}$

D. $2 \times 10^3 \text{Ms}^{-1}$

Answer: C

 [Watch Video Solution](#)

18. For the reaction, $\text{A} + \text{B} \rightarrow \text{product}$

If concentration of A is doubled, rate increases 4 times. If concentrations of A and B both are doubled, rate increases 8 times. The differential rate equation of the reaction will be

A. $\frac{dC}{dt} = kC_A \times C_B$

B. $\frac{dC}{dt} = kC_A^2 \times C_B^3$


C. $\frac{dC}{dt} = kC_A^2 \times C_B$

D. $\frac{dC}{dt} = kC_A^2 \times C_B^2$

Answer: C

 [View Text Solution](#)

19. Consider the following reaction,



The activation energy for the forward reaction is 50 kcal. What is the activation energy for the backward reaction?

A. $-28 \text{ kcal mol}^{-1}$

B. $+28 \text{ kcal mol}^{-1}$

C. $-72 \text{ kcal mol}^{-1}$

D. $+72 \text{ kcal mol}^{-1}$

Answer: D

 [Watch Video Solution](#)

20. Consider the following reaction,




How will the rate of reaction changes when the concentration of A is doubled and that of B is triplet while C is taken in excess ?

- A. The rate reduces 8 times of its original value
- B. The rate reduces 12 times of its original value
- C. The rate increases 8 times of its original value
- D. The rate increases 12 times of its original value

Answer: D

 [Watch Video Solution](#)

21. Which of the following expression is correct for the rate of reaction given below ?



- A. $\frac{\Delta[\text{Br}^{\ominus}]}{\Delta t} = 6 \frac{\Delta[\text{H}^{\oplus}]}{\Delta t}$
- B. $\frac{\Delta[\text{Br}^{\ominus}]}{\Delta t} = \frac{6}{5} \frac{\Delta[\text{H}^{\oplus}]}{\Delta t}$
- C. $\frac{\Delta[\text{Br}^{\ominus}]}{\Delta t} = 5 \frac{\Delta[\text{H}^{\oplus}]}{\Delta t}$
- D. $\frac{\Delta[\text{Br}]}{\Delta t} = \frac{5}{6} \frac{\Delta[\text{H}^{\oplus}]}{\Delta t}$

Answer: D

 [Watch Video Solution](#)

22. The reaction $\text{A} \rightarrow \text{B}$ follows first order kinetics. The time taken for 0.8 mol of A to produce 0.6 mol of B is 1 hr . What is the time taken for the conversion of 9.0 mol of A to Product 0.675 mol of B ?

- A. 0.5 h
- B. 0.25 h
- C. 1 h
- D. 2 h

Answer: C

 [Watch Video Solution](#)

23. The rate of formation of SO_3 in the following reaction is 100 g min^{-1} .
 $2\text{SO}_2 + \text{O}_2 \rightarrow 2\text{SO}_3$


The rate of disappearance of O_2 is

- A. 29 g min^{-1}
- B. 20 g min^{-1}
- C. 200 g min^{-1}
- D. 50 g min^{-1}

Answer: B

 [Watch Video Solution](#)

24. Consider the following reaction,



$\frac{d[O_2]}{dt} = k_3[N_2O_5]$ and $\frac{d[N_2O_5]}{dt} = -k_1$

The relation between k_1 , k_2 and k_3 is

- A. $k_1 = k_2 = k_3$
- B. $2k_1 = k_2 = 4k_3$
- C. $2k_1 = 4k_2 = k_3$
- D. None of these

Answer: B

 [Watch Video Solution](#)

25. For a hypothetical reaction, $A \rightarrow P$


the rate constant is 0.12 s^{-1} . which of the following relation is correct, if $[R]_0$ is the initial concentration?

- A. $t_{1/2} = \frac{[R]_0}{0.12 \times 2}$
- B. $t_{1/2} = \frac{3}{2(0.12)[R]_0^2}$
- C. $t_{1/2} = \frac{0.693}{0.12}$
- D. $t_{1/2} = \frac{0.693}{0.12 \times 3}$

Answer: C

 [Watch Video Solution](#)

26. The initial concentration of N_2O_5 in the following first order reaction:



was $1.24 \times 10^{-2} \text{ mol L}^{-1}$ at 318 K . The concentration of N_2O_5 after 60 min was $0.20 \times 10^{-2} \text{ mol L}^{-1}$. Calculate the rate constant of the reaction at 318 K .

A. 0.0104 min^{-1}

B. 0.0204 min^{-1}


C. 0.0304 min^{-1}

D. 0.0404 min^{-1}

Answer: C

 [Watch Video Solution](#)

27. A following mechanism has been proposed for a reaction



The rate law expression for the reaction by RDS method is:

A. $r = k[\text{P}][\text{Q}]$

B. $r = k[\text{P}]^2$

C. $r = k[\text{P}][\text{T}]$

D. $r = k[\text{P}]^2[\text{Q}]$

Answer: A

 [Watch Video Solution](#)

28. The half-time of the following first order decomposition of nitramide is 2.1 h at 15°C :



If 6.2 g of nitramide is allowed to decompose then time taken for it to decompose 99%, will be

- A. 2.1 h
- B. 12 h
- C. 13.96 h
- D. 33 h

Answer: C

 [Watch Video Solution](#)

29. Consider a reaction $a\text{G} + b\text{H} \rightarrow \text{Products}$. When concentration of both the reactants G and H is doubled, the rate increases eight times. However, when the concentration of G is doubled, keeping the concentration of H fixed, the rate is doubled. The overall order of reaction is

- A. 3
- B. 2
- C. 1
- D. 0

Answer: A

 [Watch Video Solution](#)

30. Find out two-third $\frac{2}{3}$ life of a first order reaction in which $k = 5.48 \times 10^{-14} \text{ s}^{-1}$

A. $2.01 \times 10^{11} \text{ s}$

B. $2.01 \times 10^{13} \text{ s}$

C. $8.08 \times 10^{13} \text{ s}$

D. $16.04 \times 10^{11} \text{ s}$

Answer: B

 [Watch Video Solution](#)

31. Rate constant $k = 1.2 \times 10^3 \text{ mol}^{-1} \text{ L s}^{-1}$ and $E_a = 2.0 \times 10^2 \text{ kJ mol}^{-1}$. When $T \rightarrow \infty$:

A. $A = 2.0 \times 10^2 \text{ kJ mol}^{-1}$

B. $A = 1.2 \times 10^3 \text{ mol}^{-1} \text{ L s}^{-1}$

C. $A = 1.2 \times 10^3 \text{ mol}^{-1} \text{ L s}^{-1}$

D. $A = 2.4 \times 10^3 \text{ kJ mol}^{-1} \text{ s}^{-1}$

Answer: C

 [Watch Video Solution](#)

32. A given sample of milk turns sour at room temperature (27°C) in five hours. In a refrigerator at -3°C , it can be stored 10 times longer. The energy of activation for the souring of milk is

- A. $2.303 \times 5 \text{RkJ mol}^{-1}$
- B. $2.303 \times 3 \text{RkJ mol}^{-1}$
- C. $2.303 \times 2.7 \text{RkJ mol}^{-1}$
- D. $2.303 \times 10 \text{RkJ mol}^{-1}$

Answer: C

 [Watch Video Solution](#)

33. The rate law for a reaction between the substances A and B is given by rate $= k[A]^n [B]^m$. On doubling the concentration of A and having the concentration of B, the ratio of the new rate to the earlier rate of the reaction will be:

- A. $(1)/(2^{(m)+n})$
- B. $(m+n)$
- C. $(n-m)$
- D. $2^{((n-m))}$

Answer: D

 [Watch Video Solution](#)

34. The time required for 100% completion of a zero order reaction is

A. ak

B. $\frac{a}{2k}$

C. $\frac{a}{k}$

D. $\frac{2k}{a}$

Answer: C

 [Watch Video Solution](#)

35. For a reaction $2A \rightarrow 3B$, if the rate of formation of B is $x \text{ mol/L}$, the rate of consumption of A is

A. x

B. $\frac{3x}{2}$

C. $3x$

D. $\frac{2x}{3}$

Answer: D

 [Watch Video Solution](#)

36.A..... Is the expression in which reaction rate is given in term of molar concentration of reactants with each term raised to some power, which may or may not be same as the stoichiometric coefficient of the reacting species in a balanced chemical equation. Here, A is

A. Rate law

- B. Rate equation
- C. Differential rate equation
- D. None of these

Answer: A

 [Watch Video Solution](#)

37. Consider some facts about decomposition of H_2O_2 .

It is catalysed by iodide ion in acidic medium.

It is second order reaction with respect to both H_2O_2 and I^- .

Rate equation of this particular reaction will be

$$\text{Rate} = -\frac{d[\text{H}_2\text{O}_2]}{dt} = k[\text{H}_2\text{O}_2]^2[\text{I}^-]$$

It completes in two steps and both steps are unimolecular elementary reactions.

Which of the above written facts are correct, regarding decomposition of H_2O_2 ? Choose the correct option.

- A. I and II
- B. II and III
- C. III and IV
- D. I and IV

Answer: D

 [View Text Solution](#)

38. If rate constant is numerically the same for three reaction of first, second and third order respectively, then which of the following is correct?

- A. $r_1 > r_2 > r_3$
- B. $r_1 = r_2 = r_3$
- C. $r_1 < r_2 < r_3$
- D. All of these

Answer: D

 [Watch Video Solution](#)

39. The rate constant of a reaction is given as

$$k = 2.1 \times 10^{10} e^{-(2700/RT)}$$

It means that

"log k" vs "1/T" will be a straight line with intercept on "log k" axis = $\log 2.1 \times 10^{10}$.

Number of effective collisions of temperature are $2.1 \times 10^{10} \text{ cm}^{-3} \text{ s}^{-1}$.

Half-life of a reaction increases of temperature.

"log k" vs "1/T" will be a straight line with slope = $-(2700)/(2.303R)$.

Which of the above statements are true? Choose the correct option.

- A. I and II
- B. II and III
- C. III and IV
- D. I and IV

Answer: d

 [Watch Video Solution](#)

40. On mixing 1 dm³ of 3M ethanol with 1 dm³ of 2 M ethanoic acid, an ester is formed.



If each solution is diluted with an equal volume of water, the decrease in the initial rate would be

- A. 0.5 times
- B. 4 times
- C. 0.25 times
- D. 2 times

Answer: C

 [Watch Video Solution](#)

41. In a reversible reaction $2\text{NO}_2 \xrightleftharpoons[k_{-2}]{k_1} \text{N}_2\text{O}_4$, the rate of disappearance of NO_2 is equal to

- A. $\frac{2k_1}{k_2}[\text{NO}_2]^2$
- B. $2k_1[\text{NO}_2]^2 - 2k_2[\text{N}_2\text{O}_4]$
- C. $2k_2[\text{NO}_2]^2 - k_1[\text{N}_2\text{O}_4]$
- D. $(2k_1 - k_2)[\text{NO}_2]$

Answer: B

 [Watch Video Solution](#)


42. The conversion of molecules 'X' to 'Y' follows second order kinetics. If the concentration of 'X' is increased to three times, how will it affect the rate of formation of 'Y' ?

- A. increased by three times
- B. decreased by three times
- C. increases by nine times
- D. decrease by nine times

Answer: C

 [Watch Video Solution](#)

43. In the sequence of reaction,



$$k_3 > k_2 > k_1$$

The rate determining step of the reaction is :

- A. $A \rightarrow B$
- B. $B \rightarrow C$
- C. $C \rightarrow D$
- D. $A \rightarrow D$

Answer: A

 [Watch Video Solution](#)

44. Half-life of a reaction is found to be inversely proportional to the cube of its initial concentration. The order of reaction is

- A. 4
- B. 3
- C. 5
- D. 2

Answer: A

 [Watch Video Solution](#)

45. For an exothermic chemical process occurring in two as

(i) $A + B \rightarrow X$ (slow)

(ii) $X \rightarrow AB$ (fast)

The process of the reaction can be best described by



A.



B.



C.

D. All are correct

Answer: C

 [Watch Video Solution](#)

46. A reactant (A) forms two products

$A \xrightarrow{k_1} B$, Activation energy E_{a1}

$A \xrightarrow{k_2} C$, Activation energy E_{a2}

If $E_{a2} = 2E_{a1}$ then k_1 and k_2 are related as

A. $k_1 = 2k_2 e^{(E_{a2} - E_{a1})/RT}$

B. $k_1 = k_2 e^{(2E_{a1})/RT}$

C. $k_2 = k_1 e^{(E_{a2} - E_{a1})/RT}$

D. $k_1 = k_2 e^{(E_{a2} - E_{a1})/RT}$

Answer: B

 [Watch Video Solution](#)

47. In a first order reaction, $A \rightarrow P$, the ratio of $a/(a-x)$ was found to be 8 after 60 min. If the concentration is 0.1 M then the rate of reaction is

A. $2.226 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$

B. $4.455 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$

C. $3.466 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$

D. $5.532 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$

Answer: C

 [Watch Video Solution](#)

48. The rate of a chemical reaction becomes double for every 10° rise in temperature. If the temperature is raised by 50° C, the rate of reaction increases by about:

- A. 10 times
- B. 24 times
- C. 32 times
- D. 64 times

Answer: C

 [Watch Video Solution](#)

49. For a reaction $\frac{1}{2}A \rightarrow 2B$, rate of disappearance of A is related to the rate of appearance of B by the expression:

- A. $-(d[A])/dt = (1/2)(d[B])/dt$
- B. $-(d[A])/dt = (1/4)(d[B])/dt$
- C. $-(d[A])/dt = (d[B])/dt$
- D. $(d[A])/dt = 4(d[B])/dt$

Answer: B

 [Watch Video Solution](#)

50. The rate of a reaction doubles when its temperature changes from 300 K to 310 K . Activation energy of such a reaction will be:

($R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ and $\log 2 = 0.301$)

- A. 53.6 kJ mol^{-1}
- B. 48.6 kJ mol^{-1}
- C. 58.5 kJ mol^{-1}
- D. 60.5 kJ mol^{-1}

Answer: A

 [Watch Video Solution](#)

51. A plot of $\ln k$ vs $1/T$ for a reaction gives the slope $-1 \times 10^4 \text{ K}$. The energy of activation for the reaction is (Given, $R = 8.314 \text{ k}^{-1} \text{ mol}^{-1}$)

- A. 8314 kJ mol^{-1}
- B. $1.202 \text{ kJ mol}^{-1}$
- C. 1202 kJ mol^{-1}
- D. $83.14 \text{ kJ mol}^{-1}$

Answer: D

 [Watch Video Solution](#)

52. In a zero-order reaction for every 10° rise of temperature, the rate is doubled. If the temperature is increased from 10°C to 100°C , the rate of the reaction will become

- A. 256 times
- B. 512 times
- C. 64 times
- D. 128 times

Answer: B

 [Watch Video Solution](#)

53. The inversion of cane sugar is first order in [sugar] and proceeds with half-life of 600 min at pH=4 for a given concentration of sugar. However , if pH=5, the half-life changes to 60 min. The rate law expression for the sugar inversion can be written as

- A. $\text{rate} = k[\text{sugar}]^1[\text{H}^+]^2$
- B. $\text{rate} = k[\text{sugar}][\text{H}^+]^1$
- C. $\text{rate} = k[\text{sugar}][\text{H}^+]^4$
- D. $\text{rate} = k[\text{sugar}][\text{H}^+]^0$

Answer: D

 [View Text Solution](#)

54. The initial rates of reaction $3A + 2B + C \rightarrow \text{products}$ at different initial concentration are given below

{:("Initial rate, MS⁻¹),|A_(0)|M,|B_(0)|M,|C_(0)|M),(5.0xx10⁻³),0.010,0.005,0.010),

$(5.0 \times 10^{-3}, 0.010, 0.005, 0.015), (1.0 \times 10^{-2}, 0.010, 0.010, 0.010), (1.25 \times 10^{-3}, 0.005, 0.005, 0.010):$

The order of reaction with respect to the reacts A, B and C are respectively.

- A. 3,2,0
- B. 3,2,1
- C. 2,2,0
- D. 2,1,0

Answer: D

 [Watch Video Solution](#)

55. For a reaction , $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$, the rate is directly proportional to $[\text{N}_2\text{O}_5]$. At 45°C , 90% of the N_2O_5 react in 3600 s. The value of the rate constant is

- A. $3.2 \times 10^{-4} \text{s}^{-1}$
- B. $6.4 \times 10^{-4} \text{s}^{-1}$
- C. $8.5 \times 10^{-4} \text{s}^{-1}$
- D. $12.8 \times 10^{-4} \text{s}^{-1}$

Answer: B

 [Watch Video Solution](#)

56. At 500 K, the half-life period of a gaseous reaction at the initial pressure of 80 kPa is 350 sec. When the pressure is 40 kPa, the half life period is 175 sec. The order of reaction is

- A. zero
- B. one
- C. two
- D. three

Answer: A

 [Watch Video Solution](#)

57. The energies of activation for forward and reverse reaction for $A_2 + B_2 \rightleftharpoons 2AB$ are 180 kJ mol^{-1} and 200 kJ mol^{-1} respectively. The presence of catalyst lowers the activation energy of both (forward and reverse) reactions by 100 kJ mol^{-1} . The enthalpy change of the reaction $A_2 + B_2 \rightleftharpoons 2AB$ in the presence of catalyst will be (in kJ mol^{-1}):

- A. 300
- B. 120
- C. 280
- D. -20

Answer: D

 [Watch Video Solution](#)


58. A reaction was found to be second order with respect to the concentration of carbon monoxide. If the concentration of carbon monoxide is doubled, with everything else kept the same, the rate of reaction will:

- A. remain unchanged
- B. triple
- C. increase by a factor of four
- D. double

Answer: C

 [Watch Video Solution](#)

59. For the second order reaction,



when a moles of A react with b moles of B, the rate equation is given by

$$k_2 t = \frac{1}{a-b} \ln \frac{b(a-x)}{a(b-x)}$$

when $a = b$, the rate expression becomes that of

- A. first order
- B. zero order
- C. unchanged, second order
- D. third order

Answer: A

 [Watch Video Solution](#)

60. $A(g) \rightarrow P(g) + Q(g) + R(g)$ follows first order kinetics with a half-life of 69.3 s at 500°C . Starting from the gas A enclosed in a container at 500°C and at a pressure of 0.4 atm, the total pressure of the system after 230 s will be

- A. 1.15 atm
- B. 1.32 atm
- C. 1.22 atm
- D. 1.12 atm

Answer: D

 [Watch Video Solution](#)

61. The value of rate constant for a first order reaction is $2.303 \times 10^{-2} \text{ sec}^{-1}$. What will be time required to reduce the concentration to $\frac{1}{10}$ th of its initial concentration ?

- A. 100 s
- B. 10 s
- C. 2303 s
- D. 230.3 s

Answer: A

 [Watch Video Solution](#)

62. The rate of a certain reaction is given by , rate $=K [H^{(+)}]^{(n)}$. The rate increases 100 times when the pH changes from 3 to 1 . The order (n) of the reaction is _____.

- A. 3
- B. 0
- C. 1
- D. 1.5

Answer: C

 [Watch Video Solution](#)

63. For the elementary reaction $M \rightarrow N$, the rate of disappearance of M increases by a factor of 8 upon doubling the concentration of M . The order of the reaction with respect to M is

- A. 4
- B. 3
- C. 2
- D. 1

Answer: B

 [Watch Video Solution](#)

64. The initial rate, $-(d[A])/dt$ at $t=0$ was found to be 2.6×10^{-2} mol $L^{-1} s^{-1}$ for the reaction $A + 2B \rightarrow \text{products}$

The initial rate, $-(d[B])/dt$, at $t=0$ is

- A. $0.10 \text{ mol L}^{-1} \text{ s}^{-1}$
- B. $2.6 \times 10^2 \text{ mol L}^{-1} \text{ s}^{-1}$
- C. $5.2 \times 10^2 \text{ mol L}^{-1} \text{ s}^{-1}$
- D. $6.5 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$

Answer: C

 [Watch Video Solution](#)

65. A chemical reaction was carried out at 300 K and 280 K. The rate constants were found to be k_1 and k_2 respectively. Then

- A. $k_2 = 0.25 k_1$
- B. $k_2 = 0.5 k_1$
- C. $k_2 = 4 k_1$
- D. $k_2 = 2 k_1$

Answer: A

 [Watch Video Solution](#)

66. 75 % of first order reaction is complete in 30 minutes. What is the time required for 93.75 % of the reaction (in minutes) ?

- A. 45
- B. 120
- C. 90
- D. 60

Answer: D

 [Watch Video Solution](#)

67. The half-life of a reaction is halved as the initial concentration of the reaction is doubled. The order of the reaction is

- A. 0.5
- B. 1
- C. 2
- D. 0

Answer: C

 [Watch Video Solution](#)

68. The half-life of 2 sample are 0.1 and 0.4 seconds. Their respective concentration are 200 and 50 respectively. What is the order of the reaction

- A. 0

B. 2

C. 1

D. 4

Answer: B

 [Watch Video Solution](#)

69. A first order reaction is 60% complete in 20 min. How long will the reaction take to be 84% complete?

A. 68 min

B. 40 min

C. 76 min

D. 54 min

Answer: B

 [Watch Video Solution](#)

70. An organic compound undergoes first decomposition. The time taken for its decomposition to $\frac{1}{8}$ and $\frac{1}{10}$ of its initial concentration are $t_{(1/8)}$ and $t_{(1/10)}$, respectively. What is the value of $\frac{t_{(1/8)}}{t_{(1/10)}} \times 10^2$? ($\log_{10} 2 = 0.3$)

A. 2

B. 3

C. 3

D. 9

Answer: D

 [Watch Video Solution](#)

71. Which graph represents zero-order reaction $[A(g) \rightarrow B(g)]$?



A.



B.



C.





D.


Answer: C

 [Watch Video Solution](#)

72. The reaction, $2\text{NO} + \text{Br}_2 \rightarrow 2\text{NOBr}$, is supposed to follow the following mechanism,



suggest the rate law expression.



Answer: A

 [Watch Video Solution](#)

73. The inversion of a sugar follows first order rate equation which can be followed by noting the change in the rotation of the plane of polarization of light in the polarimeter. If r_{∞} , r_t and r_0 are the rotations at $t = \infty$, $t = t$, and $t = 0$, then the first order reaction can be written as

A. $k = \frac{1}{t} \ln \frac{r_t - r_{\infty}}{r_0 - r_{\infty}}$

B. $k = \frac{1}{t} \ln \frac{r_0 - r_{\infty}}{r_t - r_{\infty}}$

C. $k = \frac{1}{t} \ln \frac{r_0 - r_t}{r_{\infty} - r_t}$

D. $k = \frac{1}{t} \ln \frac{r_{\infty} - r_t}{r_{\infty} - r_0}$

Answer: B

 [Watch Video Solution](#)

74. The unit of the rate constant for first order reaction is

A. mol^{-1}

B. s^{-1}

C. $\text{s}^{-1} \text{mol}^{-1} \text{dm}^3$

D. $\text{mol}^{-1} \text{s}^{-1} \text{dm}^3$

Answer: B

 [Watch Video Solution](#)

75. Which of the following represent the expression for $\frac{3}{4}$ th life of first order reaction

A. $\frac{2.303}{k} \log \frac{4}{3}$

B. $\frac{2.303}{k} \log \frac{3}{4}$

C. $\frac{2.303}{k} \log 4$

D. $\frac{2.303}{k} \log 3$

Answer: A

 [Watch Video Solution](#)

MHT CET CORNER

1. The rate constant and half - life of a first order reaction are related to each other as _____.

A. $t_{1/2} = \frac{0.693}{k}$

B. $t_{1/2} = 0.693k$

C. $k = 0.693 t_{1/2}$

D. $kt_{1/2} = \frac{1}{0.693}$

Answer: A

 [Watch Video Solution](#)

2. Average rate of reaction for the following reaction, $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{SO}_3(\text{g})$ is written as

A. $\frac{\Delta[\text{SO}_2]}{\Delta t}$

B. $-\frac{\Delta[\text{O}_2]}{\Delta t}$

C. $\frac{1}{2}\frac{\Delta[\text{SO}_2]}{\Delta t}$

D. $\frac{\Delta[\text{SO}_3]}{\Delta t}$

Answer: B

 [Watch Video Solution](#)

3. For the reaction $\text{O}_3(\text{g}) + \text{O}(\text{g}) \rightarrow 2\text{O}_2(\text{g})$, if the rate law expression is, rate $= k [\text{O}_3][\text{O}]$ the molecularity and order of the reaction are respectively _____.

A. 2 and 2

B. 2 and 1.33

C. 2 and 1

D. 1 and 2

Answer: A

 [Watch Video Solution](#)

4. The relationship between rate constant and half life period of zero order reaction is given by _____.

A. $t_{1/2} = \frac{[\text{A}]_0}{2k}$

B. $t_{1/2} = \frac{0.693}{k}$

C. $t_{1/2} = ([A]_0) / (k)$

D. $t_{1/2} = ([A]_0) / (2k)$

Answer: C

 [Watch Video Solution](#)

5. Half life period of a first order reaction $A \rightarrow$ product is 6.93 hour . What is the value of rate constant ?

A. $1.596h^{-1}$

B. $0.1h^{-1}$

C. $4.802h^{-1}$

D. $10h^{-1}$

Answer: B

 [Watch Video Solution](#)

6. Rate law for the reaction $A + B \rightarrow$ product is rate $= k [A]^2 [B]$. What is the rate of reaction at a given temperature is $0.22Ms^{-1}$, when $[A]=1 M$ and $[B]=0.25 M$?

A. $3.52M^{-2}s^{-1}$

B. $0.88M^{-2}s^{-1}$

C. $1.136M^{-2}s^{-1}$

D. $0.05M^{-2}s^{-1}$

Answer: B

 [Watch Video Solution](#)

7. What is the activation energy for a reaction if its rate doubles when the temperature is raised from 20°C to 35°C ? ($R = 8.314 \text{ J mol}^{-1}\text{K}^{-1}$)

- A. 342 kJ mol^{-1}
- B. 269 kJ mol^{-1}
- C. 34.7 kJ mol^{-1}
- D. 15.1 kJ mol^{-1}

Answer: C

 [Watch Video Solution](#)

8. In a multistep reaction, the overall rate of reaction is equal to the

- A. rate of slowest step
- B. rate of fastest step
- C. average rate of various step
- D. the rate of last step

Answer: A

 [Watch Video Solution](#)

9. The first order integrated rate equation is

A. $k = x/t$

B. $k = -(2.303)/t \log(a)/(a-x)$

C. $k = (1)/t \ln(a)/(a-x)$

D. $k = (1)/t(x)/(a(a-x))$

Answer: C

 [Watch Video Solution](#)

10. If the concentration is expressed in moles per liter, the unit of the rate constant for a first-order reaction is

A. $\text{mol}^{-1} \text{L}^{-1} \text{s}^{-1}$

B. s^{-1}

C. $\text{L}^{-1} \text{mol}^{-1} \text{s}^{-1}$

D. $\text{L}^{-2} \text{mol}^{-2} \text{s}^{-1}$

Answer: B

 [Watch Video Solution](#)

11. The unit of rate constant for a zero order reaction is s^{-1} .

- A. Zero order
- B. First order
- C. Second order
- D. third order

Answer: B

 [Watch Video Solution](#)

12. Which is a correct integrated rate equation?

- A. $k = -\frac{2.303}{t} \log \frac{a}{a-x}$
- B. $k = -\frac{2.303}{t} \log \frac{a-x}{a}$
- C. $-d(a-x) = kdt$
- D. All are integrated rate equations

Answer: B

 [Watch Video Solution](#)

13. After how many seconds will the concentration of the reactant in a first order reaction be halved if the rate constant is $1.155 \times 10^{-3} \text{ s}^{-1}$?

- A. 600
- B. 100

C. 60

D. 10

Answer: A

 [Watch Video Solution](#)

14. A certain nuclide has a half life period of 30 min. If a sample containing 600 atoms is allowed to decay for 90 min, how many atoms will remain?

A. 200 atoms

B. 450 atoms

C. 75 atoms

D. 150 atoms

Answer: C

 [Watch Video Solution](#)

15. If 50% of a radioactive substance dissociates in 15 min, then the time taken by substance to dissociate 99% will be

A. 50 min

B. 100 min

C. 99 min

D. 150 min

Answer: C

 [Watch Video Solution](#)

16. The disintegration constant of radium with half-life 1600 yr is

A. $2.12 \times 10^{-4} \text{ yr}^{-1}$

B. $4.33 \times 10^{-4} \text{ yr}^{-1}$

C. $3.26 \times 10^{-3} \text{ yr}^{-1}$

D. $4.33 \times 10^{-12} \text{ yr}^{-1}$

Answer: B

 [Watch Video Solution](#)

17. The half-life period of a radioactive element is 1 h. After 3 h, what fraction of it will remain?

A. $\frac{1}{8}$

B. $\frac{1}{16}$

C. $\frac{1}{64}$

D. $\frac{1}{9}$

Answer: A

 [Watch Video Solution](#)

18. If the half-life period of a first order reaction is 138.6 min, then the value of decay constant for the reaction will be (in min^{-1})

- A. 5
- B. 0.5
- C. 0.05
- D. 0.005

Answer: D

 [Watch Video Solution](#)

19. 20 mg of C-14 has half-life of 5760 yr. 100 mg of sample containing C-14 is reduced to 25 mg in

- A. 280 yr
- B. 1440 yr
- C. 2880 yr
- D. 11520 yr

Answer: D

 [Watch Video Solution](#)

20. If a is the initial concentration then time required to decompose half of the substance for n th order is inversely proportional to:

- A. $a^{(n-2)}$
- B. $a^{(1-n)}$
- C. $a^{(n-1)}$
- D. $a^{(n)}$

Answer: B

 [Watch Video Solution](#)

EXAMPLE

1. A reaction is of second order with respect to a reactant . How is its effected if the concentration of the reactant is (i)doubled (ii) reduced to half ?

- A. Becomes 2 times and $(1/2)$ times respectively
- B. Becomes $(1/2)$ times and $(1/2)$ times respectively
- C. Becomes 2 times and 4 times, respectively
- D. Becomes 4 time and 2 times, respectively


 [Watch Video Solution](#)

2. The decomposition of NH_3 on platinum surface is zero order reaction the rate of production of H_2 is $(k=2.5 \times 10^{-4} \text{ M s}^{-1})$

- A. 3.35×10^{-4} and 1.25×10^{-4} , respectively
- B. 1.25×10^{-4} and 3.75×10^{-4} , respectively
- C. 3.75×10^{-3} and 2.45×10^{-3} , respectively
- D. 1.25×10^{-3} and 3.25×10^{-3} , respectively

 [Watch Video Solution](#)

3. The initial concentration of N_2O_5 in the following first order reaction:




was $1.24 \times 10^{-2} \text{ mol L}^{-1}$ at 318 K . The concentration of N_2O_5 after 60 min was $0.20 \times 10^{-2} \text{ mol L}^{-1}$. Calculate the rate constant of the reaction at 318 K .

- A. 0.0304 min^{-1}
- B. 0.0204 min^{-1}
- C. 0.0034 min^{-1}
- D. 1.0304 min^{-1}

 [Watch Video Solution](#)

4. The following data were obtained during the first thermal decomposition of $\text{N}_2\text{O}_5(\text{g})$ at constant volume.



{("S.No.", "Time (s)", "Total pressure (atm)"), (i., 0, 0.5), (ii., 100, 0.512):}

Calculate the rate constant.

A. $3.39 \times 10^{-4} \text{ s}^{-1}$

B. $1.39 \times 10^{-5} \text{ s}^{-1}$

C. $5.45 \times 10^{-4} \text{ s}^{-1}$

D. $4.91 \times 10^{-4} \text{ s}^{-1}$

 [Watch Video Solution](#)

5. A first order reaction takes 40 min for 30% decomposition. Calculate $t_{1/2}$. (Given $\log 7 = 0.845$)

A. 77.78 min

B. 78.34 min

C. 84.36 min

D. 65.34 min

 [Watch Video Solution](#)