



# MATHS

## BOOKS - NCERT MATHS (ENGLISH)

### STATISTICS

#### Short Answer Type Questions

1. Find the mean deviation about the mean of the distributon .

|                  |    |    |    |    |    |
|------------------|----|----|----|----|----|
| <b>Size</b>      | 20 | 21 | 22 | 23 | 24 |
| <b>Frequency</b> | 6  | 4  | 5  | 1  | 4  |



Watch Video Solution

2. Find the mean deviation about the mean of the distribution .

|                           |    |    |    |    |    |
|---------------------------|----|----|----|----|----|
| <b>Marks obtained</b>     | 10 | 11 | 12 | 14 | 15 |
| <b>Number of students</b> | 2  | 3  | 8  | 3  | 4  |



Watch Video Solution

3. Calculate the mean deviation about the mean of the set of first  $n$  natural numbers when  $n$  is odd natural number.

 [View Text Solution](#)

4. Calculate the mean deviation about the mean of the set of first  $n$  natural numbers when  $n$  is even natural number.

 [Watch Video Solution](#)

5. Find the standard deviation of first  $n$  natural numbers.

|         |   |   |   |    |    |     |     |       |
|---------|---|---|---|----|----|-----|-----|-------|
| $x_j$   | 1 | 2 | 3 | 4  | 5  | ... | ... | $n$   |
| $x_j^2$ | 1 | 4 | 9 | 16 | 25 | ... | ... | $n^2$ |

6. The mean and standard deviation of some data for the time taken to complete a test are calculated with the following results

Number of observations = 25, mean = 18.2 s, standard deviation = 3.25 s. Further another set of 15 observations  $x_1, x_2, \dots, x_{15}$  also in seconds is now available and we have

$\sum_{i=1}^{15} x_i = 279$  and  $\sum_{i=1}^{15} x_i^2 = 5524$ . Calculate the standard deviation based on all 40 observations.



Watch Video Solution

7. The mean and standard deviation of a set of  $n_1$  observations are  $\bar{x}_1$  and  $s_1$  respectively while the mean and standard deviation of another set of  $n_2$  observations are  $\bar{x}_2$  and  $s_2$  respectively. Show that the standard deviation of the combined set of  $(n_1 + n_2)$  observations

is given by

$$SD = \sqrt{\frac{n_1(s_1)^2 + n_2(s_2)^2}{n_1 + n_2} + \frac{n_1 n_2 (\bar{x}_1 - \bar{x}_2)^2}{(n_1 + n_2)^2}}$$



View Text Solution

8. Two sets each of 20 observations have the same standard deviation 5. The first set has a mean 17 and second a mean 22. Then the standard deviation of the set obtained by combining the given two sets.



[View Text Solution](#)

9. The frequency distribution

|     |   |    |    |    |    |    |
|-----|---|----|----|----|----|----|
| $x$ | A | 2A | 3A | 4A | 5A | 6A |
| $f$ | 2 | 1  | 1  | 1  | 1  | 1  |



[View Text Solution](#)

[View Text Solution](#)

10. For the frequency distribution

|     |   |   |    |    |    |   |
|-----|---|---|----|----|----|---|
| $x$ | 2 | 3 | 4  | 5  | 6  | 7 |
| $f$ | 4 | 9 | 16 | 14 | 11 | 6 |



[View Text Solution](#)

11. For the frequency distribution

|                  |       |     |       |           |      |       |
|------------------|-------|-----|-------|-----------|------|-------|
| <b>Marks</b>     | 0     | 1   | 2     | 3         | 4    | 5     |
| <b>Frequency</b> | $x-2$ | $x$ | $x^2$ | $(x+1)^2$ | $2x$ | $x+1$ |



[View Text Solution](#)

**12.** The mean life of a sample of 60 bulbs was 650 h and the standard deviation was 8 h, If a second sample of 80 bulbs has a mean life of 660 h and standard deviation 7 h then find the over all standard deviation



[View Text Solution](#)

**13.** If mean and standard deviation of 100 items are 50 and 4 respectively the find the sum of all the item and the sum of the squares of item.





[View Text Solution](#)

14. If for distribution of 18 observations

$$\sum (x_i - 5) = 3 \text{ and } \sum (x_i - 5)^2 = 43, \text{ find}$$

the mean and standard deviation.



[View Text Solution](#)

15. Find the mean and variance of the frequency distribution given below

|     |                   |                   |                   |                    |
|-----|-------------------|-------------------|-------------------|--------------------|
| $x$ | $1 \leq x \leq 3$ | $3 \leq x \leq 5$ | $5 \leq x \leq 7$ | $7 \leq x \leq 10$ |
| $f$ | 6                 | 4                 | 5                 | 1                  |



[View Text Solution](#)

## Long Answer Type Questions

1. Calculate the mean deviation about the mean for the following frequency distribution

|                       |     |     |      |       |       |
|-----------------------|-----|-----|------|-------|-------|
| <b>Class interval</b> | 0-4 | 4-8 | 8-12 | 12-16 | 16-20 |
| <b>Frequency</b>      | 4   | 6   | 8    | 5     | 2     |



[View Text Solution](#)

2. Calculate the mean deviation from the median of the following data

|                       |     |      |       |       |       |
|-----------------------|-----|------|-------|-------|-------|
| <b>Class interval</b> | 0-6 | 6-12 | 12-18 | 18-24 | 24-30 |
| <b>Frequency</b>      | 4   | 5    | 3     | 6     | 2     |



[View Text Solution](#)

3. Determine the mean and standard deviation for the following distribution

|                  |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |
|------------------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|
| <b>Marks</b>     | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 |
| <b>Frequency</b> | 1 | 6 | 6 | 8 | 8 | 2 | 2 | 3 | 0  | 2  | 1  | 0  | 0  | 0  | 1  |



[View Text Solution](#)

4. The weights of coffee in 70 jars is shown in the following table

| <b>Weight (in g)</b> | <b>Frequency</b> |
|----------------------|------------------|
| 200-201              | 13               |
| 201-202              | 27               |
| 202-203              | 18               |
| 203-204              | 10               |
| 204-205              | 1                |
| 205-206              | 1                |

Determine variance and standard deviation of the above distribution

 [View Text Solution](#)

5. Determin mean and standard deviation of first n terms of an AP whose first term is a and common difference is d.

| $x_i$                                    | $x_i - a$ | $(x_i - a)^2$ |
|--|-----------|---------------|
| $a$                                      | $0$       | $0$           |
| $a + d$                                  | $d$       | $d^2$         |
| $a + 2d$                                 | $2d$      | $4d^2$        |
| .....                                    | .....     | $9d^2$        |
| .....                                    | .....     | .....         |
| .....                                    | .....     | .....         |
| $a + (n-1)d$                             | $(n-1)d$  | $(n-1)^2 d^2$ |
| $\Sigma x_i = \frac{n}{2} [2a + (n-1)d]$ |           |               |



**Watch Video Solution**

6. Following are the marks obtained ,out of 100, by two student Ravi and Hashina in 10

## tests

|                |    |    |    |    |    |    |    |    |    |    |
|----------------|----|----|----|----|----|----|----|----|----|----|
| <b>Ravi</b>    | 25 | 50 | 45 | 30 | 70 | 42 | 36 | 48 | 35 | 60 |
| <b>Hashina</b> | 10 | 70 | 50 | 20 | 95 | 55 | 42 | 60 | 48 | 80 |

Who is more intellogent and who is more consistent ?



[View Text Solution](#)

7. Mean and standard deviation of 100 observations were found to be 40 and 10 respecitly .If at the time of calculation two observations were wrongly taken as 30 ans 70

in place of 3 and 27 respectively , then find the correct standard deviation.



[Watch Video Solution](#)

8. While calculating the mean and variance of 10 readings, a student wrongly used the reading 52 for the correct reading 25. He obtained the mean and variance as 45 and 16 respectively. Find the correct mean and the variance .



[Watch Video Solution](#)

## Objective Type Question

1. The mean deviation of the data 3,10,10,4,7,10,5 from the mean is

- A. 2
- B. 2.57
- C. 3
- D. 3.75

**Answer: B**



**Watch Video Solution**



2. Mean deviation for n observation

$x_1, x_2, \dots, x_n$  from their mean  $\bar{x}$  is given by

A.  $\sum_{i=1}^n (x_i - \bar{x})$

B.  $\frac{1}{n} \sum_{i=1}^n |x_i - \bar{x}|$

C.  $\sum_{i=1}^n (x_i - \bar{x})^2$

D.  $\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$

**Answer: B**



**Watch Video Solution**

3. when tested the lives (in hours) of 5 bulbs were noted as follows 1357,1090,1666,1494,1623

The mean deviations (in hours ) from their mean is

A. 178

B. 179

C. 220

D. 356

**Answer: A**



**Watch Video Solution**

4. Following are the marks obtained by 9 student in a mathematics test

50,69,20,33,53,39,40,65,59,

The mean deviation from the median is

A. 9

B. 10.5

C. 12.67

D. 14.76

**Answer: C**



**Watch Video Solution**

5. The standard deviation of data 6,5,9,13,12,8 and 10 is

A.  $\sqrt{\frac{52}{7}}$

B.  $52/7$

C.  $\sqrt{6}$

D. 6

**Answer: A**



**Watch Video Solution**

**6.** If  $x_1, x_2, \dots, x_n$  be  $n$  observations and  $\bar{x}$  be their arithmetic mean. Then formula of the standard deviation is given by

A.  $\Sigma(x_i - \bar{x})^2$

B.  $\frac{\Sigma(x_i - \bar{x})^2}{n}$

C.  $\sqrt{\frac{\Sigma(x_i - \bar{x})^2}{n}}$

D.  $\sqrt{\frac{\sum x^2 i}{n} + \bar{x}^{-2}}$

**Answer: C**



**Watch Video Solution**

7. If the mean of 100 observations is 50 and their standard deviations is 5, then the sum of all squares of all the observations is

A. 50000

B. 250000

C. 252500

D. 255000

**Answer: C**



**Watch Video Solution**

**8.** Let  $a, b, c, d, e$ , be the observations with mean  $m$  and standard deviation  $s$ . The standard deviation of the observations  $a+k, b+k, c+k, d+k, e+k$  is (a)  $s$  (b)  $ks$  (c)  $s + k$  (d)  $\frac{s}{k}$

A.  $s$

B.  $ks$

C.  $s+k$

D.  $\frac{s}{k}$

**Answer: A**



**Watch Video Solution**

9. If,  $s$  is the standard deviation of the observations  $x_1, x_2, x_3, x_4$  and  $x_5$  then the



standard deviation of the observations

$kx_1, kx_2, kx_3, kx_4$  and  $kx_5$  is

A.  $k + s$

B.  $\frac{s}{k}$

C.  $ks$

D.  $s$

**Answer: C**



**Watch Video Solution**

**10.** Let  $x_1, x_2, \dots, x_n$  be  $n$  observations. Let  $w_i = lx_i + k$  for  $i = 1, 2, \dots, n$ , where  $l$  and  $k$  are constants. If the mean of  $x_i$  is 48 and their standard deviation is 12 the mean of  $w_i$ 's is 55 and standard deviation of  $w_i$  is 15 then the value of  $l$  and  $k$  should be

A.  $l=1.25, k=-5$

B.  $l=-1.25, k=5$

C.  $l=2.5, k=-5$

D.  $l=2.5, k=5$

**Answer: A**



**Watch Video Solution**

**11.** The standard deviation for first 10 natural number is

A. 5.5

B. 3.87

C. 2.97

D. 2.87

**Answer: D**



**Watch Video Solution**

**12.** Consider the number 1,2,3,4,5,6,7,8,9 and 10. If 1 is added to each number the variance of the number so obtained is

A. 6.5

B. 2.87

C. 3.87

D. 8.25

**Answer: D**



**Watch Video Solution**

**13.** Consider the first 10 positive integers. If we multiply each number by  $-1$  and then add 1 to each number, the variance of the number so obtained

A. 8.25

B. 6.5

C. 3.87

D. 2.87

**Answer: A**



**Watch Video Solution**

**14.** If for a sample of size 60, we have the following information  $\sum (x_i)^2 = 18000$  and  $\sum x_i = 960$ , then the variance is

A. 6.63

B. 16

C. 22

D. 44

**Answer: D**



**Watch Video Solution**

**15.** If the coefficient of variation of two distribution are 50 ,60 and their arithmetic means are 30 and 25 respectively then the difference of their standard deviation is

A. 0

B. 1

C. 1.5

D. 2.5

**Answer: A**



**Watch Video Solution**

**16.** The standard deviation of some temperature data in  $^{\circ}C$  is 5. If the data were converted into  $^{\circ}F$  then variance would be



A. 81

B. 57

C. 36

D. 25

**Answer: A**



**Watch Video Solution**

**Fillers**

1. Coefficient of variation =  $\frac{\dots}{\text{Mean}} \times 100$

A. SD

B. MD

C. MEDIAN

D. none of these

**Answer: A**



**Watch Video Solution**

2. If  $\bar{x}$  is the mean of  $n$  values of  $x$ , then

$\sum_{i=1}^n (x_i - \bar{x}) = 0$  and if  $a$  has a value other

than

$\bar{x}$  then  $\sum_{i=1}^n (x_i - \bar{x})^2$  is less than  $\sum (x_i - a)^2$



[Watch Video Solution](#)

3. If the variance of a data is 121, then the standard deviation of the data is .....

A. 11

B. 12

C. 10

D. 9

**Answer: A**



**Watch Video Solution**

4. The standard deviation of a data is .... Of any change in origin but is ....of change of scale .



**Watch Video Solution**

5. The sum of squares of the deviation of the values of the variable is \_\_\_\_\_ when taken about their arithmetic mean



[Watch Video Solution](#)

6. The mean deviation of the data is .... When measured from the median



[Watch Video Solution](#)

7. The standard deviation is .... To the mean deviation taken from the arithmetic mean



**Watch Video Solution**