





CHEMISTRY

BOOKS - NTA MOCK TESTS

JEE MOCK TEST 5



1. Among the following compounds the correct order of

basicity is



$$A.$$







Answer: C



2. Benzene and naphthalene form ideal solution over the entire range of composition. The vapour pressure of pure benzene and naphthalene at 300 K are 50.71 mm Hg and 32.06 mm Hg respectively. Calculate the mole fraction of benzene in vapour phase if 80 g of benzene is mixed with 100 g of naphthalene.

A. 0.0675

B. 0.675

C. 0.35

D. 0.5

Answer: B





3. The pressure of H_2 required to make the potential of

 $H_{2^-}\,$ electrode zero in pure water at $298K\,{\rm is}$

A. $10^{-14} \ \mathrm{atm}$

B. 10^{-12} atm

C. 10^{-10} atm

D. 10^{-4} atm



4. Calculate ΔH_f° for chloride ion from the following data :

$$egin{aligned} &rac{1}{2} H_{2\,(\,g\,)} \,+\, rac{1}{2} Cl_{2\,(\,g\,)} \, o HCl_{\,(\,g\,)}\,, \Delta H_{f}^{\,\circ} \,=\,\, -\,\, 92.4 kJ \ &HCl_{\,(\,g\,)} \,+\, nH_{2}O \, o \, H_{\,(\,aq\,.\,\,)}^{\,+} \,+\, Cl_{\,(\,aq\,.\,\,)}^{\,-}\,, \Delta H^{\,\circ} \,=\,\, -\,\, 74.8 kJ \ &\Delta H_{f}^{\,\circ}\, H_{\,(\,aq\,.\,\,)}^{\,+} \,=\, 0.0 kJ \end{aligned}$$

 $\mathrm{A.}-167.2~\mathrm{kJ}$

 $\mathrm{B.}-165.2\,\mathrm{kJ}$

 $\mathrm{C.}-157.2~\mathrm{kJ}$

 $\mathrm{D.}-147.2~\mathrm{kJ}$



5. Complete the following chemical reaction equations :

- (i) $NaOH_{(ext{cold and dilute})} + Cl_2
 ightarrow$
- (ii) $XeF_6 + H_2O
 ightarrow (ext{excess})$

A. NaOH and XeO_3

- B. $HClO_3$ and XeO_2F_2
- C. $NaClO_3$ and XeO_3
- D. None of these

Answer: C



6. Select the nature or type of redox change in the following reaction - $Cl_2 \rightarrow ClO^- + Cl^-$

A. Disproportionation

B. Intramolecular redox

C. Intermolecular redox

D. None of the above



7. The K_{sp} of $FeS=4 imes10^{-19}$ at 298 K. The minimum concentration of H^+ ions required to prevent the precipitation of FeS from a 0.01 M solution Fe^{2+} salt by passing $H_2S(0.1M)$ (Given $H_2Sk_{a_1} imes k_{b_1}=10^{-21}$)

A.
$$1.6 imes 10^{-3}$$
 M

B.
$$2.5 imes 10^{-4}$$
 M

C.
$$2.0 imes 10^{-2}$$
 M

D.
$$1.2 imes 10^{-4}$$
 M

Answer: A

Watch Video Solution

8. Pb and Sn are extracted from their chief ore by :

A. Carbon reduction and self reduction

B. self reduction and carbon reduction

C. Electryloysis and self reduction.

D. Self reduction and electrolysis.

Answer: B

Watch Video Solution

9. The carbonate of which of the following cation is soluble in water ?

A. Na^+

 $\mathsf{B.}\,K^{\,+}$

C. NH_4^+

D. Ca^{2+}

Answer: D

Watch Video Solution

10. The lewis structure of allene is



Which of the following statements correctly gives answers for all 3 parts:

(a) Is the molecule planar?

(b) Does 1,3- dichloro propadiene show geometrical isomerism?

(c) is the molecule 1,3 - dichloro propadiene polar?

A. (i) Non-planar, (ii) No geometrical isomersim , (iii) polar

B. (i) planar, (ii) No geometrical isomerism, (iii) polar

C. (i) planar, (ii) Yes geometrical isomerism, (iii) polar

D. (i) Non planar, (ii) No geometrical isomerism, (iii)

Not polar



11. In orthorhombic , the value of a, b and c are respectively 4.2\AA , 8.6\AA and 8.3\AA .Given the molecular mass of the solute is 155gm/mol and that of density is $3.3gm/cm^3$ the number of formula unit per unit cell is

A. 2

B. 3

C. 4

D. 6

Answer: C



12. Determine the number of planes of symmetry of the

given compound



A. 1

B. 2

C. 3

D. 4



13. Gaseous benzene reacts with hydrogen gas in the presence of nickel catalyst to give gaseous cyclohexane. A mixture of benzene vapour and hydrogen had a pressure of 60 mm Hg in vessel. After all benzene converted to cyclohexane, the pressure of the gas was 30 mm Hg in the same volume and at the same temperature. What fraction (by mole) of the original mixture was benzene?

A. 0.167

B. 0.333

C. 0.666

D. 1

Answer: A

14. Correct acidic strength of given acids is:-

$$H_{3}C - C_{I}^{||} - OH$$

$$H_{3}C - C_{I}^{||} - OH$$

$$(II) HO - C - C - OH$$

$$(III) HO - C - CH_{2} - OH$$

 $\mathsf{A.I} \ > \ \mathsf{II} \ > \ \mathsf{III}$

 $\mathsf{B}.\,II>III>I$

 $\mathsf{C}.\,II>I>III$

D. III > II > I

Answer: B



15. Larger number of oxidation states are exhibited by the actinoids then those by the lanthanoids, the main reason being

A. More reactive nature of the actinides than the lanthanides

B. 4f orbitals more diffused than the 5f orbitals

C. More energy difference between 5f and 6d than

between 4f and 5d orbitals

D. lesser energy difference between 5f and 6d than

between 4f and 5d orbitals.

Answer: D

Α.

Watch Video Solution

16. Hydroboration oxidation and acid hydration will yield

the same product in case of:





 $\mathsf{C}.\,CH_2=CH_2$

D. $CH_3CH = CH - CH_3$

Answer: A

Β.

Watch Video Solution

17. According to kinetic theory of gases in an ideal gas between two successive collisions a gas molecule travels

A. In a circular path

B. In a wavy path

C. In a straight line path

D. With an accelerated velocity

Answer: C

Watch Video Solution

18. The equilibrium constant for a reaction $A + B \Leftrightarrow C + D$ is 1×10^{-2} at 298K and is 2 at 273K. The chemical process resulting in the formation of C and D is

A. Exothermic

B. Endothermic

C. Unpredictable

D. There is no relationship between ΔH and K

Answer: A



A. 1-Ethoxy -1-ethyl-1-aminopropane

B. 1-Ethoxy-1-amino-1-ethylpropane

C. 1-Ethoxy-2-butanol

D. 2-Ethoxybutan-2-amine

Answer: D



20. Second ionization potential value is very low for

A. sodium

B. magnesium

C. fluorine

D. oxygen

Answer: B

21. Adsorption of a gas follows Freundlich adsorption isotherm. In the given plot,x is the mass of the gas adsorbed on mass m of the adsorbent at pressure P. If $\frac{x}{m} \propto P^{\frac{1}{y}}$ find the magnitude of y is:



22. Washing soda is Na_2CO_3 . xH_2O . The value of x is



23. How many among the following can exhibit linkage

isomerism?





25. The highest oxidation state exhibited by a transition

metal is

