



MATHS

BOOKS - NTA MOCK TESTS

NTA JEE MOCK TEST 101

Mathematics

1. If the area covered by $y = \frac{2}{x}$ and $y = \frac{2}{2x - 1}$ from $x = 1$ or $x = e$ is

In (a) sq. units, then $(2e - 1)^2 a^2$ is equal to

A. $\frac{e}{2}$

B. e^4

C. 1

D. e

Answer: B



Watch Video Solution

2. If a, b, c are real numbers satisfying the condition $a + b + c = 0$ then the roots of the quadratic equation $3ax^2 + 5bx + 7c = 0$ are

- A. Positive
- B. Negative
- C. real and equal
- D. distinct but not imaginary

Answer: D



Watch Video Solution

3. The indefinite integral $I = \int \frac{(\sin^2 x - \cos^2 x)^{2019}}{(\sin x)^{2021} (\cos x)^{2021}} dx$ simplifies to

(where c is an integration constant)

A. $\frac{(\sin^2 x - \cos^2 x)^{2020}}{2020} + c$

$$B. \frac{(\tan x - \cot x)^{2020}}{2020} + c$$

$$C. \frac{(\sin x - \cos x)^{2020}}{2020} + c$$

$$D. \frac{(\tan^2 x + \cot^2 x)^{2020}}{2020} + c$$

Answer: B

 [Watch Video Solution](#)

4. Coefficient of α^t in the expansion of $(\alpha + p)^{m-1} + (\alpha + p)^{m-2}(\alpha + q) + (\alpha + p)^{m-3}(\alpha + q)^2 + \dots + (\alpha + p)$, where $\alpha \neq -q$ and $p \neq q$ is

$$A. \frac{{}^m C_t (t^t - q^t)}{p - q}$$

$$B. \frac{{}^m C_t (p^{m-1} - q^{m-1})}{p - q}$$

$$C. \frac{{}^m C_t (p^t - q^t)}{p - q}$$

$$D. \frac{{}^m C_t (p^{m-t} + q^{m-t})}{p - q}$$

Answer: B



Watch Video Solution

5. The solution of the differential equation $\frac{dy}{dx} = \frac{2x - y}{x - 6y}$ is (where c is an arbitrary constant)

A. $4xy = x^2 - 3y + c$

B. $2xy = x^2 + 3y^2 + c$

C. $xy = x^2 + 3y^2 + c$

D. $xy = x^2 + c$

Answer: C



Watch Video Solution

6. Let a_1, a_2, a_3 be three positive numbers which are in geometric progression with common ratio r . The inequality $a_3 > a_2 + 2a_1$ holds true if r is equal to

A. 2

B. 1.5

C. 0.5

D. 2.5

Answer: D



Watch Video Solution

7. The arithmetic mean of the numbers

$2\sin 2^\circ, 4\sin 4^\circ, 6\sin 6^\circ, \dots, 178\sin 178^\circ, 180\sin 180^\circ$

A. $\sin 1^\circ$

B. $\cot 1^\circ$

C. $\tan 1^\circ$

D. $\cos 1^\circ$

Answer: B

 [Watch Video Solution](#)

8. If PQ is the focal chord of the parabola $y^2 = -x$ and P is $(-4, 2)$, then the ordinate of the point of intersection of the tangents at P and Q is

A. $\frac{15}{16}$

B. $\frac{7}{8}$

C. 4

D. $\frac{17}{18}$

Answer: A

 [Watch Video Solution](#)

9. Consider $f(x) = \begin{cases} [x] + [-x] & x \neq 2 \\ \lambda & x = 2 \end{cases}$ where $[.]$ denotes the greatest integer function. If $f(x)$ is continuous at $x = 2$ then the value of λ is equal to

A. -1

B. 0

C. 1

D. No value of possible

Answer: A



[Watch Video Solution](#)

10. Consider the statement p : If a hexagon is regular then all its sides and angles are equal. The contrapositive of statement p is

A. If all sides and angles of a hexagon are not equal then it is not regular

B. If all sides and angles of a hexagon are equal then it is regular

C. If all sides and angles of a hexagon are not equal then it is regular

D. If all sides and angles of a hexagon are equal then it is not regular

Answer: A



Watch Video Solution

11. If $\sin(\cot^{-1}(1 - x)) = \cos(\tan^{-1}(-x))$, then x is

A. 1

B. $\frac{1}{2}$

C. 0

D. $-\frac{1}{2}$

Answer: B



Watch Video Solution

12. A circle C_1 has radius 2 units and a circles C_2 has radius 3 units. The distance between the centres of C_1 and C_2 is 7 units. If two points, one tangent to both circles and the other passing through the centre of both

circles, intersect at point P which lies between the centers of C_1 and C_2 ,

then the distance between P and the centre of C_2 is

A. $\frac{9}{4}$ units

B. $\frac{7}{3}$ units

C. $\frac{21}{5}$ units

D. $\frac{14}{5}$ units

Answer: C



Watch Video Solution

13. The number of values of x in the interval $[0, 3\pi]$ satisfying the equation $3\sin^2 x - 7\sin x + 2 = 0$ is

A. 0

B. 5

C. 4

D. 10

Answer: C



[Watch Video Solution](#)

14. From the point $P(3, 4)$ pair of tangents PA and PB are drawn to the ellipse $\frac{x^2}{16} + \frac{y^2}{9} = 1$. If AB intersects y - axis at C and x - axis at D, then OC. OD is equal to (where O is the origin)

A. 9

B. 18

C. 8

D. 12

Answer: D



[Watch Video Solution](#)

15. For two non - zero complex numbers A and B, if $A + \frac{1}{B} = \bar{A}$ and $\frac{1}{A} + B = \bar{B}$, then

A. A is purely real

B. B is purely real

C. $|AB| = \frac{1}{2}$

D. $|A| = |B|$

Answer: C



Watch Video Solution

16. The value of the determinant $\Delta = \begin{vmatrix} \sqrt{13} + \sqrt{3} & 2\sqrt{5} & \sqrt{5} \\ \sqrt{15} + \sqrt{26} & 5 & \sqrt{10} \\ 3 + \sqrt{65} & \sqrt{15} & 5 \end{vmatrix}$ is

equal to

A. $15\sqrt{2} - 25\sqrt{3}$

B. $25\sqrt{3} - 15\sqrt{2}$

C. $3\sqrt{5}$

D. $-15\sqrt{2} + 7\sqrt{3}$

Answer: A



Watch Video Solution

17. Two dice are thrown at a time, the probability that the absolute value of the difference of number on dice is 3, is

A. $\frac{3}{4}$

B. $\frac{5}{6}$

C. $\frac{2}{7}$

D. $\frac{1}{6}$

Answer: D



Watch Video Solution

18. Let $\vec{a}, \vec{b}, \vec{c}$ be three non-zero, non-coplanar vectors and $\vec{p}, \vec{q}, \vec{r}$ be three vectors given by

$$\vec{p} = \vec{a} + 2\vec{b} - 2\vec{c}, \vec{q} = 3\vec{a} + \vec{b} - 3\vec{c} \text{ and } \vec{r} = \vec{a} - 4\vec{b} + 4\vec{c}$$

. If the volume of parallelepiped determined by \vec{a}, \vec{b} and \vec{c} is v_1 cubic units and volume of tetrahedron determined by \vec{p}, \vec{q} and \vec{r} is v_2 cubic units, then $\frac{v_1}{v_2}$ is equal to

- A. $\frac{1}{3}$
- B. $\frac{3}{4}$
- C. $\frac{1}{4}$
- D. $\frac{1}{2}$

Answer: D



[Watch Video Solution](#)

19. The number of matrices X with entries $\{0, 2, 3\}$ for which the sum of all the principal diagonal elements of $X \cdot X^T$ is 28 (where X^T is the

transpose matrix of X), is

A. 12

B. 18

C. 36

D. 44

Answer: C



[Watch Video Solution](#)

20. Let $f(x) = 2 \tan^3 x - 6 \tan^2 x + 1 + \operatorname{sgn}(e^x)$, $\forall x \in \left[-\frac{\pi}{4}, \frac{\pi}{4} \right]$,

Then the positive difference between the least value and the local maximum value of the function is (where $\operatorname{sgn}(f(x))$ represents the signum function)

A. 7

B. 8

C. 9

Answer: B

 [Watch Video Solution](#)

21. If the integral $I = \int_0^{\pi} \frac{\sec^{-1}(\sec x)}{1 + \tan^8 x} dx, \forall x \neq \frac{\pi}{2}$, then the value of $[I]$ is equal to (where $[.]$ is the greatest integer function)

 [Watch Video Solution](#)

22. The value of $\lim_{x \rightarrow 0} \frac{\log(1 + 2x)}{5x} + \lim_{x \rightarrow 2} \frac{x^4 - 2^4}{x - 2}$ is equal to

 [Watch Video Solution](#)

23. The sum of square of the abscissas for all the points on the line $x + y = 4$ that lie at a unit distance from the line $4x + 3y - 10 = 0$ is

 [Watch Video Solution](#)

24. The line $\frac{x}{k} = \frac{y}{2} = \frac{z}{-12}$ makes an isosceles triangle with the planes $2x + y + 3z - 1 = 0$ and $x + 2y - 3z - 1 = 0$ then the value of k is-

 [Watch Video Solution](#)

25. If the total number of ways of selecting two numbers from the set $\{1, 2, 3, \dots, 89, 90\}$ such that their sum is divisible by 3 is k , then $\frac{k}{500}$ is

 [Watch Video Solution](#)