



# MATHS

## BOOKS - NTA MOCK TESTS

### NTA JEE MOCK TEST 103

#### Mathematics

1. The coefficient of  $x^4$  in the expansion  $(1 - x)^8(1 + x)^{12}$  is equal to

A. 1

B.  $-48$

C. 28

D.  $-19$

**Answer: D**



**Watch Video Solution**

2. For a complex number  $z$ , if  $\arg(z) \in (-\pi, \pi]$ , then

$\arg\left(1 + \cos.\frac{6\pi}{5} + I \sin.\frac{6\pi}{5}\right)$  is (here  $i^2 = -1$ )

A.  $\frac{3\pi}{5}$

B.  $\frac{2\pi}{5}$

C.  $-\frac{2\pi}{5}$

D.  $-\frac{3\pi}{5}$

**Answer: C**



**Watch Video Solution**

3. Consider the system of equations

$$x + 2y + 3z = 6,$$

$$4x + 5y + 6z = \lambda,$$

$$7x + 8y + 9z = 24.$$

Then, the value of  $\lambda$  for which the system has infinite solutions is

A. 15

B. 20

C. 16

D. 30

**Answer: A**



**Watch Video Solution**

4. Tibu and Babu are playing a game in which both the players roll a pair of dice. Tibu wins if the sum of the numbers appearing on the dice is a prime number while Babu wins if the product of the numbers appearing on the dice is prime. The chance that no one wins is (both players are allowed to win simultaneously)

A.  $\frac{19}{36}$

B.  $\frac{1}{2}$

C.  $\frac{17}{36}$

D.  $\frac{4}{9}$

**Answer: C**



**Watch Video Solution**

5. Let  $\alpha$ ,  $\beta$  and  $\gamma$  be the roots of equation  $x^3 + x + 1 = 0$ , then

$$\frac{\alpha\beta(\alpha + \beta) + \beta\gamma(\beta + \gamma) + \gamma\alpha(\gamma + \alpha)}{\alpha^2 + \beta^2 + \gamma^2}$$

is

equal to

A.  $\frac{3}{2}$

B. 1

C.  $-3$

D.  $-\frac{3}{2}$

**Answer: D**



**Watch Video Solution**

6. Let  $l$ ,  $m$  and  $n$  are three distinct numbers in arithmetic progression. Also  $l^2$ ,  $m^2$  and  $n^2$  are in geometric progression and  $l + m + n = 3$ . If  $l < m < n$ , then  $n$  is equal to

A. 1

B.  $1 - \sqrt{2}$

C.  $1 + \sqrt{2}$

D.  $2 + \sqrt{5}$

**Answer: C**



Watch Video Solution



7. The equation of incircle of the triangle formed by common tangents to the circles  $x^2 + y^2 = 4$  and  $x^2 + y^2 - 6x + 8 = 0$  is

A.  $x^2 + y^2 = 1$

B.  $x^2 + y^2 - 6x + 8 = 0$

C.  $x^2 + y^2 - 6x + 9 = 0$

D.  $x^2 - y^2 = 1$

**Answer: B**





8. The tangent to the ellipse  $\frac{x^2}{25} + \frac{y^2}{16} = 1$  at point P lying in the first quadrant meets x - axis at Q and y - axis at R. If the length QR is minimum, then the equation of this tangent is

A.  $2x + \sqrt{5}y = 6\sqrt{5}$

B.  $\sqrt{4}x + 2y = 6\sqrt{5}$

C.  $2x - \sqrt{5} = 6\sqrt{5}$

D.  $2\sqrt{5}x + y = 6\sqrt{5}$

**Answer: A**



**Watch Video Solution**

9. Let the tangents to the parabola  $y^2 = 4ax$  drawn from point P have slope  $m_1$  and  $m_2$ . If  $m_1 m_2 = 2$ , then the locus of point P is

A.  $x = a$

B.  $x = \frac{a}{2}$

C.  $x + a = 0$

D.  $x = 2a$

**Answer: B**



**Watch Video Solution**

**10.** In the triangle  $ABC$ , vertices  $A, B, C$  are  $(1, 2), (3, 1), (-1, 6)$  respectively. If the internal angle bisector of  $\angle BAC$  meets  $BC$  at  $D$ , then the coordinates of  $D$  are  $\left(\frac{5}{3}, \frac{8}{3}\right)$

A.  $\left(1, \frac{7}{2}\right)$

B.  $\left(\frac{1}{3}, \frac{13}{3}\right)$

C.  $\left(2, \frac{3}{4}\right)$

D.  $\left(2, \frac{3}{4}\right)$

**Answer: A**



**Watch Video Solution**

11. The image of the line  $\frac{x-2}{2} = \frac{y-3}{-3} = \frac{z-4}{1}$  in the plane  $x + y + z = 6$  can be

A.  $\vec{r} = \lambda(2\hat{i} - 3\hat{j} + \hat{k})$

B.  $\vec{r} = (2\hat{i} + \hat{j} + \hat{k}) + \alpha(2\hat{i} - 3\hat{j} + \hat{k})$

$$\text{C. } \vec{r} = (\hat{j} + 2\hat{k}) + \beta(2\hat{i} - 3\hat{j} + \hat{k})$$

$$\text{D. } \vec{r} = (\hat{i} + 2\hat{j}) + u(2\hat{i} - 3\hat{j} + \hat{k})$$

**Answer: C**



**Watch Video Solution**

**12.** Let  $\vec{a} = \hat{i} + \hat{j} - \hat{k}$  and  $\vec{b} = 2\hat{i} - \hat{j} + \hat{k}$ .

If  $\vec{c}$  is a non - zero vector perpendicular to

both  $\vec{a}$  and  $\vec{b}$ , such that its component

along x, y, z axes are rational numbers, then

$|\vec{c}|$  is

A. an integer

B. an irrational number

C. a rational number

D. None of these

**Answer: B**



**Watch Video Solution**

**13.**

Let

$$f: \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^4 - 8x^3 + 22x^2 - 24x + c$$

.

If sum of all extremum value of  $f(x)$  is 1, then  $c$  is equal to

A. 8

B. 9

C. 10

D. 11

**Answer: B**



**Watch Video Solution**



14. If  $f(x)$  is a continuous function such that its value  $\forall x \in R$  is a rational number and  $f(1) + f(2) = 6$ , then the value of  $f(3)$  is equal to

A. 3

B. 9

C. 2

D. 4

**Answer: A**



Watch Video Solution

15. The domain of the function

$$f(x) = \sqrt{(x^2 - 8x + 12) \cdot \ln^2(x - 3)}$$
 is

- A.  $[3, \infty)$
- B.  $[4, \infty)$
- C.  $[6, \infty) \cup \{2, 4\}$
- D.  $[6, \infty) \cup \{4\}$

**Answer: D**



**Watch Video Solution**

16. The area bounded by  $y = \sqrt{1 - x^2}$  and  $y = x^3 - x$  is divided by y - axis in the ratio  $\lambda : 1$  ( $\forall \lambda > 1$ ), then  $\lambda$  is equal to

A.  $\frac{\pi + 1}{\pi - 1}$

B.  $\frac{\pi}{\pi - 1}$

C.  $\frac{\pi + 1}{\pi}$

D.  $\frac{\pi + 2}{\pi - 2}$

**Answer: A**



**Watch Video Solution**

17. The straight line  $y = 2x$  meets  $y = f(x)$  at P, where  $f(x)$  is a solution of the differential equation  $\frac{dy}{dx} = \frac{x^2 + xy}{x^2 + y^2}$  such that  $f(1) = 3$ , then  $f'(x)$  at point P is

A.  $\frac{3}{5}$

B.  $\frac{5}{3}$

C. 2

D. 1

**Answer: A**



Watch Video Solution

18. If  $\int \frac{2x^2 + 5}{x^2 + a} dx = f(x)$ , where  $f(x)$  is a polynomial or ratio of polynomials, then the number of possible value(s) of  $a$  is equal to

A. 0

B. 1

C. 2

D. 3

**Answer: C**



**Watch Video Solution**

**19.** Negation of the statement, "I will work hard and party" is

- A. I will not work hard and party
- B. I will not work hard or I will party
- C. I will not work hard or I will not party
- D. I will not work had and I will not party

**Answer: C**



**Watch Video Solution**

**20.** The solution set of the inequality

$$(\tan^{-1} x)^2 \leq (\tan^{-1} x) + 6 \text{ is}$$

A.  $[-\tan 2, \tan 3]$

B.  $[\tan 2, -\tan 3]$

C.  $[\tan 2, \tan 3]$

D.  $(-\infty, \infty)$

**Answer: A**



**Watch Video Solution**

21. Let the matrix  $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$  and

$B = A + A^2 + A^3 + A^4$ . If

$B = \lambda A$ ,  $\forall \lambda \in R$ , then the value of  $\lambda$  is equal to



**Watch Video Solution**



22. The number of solution of the equation  $2a + 3b + 6c = 60$ , where  $a, b, c \in N$ , is equal to

 [Watch Video Solution](#)

23. The value of the integral

$$\int_0^8 \frac{x^2}{x^2 + 8x + 32} dx \text{ is equal to}$$

 [Watch Video Solution](#)

24. A person standing at point P sees angle of elevation of the top of a building, whose base is 50 meters away, to be  $60^\circ$ . Another building whose base is 20 meters away from the base of the first building and is between the observer and first building has height  $h$  meters, then the maximum possible height (in meters) of this second building is (Take  $\sqrt{3} = 1.73$ )



[Watch Video Solution](#)

**25.** The number of solutions of the equation

$$\cot x \cos x - 1 = \cot x - \cos x, \quad \forall x \in [0, 2\pi]$$

is equal to



**Watch Video Solution**