



## MATHS

### BOOKS - NTA MOCK TESTS

### NTA JEE MOCK TEST 25

#### Mathematics

1. If  $y = (1 + x)^y + \sin^{-1}(\sin^2 x)$ , then  $\frac{dy}{dx}$  at  $x = 0$  is

A. 0

B.  $\ln 2$

C. 1

D.  $\frac{1}{2}$

**Answer: C**



**Watch Video Solution**

2. The area bounded by  $f(x) = \sin^2 x$  and the x - axis from  $x = a$  to  $x = b$  , where  $f''(a) = f''(b) = 0 (\forall a, b, \in (0, \pi))$  is

A.  $\frac{\pi}{4}$

B.  $\pi + 2$

C. 2

D.  $\frac{\pi + 2}{4}$

**Answer: D**



**Watch Video Solution**

**3.** The domain of the function

$$f(x) = \frac{1}{9 - x^2} + \log_{20}(x^3 - 3x) \text{ is}$$

- A.  $(-\sqrt{3}, 0) \cup (\sqrt{3}, \infty)$
- B.  $(-\sqrt{3}, 0) \cup (\sqrt{3}, 3)$
- C.  $(-\sqrt{3}, 0) \cup (3, \infty)$
- D.  $(-\sqrt{3}, 0) \cup (\sqrt{3}, 3) \cup (3, \infty)$

**Answer: D**



**Watch Video Solution**

4. If for a sample size of 10,

$$\sum_{i=1}^{10} (x_i - 5)^2 = 350 \text{ and } \sum_{i=1}^{10} (x_i - 6) = 20, \text{ then}$$

the variance is

A. 23

B. 24

C. 25

D. 26

**Answer: D**



**Watch Video Solution**

5. If  $a$  and  $b$  are two real number lying between 0 and 1 such that  $z_1 = a + i$ ,  $z_2 = 1 + bi$  and  $z_3 = 0$  form an equilateral triangle , then

A.  $a = 2 + \sqrt{3}$

B.  $a = 4 - \sqrt{3}$

C.  $a = b$

D.  $a = 2, b = \sqrt{3}$

**Answer: C**



**Watch Video Solution**

6. If any tangent to the ellipse  $25x^2 + 9y^2 = 225$  meets the coordinate axes at A and B such that  $OA = OB$  then, the length AB is equal to (where, O is the origin)

A.  $\sqrt{17}$  units

B.  $\sqrt{34}$  units

C.  $2\sqrt{17}$  units

D.  $2\sqrt{34}$  units

**Answer: C**



**Watch Video Solution**

7. If  $\frac{1}{1!11!} + \frac{1}{3!9!} + \frac{1}{5!7!} = \frac{2^n}{m!}$  then the value of  $m + n$  is

A. 18

B. 23

C. 12

D. 22

**Answer: D**



**Watch Video Solution**

8. There are 10 seats in the first row of a theatre of which 4 are to be occupied. The number of ways of

arranging 4 persons so that no two persons sit side by side is:

A. 240

B. 480

C. 840

D. 420

**Answer: C**



[Watch Video Solution](#)

9. Let the curve  $y = f(x)$  satisfies the equation

$$\frac{dy}{dx} = 1 - \frac{1}{x^2} \text{ and passes through the point } \left(2, \frac{7}{2}\right)$$



then the value of  $f(1)$  is

A. 3

B. 2

C.  $\frac{7}{2}$

D. 1

**Answer: A**



**Watch Video Solution**

10. If a function  $F: \mathbb{R} \rightarrow \mathbb{R}$  is defined as

$$f(x) = \int \frac{x^8 + 4}{x^4 - 2x^2 + 2} dx \text{ f (0) =1 , then which of the}$$

following is correct ?

A.  $f(x)$  is an even function

B.  $f(x)$  is an onto function

C.  $f(x)$  is an odd function

D.  $f(x)$  is an many one function

**Answer: B**



**Watch Video Solution**

**11.** Six fair dice are rolled . The probability that the product of the numbers appearing on top faces is prime is

A.  $\frac{1}{2} \left( \frac{1}{6} \right)^4$

B.  $\left(\frac{1}{2}\right)^6$

C.  $\frac{1}{6^4}$

D.  $\frac{1}{2} \left(\frac{1}{6}\right)^5$

**Answer: A**



**Watch Video Solution**

12. If  $x$  satisfies the inequality

$$(\tan^{-1} x)^2 + 3(\tan^{-1} x) - 4 > 0, \text{ then the complete}$$

set of values of  $x$  is

A.  $\left(-\tan 4, \frac{\pi}{4}\right)$

B.  $(\infty, \tan 4) \cup \left(\frac{\pi}{4}, \infty\right)$

C.  $(\tan 1, \infty)$

D.  $(\tan 4, \tan 1)$

**Answer: C**



**Watch Video Solution**

**13.** Let  $2a + 2b + c = 0$  , then the equation of the straight line  $ax + by + c = 0$  which is farthest the point  $(1,1)$  is

A.  $y = x$

B.  $y + x = 2$

C.  $y + x = 4$

D.  $y = x + 2$

**Answer: C**



**Watch Video Solution**

**14.** If  $x, |x + a|, |x - 1|$  are first three terms of an A.P.,  
then the sum of its first 20 terms is

A. 90 or 175

B. 180 or 350

C. 360 or 700

D. 720 or 1400

**Answer: B**



**Watch Video Solution**

**15.** The difference between the greatest and the least possible value of the expression  $3 - \cos x + \sin^2 x$  is

A.  $\frac{13}{4}$

B.  $\frac{17}{4}$

C.  $\frac{9}{4}$

D.  $\frac{1}{4}$

**Answer: C**



**Watch Video Solution**

16. The value of  $\int_0^{12\pi} ([\sin t] + [-\sin t]) dt$  is equal to  
(where  $[.]$  denotes the greatest integer function )

A.  $12\pi$

B.  $-12\pi$

C.  $-10\pi$

D.  $-6\pi$

**Answer: B**



**Watch Video Solution**

17. For the function

$f(x) = \sin x + 2 \cos x, \forall x \in [0, 2\pi]$  we obtain

A. a local point of maxima at  $x = \alpha$ , where  $\alpha$  is in

1<sup>st</sup> quadrant

B. a local point of maxima at  $x = \alpha$ , where  $\alpha$  is in

3<sup>rd</sup> quadrant

C. a local point of minima at  $x = \alpha$ , where  $\alpha$  is in 1<sup>st</sup>

quadrant

D. a local point of minima at  $x = \alpha$ , where  $\alpha$  is in

2<sup>nd</sup> quadrant

**Answer: A**





Watch Video Solution

18. Find the equation of the plane containing the lines  $2x-y+z-3=0$ ,  $3x+y+z=5$  and at a distance of  $\frac{1}{\sqrt{6}}$  from the point  $(2,1,-1)$ .

A.  $x + y + z - 3 = 0$

B.  $2x - y - z - 3 = 0$

C.  $2x - y + z + 3 = 0$

D.  $62x + 29y + 19z - 105 = 0$

**Answer: D**



Watch Video Solution

19. If  $\vec{m}_a$ ,  $\vec{m}_b$  and  $\vec{m}_c$  are 3 unit vectors such that  $\vec{m}_a \cdot \vec{m}_b = \vec{m}_a \cdot \vec{m}_c = 0$  and the angle between  $\vec{m}_b$  and  $\vec{m}_c$  is  $\frac{\pi}{3}$ , then the value of  $|\vec{m}_a \times \vec{m}_b - \vec{m}_a \times \vec{m}_c|$  is equal to

A. 1

B. 2

C. 3

D. 4

**Answer: A**



**Watch Video Solution**

20. Let the points A lies on  $3x - 4y + 1 = 0$ , the point B lines on  $4x + 3y - 7 = 0$  and the point C is  $(-2,5)$  . If ABCD is a rhombus , then the locus of D is

A.  $25\left((x + 2)^2 + (y - 5)^2\right) = (3x + 4y + 1)^2$

B.  $(3x - 4y + 1)^2 + (4x - 3y - 7)^2 = 1$

C.  $(3x - 4y + 1)^2 - (4x - 3y - 7)^2 = 1$

D.  $(4x + 3y - 7)^2 + (3x - 4y + 1)^2 = 1$

**Answer: A**



**Watch Video Solution**

21.

Let

$$A + 2B = \begin{bmatrix} 2 & 4 & 0 \\ 6 & -3 & 3 \\ -5 & 3 & 5 \end{bmatrix} \text{ and } 2A - B = \begin{bmatrix} 6 & -2 & 4 \\ 6 & 1 & 5 \\ 6 & 3 & 4 \end{bmatrix}$$

, then  $\text{tr}(A) - \text{tr}(B)$  is equal to (where,  $\text{tr}(A)$  = trace of matrix A i.e. . Sum of the principle diagonal elements of matrix A)



Watch Video Solution

22.

Let

$$px^4 + qx^3 + rx^2 + sx + t = \begin{vmatrix} x^2 + 3x & x - 1 & x + 3 \\ x + 1 & -2 & x - 4 \\ x - 3 & x + 4 & 3x \end{vmatrix}$$

be an identity where p,q,r,s and t are constants, then the value of s is equal to



Watch Video Solution

23. If  $S = \sum_{r=1}^{80} \frac{r}{(r^4 + r^2 + 1)}$ , then the value of  $\frac{6481s}{1000}$

is




Watch Video Solution

24. If the equation of the tangent at the point P(3,4) on the parabola whose axis is the x - axis is  $3x - 4y + 7 = 0$ , then distance of the tangent from the focus of the parabola is



Watch Video Solution

25. In the figure  $PQ$ ,  $PO_1$  and  $O_1Q$  are the diameters of semicircles  $C_1$ ,  $C_2$  and  $C_3$  with centres at  $O_1$ ,  $O_2$  and  $O_3$  respectively.  $C_1C_2$  and  $C_3$ . If  $PQ = 24$  units and the area of the circle  $C_4$  is  $A$  sq. units, then the value of  $\frac{8\pi}{4}$  is equal to (here,  $PO_1 = O_1Q$ )



**Watch Video Solution**