



MATHS

BOOKS - NTA MOCK TESTS

NTA JEE MOCK TEST 51

Mathematics

1. find the term independent of 'x' in the expansion of $(1 + x + x^2) \left(\frac{3}{2}x^2 - \frac{1}{3x} \right)^9$

A. $\frac{1}{3}$

B. $\frac{19}{54}$

C. $\frac{17}{54}$

D. $\frac{1}{4}$

Answer: C



Watch Video Solution

2. For a complex number Z , if the argument of $(Z - a)(\bar{Z} - b)$ is $\frac{\pi}{4}$ or $\frac{-3\pi}{4}$ (where a, b are two real numbers), then the value of ab such

that the locus of Z represents a circle with

centre $\frac{3}{2} + \frac{i}{2}$ is (where, $i^2 = -1$)

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

3. The value of $\int \frac{(1 - \cos \theta)^{\frac{3}{10}}}{(1 + \cos \theta)^{\frac{13}{13}}} d\theta$ is equal to

(where, c is the constant of integration)

A. $\frac{5}{8}(\tan \theta)^{\frac{5}{8}} + c$

B. $\frac{5}{8} \left(\tan. \frac{\theta}{2} \right)^{\frac{8}{5}} + c$

C. $\frac{5}{16} \left(\tan. \frac{\theta}{2} \right)^{\frac{8}{5}} + c$

D. $\frac{5}{8}(\tan \theta)^{\frac{5}{16}} + c$

Answer: B



Watch Video Solution

4. In an increasing geometric progression, the sum of the first and the last term is 99, the product of the second and the last but one term is 288 and the sum of all the terms is 189. Then, the number of terms in the progression is equal to

A. 5

B. 6

C. 7

D. 8

Answer: B



Watch Video Solution

5. The sum of the first 10 terms of the series

$$\frac{5}{1.2.3} + \frac{7}{2.3.9} + \frac{9}{3.4.27} + \dots \text{ is}$$

A. $2 + \frac{1}{3^{10}}$

B. $1 + \frac{1}{11(3)^{10}}$

C. $2 - \frac{1}{11(3)^{10}}$

D. $1 - \frac{1}{11(3)^{10}}$

Answer: D



Watch Video Solution

6.

if

$$|\sin^2 x + 10x^2| = |9 - x^2| + 2\sin^2 x + \cos^2 x$$

, then x lies in

A. $[-8, 8]$

B. $[-3, 3]$

C. $[-\sqrt{17}, \sqrt{17}]$

D. $[-\sqrt{21}, \sqrt{21}]$

Answer: B



Watch Video Solution

7. An isosceles triangle made of wood of base 10 feet and height 8 feet is placed vertically with its base on the ground and vertex directly above it. If the triangle faces the sun whose altitude is 30° , then the tangent of the angle at the apex of the shadow is

A. $\frac{80}{\sqrt{3}}$

B. $\frac{80\sqrt{3}}{167}$

C. $\frac{89}{2\sqrt{3}}$

D. $\frac{80\sqrt{3}}{217}$

Answer: B



Watch Video Solution

8. The mean deviation of the series $a^2, a^2 + d, a^2 + 2d, \dots, a^2 + 2nd$ from its median is

A. $\frac{(n + 1)d}{(2n + 1)}$

B. $\frac{nd}{2n + 1}$

C. $\frac{n(n + 1)d}{(2n + 1)}$

D. $\frac{(2n + 1)d}{n(n + 1)}$

Answer: C



Watch Video Solution

9. The number of roots of the equation

$$\sin^{-1} x - \cos^{-1} x = \sin^{-1}(5x - 3) \text{ is/ are}$$

A. 3

B. 1

C. 2

D. 0

Answer: B



Watch Video Solution

10. The value of

$$\lim_{x \rightarrow 0} \frac{\ln(1 + 2x + 4x^2) + \ln(1 - 2x + 4x^2)}{\sec x - \cos x}$$

is equal to

A. -1

B. 1

C. 0

D. 4

Answer: D



Watch Video Solution

11. The radius of a right circular cylinder increases at the rate of 0.2 cm/sec and the height decreases at the rate of 0.1 cm/sec. The rate of change of the volume of the cylinder when the radius is 1 cm and the height is 2 cm is

A. $\frac{\pi}{10} \text{ cm}^3 / \text{sec}$

B. $\frac{9\pi}{10} \text{ cm}^3 / \text{sec}$

C. $\frac{7\pi}{10} \text{ cm}^3 / \text{sec}$

D. $\frac{\pi}{5} \text{ cm}^3 / \text{sec}$

Answer: C



Watch Video Solution

12. If the graph of the function
$$y = (a - b)^2 x^2 + 2(a + b - 2c)x + 1 (\forall a \neq b)$$

A. $a < b < c$

B. $a < c < b$

C. $b < a < c$

D. $c < b < a$

Answer: B



Watch Video Solution

13. The value of $\int_{-\pi}^{\pi} \frac{\sqrt{2} \cos x}{1 + e^x} dx$ is equal to

A. $\frac{\pi}{2}$

B. π

C. 0

D. 2π

Answer: C



Watch Video Solution

14. Let the tangents PQ and PR are drawn to $y^2 = 4ax$ from any point P on the line $x + 4a = 0$. The angle subtended by the chord of contact QR at the vertex of the parabola $y^2 = 4ax$ is

A. $\frac{\pi}{4}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{2}$

D. $\frac{\pi}{6}$

Answer: C



Watch Video Solution

15. Let $\vec{p} = \hat{i} + \hat{j} + \hat{k}$ and \vec{r} be a variable vector such that $\vec{r} \cdot \hat{i}$, $\vec{r} \cdot \hat{j}$ and $\vec{r} \cdot \hat{k}$ are even natural numbers. If $\vec{r} \cdot \vec{p} \leq 20$, then the number of values of \vec{r} is

A. 20

B. 60

C. 75

D. 120

Answer: D



Watch Video Solution

16. Let A , B and C are $n \times n$ matrices such that

$|A| = 2$, $|B| = 3$ and $|C| = 5$. If

$|(2A)^2(3B)(5C)^{-1}| = \frac{1728}{125}$, then the value

of n is equal to

A. 2

B. 3

C. 4

D. 6

Answer: A



Watch Video Solution

17. Given two independent events, if the probability that both the events occur is $\frac{8}{49}$, the probability that exactly one of them

occurs is $\frac{26}{49}$ and the probability of more probable of the two events is λ , then 14λ is equal to

A. 2

B. 4

C. 8

D. 7

Answer: C



Watch Video Solution

18. The number of integral value(s) of k such that the system of equations $kz - 2y - z = x$, $ky - z = z + 3x$ and $2x + kz = 2y - z$ has non-trivial solution, is/are

A. 0

B. 1

C. 2

D. 3

Answer: D



19. The vertices of a triangle are the points $P(-26, 17)$, $Q(30, 17)$ and $R(10, 2)$. If G and I be the centroid and incentre of the triangle PQR , then the value of $(GI)^2$ is equal to

A. $\frac{205}{9}$

B. $\frac{\sqrt{205}}{3}$

C. $3\sqrt{3}$

D. 27

Answer: A



Watch Video Solution

20. General solution of the differential

equation $(\cos x) \frac{dy}{dx} + y \cdot \sin x = 1$ is

A. $y = c \sin x + \cos x$

B. $y = \sin x + c \cos x$

C. $y = \tan x + c$

D. $y \sin x = \sin x + c$

Answer: B



Watch Video Solution

21. The sum of the y - intercepts of the tangents drawn from the point $(- 2, - 1)$ to the hyperbola $\frac{x^2}{3} - \frac{y^2}{2} = 1$ is



Watch Video Solution

22. In a class tournament where the participants were to play one game with one another, two of the class players fell ill, having played 3 games each. If the total number of games played is 24, then the number of participants at the beginning was



[Watch Video Solution](#)

23. The value of $\frac{\int_0^2 x^4 \sqrt{4-x^2} dx}{\int_0^2 x^2 \sqrt{4-x^2} dx}$ is equal to



[Watch Video Solution](#)

24. Let four circles having radii $r_1 = 5$ units, $r_2 = 5$ units, $r_3 = 8$ units and r_4 units ($r_4 < 5$) are mutually touching each other externally, then the value of $\frac{2}{r_4}$ is equal to



[Watch Video Solution](#)

25. If the distance of point $P(3, 2, 6)$ from the line $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ measured

parallel to the plane $3x - 5y + 2z = 5$ is k ,
then then the value of k^2 is equal to



Watch Video Solution