



## MATHS

### BOOKS - NTA MOCK TESTS

#### NTA JEE MOCK TEST 81

#### Mathematics

1. If the function  $f(x)$  is symmetric about the line  $x = 3$ , then the value of the integral

$$I = \int_{-2}^8 \frac{f(x)}{f(x) + f(6-x)} dx \text{ is}$$

A. 0

B. 5

C. 10

D. 16

**Answer: B**



**Watch Video Solution**

2. The normal to the parabola  $y^2 = 8ax$  at the point  $(2, 4)$  meets the parabola again at the point

A.  $(-18, -12)$

B.  $(-18, 12)$

C.  $(18, 12)$

D.  $(18, -12)$

**Answer: D**

 [Watch Video Solution](#)

3. The number of values of  $x \in [-2\pi, 2\pi]$  which satisfy the equation  $\operatorname{cosec} x = 1 + \cot x$  is equal to

A. 0

B. 2

C. 4

D. 6

**Answer: B**

4. If the integral

$$I = \int \frac{x\sqrt{x} - 3x + 3\sqrt{x} - 1}{x - 2\sqrt{x} + 1} dx = f(x) + C \quad (\text{where,}$$

$x > 0$  and  $C$  is the constant of integration) and

$f(1) = \frac{-1}{3}$ , then the value of  $f(9)$  is equal to

- A. 3
- B. 6
- C. 9
- D. 12

**Answer: C**

5. The number of ways of arranging the letters AAAAA, BBB, CCC, D, EE and F in a row, if the letters B are separated from one another, is equal to

A.  $\frac{13!}{5!3!3!2!}$

B.  $\frac{14!}{3!3!2!}$

C.  $\frac{15!}{(3!)^2 2!5!}$

D.  ${}^{13}C_3 \times \frac{12!}{5!3!2!}$

**Answer: D**



**Watch Video Solution**

6. If  $a, b, c \in R^+$  such that  $a + b + c = 27$ , then the maximum value of  $a^2 b^3 c^4$  is equal to

A.  $2^8 \cdot 3^{10}$

B.  $2^9 \cdot 3^{12}$

C.  $2^{10} \cdot 3^{12}$

D.  $2^{11} \cdot 3^{13}$

**Answer: C**



[Watch Video Solution](#)

7. Find the degrees and radians the angle between the hour hand and the minute hand of a clock at half past

three.

A.  $90^\circ$

B.  $80^\circ$

C.  $75^\circ$

D.  $60^\circ$

**Answer: C**



**Watch Video Solution**

**8.** If  $f(x) = 2 \sin x - x^2$ , then in  $x \in [0, \pi]$

A.  $f(x)$  has no local maximum

B.  $f(x)$  has one local minimum

C.  $f(x)$  has 2 local maxima

D.  $f(x)$  has one local maximum

**Answer: D**



**Watch Video Solution**

9. 15 coins are tossed. If the probability of getting at least 8 heads is equal to  $p$ , then  $\frac{8}{p}$  is equal to

A. 2

B. 4

C. 8

D. 16



**Answer: D**



**Watch Video Solution**

**10.** A normal line with positive direction cosines to the plane P makes equal angles with the coordinate axis. The distance of the point A(1, 2, 3) from the line  $\frac{x-1}{1} = \frac{y+2}{1} = \frac{z-3}{2}$  measured parallel to the plane P is equal to

- A. 3 units
- B.  $\sqrt{13}$  units
- C.  $\sqrt{14}$  units
- D.  $2\sqrt{5}$  units

**Answer: C**



**Watch Video Solution**

11. Let  $A = [a_{ij}]_{3 \times 3}$  be a scalar matrix whose elements are the roots of the equation  $x^9 - 15x^8 + 75x^7 - 125x^6 = 0$ .

If  $|A \cdot adj A| = k$ , then the value of  $k$  is equal to

A.  $5^{12}$

B.  $5^9$

C.  $3^{12}$

D.  $3^9$

**Answer: B**



**Watch Video Solution**

12. For three non - zero vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$ , if

$$\left[ \begin{array}{ccc} \vec{a} & \vec{b} & \vec{c} \end{array} \right] = 4, \quad \text{then}$$

$$\left[ \vec{a} \times \left( \vec{b} + 2\vec{c} \right) \quad \vec{b} \times \left( \vec{c} - 3\vec{a} \right) \quad \vec{c} \times \left( 3\vec{a} + \vec{b} \right) \right]$$

is equal to

A. 12

B. 16

C. 84

D. 144

**Answer: D**



**Watch Video Solution**

13. Let  $f: [-1, 1] \Rightarrow B$  be a function defined as

$$f(x) = \cot^{-1} \left( \cot \left( \frac{2x}{\sqrt{3}(1+x^2)} \right) \right). \text{ If } f \text{ is both one -}$$

one and onto, then B is the interval

A.  $\left(0, \frac{\pi}{3}\right)$

B.  $\left[0, \frac{2\pi}{3}\right)$

C.  $\left[\frac{\pi}{3}, \frac{2\pi}{3}\right]$

D.  $\left(\frac{\pi}{3}, \pi\right)$

**Answer: C**



**Watch Video Solution**

14. If  $p$ ,  $q$  and  $r$  are three logical statements, then the truth value of the statement  $(p \wedge q) \vee (\sim q \rightarrow r)$ , where  $p$  is true, is

- A. True if  $q$  is false
- B. False if  $q$  is false
- C. True if  $q$  is true
- D. False if  $q$  is true

**Answer: C**



**Watch Video Solution**

15. If  $f(x) = \begin{cases} \frac{e^{[2x]+2x+1}-1}{[2x]+2x+1} & : x \neq 0 \\ 1 & : x = 0 \end{cases}$ , then (where  $[.]$

represents the greatest integer function)

A.  $\lim_{x \rightarrow 0^+} f(x) = 1$

B.  $\lim_{x \rightarrow 0^-} f(x) = e - 1$

C.  $f(x)$  is continuous at  $x = 0$

D.  $f(x)$  is discontinuous at  $x = 0$

**Answer: D**



**Watch Video Solution**

16. The set  $(A \cup B \cup C) \cap (A \cap B' \cap C')$  is equal to

A.  $A \cap B$

B.  $A \cap C'$

C.  $B \cup C$

D.  $B \cap C$

**Answer: C**



**Watch Video Solution**

17. Let the circumcentre of  $\Delta ABC$  is  $S(-1, 0)$  and the midpoints of the sides  $AB$  and  $AC$  are  $E(1, -2)$  and  $F(-2, -1)$  respectively, then the coordinates of  $A$  are

A.  $(0, -3)$

B.  $(0, 3)$

C.  $(-3, 0)$

D.  $(3, 0)$

**Answer: B**



**Watch Video Solution**

18. For a  $\triangle ABC$  the vertices are  $A(0, 3)$ ,  $B(0, 12)$  and  $C(x, 0)$ . If the circumcircle of  $\triangle ABC$  touches the  $x$  - axis, then the area (in sq. units) of the  $\triangle ABC$  is

A. 36

B. 27

C. 30



**Answer: B**

 [Watch Video Solution](#)

**19.** The solution of the differential equation

$$\left(\frac{dy}{dx}\right)^4 - \left(\frac{dy}{dx}\right)^2 - 2 = 0 \text{ is } y = \pm \sqrt{\lambda x} + C \text{ (where,}$$

$C$  is an arbitrary constant). Then,  $\lambda^2$  is equal to

A. 2

B. 4

C. 8

D. 16

**Answer: B**

 [Watch Video Solution](#)

20. For the complex number  $z$  satisfying the condition

$$\left| z + \frac{2}{z} \right| = 2, \text{ the maximum value of } |z| \text{ is}$$

A.  $\sqrt{3} - 1$

B.  $\sqrt{3} + 1$

C.  $\sqrt{2} + \sqrt{3}$

D.  $\sqrt{3}$

**Answer: B**

 [Watch Video Solution](#)

21. If the area bounded by  $y \leq e - |x - e|$  and  $y \geq 0$  is  $A$  sq. units, then  $\log_e(A)$  is equal is

 [Watch Video Solution](#)

22. If the middle term in the expansion of  $\left(\frac{1}{x} + x \sin x\right)^{10}$  is equal to  $7\frac{7}{8}$ , then the number of values of  $x$  in  $[0, 2\pi]$  is equal to

 [Watch Video Solution](#)

23. Let  $A = \begin{bmatrix} 2 & -1 & 1 \\ -2 & 3 & -1 \\ -4 & 4 & -x \end{bmatrix}$  be a matrix. If  $A^2 = A$ ,

then the value of  $x$  is equal to

 [Watch Video Solution](#)

24. The value of  $\lim_{x \rightarrow 0} (\cos x + \sin x)^{\frac{1}{x}}$  is equal to to (take  $e = 2.71$ )

 [Watch Video Solution](#)

25. A tangent of slope 2 of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{1} = 1$  passes through  $(-2, 0)$ . Then, three times the square of the eccentricity of the ellipse is equal to

 [Watch Video Solution](#)