



## PHYSICS


### BOOKS - NCERT PHYSICS (ENGLISH)

#### ELECTRIC CHARGES AND FIELD

##### Mcq 5

1. In figure two positive charges  $q_2$  and  $q_3$  fixed along the  $y$ -axis, exert a net electric force in the  $+x$  direction on a charge  $q_1$  fixed along the  $x$ -axis if a positive charge  $Q$  is

added at  $(x, 0)$  the force on  $q_1$



A. shall increase along the positive x-axis

B. shall decrease along the positive x-axis.

C. shall point along the negative x-axis

D. shall increase but the direction changes because of





the intersection of Q with  $q_2$  and  $q_3$ .

**Answer: A**




**Watch Video Solution**

2. A point positive charge is brought near an isolated conducting sphere as shown in figure the electric field is best given by





A. (a)



(b)

B.



(c)

C.




(d)

D.

Answer: a

 Watch Video Solution

3. The electric flux through the surface



A. in fig. (iv) is the largest

B. in fig. (iii) is the least

C. in fig. (ii) is same as fig.(iii) but is smaller than fig.(iv)

D. Is the same for all the figures.

**Answer: D**


 [Watch Video Solution](#)

4. five charge  $q_1, q_2, q_3, q_4$  and  $q_5$  are fixed at their positions as shown in figure .s is Gaussian surface .The Gauss's law is

given by 
$$\oint \vec{E} \cdot \vec{ds} = \frac{q}{\epsilon_0}$$



# Gaussian surface



Which of the following statement is correct?

A. E on the LHS of the above equation will have a contribution from  $q_1$ ,  $q_5$  and  $q_3$  while q on the RHS will have a contribution from  $q_2$  and  $q_4$  only.

B. E on the LHS of the above equation will have a contribution from all charges while  $q$  on the RHS will have a contribution from  $q_2$  and  $q_4$  only.

C. E on the LHS of the above equation will have a contribution from all charges while  $q$  on the RHS will have a contribution from  $q_1$ ,  $q_3$  and  $q_5$  only


D. Both E on the LHS and  $q$  on the RHS will have contributions from  $q_2$  and  $q_4$  only.

**Answer: B**



**Watch Video Solution**

5. Figure shows electric field lines in which an electric dipole  $p$  is placed as shown. Which of the following statements is correct ?



- A. The dipole will not experience any force
- B. The dipole will experience a force towards right
- C. The dipole will experience a force towards left
- D. The dipole will experience a force upwards.

**Answer: C**



**Watch Video Solution**

**6.** A point charge  $+q$  is placed at a distance  $d$  from an isolated conducting plane. The field at a point  $P$  on the other side of plane is

- A. directed perpendicular to the plane and away from the plane.
- B. directed perpendicular to the plane but towards the plane.
- C. directed radially away from the point charge
- D. directed radially towards the point charge.

**Answer: A**



**Watch Video Solution**

7. A hemispherical shell is uniformly charge positively. The electric field at point on a diameter away from the center is directed

- A. perpendicular to the diameter
- B. parallel to the diameter
- C. at an angle tilted towards the diameter
- D. at an angle tilted away from the diameter.

**Answer: A**



**Watch Video Solution**

## Mcq 5 More Than One Options

1. If  $\oint_s E \cdot ds = 0$  Over a surface, then

- A. The electric field inside the surface and on it is zero.
- B. the electric field inside the surface is necessarily uniform
- C. the number of flux lines entering the surface must be equal to the number of flux lines leaving it.
- D. all charges must necessarily be outside the surface.

**Answer: C::D**



**Watch Video Solution**

2. The Electric field at a point is

A. always continuous

B. continuous if there is no charge at that point.

C. discontinuous only if there is a negative charge at that point

D. None of these

**Answer: B**



**Watch Video Solution**

3. If there were only one type of charge in the universe then

- A.  $\oint_S E \cdot dS \neq 0$  on any surface
- B.  $\oint_S E \cdot dS = 0$  if the charge is outside the surface.
- C.  $\oint_S E \cdot dS$  could not be defined.
- D.  $\oint_S E \cdot dS = \frac{q}{\epsilon_0}$  if charges of magnitude  $q$  were inside the surface.

**Answer: C::D**

 [Watch Video Solution](#)

4. Consider a region inside which, there are various types of charges but the total charge is zero. At points outside the region

A. the electric field is necessarily zero.



B. The electric field is due to the dipole moment of the charge distribution only.

C. The dominant electric field is  $\propto \frac{1}{r^3}$ , for large  $r$ ,


where  $r$  is the distance from a origin in this regions.

D. the work done to move a charged particle along a closed path, away from the region, will be zero.

**Answer: C::D**

 [Watch Video Solution](#)

5. Refer to the arrangement of charges in Fig and a Gaussian surface of radius  $R$  with  $Q$  at the centre. Then




- A. total flux through the surface of the sphere is  $\frac{-Q}{\epsilon_0}$
- B. field on the surface of the sphere is  $\frac{-Q}{4\pi\epsilon_0 R^2}$
- C. flux through the surface of sphere due to 5Q is zero
- D. field on the surface of sphere due to -2Q is same everywhere

Answer: A::C



Watch Video Solution

6. A positive charge  $Q$  is uniformly distributed along a circular ring of radius  $R$ . a small test charge  $q$  is placed at the centre of the ring.



A. if  $q > 0$  and is displaced away from the centre in the place of the ring, it will be pushed back towards the

centre.

B. if  $q < 0$  and is displaced away from the centre in the plane of the ring, it will never return to the centre and will continue moving till it hits the ring.

C. if  $q < 0$ , it will perform SHM for small displacement along the axis

D.  $q$  at the centre of the ring is in an unstable equilibrium within the plane of the ring for  $q > 0$

**Answer: A::B::C**



**Watch Video Solution**

**Very Short Type Question**

1. An arbitrary surface encloses a dipole. What is the electric flux through this surface ?

 [Watch Video Solution](#)

2. A metallic spherical shell has an inner radius  $R_1$  and outer radius  $R_2$ . A charge  $Q$  is placed at the centre of the shell. What will be the surface charge density on the (i) inner surface, and (ii) outer surface of the shell?

 [Watch Video Solution](#)

3. The dimensions of an atom are of the order of an Angstrom. Thus there must be large electric fields between



the protons and electrons. Why, then is the electrostatic field inside a conductor zero?

 [Watch Video Solution](#)


4. If the total charge enclosed by a surface is zero, does it imply that the electric field everywhere on the surface is zero? Conversely, if the electric field everywhere on a surface is zero, does it imply that net charge inside is zero.

 [Watch Video Solution](#)

5. Sketch the electric field lines for a uniformly charged hollow cylinder shown in Fig.



Side View



Top View

 [Watch Video Solution](#)

6. What will be the total flux through the faces of the cube, with side of length  $a$  if a charge  $q$  is placed at (a) A : a corner of the cube (b) B : mid-point of an edge of the cube (c) C : center of a face of the cube (d) D : mid-point of B and C.

 [Watch Video Solution](#)

## Short Answer Type Question

1. A paisa coin is made up Al.Mg alloy and weighs 0.75 g. It has a square shape and its diagonal measures 17mm. It is electrically neutral and contains equal amounts of positive and negative charges . Treating the paisa coin made up of only Al, find the magnitude of equal number of positive and negative charges. What conclusion do you draw from this magnitude ?



[Watch Video Solution](#)

2. Consider a coin of Question 20 . It is electrically neutral and contains equal amounts of positive and negative charge of magnitude 34.8 kC. Suppose that these equal



charges were concentrated in two point charges separated

by

(i)  $1\text{cm} \left( \sim \frac{1}{2} \times \text{diagonal of the one paisa coin} \right)$

(ii)  $100\text{ m}$  ( $\sim$  length of a long building)

(iii)  $10^6\text{m}$  (radius of the earth). find the force on each such point charge in each of the three cases. what do you conclude from these results?




[Watch Video Solution](#)

**3.** Figure represents a crystal unit of cesium chloride, CsCl.

The cesium atoms, represented by open circles are situated at the corners of a cube of side  $0.40\text{ nm}$ , whereas a Cl atom is situated at the centre of the cube. The Cs atoms are deficient in one electron while the Cl atom carries an excess

electron. (i). What is the net electric field on the Cl atom due to eight Cs atoms?

(ii) Suppose that the Cs atom at the corner A is missing. what is the net force now on the Cl atom due to seven remaining Cs atoms?



 [Watch Video Solution](#)

4. Two charges  $q$  and  $-3q$  are placed fixed on  $x$ -axis separated by distance  $d$ . where should a third charge  $2q$  be placed such that it will not experience any force?

 [Watch Video Solution](#)

5. Fig shows the electric field lines around three points charges A,B,C.

(a) Which charges are positive ?


(b) Which charge has the largest magnitude ? Why ?

(c ) In which region or regions of the picture could be the electric field be zero? Justify your answer.

(i) near A, (ii) near B, (iii) near C, (iv) nowhere.

 [Watch Video Solution](#)

6. Five charges ,  $q$  each are placed at the corners of a regular pentagon of side.



(a) (i) What will be the electric field at  $O$ , the centre of the pentagon?

(ii) What will be the electric field at  $O$  if the charge from one of the corners (say  $A$ ) is removed?

(iii). What will be the electric field at  $O$  if the charge  $q$  at  $A$  is

replaced by n-sided regular polygon with charge  $q$  at each of its corner?



[View Text Solution](#)

## Long Answer Type Question

1. In 1959 Lyttleton and Bondi suggested that the expansion of the universe could be explained if matter carried a net charge. Suppose that the universe is made up of hydrogen atoms with a number density  $N$ , which is maintained a constant. Let the charge on the proton be  $e_p = -(1 + y)e$  where  $e$  is the electronic charge.

(a). find the critical value of  $y$  such that expansion may

start. ItBrgt (b) show that the velocity of expansion is proportional tot he distance from the centre.

 [View Text Solution](#)

2. Consider a sphere of radius  $R$  with harge density distributed as  $p(r)=kr$  fro  $r \leq R = 0$  for  $r > R$ .

(a). Find the electric field as all points  $r$ .

(b) Suppose the total charge on the sphere is  $2e$  where  $e$  is the electron charge. Where can two protons be embedded such that the force on each of them is zero. Assume that the introduction of the proton does not alter the negative charge distribution.


 [Watch Video Solution](#)

3. Two fixed, identical conducting plates ( $\alpha$  and  $B$ ), each of surface area  $S$  are charged to  $-Q$  and  $q$ , respectively, where  $Q > q > 0$ . A third identical plate ( $\gamma$ ), free to move is located on the other side of the plate with charge  $q$  at a distance  $d$  (figure). The third plate is released and collides with the plate  $\beta$ . Assume the collision is elastic and the time of collision is sufficient to redistribute charge amongst  $\beta$  and  $\gamma$ .

(a). Find the electric field acting on the plate  $\gamma$  before collision. (b) Find the charges on  $\beta$  and  $\gamma$  after the collision.

(c) Find the velocity of the plate  $\gamma$  after the collision and at a

distance  $d$  from the plate  $\beta$ .



[View Text Solution](#)



4. There is another useful system of units, besides the SI/MKS. A system, called the CGS (centimeter-gram-second) system. In this system, coulomb's law is given by  $F = \frac{Qq}{r^2} \hat{r}$  where the distance  $r$  is measured in  $cm$  ( $= 10^{-2} \mu$ ),  $F$  in dynes ( $= 10^{-5} N$ ) and the charges in electrostatic units (esu units), where 1 esu unit of charge  $= \frac{1}{[3]} \times 10^{-9} C$ . The number [3] actually arises from the speed of light in vacuum which is now taken to be exactly given by  $c = 2.99792458 \times 10^8 m/s$ . An approximate value of  $c$ , then is  $c = 3 \times 10^8 m/s$ .

(i). Show that the coulomb's law in CGS units yields 1 esu of charge  $= 1(\text{dyne})^{1/2} \text{cm}$ . Obtain the dimensions of units of charge in terms of mass  $M$ , length  $L$  and time  $T$ . Show that it is given in terms of fractional powers of  $M$  and  $L$ .

(ii) Write 1 esu of charge  $= xC$ , where  $x$  is a dimensionless

number. Show that this gives  $\frac{1}{4\epsilon_0} = \frac{10^{-9}}{x^2} \frac{Nm^2}{C^2}$ . With

$x = \frac{1}{[3]} \times 10^{-9}$ , we have

$$\frac{1}{4\pi\epsilon_0} = [3]^2 \times 10^9 \frac{Nm^2}{C^2}, \quad \frac{1}{4\pi\epsilon_0} = (2.99792458)^2 \times 10^9 \frac{Nm^2}{C^2}$$

(exactly)



[View Text Solution](#)

5. Two charges  $-q$  each are fixed separated by distance  $2d$ .


A third charge  $q$  of mass  $m$  placed at the mid-point is

displaced slightly by  $x$  ( $x \ll d$ ) perpendicular to the line

joining the two fixed charges as shown in Fig. Show that  $q$

will perform simple harmonic oscillation of time period.

$$T = \left[ \frac{8\pi^3 \epsilon_0 m d^3}{q^2} \right]^{1/2}$$




[▶ Watch Video Solution](#)

6. Total charge  $-Q$  is uniformly spread along length of a ring of radius  $R$ . A small test  $+q$  of mass  $m$  is kept at the center of the ring .

(a) Show that the particle executes a simple harmonic oscillation.

(b) Obtain its time period.



[Watch Video Solution](#)