



India's Number 1 Education App

MATHS

BOOKS - VK JAISWAL MATHS (HINGLISH)

INDEFINITE AND DEFINITE INTEGRATION

Exercise Single Choice Problems

1. $\int a^x \left(\ln x + \ln a \cdot \ln \left(\frac{x}{e} \right)^x \right) dx =$

A. $a^x \ln \left(\frac{e}{x} \right)^{2x} + C$

B. $a^x \ln \left(\frac{x}{e} \right)^x + C$

C. $a^x + \ln \left(\frac{x}{e} \right)^x + C$

D. None of these

Answer: B



Watch Video Solution

2. The value of :

$$\lim_{n \rightarrow \infty} \left(\frac{1}{\sqrt{n}\sqrt{n+1}} + \frac{1}{\sqrt{n}\sqrt{n+2}} + \frac{1}{\sqrt{n}\sqrt{n+3}} + \dots + \frac{1}{\sqrt{n}\sqrt{2n}} \right)$$

is:

A. $\sqrt{2} - 1$

B. $2(\sqrt{2} - 1)$

C. $\sqrt{2} + 1$

D. $2(\sqrt{2} + 1)$

Answer: B



Watch Video Solution

3. If $\int \frac{\sin x}{\sin(x-a)} dx = Ax + B \log \sin(x-\alpha) + C$, then value of (A,B)

is:

A. $(\sin \alpha, \cos \alpha)$

B. $(\cos \alpha, \sin \alpha)$

C. $(-\sin \alpha, \cos \alpha)$

D. $(-\cos \alpha, \sin \alpha)$

Answer: B



Watch Video Solution

4. The value of the integral $\int_0^2 \frac{\log(x^2 + 2)}{(x + 2)^2} dx$ is

A. $\frac{\sqrt{2}}{3} \tan^{-1} \sqrt{2} + \frac{5}{12} \log 2 - \frac{1}{4} \log 3$

B. $\frac{\sqrt{2}}{3} \tan^{-1} \sqrt{2} - \frac{5}{12} \log 2 - \frac{1}{4} \log 3$

C. $\frac{\sqrt{2}}{3} \tan^{-1} \sqrt{2} + \frac{5}{12} \log 2 + \frac{1}{4} \log 3$

D. $\frac{\sqrt{2}}{3} \tan^{-1} \sqrt{2} - \frac{5}{12} \log 2 + \frac{1}{4} \log 3$

Answer: D



Watch Video Solution

5. If $I_1 = \int_0^1 \frac{1+x^8}{1+x^4} dx$ and $I_2 = \int_0^1 \frac{1+x^9}{1+x^2} dx$, then:

A. $I_1 > 1, I_2 < 1$

B. $I_1 < 1, I_2 > 1$

C. $1 < I_1 < I_2$

D. $I_2 < I_1 < 1$

Answer: D



Watch Video Solution

6. Let $f: (0, 1) \rightarrow (0, 1)$ be a differentiable function such that $f(x) \neq 0$

for all $x \in (0, 1)$ and $f\left(\frac{1}{2}\right) = \frac{\sqrt{3}}{2}$. Suppose for all x ,

$$\lim_{x \rightarrow x} \frac{\int_0^1 \sqrt{1(f(s))^2} dx \int_0^x \sqrt{1(f(s))^2} ds}{f(t) - f(x)} = f(x)$$

Then, the value of $f\left(\frac{1}{4}\right)$ belongs to

- A. $\left\{ \frac{\sqrt{7}}{4}, \frac{\sqrt{15}}{4} \right\}$
- B. $\left\{ \frac{\sqrt{7}}{3}, \frac{\sqrt{15}}{3} \right\}$
- C. $\left\{ \frac{\sqrt{7}}{2}, \frac{\sqrt{15}}{2} \right\}$
- D. $\{\sqrt{7}, \sqrt{15}\}$

Answer: A



Watch Video Solution

7. If $f(\theta) = \frac{4}{3}(1 - \cos^6 \theta - \sin^6 \theta)$, then

$$\lim_{x \rightarrow o} \frac{1}{n} \left[\sqrt{f\left(\frac{1}{n}\right)} + \sqrt{f\left(\frac{2}{n}\right)} + \sqrt{f\left(\frac{n}{n}\right)} \right] =$$

A. $\frac{1 - \cos 1}{2}$

B. $1 - \cos 2$

C. $\frac{\sin 2}{2}$

D. $\frac{1 - \cos 2}{2}$

Answer: D



Watch Video Solution

8. The value of $\int_0^1 \frac{(x^6 - x^3)}{(2x^3 + 1)^3} dx$ is equal to :

A. $-\frac{1}{6}$

B. $-\frac{1}{12}$

C. $-\frac{1}{18}$

D. $-\frac{1}{36}$

Answer: D



Watch Video Solution

9. $2 \int_0^{\frac{1}{\sqrt{2}}} \frac{\sin^{-1} x}{x} dx - \int_0^1 \frac{\tan^{-1} x}{x} dx =$

A. $\frac{\pi}{8} \ln 2$

B. $\frac{\pi}{4} \ln 2$

C. $\frac{\pi}{2\sqrt{2}} \ln 2$

D. $\frac{\pi}{2} \ln 2$

Answer: B



Watch Video Solution

10. Let $f(x)$ be a differentiable function such that

$$f(x) = x^2 + \int_0^x e^{-t} f(x-t) dt \text{ then } \int_0^1 f(x) dx =$$

A. $\frac{1}{3}$

B. $\frac{1}{4}$

C. $\frac{7}{12}$

D. $\frac{5}{12}$

Answer: D



Watch Video Solution

11. If $f'(x) = f(x) + \int_0^1 f(x) dx$ and given $f(0) = 1$, then $\int f(x) dx$ is equal to :

- A. $\frac{2}{3-e}e^x + \left(\frac{3-e}{1-e}\right)x + C$
- B. $\frac{2}{3-e}e^x + \left(\frac{1-e}{3-e}\right)x + C$
- C. $\frac{3}{2-e}e^x + \left(\frac{1+e}{3+e}\right)x + C$
- D. $\frac{2}{2-e}e^x + \left(\frac{1-e}{3+e}\right)x + C$

Answer: B



Watch Video Solution

12. For any $x \in R$, and f be a continuous function Let
 $I_1 = \int_{\sin^2 x}^{1 - \cos^2 x} t f(t(2-t)) dt$, $I_2 = \int_{\sin^2 x}^{1 + \cos^2 x} f(t(2-t)) dt$, then $I_1 =$

A. I_2

B. $\frac{1}{2}I_2$

C. $2I_2$

D. $3I_2$

Answer: A



Watch Video Solution

13. If the integral $\int \frac{5 \tan x dx}{\tan x - 2} = x + a \ln|\sin x - 2 \cos x| + C$, then 'a' is equal to :

A. 1

B. 2

C. - 1

D. - 2

Answer: B



Watch Video Solution

14. $\int \frac{(2 + \sqrt{x}) dx}{(x + 1 + \sqrt{x})^2}$ is equal to:

A. $\frac{x}{x + \sqrt{x} + 1} + C$

B. $\frac{2x}{x + \sqrt{x} + 1} + C$

C. $\frac{-2x}{x + \sqrt{x} + 1} + C$

D. $\frac{-x}{x + \sqrt{x} + 1} + C$

Answer: B



Watch Video Solution

15. Evaluate $\int \frac{\left(3\sqrt[3]{x + \sqrt{2 - x^2}}\right) \left(6\sqrt{1 - x}\sqrt{2 - x^2}\right) dx}{3\sqrt[3]{1 - x^2}}$, $x \in (0, 1)$:

A. $2^{\frac{1}{6}}x + C$

B. $2^{\frac{1}{12}}x + C$

C. $2^{\frac{1}{3}}x + C$

D. None of these

Answer: A



Watch Video Solution

16. $\int \frac{dx}{\sqrt{1 - \tan^2 x}} = \frac{1}{\lambda} \sin^{-1}(\lambda \sin x) + C, \text{ then } \lambda =$

A. $\sqrt{2}$

B. $\sqrt{3}$

C. 2

D. $\sqrt{5}$

Answer: A



Watch Video Solution

17. $\int \frac{dx}{\sqrt[3]{x^{5/2}(x+1)^{7/2}}} \text{ is equal to:}$

A. $-\left(\frac{x-1}{x}\right)^{1/6} + C$

B. $6\left(\frac{x+1}{x}\right)^{-1/6} + C$

C. $\left(\frac{x}{x+1}\right)^{5/6} + C$

D. $-\left(\frac{x}{x+1}\right)^{5/6} + C$

Answer: B



Watch Video Solution

18. If $I_n = \int (\sin x)^n dx$, $n \in N$, then $5I_4 - 6I_6$ is equal to

A. $\sin x \cdot (\cos x)^5 + C$

B. $\sin 2x \cos 2x + C$

C. $\frac{\sin 2x}{8} [1 + \cos^2 2x - 2 \cos 2x] + C$

D. $\frac{\sin 2x}{8} [1 + \cos^2 2x + 2 \cos 2x] + C$

Answer: C



Watch Video Solution

$$19. \int \frac{x^2 dx}{(a + bx)^2} =$$

- A. $\frac{1}{b^3} \left(a + bx - a \ln|a + bx| - \frac{a^3}{a + bx} \right) + C$
- B. $\frac{1}{b^3} \left(a + bx - 2a \ln|a + bx| - \frac{a^3}{a + bx} \right) + C$
- C. $\frac{1}{b^3} \left(a + bx + 2n \ln|a + bx| \frac{a^2}{a + bx} \right) + C$
- D. $\frac{1}{b^3} \left(a + bx + 2n \ln|a - bx| \frac{a^2}{a + bx} \right) + C$

Answer: B



Watch Video Solution

$$20. \int \frac{8x^{43} + 13x^{38}}{(x^{13} + x^5 + 1)^4} dx =$$

- A. $\frac{x^{39}}{3(x^{13} + x^5 + 1)^3} + C$
- B. $\frac{x^{39}}{(x^{13} - x^5 + 1)^3} + C$
- C. $\frac{x^{39}}{5(x^{13} - x^5 + 1)^5} + C$
- D. None of these

Answer: A



Watch Video Solution

21. $\int \left(\frac{\cos 6x + 6 \cos 4x + 15 \cos 2x + 10}{10 \cos^2 x + 5 \cos x \cos 3x + \cos x \cos 5x} \right) dx = f(x) + C,$ then
 $f(10)$ is equal to:

A. 20

B. 10

C. $2 \sin 10$

D. $2 \cos 10$

Answer: A



Watch Video Solution

22. $\int (1 + x - x^{-1}) e^{x+x^{-1}} dx =$

A. $(x + 1)e^{x+x^{-1}} + C$

B. $(x - 1)e^{x+x^{-1}} + C$

C. $-xe^{x+x^{-1}} + C$

D. $xe^{x+x^{-1}} + C$

Answer: D



Watch Video Solution

23. If $\int e^x \left(\frac{2 \tan x}{1 + \tan x} + \cos x c^2 \left(x + \frac{\pi}{4} \right) \right) dx = e^x \cdot g(x) + k$, then
 $g\left(\frac{5\pi}{4}\right) =$

A. 0

B. 1

C. -1

D. 2

Answer: B



Watch Video Solution

24. $\int e^{x \sin x + \cos x} \left(\frac{x^4 \cos^3 x - x \sin x + \cos x}{x^2 \cos^2 x} \right) dx =$

A. $e^{x \sin x + \cos x} \left(x - \frac{1}{\cos x} \right) C$

B. $e^{x \sin x + \cos x} \left(x - \frac{1}{x \cos x} \right) C$

C. $e^{x \sin x + \cos x} \left(1 - \frac{1}{x \cos x} \right) C$

D. $e^{x \sin x + \cos x} \left(1 - \frac{1}{\cos x} \right) C$

Answer: B



Watch Video Solution

25. $\int \frac{1+x+\sqrt{x+x^2}}{\sqrt{x}+\sqrt{1+x}} dx =$

A. $\frac{1}{3} \left(2^{1/2} - 1 \right)$

B. $\frac{2}{3} \left(2^{1/2} - 1 \right)$

C. $\frac{2}{3} \left(2^{2/3} - 1 \right)$

D. $\frac{1}{3} \left(2^{3/2} - 1 \right)$

Answer: C



Watch Video Solution

26. $\int x^{x^2+1} (2 \ln x + 1) dx$

A. $x^{2x} + C$

B. $x^2 \ln x + C$

C. $x^{(x^x)} + C$

D. $(x^3)^x + C$

Answer: D



Watch Video Solution

27. If $\int \frac{\cos ec^2 x - 2010}{\cos^{2010} x} dx = \frac{f(x)}{(g(x))^{2010}} + C$, where $f\left(\frac{\pi}{4}\right) = 1$,

then the number of solution of the equation $\frac{f(x)}{g(x)} = \{x\}$ in $[0, 2\pi]$

is/are: (where $\{.\}$ represents fractional part function)

A. 0

B. 1

C. 2

D. 3

Answer: A



Watch Video Solution

28. $\int x^x \left((\ln x)^2 - \frac{1}{x} \right) dx$ is equal to:

A. $x^2 \left((\ln x)^2 - \frac{1}{x} \right) + C$

B. $x^x (\ln x - x) + C$

C. $x^x \frac{(\ln x)^2}{2} + C$

D. $x^x \ln x + C$

Answer: D



Watch Video Solution

29. $\int \frac{x^2 - 1}{x^3 \sqrt{2x^4 - 2x^2 + 1}} dx$ is equal to

A. $\frac{\sqrt{2x^4 - 2x^2 + 1}}{x^2} + C$

B. $\frac{\sqrt{2x^4 - 2x^2 + 1}}{x} + C$

C. $\frac{\sqrt{2x^4 - 2x^2 + 1}}{x} + C$

D. $\frac{\sqrt{2x^4 - 2x^2 + 1}}{2x^2} + C$

Answer: D



Watch Video Solution

30. $\int \left\{ \frac{(\ln x - 1)}{1 + (\ln x)^2} \right\}^2 dx$ is equal to

- A. $\frac{x}{x^2 + 1} + C$
- B. $\frac{\ln x}{(\ln x)^2 + 1} + C$
- C. $\frac{x}{1 + (\ln x)^2} + C$
- D. $e^x \left(\frac{x}{x^2 + 1} \right) + C$

Answer: C



Watch Video Solution

31. $I = \int \frac{dx}{\sqrt[4]{(x-1)^3(x+2)^5}} = k^4 \sqrt{\frac{x-1}{x+2}} + C$, then 'k' is equal to:

- A. $\frac{1}{3}$
- B. $\frac{2}{3}$
- C. $\frac{3}{4}$
- D. $\frac{4}{3}$

Answer: D



Watch Video Solution

32. If : $\int \frac{1 - x^7}{x(1 + x^7)} dx = a \cdot \log|x| + b \cdot \log|x^7 + 1| + c$, then : $(a, b) \equiv$

A. $2P - 7Q = 0$

B. $2P + 7Q = 0$

C. $7P + 2Q = 0$

D. $7P - 2Q = 1$

Answer: B



Watch Video Solution

33. $\int \frac{\sin^8 x - \cos^8 x}{1 - 2 \sin^2 x \cos^2 x} dx =$

A. $\sin 2x + C$

B. $\frac{\sin 2x}{2} + C$

C. $\frac{-\sin 2x}{2} + C$

D. $-2 \sin 2x + C$

Answer: C



Watch Video Solution

34. $\int \frac{(\sin 2x)^{1/3} d(\sqrt[3]{\tan x})}{\sin^{2/3} x + \cos^{2/3} x}$ is equal to

A. $\frac{1}{2^{2/3}} \ln(1 + \tan^{1/3} x) + C$

B. $\ln(1 + \tan^{2/3} x) + C$

C. $2^{1/3} \ln(1 + \tan^{2/3} x) + C$

D. $\frac{1}{2^{2/3}} \ln(1 + \tan^{2/3} x) + C$

Answer: D



Watch Video Solution

$$35. \int \sqrt{\frac{(2012)^{2x}}{1 - (2012)^{2x}}} (2012)^{\sin^{-1}(2012)^x} dx =$$

- A. $(\log_{2012} e)^2 (2012)^{\sin^{-1}(2012)^x} + C$
- B. $(\log_{2012} e)^2 (2012)^{x - \sin^{-1}(2012)^x} + C$
- C. $(\log_{2012} e)^2 (2012)^{\sin^{-1}(2012)^x} + C$
- D. $\frac{(2012)^{\sin^{-1}(2012)^x}}{(\log_{2012} e)^2} + C$

Answer: C



Watch Video Solution

$$36. \int \frac{x+2}{(x^2+3x+3)\sqrt{x+1}} dx \text{ is equal to}$$

- A. $\frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{x}{\sqrt{3(x+1)}} \right) + C$
- B. $\frac{2}{\sqrt{3}} \tan^{-1} \left(\sqrt{\frac{x}{3(x+1)}} \right) + C$
- C. $\frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{x}{3(x+1)} \right) + C$

$$D. \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{x}{\sqrt{3}(x+1)} \right) + C$$

Answer: A



Watch Video Solution

37. $\int \left(\frac{f(x)g'(x) - f'(x)g(x)}{f(x)g(x)} \right) \cdot (\log(g(x)) - \log(f(x))) dx$ is

A. $\log \left(\frac{g(x)}{f(x)} \right) + C$

B. $\frac{1}{2} \left(\frac{g(x)}{f(x)} \right)^2 + C$

C. $\frac{1}{2} \left(\log \left(\frac{g(x)}{f(x)} \right) \right)^2 + C$

D. $\log \left(\left(\frac{g(x)}{f(x)} \right)^2 \right) + C$

Answer: C



Watch Video Solution

38. $\int \left(\int e^x \left(\ln x + \frac{2}{x} - \frac{1}{x^2} \right) dx \right) dx =$

A. $e^x \ln x + C_1 x + C_2$

B. $e^x \ln x + \frac{1}{x} + C_1 x + C_2$

C. $\frac{\ln x}{x} + C_1 x + C_2$

D. None of these

Answer: A



Watch Video Solution

39. Maximum value of the function $f(x) = \pi^2 \int_0^2 t \sin(x + \pi t) dt$ over all

real numbers x:

A. $\sqrt{\pi^2 + 1}$

B. $\sqrt{\pi^2 + 2}$

C. $\sqrt{\pi^2 + 3}$

D. $\sqrt{\pi^2 + 4}$

Answer: D



Watch Video Solution

40. Let 'f' is a function, continuous on $[0, 1]$ such that $f(x) \leq \sqrt{5} \forall x \in [0, 1]$ and $f(x) \leq \frac{2}{x} \forall x \in \left[\frac{1}{2}, 1\right]$ then smallest 'a' for which $\int_0^1 f(x) dx \leq a$ holds for all 'f' is

A. $\sqrt{5}$

B. $\frac{\sqrt{5}}{2} + 2 \ln 2$

C. $2 + \ln\left(\frac{\sqrt{5}}{2}\right)$

D. $2 + 2 \ln\left(\frac{\sqrt{5}}{2}\right)$

Answer: D



Watch Video Solution

41. Let $I_n = \int_1^{e^2} (\ln x)^n dt(x^2)$, then the value of $2I_n + nI_{n-1}$ equals to:

A. 0

B. $2e^2$

C. e^2

D. 1

Answer: B



[View Text Solution](#)

42. Let a function $f: R \rightarrow R$ be defined as $f(x) = x + \sin x$. The value of $\int_0^{2\pi} f^{-1}(x) dx$ will be:

A. $2\pi^2$

B. $2\pi^2 - 2$

C. $2x^2 + 2$

D. π^2

Answer: A



[Watch Video Solution](#)

43. The value of the definite integral

$$\int e^{-x^4} \left(2 + \ln(x + \sqrt{x^2 + 1}) + 5x^3 - 8x^4 \right) dx$$
 is equal to

A. $4e$

B. $\frac{4}{e}$

C. $2e$

D. $\frac{2}{e}$

Answer: B



Watch Video Solution

44. $\int_{-10}^0 \left(\left| \frac{2[x]}{3x - [x]} \right| / \frac{2[x]}{3x - [x]} \right) dx$ is equal to (where $[*]$ denotes greatest integer function.) is equal to (where $[*]$ denotes greatest integer function.)

A. $\frac{28}{3}$

B. $\frac{1}{3}$

C. 0

D. None of these

Answer: A



Watch Video Solution

45. If $f(x) = \frac{x}{1 + (\ln x)(\ln x)} \forall x \in [1, \infty) \dots \infty$ then $\int_1^{2e} f(x)dx$ equals:

A. $\frac{e^2 - 1}{2}$

B. $\frac{e^2 + 1}{2}$

C. $\frac{e^2 - 2e}{2}$

D. None of these

Answer: A



Watch Video Solution

46. $\int_0^4 \frac{(y^2 - 4y + 5)\sin(y - 2)dy}{[2y^2 - 9y + 11]}$ is equal to
(a) 0 (b) 2 (c) -2 (d) none of

these

A. 0

B. 2

C. -2

D. None of these

Answer: A



Watch Video Solution

47. Let $\frac{d}{dx}F(x) = \frac{e^{\sin x}}{x}$, $x > 0$,

If $\int_1^4 \frac{3}{x} e^{\sin x^3} dx = F(k) - F(1)$, then one of the possible value of k is -

A. 15

B. 16

C. 63

D. 64

Answer: D



Watch Video Solution

48. Value of $\lim_{h \rightarrow 0} \frac{\int_0^{x - he^{-1/h}} dx - \int_0^{\pi} x^2 e^{-x^2} dx}{he^{-1/h}}$ in equal to:

A. $\pi(1 - \pi^2)e^{-\pi^2}$

B. $2\pi(1 - \pi^2)e^{-\pi^2}$

C. $\pi(1 - \pi)e^{-\pi}$

D. $\pi^2 e^{-\pi^2}$

Answer: D



View Text Solution

49. Let $f: R^+ \rightarrow R$ be a differentiable function with $f(1) = 3$ and satisfying :

$$\int_1^{xy} f(t) dt = y \int_1^x f(t) dt + x \int_1^y f(t) dt \quad \forall x, y \in R^+, \text{ then } f(e) =$$

A. 3

B. 4

C. 1

D. None of these

Answer: D



[View Text Solution](#)

50. If $[.]$ denotes the greatest integer function, then the integral

$$\int_0^{x/2} \frac{e^{\sin x - [\sin x]} d(\sin^2 x - [\sin^2 x])}{\sin x - [\sin x]} \text{ is } \lambda, \text{ then } [\lambda - 1] \text{ is equal to:}$$

A. 0

B. 1

C. 2

D. 3

Answer: C



View Text Solution

51. Calculate the reciprocal of the limit $\lim_{x \rightarrow \infty} \int_0^x xe^{t^2 - x^2} dt$

A. 0

B. 1

C. 2

D. 3

Answer: A



Watch Video Solution

52.

Let

$$L = \lim_{n \rightarrow \infty} \left(\frac{(2.1 + n)}{1^2 + n.1 + n^2} + \frac{(2.2 + n)}{2^2 + n.2 + n^2} + \frac{(2.3 + n)}{3^2 + n.3 + n^2} + \dots \dots + \right)$$

then value of e^L is:

A. 2

B. 3

C. 4

D. $\frac{3}{2}$

Answer: B



[View Text Solution](#)

53. The value of the definite integral $\int_0^2 \left(\sqrt[3]{1+x^3} + \sqrt[3]{x^2+2x} \right) dx$ is :

A. 4

B. 5

C. 6

D. 7

Answer: C



Watch Video Solution

54. The value of the definite integral $\int_0^{\infty} \frac{\ln x}{x^2 + 4} dx$ is:

A. $\frac{\pi \ln 3}{2}$

B. $\frac{\pi \ln 2}{3}$

C. $\frac{\pi \ln 2}{4}$

D. $\frac{\pi \ln 4}{3}$

Answer: C



Watch Video Solution

55. The value of the definite integral

$$\int_0^{10} \left((x - 5) + (x - 5)^2 + (x - 5)^3 \right) dx \text{ is:}$$

A. $\frac{125}{3}$

B. $\frac{250}{3}$

C. $\frac{125}{6}$

D. $\frac{250}{4}$

Answer: B



Watch Video Solution

56. The value of definite integral $\int_0^{\infty} \frac{dx}{(1+x^9)(1+x^2)}$ equal to:

A. $\frac{\pi}{16}$

B. $\frac{\pi}{8}$

C. $\frac{\pi}{4}$

D. $\frac{\pi}{2}$

Answer: C



Watch Video Solution

57. The value of the definite integral $\int_0^{\pi/2} \left(\frac{1 + \sin 3x}{1 + 2 \sin x} \right) dx$ equals to:

A. $\frac{\pi}{2}$

B. 1

C. $\frac{1}{2}$

D. $\frac{\pi}{4}$

Answer: B



Watch Video Solution

58. The value of $\lim_{x \rightarrow \infty} \frac{\int_0^x (\tan^{-1} x)^2}{\sqrt{x^2 + 1}} dx$

A. $\frac{\pi^2}{16}$

B. $\frac{\pi^2}{4}$

C. $\frac{\pi^2}{2}$

D. None of these

Answer: B



Watch Video Solution

59.

If

$$\int_0^1 \left(\sum_{r=1}^{2013} \frac{x}{x^2 + r^2} \prod_{r=1}^{2013} (x^2 + r^2) \right) dx = \frac{1}{2} \left[\left(\prod_{r=1}^{2013} (1 + r^2) \right) - k^2 \right]$$

then $k =$

A. 2013

B. $2013!$

C. 2013^2

D. 2013^{2013}

Answer: B



Watch Video Solution

60. $f(x) = 2x - \tan^{-1} x - \ln\left(x + \sqrt{1 + x^2}\right)$

- A. strictly increases $\forall x \in R$
- B. strictly increases only in $(0, \infty)$
- C. strictly decreases $\forall x \in R$
- D. strictly decreases in $(0, \infty)$ and strictly increases in $(-\infty, 0)$

Answer: A



Watch Video Solution

61. The value of the definite integral

$$\int_0^{\frac{\pi}{2}} \frac{dx}{\tan x + \cot x + \cos ex + \sec x}$$
 is

A. $1 - \frac{\pi}{4}$

B. $\frac{\pi}{4} + 1$

C. $\pi + \frac{1}{4}$

D. None of these

Answer: A



Watch Video Solution

62. The value of the definite integral $\int_3^7 \frac{\cos x^2}{\cos x^2 + \cos((10) - x)^2} dx$ is:

A. 2

B. 1

C. $\frac{1}{2}$

D. None of these

Answer: A



Watch Video Solution

63. The value of the integral $\int_{e^{-1}}^{e^2} \left| \frac{\ln x}{x} \right| dx$ is:

A. $\frac{3}{2}$

B. $\frac{5}{2}$

C. $\frac{3}{2}$

D. 5

Answer: B



Watch Video Solution

64. The value of $\lim_{x \rightarrow \frac{\pi}{4}} \frac{\int_2^{\cos e^{2x}} tg(t) dt}{x^2 - \frac{\pi^2}{16}}$ is:

A. $\frac{2}{\pi}g(2)$

B. $-\frac{4}{\pi}g(2)$

C. $-\frac{16}{\pi}g(2)$

$$\text{D. } -\frac{16}{\pi}g(2)$$

Answer: C



Watch Video Solution

65. The value of $\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{n-k}{n} \cos\left(\frac{4k}{n}\right)$ equals to

A. $\frac{1}{4}\sin 4 + \frac{1}{16}\cos 4 - \frac{1}{16}$

B. $\frac{1}{4}\sin 4 - \frac{1}{16}\cos 4 + \frac{1}{16}$

C. $\frac{1}{16}(1 - \sin 4)$

D. $\frac{1}{16}(1 - \cos 4)$

Answer: D



Watch Video Solution

66. For each ositive integer n , define a function f_n on $[0, 1]$ as follows:

$$f_n(x) = \begin{cases} 0 & \text{if } x=0 \\ \sin \frac{\pi}{2n} & \text{if } 0 < x \leq \frac{1}{n} \\ \sin \frac{2\pi}{2n} & \text{if } \frac{1}{n} < x \leq \frac{2}{n} \\ \sin \frac{3\pi}{2n} & \text{if } \frac{2}{n} < x \leq \frac{3}{n} \\ \sin \frac{n\pi}{2n} & \text{if } \frac{n-1}{n} < x \leq 1 \end{cases}$$

Then the value of $\lim_{x \rightarrow \infty} \int_0^1 f_n(x) dx$ is:

A. π

B. $\frac{\pi}{2}$

C. $\frac{1}{\pi}$

D. $\frac{2}{\pi}$

Answer: D



Watch Video Solution

67. $\int_0^{n+1} \min \{(x-1), |x-2|, |x-3|, \dots, |x-n|\} dx$

equals to (where n be a positive integer)

A. $\frac{(n+1)}{4}$

- B. $\frac{(n+2)}{4}$
- C. $\frac{(n+3)}{4}$
- D. $\frac{(n+4)}{4}$

Answer: A



Watch Video Solution

68. For positive integers $k=1,2,3,\dots,n$, let S_k denotes the area of $\triangle AOB_k$ such that $\angle AOB_k = \frac{k\pi}{2n}$, $OA=1$ and $OB_k = k$. The value of the

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} \sum_{k=1}^n S_k$$

A. $\frac{2}{\pi^2}$

B. $(4) / (\pi^2)$

C. $\frac{8}{\pi^2}$

D. $\frac{1}{2\pi^2}$

Answer: D



Watch Video Solution

69. If $A = \int_0^1 \prod_{r=4}^{2014} (r - x) dx$ and $B = \int_0^1 \prod_{r=0}^{2013} (r + x) dx$, then:

A. $A = 2B$

B. $2A = B$

C. $A + B = 0$

D. $A = B$

Answer: D



Watch Video Solution

70. If $f(x) = \left[\frac{x}{120} + \frac{x^3}{30} \right]$ defined in $[0, 3]$ then $\int_0^1 ((f(x)) + 2) dx =$
(where $[.]$ denotes greatest integer function)

A. 0

B. 1

C. 2

D. 4

Answer: B



Watch Video Solution

71. If $f(x) = \int_0^{g(x)} \frac{dt}{\sqrt{1+t^3}}$, $g(x) = \int_0^{\cos x} (1+\sin t)^2 dt$, then the value of $f' \left(\frac{\pi}{2} \right)$ is equal to:

A. 1

B. -1

C. 0

D. $\frac{1}{2}$

Answer: D



Watch Video Solution

72. Let $f(x) = \frac{1}{x^2} \int_4^x (4t^2 - 2f'(t))dt$, find $9f'(4)$

A. 16

B. 4

C. 8

D. 32

Answer: D



Watch Video Solution

73. Evaluate $\lim_{x \rightarrow \infty} \left(\frac{1^2}{n^3 + 1^3} + \frac{2^2}{n^3 + 2^3} + \frac{3^2}{n^3 + 3^3} + \dots + \frac{4}{9n} \right)$.

A. $\frac{1}{3} \ln 3$

B. $\frac{\ln 9}{3}$

C. $\frac{\ln 4}{3}$

D. $\frac{\ln 6}{3}$

Answer: A



Watch Video Solution

74. The value of $\int_0^{2\pi} \cos^{-1} \left(\frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} \right) dx$ is:

A. π^2

B. $\frac{\pi^2}{2}$

C. $2\pi^2$

D. π^3

Answer: D



Watch Video Solution

75. Given a function 'g' continuous everywhere such that $\int_0^1 g(t)dt = 2$ and $g(1) = 5$. If $f(x) = \frac{1}{2} \int_0^x (x-t)^2 g(t)dt$, then the value of $f'(1) - f'(1)$ is:

A. 0

B. 1

C. 2

D. 3

Answer: B



Watch Video Solution

76. If $\int_0^x \frac{x^3 \cos^4 x \sin^2 x}{\pi^2 - 3\pi x + 3x^2} dx = \lambda \int_0^{\pi/2} \sin^2 x dx$, then the value of λ is:

A. $\frac{\pi}{12}$

B. $\frac{\pi}{(8)}$

C. $\frac{\pi}{4}$

D. $\frac{\pi}{3}$

Answer: A



[View Text Solution](#)

77. $\int_0^{\sqrt{3}} \left(\frac{1}{2} \frac{d}{dx} \left(\tan^{-1} \frac{2x}{1-x^2} \right) \right) dx$ equals to:

A. $\frac{\pi}{3}$

B. $-\frac{\pi}{6}$

C. $\frac{\pi}{2}$

D. None of these

Answer: B



[Watch Video Solution](#)

78. Let $y = \{x\}^{\{x\}}$ then the value of $\int_0^3 y dx$ equals to (where $\{.\}$ and $[.]$ denote fractional part and integerpart function respectively)

A. 1

B. $\frac{11}{6}$

C. 3

D. $\frac{5}{6}$

Answer: C



Watch Video Solution

79. $\int_0^1 \frac{\tan^{-1}}{x} dx =$

A. $\int_0^{\pi/4} \frac{\sin x}{x} dx$

B. $\int_0^{\pi/2} \frac{x}{\sin x} dx$

C. $\frac{1}{2} \int_0^{\pi/2} \frac{x}{\sin x} dx$

D. $\frac{1}{2} \int_0^{\pi/2} \frac{x}{\sin x} dx$

Answer: C



Watch Video Solution

80. The value of $\int_0^{4/\pi} \left(3x^2 \sin \frac{1}{x} - x \cos \frac{1}{x} \right) dx$ is:

A. $\frac{8\sqrt{2}}{\pi^3}$

B. $\frac{24\sqrt{2}}{\pi^3}$

C. $\frac{32\sqrt{2}}{\pi^3}$

D. None of these

Answer: C



Watch Video Solution

81. Number of values of x satisfying the equation

$$\int_{-1}^x \left(8t^2 + \frac{28}{3}t + 4\right) dt = \frac{\left(\frac{3}{2}\right)x + 1}{\log_{x+1} \sqrt{x+1}}, \text{ is}$$

A. 0

B. 1

C. 2

D. 3

Answer: B



Watch Video Solution

82. The value of

$$\left[\lim_{n \rightarrow \infty} \frac{1 + 2^4 + 3^4 + \dots + n^4}{n^5} - \lim_{n \rightarrow \infty} \frac{1 + 2^3 + 3^3 + \dots + n^3}{n^5} \right] \text{ is}$$

equal to -

A. $\frac{1}{30}$

B. zero

C. $\frac{1}{4}$

D. $\frac{1}{5}$

Answer: D



Watch Video Solution

83. The value of $\lim_{x \rightarrow 0^+} \frac{\int_1^{\cos x} (\cos^{-1} t) dt}{2x - \sin(2x)}$ is equal to

A. 0

B. -1

C. $\frac{2}{3}$

D. $-\frac{1}{4}$

Answer: D



Watch Video Solution

84. Consider a parabola $y = \frac{x^2}{4}$ and the point $F(0, 1)$.

Let $A_1(x_1, y_1), A_2(x_2, y_2), A_3(x_3, y_3), \dots, A_n(x_n, y_n)$ are 'n' points on the parabola such $x_k > -$ and $\angle OFA_k = \frac{k\pi}{2\pi} (k = 1, 2, 3, \dots, n)$.

Then the value of $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n FA_k$, is equal to :

A. $\frac{2}{\pi}$

B. $\frac{4}{\pi}$

C. $\frac{8}{\pi}$

D. None of these

Answer: B



[View Text Solution](#)

85. The minimum value of $f(x) = \int_0^4 e^{|x-t|} dt$ where $x \in [0, 3]$ is : (A)

$2e^2 - 1$ (B) $e^4 - 1$ (C) $2(e^2 - 1)$ (D) $e^2 - 1$

A. $2e^2 - 1$

B. $e^4 - 1$

C. $e(e^2 - 1)$

D. $e^2 - 1$

Answer: C



Watch Video Solution

86. If $\int_0^\infty \frac{\cos x}{x} dx = \frac{\pi}{2}$, then $\int_0^\infty \frac{\cos^3 x}{x} dx$ is equals to:

A. $\frac{\pi}{2}$

B. $\frac{\pi}{4}$

C. π

D. $\frac{3\pi}{2}$

Answer: A



Watch Video Solution

87. $\int \sqrt{1 + \sin x} \left(\cos \frac{x}{2} - \sin \frac{x}{2} \right) dx$ is:

- A. $\frac{1 + \sin x}{2} + C$
- B. $(1 + \sin x)^2$
- C. $\frac{1}{\sqrt{1 + \sin x}} + C$
- D. $\sin x + C$

Answer: D



Watch Video Solution

88. If $I_n = \int_0^\pi \frac{\sin(2nx)}{\sin 2x} dx$, then the value of $I_{n+\frac{1}{2}}$ is equal to ($n \in I$):

- A. $\frac{n\pi}{2}$
- B. π
- C. $\frac{\pi}{2}$
- D. 0

Answer: D



Watch Video Solution

89. Find the value of the function

$$f(x) = 1 + x + \int_1^x \left((\ln t)^2 + 2 \ln t \right) dt \text{ where } f'(x) \text{ vanishes}$$

A. $\frac{1}{e}$

B. 0

C. $\frac{2}{e}$

D. $1 + \frac{2}{e}$

Answer: D



Watch Video Solution

90. Let $f: R \rightarrow R$ be a differentiable function satisfying

$$f(x) = x^2 + 3 \int_0^{x^3} e^{-t^3} \cdot t^2 \cdot f(x - t^3) dt. \text{ Then find } f(x).$$

A. $\frac{1}{3}$

B. $\frac{1}{4}$

C. $\frac{7}{12}$

D. $\frac{5}{12}$

Answer: D



Watch Video Solution

91. The value of the definite integral $\int_{-(\pi/2)}^{\pi/2} \frac{\cos^2 x}{1 + 5^x}$ equal to:

A. $\frac{3n}{4}$

B. π

C. $\frac{\pi}{2}$

D. $\frac{\pi}{4}$

Answer: D



Watch Video Solution

$$92. \int \left(\frac{x^2 - x + 1}{x^2 + 1} \right) e^{\cot^{-1}(x)} dx = f(x) \cdot e^{\cot^{-1}(x)} + C$$

where C is constant of integration. Then f(x) is equal to:

A. $-x$

B. $\sqrt{1-x}$

C. x

D. $\sqrt{1+x}$

Answer: C



Watch Video Solution

$$93. \lim_{x \rightarrow \infty} \frac{1}{n^3} \left(\sqrt{n^2 + 1} + 2\sqrt{n^2 + 2^2} + \dots + n\sqrt{(n^2 + n^2)} \right) = :$$

A. $\frac{3\sqrt{2} - 1}{2}$

B. $\frac{2\sqrt{2} - 1}{3}$

C. $\frac{3\sqrt{3} - 1}{3}$

D. $\frac{4\sqrt{2} - 1}{2}$

Answer: B



Watch Video Solution

94. $\int \frac{x^3 - 1}{(x^4 + 1)(x + 1)} dx$ is

A. $\frac{1}{4}\ln(1 + x^4) + \frac{1}{3}\ln(1 + x^3) + c$

B. $\frac{1}{4}\ln(1 + x^4) - \frac{1}{3}\ln(1 + x^3) + c$

C. $\frac{1}{4}\ln(1 + x^4) - \ln(1 + x) + c$

D. $\frac{1}{4}\ln(1 + x^4) + \ln(1 + x) + c$

Answer: C



Watch Video Solution

95. The value of $\lim_{x \rightarrow 0^+} \frac{\int_1^{\cos x} (\cos^{-1} t) dt}{2x - \sin 2x}$ is equal to:

A. 0

B. -1

C. $\frac{2}{3}$

D. $-\frac{1}{4}$

Answer: D



[Watch Video Solution](#)

96. Let $f(x) = \lim_{n \rightarrow \infty} \frac{\cos x}{(1 + \tan^{-1} x)^n}$, then $\int_0^\infty f(x) dx =$

A. $\tan(\sin 1)$

B. $\sin(\tan 1)$

C. 0

D. $\sin\left(\frac{\tan 1}{2}\right)$

Answer: B



Watch Video Solution

97. The value of $\lim_{x \rightarrow \infty} \sum_{k=1}^n \left(\frac{k}{n^2 + n + 2k} \right) =$

A. $\frac{1}{4}$

B. $\frac{1}{3}$

C. $\frac{1}{2}$

D. 1

Answer: C



Watch Video Solution

98. The value of $\lim_{y \rightarrow 1^+} \frac{\int_1^y |t - 1| dt}{\tan(y - 1)}$ is:

A. 0

B. 1

C. 2

D. does not exist

Answer: A



Watch Video Solution

99.

Given

that

$$\int \frac{dx}{(1+x^2)} = \frac{x}{2(n-1)(1+x^2)^{n-1}} + \frac{(2n-3)}{2(n-1)} \int \frac{dx}{(1+x^2)^{n-1}}. \quad \text{Find}$$

the value of $\int_0^1 \frac{dx}{(1+x^2)^4}$, (you may or may not use reduction formula

given)

A. $\frac{11}{48} + \frac{5\pi}{64}$

B. $\frac{11}{48} + \frac{5\pi}{32}$

C. $\frac{1}{24} + \frac{5\pi}{64}$

D. $\frac{1}{96} + \frac{5\pi}{32}$

Answer: A



Watch Video Solution

100. Find the value of $\int_0^{\pi/4} (\sin x)^4 \, dx :$

A. $\frac{3\pi}{16}$

B. $\frac{3\pi}{32} - \frac{1}{4}$

C. $\frac{3\pi}{32} - \frac{3}{4}$

D. $\frac{3\pi}{16} - \frac{7}{8}$

Answer: B



Watch Video Solution

101. $\int \frac{\cos 9x + \cos 6x}{2 \cos 5x - 1} dx = A \sin 4x + B \sin + C,$ then $A + B$ is equal

to:

(Where C is constant of integration)

A. $\frac{1}{2}$

B. $\frac{3}{4}$

C. 2

D. $\frac{5}{4}$

Answer: D



Watch Video Solution

102. $\int \frac{dx}{x^{2014} + x} = \frac{1}{p} \ln\left(\frac{x^q}{1+x^r}\right) + C$ where $p, qr \in N$ then the value of $(p + q + r)$ equals (Where C is constant of integration)

A. 6039

B. 6048

C. 6047

D. 6021

Answer: A



Watch Video Solution

103. If $\int_0^1 e^{-x^2} dx = 0$, then $\int_0^1 x^2 e^{-x^2} dx$ is equal to

A. $\frac{1}{2e}(e - 1)$

B. $\frac{1}{2e}(e + 1)$

C. $\frac{1}{e}(e - 1)$

D. $\frac{1}{e}(e + 1)$

Answer: A



Watch Video Solution

104. $f(x)$ is a continuous function for all real values of x and satisfies

$\int_n^{n+1} f(x) dx = \frac{n^2}{2} \forall n \in I$. Then $\int_{-3}^5 f(|x|) dx$ is equal to (b) $\frac{35}{2}$

(c) $\frac{17}{2}$ (d) none of these

A. $\frac{19}{2}$

B. $\frac{35}{2}$

C. $\frac{17}{2}$

D. $\frac{37}{2}$

Answer: B



Watch Video Solution

105. If $\int \frac{dx}{x^4(1+x^2)} = a \ln \left| \frac{1+x^3}{x^3} \right| + \frac{b}{x^3} + \frac{c}{1+x^2} + d$. then (where d is arbitrary constant)

A. $a = \frac{1}{3}, b = \frac{1}{3}, x = \frac{1}{3}$

B. $a = \frac{2}{3}, b = -\frac{1}{3}, c = \frac{1}{3}$

C. $a = \frac{2}{3}, b = -\frac{1}{3}, c = -\frac{1}{3}$

D. $a = \frac{2}{3}, b = \frac{1}{3}, c = -\frac{1}{3}$

Answer: C



Watch Video Solution

106. $\lim_{x \rightarrow \infty} \frac{1}{\sqrt{n}\sqrt{n+1}} + \frac{1}{\sqrt{n}\sqrt{n+2}}$ is equal to :

A. 2

B. 4

C. $2(\sqrt{2} - 1)$

D. $2\sqrt{2} - 1$

Answer: A



Watch Video Solution

107. Let $f(x) = \int_z^2 \frac{dy}{\sqrt{1+y^3}}$. The value of the integral $\int_0^2 xf(x)dx$ is equal to:

A. 1

B. $\frac{1}{3}$

C. $\frac{4}{3}$

D. $\frac{2}{3}$

Answer: D



Watch Video Solution

108. The value of the definite integral $\int_0^{\pi/3} \ln(1 + \sqrt{3} \tan x) dx$ equals

A. $\frac{\pi}{23} \ln 2$

B. $\frac{\pi}{3}$

C. $\frac{\pi^2}{6} \ln 2$

D. $\frac{\pi}{2} \ln 2$

Answer: A



Watch Video Solution

109. Q. if $\int_0^{100} (f(x)dx = a)$, then $\sum_{r=1}^{100} \left(\int_0^1 (f(r-1+x)dx) \right) =$

A. 100 a

B. a

C. 0

D. 0.4

Answer: B



Watch Video Solution

110. The value of $\int_0^1 \lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{x^{k+2} 2^k}{k!} dx$ is:

A. $e^2 - 1$

B. 2

C. $\frac{e^2 - 1}{2}$

D. $\frac{e^2 - 1}{4}$

Answer: D



Watch Video Solution

111. Evaluate: $\int x^5 \sqrt{1+x^3} dx$.

A. $\frac{1}{15}(1+x^2)^{5/2} - \frac{1}{9}(1+x)^3\Big)^{3/2} + xc$

B. $\frac{2}{15}(1+x^3)^{5/2} - \frac{1}{9}(1+x^3)^{3/2} + c$

C. $\frac{2}{15}(1+x^2)^{5/2} - \frac{2}{9}(1+x^3)^{3/2} + c$

D. $\frac{1}{15}(1+x^3)^{5/2} - \frac{2}{9}(1+x^3)^{3/2} + c$

Answer: C



Watch Video Solution

112. If $f(x) = \int_0^x \sin t/t dt$, which of the following is true?

A. $f(0) > f(1.1)$

B. $f(0) < f(1.1) > f(2.1)$

C. $f(0) < f(1.1) < f(2.1) > f(3.1)$

D. $f(0) < f(1.1) < f(2.1) < f(3.1) > f(4.1)$

Answer: D



Watch Video Solution

113. Evaluate : $\int \frac{x^3 + 3x^2 + x + 9}{(x^2 + 1)(x^2 + 3)} dx.$

A. $\ln|x^2 + 3| + 3\tan^{-1}x + c$

B. $\frac{1}{2}\ln|x^2 + 3| + \tan^{-1}x + c$

C. $\frac{1}{2}\ln|x^2 + 3| + 3\tan^{-1}x + c$

D. $\ln|x^2 + 3| - \tan^{-1}x + c$

Answer: C



Watch Video Solution

114. $\int \frac{\sqrt{\sec^5 x}}{\sqrt{\sin^3 x}} dx$ equals to:

- A. $(\tan x)^{3/2} - \sqrt{\tan x} + C$
- B. $2\left(\frac{1}{3}(\tan x)^{3/2} - \frac{1}{\sqrt{\tan x}}\right) + C$
- C. $\frac{1}{3}(\tan x)^{3/2} - \sqrt{\tan x} + C$
- D. $\sqrt{\sin x} + \sqrt{\cos x} + C$

Answer: B



Watch Video Solution

115. $\lim_{x \rightarrow 0} \int_0^x \frac{e^{\sin(tx)}}{x} dt$ equals to :

- A. 1
- B. 2
- C. e
- D. does not exist

Answer: A



Watch Video Solution

116. If $A = \int_0^{\pi} \frac{\sin x}{x^2} dx$, then $\int_0^{\pi/2} \frac{\cos 2x}{x} dt$ is equal to:

A. $1 - A$

B. $\frac{3}{2} - A$

C. $A - 1$

D. $1 + A$

Answer: C



Watch Video Solution

Exercise One Or More Than One Answer Is Are Correct

$$1. \int \frac{dx}{(1 + \sqrt{x})^8} = -\frac{1}{3(1 + \sqrt{x})^{k_2}} + \frac{2}{7(1 + \sqrt{x})^{k_2}} + C, \text{ then:}$$

A. $K_1 = 5$

B. $k_1 = 6$

C. $k_2 = 7$

D. $k_2 = 8$

Answer: B::C



Watch Video Solution

2. If $\int_{-\alpha}^{\alpha} \left(e^x + \cos x \ln(x + \sqrt{1 + x^2}) \right) dx > \frac{3}{2}$, then possible value of α can be:

A. 1

B. 2

C. 3

D. 4

Answer: A::B::C::D



Watch Video Solution

3. For $a > 0$, if $I = \int \sqrt{\frac{x}{a^3 - x^3}} dx = A \sin^{-1} \left(\frac{x^{3/2}}{B} \right) + C$, where C is any arbitrary constant, then:

A. $A = \frac{2}{3}$

B. $B = a^{3/2}$

C. $A = \frac{1}{3}$

D. $B = a^{1/2}$

Answer: A::B



Watch Video Solution

4. Let $\int x \sin x \sec^3 x dx = \frac{1}{2}(x \cdot f(x) - g(x)) + k$, then

- A. $f(x) \not\in (-1, 1)$
- B. $g(x) = \sin x$ has 6 solution for $x \in [-\pi, 2\pi]$
- C. $g'(x) = f(x), \forall x \in R$
- D. $f(x)g(x)$ has no solution

Answer: A::C::D



Watch Video Solution

5. If $\int (\sin 3\theta + \sin \theta) \cos \theta e^{\sin \theta} d\theta = (A \sin^3 \theta + B \cos^2 \theta + C \sin \theta + D \cos \theta + E) e^{\sin \theta} + F$, then

- A. $A = -4$
- B. $B = -12$
- C. $C = -20$
- D. None of these

Answer: A::B::C



Watch Video Solution

6. For $a > 0$, if $I = \int \sqrt{\frac{x}{a^3 - x^3}} dx = A \sin^{-1} \left(\frac{x^{3/2}}{B} \right) + C$, where C is any arbitrary constant, then:

A. $A = \frac{2}{3}$

B. $B = a^{3/2}$

C. $A = \frac{1}{3}$

D. $B = a^{1/2}$

Answer: A::B



Watch Video Solution

7. If $f(\theta) = \lim_{x \rightarrow \infty} \sum_{r=0}^{n\theta} \frac{2r}{n\sqrt{(3\theta n - 2r)(n\theta + 2r)}}$ then:

A. $f(1) = \frac{\pi}{6}$

B. $f(\theta) = \frac{\theta}{2} \int_0^\theta \frac{dx}{\sqrt{\theta^2 - \left(x - \frac{\theta}{2}\right)^2}}$

C. $f(\theta)$ is constnat function

D. $y = f(\theta)$ is invertible

Answer: A::B::D



[View Text Solution](#)

8. If $f(x + y) = f(x)f(y)$ for all x, y and

$f(0) \neq 0$, and $F(x) = \frac{f(x)}{1 + (f(x))^2}$ then:

A. $\int_{-2010}^{2011} F(x)dx = \int_0^{2011} F(x)dx$

B. $\int_{-2010}^{2011} F(x)dx = \int_0^{2011} F(x)dx = \int_0^{2011} F(x)dx$

C. $\int_{-2010}^{2011} F(x)dx = 0$

D. $\int_{-2010}^{2010} (2F(-x) - F(x))dx = e \int_0^{2010} F(x)dx$

Answer: B::D



Watch Video Solution

9. Let $J = \int_{-1}^2 \left(\cot^{-1} \frac{1}{x} + \cot^{-1} x \right) dx$, $K = \int_{-2\pi}^{7\pi} \frac{\sin x}{|\sin x|} dx$. Then which of the following alternative (s) is/are correct ?

- A. $2J + 3K = 8\pi$
- B. $4J^2 + K^2 = 26\pi^2$
- C. $2J = K = 3\pi$
- D. $\frac{J}{K} = \frac{2}{5}$

Answer: A::B



Watch Video Solution

10. Which of the following function (s) is/are even ?

$$\text{A. } f(x) = \int_0^x \ln\left(t + \sqrt{1 + t^2}\right) dt$$

$$\text{B. } g(x) = \int_0^x \frac{(2^t + 1)t}{2^t - 1} dt$$

$$\text{C. } h(x) = \int_0^x \left(\sqrt{1 + t + t^2} - \sqrt{1 - t + t^2} \right) dt$$

$$\text{D. } l(x) = \int_0^x \ln\left(\frac{1-t}{1+t}\right) dt$$

Answer: A::B::C::D



[View Text Solution](#)

11. Let $\lim_{x \rightarrow \infty} \sqrt{\frac{x - \cos^2 x}{x + \sin x}}$ and $I_2 = \lim_{h \rightarrow 0^+} \int_{-1}^1 h \frac{dx}{h^2 + x^2}$. Then

A. Both l_1 and l_2 are less than $22/7$

B. One of the two limits is rational and other irrational

C. $l_2 > l_1$

D. l_2 is greater than 3 times of l_1

Answer: A::B::C::D



[Watch Video Solution](#)

12. For $a > 0$, if $I = \int \sqrt{\frac{x}{a^3 - x^3}} dx = A \sin^{-1} \left(\frac{x^{3/2}}{B} \right) + C$,

where C is any arbitrary constant, then :

A. $A = \frac{2}{3}$

B. $B = a^{3/2}$

C. $A = \frac{1}{3}$

D. $B = a^{1/2}$

Answer: A::B



Watch Video Solution

13. If $\int \frac{dx}{1 - \sin^4 x} = a \tan x + b \tan^{-1}(c \tan x) + D$, then:

A. $a = \frac{1}{2}$

B. $b = \sqrt{2}$

C. $c = \sqrt{2}$

D. $b = \frac{1}{2\sqrt{2}}$

Answer: A::C



Watch Video Solution

14. The value of definite integral: $\int_{-2014}^{2014} \frac{dx}{1 + \sin^{2015} x + \sqrt{1 + \sin^{4030} x}}$
equals :

A. 0

B. 2014

C. $(2014)^{(2)}$

D. 4028

Answer: B



Watch Video Solution

15. Let $L = \lim_{n \rightarrow \infty} \int_a^{\infty} \frac{ndx}{1 + n^2x^2}$, where $a \in R$, then L can be

A. π

B. $\frac{\pi}{2}$

C. 0

D. $\frac{\pi}{3}$

Answer: C



Watch Video Solution

16. Let $I = \int_0^1 \sqrt{\frac{1 - \sqrt{x}}{1 + \sqrt{x}}} dx$ then correct statement (s) is/are:

A. $I = 2$

B. $I = \pi$

C. $I = \frac{2 + \pi}{2}$

D. $I = \frac{4 - \pi}{2}$

Answer: B::C



Watch Video Solution

Exercise Comprehension Type Problems

1. Let $f(x) = \int x^2 \cos^2 x (2x + 6 \tan x - 2x \tan^2 x) dx$ and $f(x)$ passes through the point $(\pi, 0)$

If $f: R - (2n + 1) \frac{\pi}{2} \rightarrow R$ then $f(x)$ be a :

- A. even function
- B. odd function
- C. neither even nor odd
- D. even as well as odd both

Answer: A



Watch Video Solution

2. Let $f(x) = \int x^2 \cos^2 x (2x + 6 \tan x - 2x \tan^2 x) dx$ and $f(x)$ passes through the point $(\pi, 0)$

The number of solution (s) of the equation $f(x) = x^3$ in $[0, 2\pi]$ be:

- A. 0
- B. 3
- C. 2
- D. None of these

Answer: B



Watch Video Solution

3. Let $f(x)$ be a non-constant twice differentiable function defined on $(-\infty, \infty)$ such that $f(x) = f(1-x)$ and $f''(1/4) = 0$. Then

- A. 2
- B. 3

C. 4

D. 5

Answer: C



Watch Video Solution

4. Let $f(x)$ be a twice differentiable function defined on $(-\infty, \infty)$ such that $f(x) = f(2 - x)$ and $f'\left(\frac{1}{2}\right) = f'\left(\frac{1}{4}\right) = 0$. Then $\int_{-1}^1 f'(1 + x)x^2 e^{x^2} dx$ is equal to :

A. 1

B. π

C. 2

D. 0

Answer: D



View Text Solution

5. Let $f(x)$ be a non-constant twice differentiable function defined on $(-\infty, \infty)$ such that $f(x) = f(1-x)$ and $f'\left(\frac{1}{4}\right) = 0$. Then $f'(x)$ vanishes at least twice on $[0, 1]$ $f'\left(\frac{1}{2}\right) = 0$

$$\int_{-\frac{1}{2}}^{\frac{1}{2}} f\left(x + \frac{1}{2}\right) \sin x dx = 0 \int_{-\frac{1}{2}}^{\frac{1}{2}} f(t) e^{\sin \pi t} dt = \int_{\frac{1}{2}}^1 f(1-t) e^{\sin \pi t} dt$$

A. $\int_0^2 f'(t) e^{\cos \pi t} dt$

B. 1

C. 2

D. π

Answer: A



[Watch Video Solution](#)

6. Consider the function $f(x)$ and $g(x)$, both defined from $R \rightarrow R$

$$f(x) = \frac{x^3}{2} + 1 - x \int_0^x g(t) dt \text{ and } g(x) = x - \int_0^1 f(t) dt, \text{ then}$$

minimum value of $f(x)$ is:

- A. 0
- B. 1
- C. $\frac{3}{2}$
- D. does not exist

Answer: B



Watch Video Solution

7. Consider the function $f(x)$ and $g(x)$, both defined from $R \rightarrow R$

$$f(x) = \frac{x^3}{2} + 1 - x \int_0^x g(t)dt \text{ and } g(x) = x - \int_0^1 f(t)dt, \text{ then}$$

The number of points of intersection of $f(x)$ and $g(x)$ is/are:

- A. 0
- B. 1
- C. 2

D. 3

Answer: A



Watch Video Solution

8. Consider the function $f(x)$ and $g(x)$, both defined from $R \rightarrow R$

$$f(x) = \frac{x^3}{2} + 1 - x \int_0^x g(t)dt \text{ and } g(x) = x - \int_0^1 f(t)dt, \text{ then}$$

minimum value of $f(x)$ is:

A. $\frac{9}{4}$

B. $\frac{9}{2}$

C. $\frac{9}{8}$

D. None of these

Answer: C



Watch Video Solution

9. Let $f(x)$ be function defined on $[0, 1]$ such that $f(1) = 0$ and for any $a \in (0, 1]$, $\int_0^a f(x)dx - \int_a^1 f(x)dx = 2f(a) + 3a + b$ where b is constant.

$b =$

A. $\frac{3}{2e} - 3$

B. $\frac{3}{2e} - \frac{3}{2}$

C. $\frac{3}{2e} + 3$

D. $\frac{3}{2e} + \frac{3}{2}$

Answer: A



Watch Video Solution

10. Let $f(x)$ be function defined on $[0, 1]$ such that $f(1) = 0$ and for any $a \in (0, 1]$, $\int_0^a f(x)dx - \int_a^1 f(x)dx = 2f(a) + 3a + b$ where b is constant.

The length of the subtangent of the curve $y = f(x)$ at $x = 1/2$ is:

A. $\sqrt{e} - 1$

B. $\frac{\sqrt{e} - 1}{2}$

C. $\sqrt{e} + 1$

D. $\frac{\sqrt{e} + 1}{2}$

Answer: A



Watch Video Solution

11. Let $f(x)$ be function defined on $[0, 1]$ such that $f(1) = 0$ and for any

$a \in (0, 1]$, $\int_0^a f(x)dx - \int_a^1 f(x)dx = 2f(a) + 3a + b$ where b is constant.

$$\int_0^1 f(x)dx =$$

A. $\frac{1}{e}$

B. $\frac{1}{2e}$

C. $\frac{3}{2e}$

D. $\frac{2}{e}$

Answer: C



Watch Video Solution

12. Let $f_a(x) = \ln x$ and for $n \geq 0$ and $x > 0$

Let $f_a(x) = \int_0^x f_a(t) dt$ then:

$f_a(x)$ equals :

A. $\frac{x^3}{3} \left(\ln x - \frac{5}{6} \right)$

B. $\frac{x^3}{3} \left(\ln x - \frac{11}{6} \right)$

C. $\frac{x^3}{3} \left(\ln x - \frac{11}{6} \right)$

D. $\frac{x^3}{3} \left(\ln x - \frac{5}{6} \right)$

Answer: C



View Text Solution

13. $\int \frac{x^2 + 4}{x^2 + 1} dx$



Watch Video Solution

14. Let $f: R \rightarrow \left[\frac{3}{4}, \infty\right)$ be a surjective quadratic function with line of symmetry $2x - 1 = 0$ and $f(1) = 1$

If $g(x) = \frac{f(x) + f(-x)}{2}$ then $\int \frac{dx}{\sqrt{g(e^x) - 2}}$ is equal to:

A. $\sec^{-1}(e^{-x}) + C$

B. $\sec^{-1}(e^x) + C$

C. $\sin^{-1}(e^{-x}) + C$

D. $\sin^{-1}(e^x) + C$

Answer: B



Watch Video Solution

15. Let $f: R \rightarrow \left[\frac{3}{4}, \infty\right)$ be a surjective quadratic function with line of symmetry $2x - 1 = 0$ and $f(1) = 1$

$$\int \frac{e^x}{f(e^x)} dx$$

- A. $\cot^{-1}\left(\frac{2x^2 - 1}{\sqrt{3}}\right) + C$
- B. $\frac{2}{\sqrt{3}}\cot^{-1}\left(\frac{2x^2 - 1}{\sqrt{3}}\right) + C$
- C. $\tan^{-1}\left(\frac{2x^2 - 1}{\sqrt{3}}\right) + C$
- D. $\frac{2}{\sqrt{3}}\tan^{-1}\left(\frac{2x^2 - 1}{\sqrt{3}}\right) + C$

Answer: D



Watch Video Solution

16.

Let

$$g(x) = x^C e^{Cx} \text{ and } f(x) = \int_0^x te^{2r}(1+3t^2)^{1/2} dt. \text{ If } L = \lim_{x \rightarrow \infty} \frac{f'(x)}{g'(x)}$$

is non-zero finite number then :

The value of C is:

A. 7

B. $\frac{3}{2}$

C. 2

D. 3

Answer: C



Watch Video Solution

17. Let $g(x) = x^C e^{Cx}$ and $f(x) = \int_0^x t e^{2t} (1 + 3t^2)^{1/2} dt$. If $L = \lim_{x \rightarrow \infty} \frac{f'(x)}{g'(x)}$ is non-zero finite number then :

The value of L . Is :

A. $\frac{2}{7}$

B. $\frac{1}{2}$

C. $\frac{\sqrt{3}}{4}$

D. $\frac{\sqrt{3}}{2}$

Answer: D



Watch Video Solution

Exercise Matching Type Problems

	Column-I	Column-II
(A)	$\lim_{n \rightarrow \infty} 4 \left[\frac{\frac{1}{e^n}}{n^2} + \frac{2}{n^2} e^{-n} + \frac{3}{n^2} e^{-2n} + \dots + \frac{1}{n} e^{-n} \right] =$	(P) 0
(B)	$\int_0^1 \ln\left(\frac{1}{x} - 1\right) dx =$	(Q) 1
(C)	$\int_0^{10\pi} \left(\lim_{x \rightarrow y} \left(\frac{\sin x - \sin y}{x - y} \right) \right) dy =$	(R) 2
(D)	$\int_0^{\infty} \frac{\ln\left(x + \frac{1}{x}\right)}{(1+x^2)} dx = \frac{\pi}{2} \ln a, \text{ then } a =$	(S) 4 (T) 5

1.



View Text Solution

2. Match the following $\int f(x) dx$ is equal to, if

	Column-I	Column-II
(A)	$f(x) = \frac{1}{(x^2 + 1)\sqrt{x^2 + 2}}$	(P) $\frac{x^5}{5(1-x^4)^{5/2}} + C$
(B)	$f(x) = \frac{1}{(x+2)\sqrt{x^2 + 6x + 7}}$	(Q) $\sin^{-1}\left(\frac{x+1}{(x+2)\sqrt{2}}\right) + C$
(C)	$f(x) = \frac{x^4 + x^8}{(1-x^4)^{7/2}}$	(R) $(\sqrt{x}-2)\sqrt{1-x} + \cos^{-1}\sqrt{x} + C$
(D)	$f(x) = \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}$	(S) $-\tan^{-1}\sqrt{1+\frac{2}{x^2}} + C$ (T) $\frac{x^6}{6(1-x^4)^{5/2}} + C$



View Text Solution

Column-I		Column-II	
(A)	$\int_0^{\pi/2} \frac{\cos x}{(1 + \sin x)(2 + \sin x)} dx =$	(P)	$\frac{\pi}{6}$
(B)	$\int_0^{41\pi/4} \cos x dx =$	(Q)	$20 + \frac{1}{\sqrt{2}}$
(C)	$\int_{-1/2}^{1/2} \left[[x] + \ln\left(\frac{1+x}{1-x}\right) \right] dx =$ where $[.]$ greatest integer function	(R)	$\ln 4 - \ln 3$
(D)	$\int_0^{\pi/2} \frac{2\sqrt{\cos \theta}}{3(\sqrt{\sin \theta} + \sqrt{\cos \theta})} d\theta =$	(S)	$-\frac{1}{2}$

3.



View Text Solution

Column-I		Column-II	
(A)	If quadratic equation $3x^2 + ax + 1 = 0$ and $2x^2 + bx + 1 = 0$ have a common root then value of $5ab - 2a^2 - 3b^2 =$	(P)	6
(B)	Number of solution of $x^4 - 2x^2 \sin^2 \frac{\pi x}{2} + 1 = 0$ is/are	(Q)	1
(C)	Number of points of discontinuity $y = \frac{1}{u^2 + u - 2}$ where $u = \frac{1}{x-1}$ is/are	(R)	2
(D)	$\int \frac{dx}{\sqrt[3]{x^5(1+x)^7/2}} = A \left(\frac{x+1}{x} \right)^{-1/A} + C$ (Where C is integration constant), then A =	(S)	3

4.



View Text Solution

(A)	$\int_0^{1.5} [x^2] dx$	(P)	-π
(B)	$\int_0^4 \{\sqrt{x}\} dx$	(Q)	$4(\sqrt{2} - 1)$
	where $\{x\}$ denotes the fractional part of x		
(C)	$\int_0^{2\pi} [\sin x + \cos x] dx$	(R)	$\frac{7}{3}$
(D)	$\int_0^{\pi} \sin x - \cos x dx$	(S)	$2 - \sqrt{2}$

5.



[View Text Solution](#)

Exercise Subjective Type Problems

1.
$$\int \frac{x + (\cos^{-1} 3x)^2}{\sqrt{1 - 9x^2}} dx = \frac{1}{k_1} \left(\sqrt{1 - 9x^2} + (\cos^{-1} 3x)^{k_2} \right) + C, \text{ then}$$

$k_1^2 + k_2^2 =$ (where C is an arbitrary constnat.)



[Watch Video Solution](#)

2. If $\int_0^{\infty} \frac{x^3}{(a^2 + x^2)} dx = \frac{1}{ka^6}$, then find the value of $\frac{k}{8}$.



[Watch Video Solution](#)

3. $\int \frac{(x^2 + 1)dx}{x}$



Watch Video Solution

4.

If

$$\int (x^6 + x^4 + x^2) \sqrt{2x^4 + 3x^2 + 6} dx = \frac{(ax^6 + \beta x^4 + \gamma x^2)^{3/2}}{18} + C$$

where C is constnat, then find the value of $(\beta + \gamma - \alpha)$.



Watch Video Solution

5. $\int \frac{x^2 + 3}{x^2 + 2} dx$



Watch Video Solution

6. The value of

$$\int \frac{\tan x}{\tan^2 x + \tan x + 1} dx = x - \frac{2}{\sqrt{A}} \tan^{-1} \left(\frac{2 \tan x + 1}{\sqrt{A}} \right) + C \quad \text{Then}$$

the value of A is:



Watch Video Solution

7. Let $\int_0^1 \frac{4x^3 \left(1 + (x^4)^{2010}\right)}{(1 + x^4)^{2012}} dx = \frac{\lambda}{\mu}$

where λ and μ are relatively prime positive integers. Find unit digit of $\lambda \mu$.



Watch Video Solution

8. Let $\int_1^{\sqrt{5}} \left(x^{2x^2+1} + \ln(x^{2x^{2x^2+1}}) \right) dx = N$. Find the value of $(N - 6)$.



Watch Video Solution

9. If $\int \frac{dx}{\cos^3 x - \sin^3} = A \tan^{-1}(f(x)) + b \ln \left| \frac{\sqrt{2} + f(x)}{\sqrt{2} - f(x)} \right| + C$ where $f(x) = \sin x + \cos x$ find the value of $(12A + 9\sqrt{2}V) - 3$.



Watch Video Solution

10. Find the value of $|a|$ for which the area of triangle included between the coordinate axes and any tangent to the curve $x^a y = \lambda^a$ is constant (where λ is constnat.),



Watch Video Solution

11. Let $I = \int_0^\pi x^6 (\pi - x)^8 dx$, then $\frac{\pi^{15}}{(15C_9)I} =$



Watch Video Solution

12. $\int \frac{x^3}{x^2 - 3} dx$



Watch Video Solution

13. If a continuous function f on $[0, a]$ satisfies $f(x)f(a - x) = 1$, $a > 0$, then find the value of $\int_0^a \frac{dx}{1 + f(x)}$.

 Watch Video Solution

14. If $\{x\}$ denotes the fractional part of x , then $I = \int_0^{100} (\sqrt{x}) dx$, then the value of $\frac{9f}{155}$ is:

 Watch Video Solution

$$15. \int \frac{x^3}{x^2 - 2} dx$$

 Watch Video Solution

16. If M be the maximum value of $72 \int_0^y \sqrt{x^4 + (y - y^2)^2} dx$ for $y \in [0, 1]$, then find $M/6$.



Watch Video Solution

17. Find the number points where $f(\theta) = \int_{-1}^1 \frac{\sin \theta dx}{2x \cos \theta + x^2}$ is discontinuous where $\theta \in [0, 2\pi]$.



Watch Video Solution

18. $\lim_{n \rightarrow \infty} \left[\frac{1}{\sqrt{n}} + \frac{1}{\sqrt{2n}} + \frac{1}{\sqrt{3n}} + \dots + \frac{1}{\sqrt{n}} \right]$



Watch Video Solution

19. The maximum value of $\int_{-\pi/2}^{2\pi/2} \sin x \cdot f(x) dx$, subject to the condition $|f(x)| \leq 5$ is M, then 'M/10' is equal to :



Watch Video Solution

20. Given a function g , continuous everywhere such that $g(1) = 5$ and $\int_0^1 g(t)dt = 2$. If $f(x) = \frac{1}{2} \int_0^x (x-t)^2 g(t)dt$, then find the value of $f''(1) + f'(1)$.



Watch Video Solution

21. If $f(n) = \frac{1}{\pi} \int_0^{\pi/2} \frac{\sin^2(n\theta)d\theta}{\sin^2\theta}$, $n \in N$, then evaluate $\frac{f(15) + f(3)}{f(12) - f(10)}$.



Watch Video Solution

22. Let $f(2-x) = f(2+x)$ and $f(4-x) = f(4+x)$. Function $f(x)$ satisfies $\int_0^2 f(x)dx = 5$. if $\int_0^{50} f(x)dx = l$. Find $[\sqrt{l} - 2]$. (where $[.]$ denotes greatest integer function.)



Watch Video Solution

23.

Let

$$l_n = \int_{-1}^1 |x| \left(1 + x + \frac{x^2}{2} + \frac{x^2}{3} + \dots + \frac{x^{2n}}{2n} \right) dx \quad \text{if } \lim_{x \rightarrow \infty} l_n \text{ can be expressed as rational } \frac{p}{q} \text{ in this lowest form, then find the value of } \frac{pq(p+q)}{10}$$



Watch Video Solution

24. $\int \sqrt{x^2 + 4} dx$



Watch Video Solution

25. If $\int_a^b |\sin x| dx = 8$ and $\int_0^{1+b} |\cos x| dx = 9$ then the value of $\frac{1}{\sqrt{2x}} \left| \int_a^b x \sin x dx \right|$ is:



Watch Video Solution

26. If $f(x)$, $g(x)$, $h(x)$ and $\phi(x)$ are polynomial in x ,
 $\left(\int_1^x f(x)h(x)dx\right)\left(\int_1^x g(x)\phi(x)dx\right) - \left(\int_1^x g(x)h(x)dx\right)$ is divisible by $(x - 1)^2$. Find maximum value of λ .



[Watch Video Solution](#)

27. If $\int_0^2 (3x^2 - 3x + 1) \cos(x^3 - 3x^2 + 4x - 2) dx = a \sin(b)$, where a and b are positive integers. Find the value of $(a + b)$.



[Watch Video Solution](#)

28. let $f(x) = \int_0^x e^{x-y} f'(y) dy - (x^2 - x + 1)e^x$
Find the number of roots of the equation $f(x) = 0$.



[Watch Video Solution](#)

29. For a positive integer n , let $I_n \int_{-\pi}^{\pi} \left(\frac{\pi}{2} - |x| \right) \cos nx dx$

Find the value of $[I_1 + I_3 + I_4]$ where $[.]$ denotes greatest integer function.



[Watch Video Solution](#)