



## MATHS

### BOOKS - VIKAS GUPTA MATHS (HINGLISH)

## CIRCLE

#### Exercise 1 Single Choice Problems

1. The locus of mid-points of the chords of the circle

$x^2 - 2x + y^2 - 2y + 1 = 0$  which are of unit length is :

A.  $(x - 1)^2 + (y - 1)^2 = \frac{3}{4}$

B.  $(x - 1)^2 + (y - 1)^2 = 2$

C.  $(x - 1)^2 + (y - 1)^2 = \frac{1}{4}$

D.  $(x - 1)^2 + (y - 1)^2 = \frac{2}{3}$

**Answer: A**

[Watch Video Solution](#)

2. The length of a common internal tangent to two circles is 5 and a common external tangent is 15, then the product of the radii of the two circles is :

A. 25

B. 50

C. 75

D. 30

**Answer: B**

[Watch Video Solution](#)

3. A circle with center  $(2, 2)$  touches the coordinate axes and a straight line AB where A and B are on the direction of coordinate axes such that the line lies

between and the line AB be the origin then the locus of circumcenter of

$\triangle OAB$  will be:

A.  $xy = x + y + \sqrt{x^2 + y^2}$

B.  $xy = x + y - \sqrt{x^2 + y^2}$

C.  $xy + x + y = \sqrt{x^2 + y^2}$

D.  $xy + x + y + \sqrt{x^2 + y^2} = 0$

**Answer: A**



**Watch Video Solution**

4. Length of chord of contact of point (4,4) with respect to the circle

$x^2 + y^2 - 2x - 2y - 7 = 0$  is

A.  $\frac{3}{\sqrt{2}}$

B.  $3\sqrt{2}$

C. 3

D. 6

**Answer: B**



**Watch Video Solution**

5. Let P, Q, R, S be the feet of the perpendiculars drawn from a point (1, 1) upon the lines  $x + 4y = 12$ ,  $x - 4y + 4 = 0$  and their angle bisectors respectively, then equation of the circle which passes through Q, R, S is :

A.  $x^2 + y^2 - 5x + 3y - 6 = 0$

B.  $x^2 + y^2 - 5x - 3y + 6 = 0$

C.  $x^2 + y^2 - 5x - 3y - 6 = 0$

D. None of these

**Answer: B**



**Watch Video Solution**

6. From a point  $P$  on the line  $2x + y = 4$  which is nearest to the circle  $x^2 + y^2 - 12y + 35 = 0$ . tangents are drawn to give circle. The area of quadrilateral PACB (where 'C' is the center of circle and PA & PB re the tangents.) is

A. 8

B.  $\sqrt{110}$

C.  $\sqrt{19}$

D. None of these

**Answer: C**



**Watch Video Solution**

7. The line  $2x - y + 1 = 0$  is tangent to the circle at the point (2,5) and the centre of the circles lies on  $x-2y = 4$ . The radius of the circle is :

A.  $3\sqrt{5}$

B.  $5\sqrt{3}$

C.  $2\sqrt{5}$

D.  $5\sqrt{2}$

**Answer: A**



**Watch Video Solution**

8. If  $A(\cos \alpha, \sin \alpha)$ ,  $B(\sin \alpha, -\cos \alpha)$ ,  $C(1, 2)$  are the vertices of a triangle, then as  $\alpha$  varies the locus of centroid of the  $\triangle ABC$  is a circle whose radius is :

A.  $\frac{2\sqrt{2}}{3}$

B.  $\sqrt{\frac{4}{3}}$

C.  $\frac{2}{3}$

D.  $\sqrt{\frac{2}{9}}$

**Answer: D**



**Watch Video Solution**

9. Tangents drawn to circle  $(x - 1)^2 + (y - 1)^2 = 5$  at point P meet the line  $2x + y + 6 = 0$  at Q on the axis. Length PQ is equal to

A.  $\sqrt{12}$

B.  $\sqrt{10}$

C. 4

D.  $\sqrt{15}$

**Answer: A**



**Watch Video Solution**

10. ABCD is square in which A lies on positive y-axis and B lies on the positive x-axis. If D is the point (12, 17), then co-ordinate axis. of C is

A. (17, 12)

B. (17, 5)

C. (17, 16)

D. (15, 3)

**Answer: B**



**Watch Video Solution**

**11. Statement-1:** The lines  $y = mx + 1 - m$  for all values of  $m$  is a normal to the circle  $x^2 + y^2 - 2x - 2y = 0$ .

**Statement-2:** The line  $L$  passes through the centre of the circle.

A. Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.

B. Statement-1 is true, statement-2 is true and statement-2 is not the correct explanation for statement-1.

C. Statement-1 is true, statement-2 is false.

D. Statement-1 is false, statement-2 is true.



**Answer: A**



**Watch Video Solution**

**12.** A(1, 0) and 0, 1) are two fixed points on the circle  $x^2 + y^2 = 1$ . C is a variable point on this circle. As C moves, the locus of the orthocentre of the triangle ABC is

A.  $x^2 + y^2 - 2x - 2y + 1 = 0$

B.  $x^2 + y^2 - x - y = 0$

C.  $x^2 + y^2 = 4$

D.  $x^2 + y^2 + 2x - 2y + 1 = 0$

**Answer: A**



**Watch Video Solution**

13. Equation of a circle passing through (1, 2) and (2, 1) and for which line  $x + y = 2$  is a diameter, is :

A.  $x^2 + y^2 + 2x + 2y - 11 = 0$

B.  $x^2 + y^2 - 2x - 2y - 1 = 0$

C.  $x^2 + y^2 - 2x - 2y + 1 = 0$

D. None of these

**Answer: C**



**Watch Video Solution**

14. The area of an equilateral triangle inscribed in a circle of radius  $4cm$ , is

A.  $12cm^2$

B.  $9\sqrt{3}cm^2$

C.  $8\sqrt{3}cm^2$

D.  $12\sqrt{3}cm^2$

**Answer: D**



**Watch Video Solution**

15. let all the points on the curve  $x^2 + y^2 - 10x = 0$  are reflected about the line  $y = x + 3$ . The locus of the reflected points is in the form  $x^2 + y^2 + gx + fy + c = 0$ . The value of  $g + f + c$  is equal to

A. 28

B. -28

C. 38

D. -38

**Answer: C**



**Watch Video Solution**

16. The shortest distance from the line  $3x + 4y = 25$  to the circle  $x^2 + y^2 = 6x - 8y$  is equal to :

A.  $7/5$

B.  $9/5$

C.  $11/5$

D.  $32/5$

**Answer: A**



**Watch Video Solution**

17. Q.ys In the xy-plane, the length of the shortest path from (0.0) to (12.16) that does not go inside the circle  $(y-8)^2 = 25$  is (D) 10 (B) 10 5 (A) 10

Dps' on Circle

A.  $10\sqrt{3}$

B.  $10\sqrt{5}$

C.  $10\sqrt{3} + \frac{5\pi}{3}$

D.  $10 + 5\pi$

**Answer: C**



**Watch Video Solution**

**18.** A circle is inscribed in an equilateral triangle with side lengths 6 unit. Another circle is drawn inside the triangle (but outside the first circle), tangent to the first circle and two of the sides of the triangle. The radius of the smaller circle is

A.  $\frac{1}{\sqrt{3}}$

B.  $\frac{2}{3}$

C.  $\frac{1}{2}$

D. 1

**Answer: A**



**Watch Video Solution**

19. The equation of tangent to the circle  $x^2 + y^2 - 4x = 0$  which is perpendicular to the normal drawn through the origin can be : (A)  $x = 0$   
(B)  $x = 4$  (C)  $x + y = 2$  (D) none of these

A.  $x = 1$

B.  $x = 2$

C.  $x + y = 2$

D.  $x = 4$

**Answer: D**



Watch Video Solution

20. the equation of the line parallel to the line  $3x + 4y = 0$  and touching the circle  $x^2 + y^2 = 9$  in the first quadrant is:

A.  $3x + 4y = 15$

B.  $3x + 4y = 45$

C.  $3x + 4y = 9$

D.  $3x + 4y = 12$

**Answer: A**



**Watch Video Solution**

21. The centres of the three circles

$$x^2 + y^2 - 10x + 9 = 0, x^2 + y^2 - 6x + 2y + 1 = 0, x^2 + y^2 - 9x - 4y + 2 = 0$$

A. lie on the straight line  $x - 2y = 5$

B. lie on circle  $x^2 + y^2 = 25$

C. do not lie on straight line

D. lie on circle  $x^2 + y^2 + x + y - 17 = 0$

**Answer: C**



**Watch Video Solution**

22. The equation of diameter of a circle  $x^2 + y^2 + 2x - 4y = 4$ , that is parallel to  $3x + 5y = 4$  is

A.  $3x + 5y = -7$

B.  $3x + 5y = 7$

C.  $3x + 5y = 9$

D.  $3x + 5y = 1$

**Answer: B**



**Watch Video Solution**

23. There are 2 circles passing through points A(-1, 2) and B(2, 3) having radius  $\sqrt{5}$ . Then the length of intercept on axis of the circle intersecting x-axis is :

A. 2



B. 3

C. 4

D. 5

**Answer: C**



**Watch Video Solution**

**24.** A square OABC is formed by line pairs  $xy = 0$  and  $xy + 1 = x + y$  where o is the origin .A circle with circle  $C_1$  inside the square is drawn to touch the line pair  $xy = 0$  and another circle with centre  $C_2$  and radius twice that  $C_1$  is drawn touch the circle  $C_1$  and the other line .The radius of the circle with centre  $C_1$

A.  $\frac{\sqrt{2}}{\sqrt{3}(\sqrt{2} + 1)}$

B.  $\frac{2\sqrt{2}}{3(\sqrt{2} + 1)}$

C.  $\frac{\sqrt{2}}{3(\sqrt{2} + 1)}$

D.  $\frac{\sqrt{2} + 1}{3\sqrt{2}}$

**Answer: C**



**Watch Video Solution**

**25.** The equation of the circle circumscribing the triangle formed by the points  $(3, 4)$ ,  $(1, 4)$  and  $(3, 2)$  is :

A.  $8x^2 + 8y^2 - 16x - 13y = 0$

B.  $x^2 + y^2 - 4x - 8y + 19 = 0$

C.  $x^2 + y^2 - 4x - 6y + 11 = 0$

D.  $x^2 + y^2 - 6x - 6y + 17 = 0$

**Answer: C**



**Watch Video Solution**

26. The equation of the tangent to circle  $x^2 + y^2 + 2gx + 2fy = 0$  at origin is :

A.  $fx + gy = 0$

B.  $gx + fy = 0$

C.  $x = 0$

D.  $y = 0$

**Answer: B**



**Watch Video Solution**

27. The line  $y = x$  is a tangent at  $(0, 0)$  to a circle of radius unity. The center of the circle is

A. either  $\left(-\frac{1}{2}, \frac{1}{2}\right)$  or  $\left(\frac{1}{2}, -\frac{1}{2}\right)$

B. either  $\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$  or  $\left(-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$

C. either  $\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$  or  $\left(-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$

D. either (1, 0) or (-1, 0)

**Answer: C**



**Watch Video Solution**

**28.** The circles  $x^2 + y^2 + 6x + 6y = 0$  and  $x^2 + y^2 - 12x - 12y = 0$

- A. cut orthogonally
- B. touch each other internally
- C. intersect in two points
- D. touch each other externally

**Answer: D**



**Watch Video Solution**

29. In a right triangle ABC, right angled at A, on the leg AC as diameter, a semicircle is described. The chord joining A with the point of intersection D of the hypotenuse and the semicircle, then the length AC equals to

A.  $\frac{AB \cdot AB}{\sqrt{AB^2 + AB^2}}$

B.  $\frac{AB \cdot AD}{AB + AD}$

C.  $\sqrt{AB \cdot AD}$

D.  $\frac{AB \cdot AD}{\sqrt{AB^2 - AD^2}}$

**Answer: D**



**Watch Video Solution**

30. 30. Radical centre of circles drawn on the sides as a diameter of triangle formed by the lines the  $3x - 4y + 6 = 0$ ,  $x - y + 2 = 0$  and  $4x + 3y - 17 = 0$  is

A. (3, 2)

B. (3, -2)

C. (2, -3)

D. (2, 3)

**Answer: D**



**Watch Video Solution**

**31.** Statement-1: A circle can be inscribed in a quadrilateral whose sides are  $3x - 4y = 0$ ,  $3x - 4y = 5$ ,  $3x + 4y = 0$  and  $3x + 4y = 7$

Statement-2: A circle can be inscribed in a parallelogram if and only if it is a rhombus (a) statement-1 is true, statement-2 is true and statement-2 is correct explanation for Statement-1. (b) Statement-1 is true, statement-2 is true and statement-2 is not the correct explanation for statement-1. (c) Statement-1 is true, statement-2 is false. d) Statement-1 is false, statement-2 is true

A. Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.

- B. Statement-1 is true, statement-2 is true and statement-2 is not the correct explanation for statement-1.
- C. Statement-1 is true, statement-2 is false.
- D. Statement-1 is false, statement-2 is true.

**Answer: D**



**Watch Video Solution**

**32.** If  $x = 3$  is the chord of the contact of the circle  $x^2 + y^2 = 81$ , then the equation of the corresponding pair of tangents, is

A.  $x^2 - 8y^2 + 54x + 729 = 0$

B.  $x^2 - 8y^2 - 54x + 729 = 0$

C.  $x^2 - 8y^2 - 54x - 729 = 0$

D.  $x^2 - 8y^2 = 729$

**Answer: B**

 [Watch Video Solution](#)

33. The shortest distance from the line  $3x + 4y = 25$  to the circle  $x^2 + y^2 = 6x - 8y$  is equal to :

A.  $\frac{7}{3}$

B.  $\frac{9}{5}$

C.  $\frac{11}{5}$

D.  $\frac{7}{5}$

**Answer: D**

 [Watch Video Solution](#)

34. The circle with equation  $x^2 + y^2 = 1$  intersects the line  $y = 7x + 5$  at two distinct points A and B. Let C be the point at which the positive x-axis intersects the circle. The angle ACB is



A.  $\tan^{-1} \frac{4}{3}$

B.  $\cot^{-1}(-1)$

C.  $\tan^{-1} 1$

D.  $\cot^{-1} \frac{4}{3}$

**Answer: C**



**Watch Video Solution**

**35.** The abscissa of the two points A and B are the roots of the equation  $x^2 + 2ax - b^2 = 0$  and their ordinates are the roots of the equation  $x^2 + 2px - q^2 = 0$ . Find the equation of the circle with AB as diameter.

Also, find its radius.

A.  $\sqrt{a^2 + b^2 + p^2 + q^2}$

B.  $\sqrt{a^2 + p^2}$

C.  $\sqrt{b^2 + q^2}$

D.  $\sqrt{a^2 + b^2 + p^2 + 1}$

**Answer: A**



**Watch Video Solution**

**36.** Let  $C$  be the circle of radius unity centred at the origin. If two positive numbers  $x_1$  and  $x_2$  are such that the line passing through  $(x_1, -1)$  and  $(x_2, 1)$  is tangent to  $C$  then  $x_1 \cdot x_2$

A.  $x_1 x_2 = 1$

B.  $x_1 x_2 = -1$

C.  $x_1 + x_2 = 1$

D.  $4x_1 x_2 = 1$

**Answer: A**



**Watch Video Solution**

37. A circle bisects the circumference of the circle  $x^2 + y^2 + 2y - 3 = 0$  and touches the line  $x = y$  at the point  $(1, 1)$ . Its radius is

A.  $\frac{3}{\sqrt{2}}$

B.  $\frac{9}{\sqrt{2}}$

C.  $4\sqrt{2}$

D.  $3\sqrt{2}$

**Answer: B**



**Watch Video Solution**

38. The distance between the chords of contact of tangents to the circle  $x^2 + y^2 + 2gx + 2fy + c = 0$  from the origin & the point  $(g, f)$  is

A.  $\sqrt{g^2 + f^2}$

B.  $\frac{\sqrt{g^2 + f^2 - c}}{2}$

C.  $\frac{g^2 + f^2 - c}{2\sqrt{g^2 + f^2}}$

D.  $\frac{\sqrt{g^2 + f^2 + c}}{2\sqrt{g^2 + f^2}}$

**Answer: C**



**Watch Video Solution**

**39.** If the tangents AB and AQ are drawn from the point A(3, -1) to the circle  $x^2 + y^2 - 3x + 2y - 7 = 0$  and C is the centre of circle, then the area of quadrilateral APCQ is :

A. 9

B. 4

C. 2

D. non-existent

**Answer: D**



**Watch Video Solution**

40. Number of integral value(s) of  $k$  for which no tangent can be drawn from the point  $(k, k + 2)$  to the circle  $x^2 + y^2 = 4$  is :

A. 0

B. 1

C. 2

D. 3

**Answer: B**



**Watch Video Solution**

41. If the length of the normal for each point on a curve is equal to the radius vector, then the curve (a) is a circle passing through origin (b) is a circle having centre at origin and radius 0 (c) is a circle having centre on x-axis and touching y-axis (iv) is a circle having centre on y-axis and touching x-axis

A. is a circle passing through origin

- B. is a circle having centre at origin and radius  $> 0$
- C. is a circle having centre on x-axis and touching y-axis
- D. is a circle having centre on y-axis and touching x-axis

**Answer: B**



**Watch Video Solution**

**42.** A circle of radius unity is centered at the origin. Two particles start moving at the same time from the point  $(1, 0)$  and move around the circle in opposite directions. One of the particles moves anticlockwise with constant speed  $v$  and the other moves clockwise with constant speed  $3v$ . After leaving  $(1, 0)$ , the two particles meet first at a point  $P$ , and continue until they meet next at point  $Q$ . The coordinates of the point  $Q$  are

- A.  $(1, 0)$
- B.  $(0, 1)$

C. (0, -1)

D. (-1, 0)

**Answer: D**



**Watch Video Solution**

**43.** A variable circle is drawn to touch the x-axis at the origin. The locus of the pole of the straight line  $lx + my + n = 0$  w.r.t the variable circle has the equation:

A.  $x(my - n) - ly^2 = 0$

B.  $x(my + n) - ly^2 = 0$

C.  $x(my - n) + ly^2 = 0$

D. None of these

**Answer: A**



**Watch Video Solution**

44. The minimum length of the chord of the circle  $x^2 + y^2 + 2x + 2y - 7 = 0$  which is passing (1,0) is :

A. 2

B. 4

C.  $2\sqrt{2}$

D.  $\sqrt{5}$

**Answer: B**



**Watch Video Solution**

45. (C) 2 45. Three concentric circles of which the biggest is  $x^2 + y^2 = 1$ , have their radii in A.P If the line  $y = x + 1$  cuts all the circles in real and distinct points. The interval in which the common difference of the A.P will lie is:

A.  $\left(0, \frac{1}{4}\right)$



B.  $\left(0, \frac{1}{2\sqrt{2}}\right)$

C.  $\left(0, \frac{2 - \sqrt{2}}{4}\right)$

D. none

**Answer: C**



**Watch Video Solution**

**46.** The locus of the point of intersection of the tangent to the circle  $x^2 + y^2 = a^2$ , which include an angle of  $45^\circ$  is the curve  $(x^2 + y^2)^2 = \lambda(a^2)(x^2 + y^2 = a^2)$ . the value of  $\lambda$  is : (a) 2 (b) 4 (c) 8 (d) 16

A. 2

B. 4

C. 8

D. 16

**Answer: C**

[Watch Video Solution](#)

47. A circle touches the line  $y = x$  at point  $(4, 4)$  on it. The length of the chord on the line  $x + y = 0$  is  $6\sqrt{2}$ . Then one of the possible equation of the circle is

A.  $x^2 + y^2 + x - y + 30 = 0$

B.  $x^2 + y^2 + 2x - 18y + 32 = 0$

C.  $x^2 + y^2 + 2x + 18y + 32 = 0$

D.  $x^2 + y^2 - 2x - 22y + 32 = 0$

**Answer: B**

[Watch Video Solution](#)

48. Point on the circle  $x^2 + y^2 - 2x + 4y - 4 = 0$  which is nearest to the line  $y = 2x + 11$  is :

A.  $\left(1 - \frac{6}{\sqrt{5}}, -2 + \frac{3}{\sqrt{5}}\right)$

B.  $\left(1 + \frac{6}{\sqrt{5}}, 2 - \frac{3}{\sqrt{5}}\right)$

C.  $\left(1 - \frac{6}{\sqrt{5}}, -2 - \frac{3}{\sqrt{5}}\right)$

D. None of these

**Answer: A**



**Watch Video Solution**

**49.** A foot of the normal from the point  $(4, 3)$  to a circle is  $(2, 1)$  and a diameter of the circle has the equation  $2x - y - 2 = 0$ . Then the equation of the circle is:

A.  $x^2 + y^2 - 4y + 2 = 0$

B.  $x^2 + y^2 - 4y + 1 = 0$

C.  $x^2 + y^2 - 2x - 1 = 0$

D.  $x^2 + y^2 - 2x + 1 = 0$

**Answer: C**



**Watch Video Solution**

50. If  $\left(a, \frac{1}{a}\right), \left(b, \frac{1}{b}\right), \left(c, \frac{1}{c}\right), \left(d, \frac{1}{d}\right)$  are four distinct points on a circle of radius 4 units then,  $abcd$  is equal to:

A. 4

B.  $1/4$

C. 1

D. 16

**Answer: C**



**Watch Video Solution**

**Exercise 2 One Or More Than One Answer Is Are Correct**

1. Number of circle touching both the axes and the line  $x + y = 4$  is greater than or equal to :

A. 1

B. 2

C. 3

D. 4

**Answer: A::B::C::D**



**Watch Video Solution**

2. Which of the following is/are true ? The circles  $x^2 + y^2 - 6x + 6y + 9 = 0$  and  $x^2 + y^2 + 6x + 6y + 9 = 0$  are such that :

A. They do not intersect

B. They touch each other

C. Their exterior common tangents are parallel

D. Their interior common tangents are perpendicular

**Answer: A::C::D**



**Watch Video Solution**

3. Let  $\alpha$  be a variable parameter, then the length of the chord of the curve:  $(x - \sin^{-1} \alpha)(x - \cos^{-1} \alpha) + (y - \sin^{-1} \alpha)(y + \cos^{-1} \alpha) = 0$  along the line  $x = \frac{\pi}{4}$  can not be equal to

A.  $\frac{\pi}{3}$

B.  $\frac{\pi}{6}$

C.  $\frac{\pi}{4}$

D.  $\frac{\pi}{2}$

**Answer: A::B::C**



**Watch Video Solution**

4. The point  $(1, 4)$  are inside the circle  $S: x^2 + y^2 - 6x - 10y + k = 0$ .

What are the possible values of  $k$  if the circle  $S$  neither touches the axes nor cut them

A.  $p < 29$

B.  $p > 25$

C.  $p > 27$

D.  $p < 27$

**Answer: A::B**



**Watch Video Solution**

5. The equation of a circle  $C_1$  is  $x^2 + y^2 = 4$ . The locus of the intersection of orthogonal tangents to the circle is the curve  $C_2$  and the locus of the intersection of perpendicular tangents to the curve  $C_2$  is the curve  $C_3$ , Then

A.  $C_2$  is a circle

B.  $C_1, C_2$  are circles having different centres

C.  $C_1, C_2$  are circles having same centres

D. area enclosed between  $C_1$  and  $C_2$  is  $8\pi$

**Answer: A::C::D**



**Watch Video Solution**

6. If two distinct chords, drawn from the point  $(p, q)$  on the circle

$x^2 + y^2 = px + qy$  (where  $pq \neq q$ ) are bisected by the x-axis, then

$p^2 = q^2$  (b)  $p^2 = 8q^2$   $p^2 < 8q^2$  (d)  $p^2 > 8q^2$

A.  $p^2 = q^2$

B.  $p^2 > q^2$

C.  $p^2 < 8q^2$

D.  $p^2 > 8q^2$



**Answer: B::D**



**Watch Video Solution**

7.

If

$$a = \max\{(x + 2)^2 + (y - 3)^2\} \text{ and } b = \min\{(x + 2)^2 + (y - 3)^2\}$$

where  $x, y$  satisfying  $x^2 + y^2 + 8x - 10y - 40 = 0$  then :

A.  $a + b = 18$

B.  $a + b = 178$

C.  $a - b = 4\sqrt{2}$

D.  $a - b = 72\sqrt{2}$

**Answer: B::D**



**Watch Video Solution**

8. The locus of points of intersection of the tangents to  $x^2 + y^2 = a^2$  at the extremities of a chord of circle  $x^2 + y^2 = a^2$  which touches the circle  $x^2 + y^2 - 2ax = 0$  is/are :

A.  $y^2 = a(a - 2x)$

B.  $x^2 = a(a - 2y)$

C.  $x^2 + y^2 = (x - a)^2$

D.  $x^2 + y^2 = (y - a)^2$

**Answer: A::C**



**Watch Video Solution**

9. A circle passes through the points  $(-1,1)$ ,  $(0,6)$  and  $(5,5)$ . The point(s) on this circle, the tangents at which is/are parallel to the straight line joining the origin to its centre is/are:

A.  $(1, -5)$

B. (5, 1)

C. (-5, -1)

D. (-1, 5)

**Answer: B::D**



**Watch Video Solution**

**10.** A square is inscribed in the circle  $x^2 + y^2 - 2x + 4y - 93 = 0$  with its sides parallel to the coordinate axes. The coordinates of its vertices are

$(-6, -9), (-6, 5), (8, -9), (8, 5)$

$(-6, -9), (-6, -5), (8, -9), (8, 5)$

$(-6, -9), (-6, 5), (8, 9), (8, 5)$

$(-6, -9), (-6, 5), (8, -9), (8, -5)$

A. (8, 5)

B. (8, 9)

C. (-6, 5)

D. (-6, -9)

Answer: A::C



Watch Video Solution

### Exercise 3 Comprehension Type Problems

1. Let each of the circles,

$$S_1 = x^2 + y^2 + 4y - 1 = 0,$$

$$S_2 = x^2 + y^2 + 6x + y + 8 = 0,$$

$$S_3 = x^2 + y^2 - 4x - 4y - 37 = 0$$

touches the other two. Let  $P_1, P_2, P_3$  be the points of contact of  $S_1$  and  $S_2, S_2$  and  $S_3, S_3$  and  $S_1$  respectively and  $C_1, C_2, C_3$  be the centres of  $S_1, S_2, S_3$  respectively.

Q. The co-ordinates of  $P_1$  are :

A. (2, -1)

B. (2, 1)

C.  $(-2, 1)$

D.  $(-2, -1)$

**Answer: D**



**Watch Video Solution**

2. Let each of the circles,

$$S_1 = x^2 + y^2 + 4y - 1 = 0,$$

$$S_2 = x^2 + y^2 + 6x + y + 8 = 0,$$

$$S_3 = x^2 + y^2 - 4x - 4y - 37 = 0$$

touches the other two. Let  $P_1, P_2, P_3$  be the points of contact of  $S_1$  and  $S_2$ ,  $S_2$  and  $S_3$ ,  $S_3$  and  $S_1$  respectively and  $C_1, C_2, C_3$  be the centres of  $S_1, S_2, S_3$  respectively.

Q. The ratio  $\frac{\text{area}(\Delta P_1 P_2 P_3)}{\text{area}(\Delta C_1 C_2 C_3)}$  is equal to :

A. 3 : 2

B. 2 : 5

C.  $5:3$

D.  $2:3$

**Answer: B**



**Watch Video Solution**

**3.** Let each of the circles,

$$S_1 = x^2 + y^2 + 4y - 1 = 0,$$

$$S_2 = x^2 + y^2 + 6x + y + 8 = 0,$$

$$S_3 = x^2 + y^2 - 4x - 4y - 37 = 0$$

touches the other two. Let  $P_1, P_2, P_3$  be the points of contact of  $S_1$  and  $S_2$ ,  $S_2$  and  $S_3$ ,  $S_3$  and  $S_1$  respectively and  $C_1, C_2, C_3$  be the centres of  $S_1, S_2, S_3$  respectively.

Q.  $P_2$  and  $P_3$  are image of each other with respect to line :

A.  $y = x + 1$

B.  $y = -x$

C.  $y = x$

D.  $y = -x + 2$

**Answer: C**



**Watch Video Solution**

4. Let A(3, 7) and B(6, 5) are two points.  $C: x^2 + y^2 - 4x - 6y - 3 = 0$  is a circle.

Q. The chords in which the circle C cuts the members of the family S of circle passing through A and B are concurrent at:

A. (2, 3)

B.  $\left(2, \frac{23}{3}\right)$

C.  $\left(3, \frac{23}{2}\right)$

D. (3, 2)

**Answer: B**



**Watch Video Solution**

5. Let A(3, 7) and B(6, 5) are two points.  $C: x^2 + y^2 - 4x - 6y - 3 = 0$  is a circle.

Q. Equation of the member of the family of circles S that bisects the circumference of C is :

A.  $x^2 + y^2 - 5x - 1 = 0$

B.  $x^2 + y^2 - 5x + 6y - 1 = 0$

C.  $x^2 + y^2 - 5x - 6y - 1 = 0$

D.  $x^2 + y^2 + 5x - 6y - 1 = 0$

**Answer: C**



**View Text Solution**

6. Let A(3, 7) and B(6, 5) are two points.  $C: x^2 + y^2 - 4x - 6y - 3 = 0$  is a circle.

Q. If O is the origin and P is the center of C, then absolute value of



difference of the squares of the lengths of the tangents from A and B to the circle C is equal to :

A.  $(AB)^2$

B.  $(OP)^2$

C.  $|(AP)^2 - (BP)^2|$

D.  $(AP)^2 + (BP)^2$

**Answer: C**



**Watch Video Solution**

7. Let the diameter of a subset S of the plane be defined as the maximum of the distance between arbitrary pairs of points of S.

Q. Let  $S = \{(x, y) : (y - x) \leq 0, x + y \geq 0, x^2 + y^2 \leq 2\}$  then the diameter of S is :

A. 2

B. 4

C.  $\sqrt{2}$

D.  $2\sqrt{2}$

**Answer: A**



**Watch Video Solution**

**8.** Let the diameter of a subset  $S$  of the plane be defined as the maximum of the distance between arbitrary pairs of points of  $S$ .

Q.

Let

$$S = \left\{ (x, y) : (\sqrt{5} - 1)x - \sqrt{10 + 2\sqrt{5}}y \geq 0, (\sqrt{5} - 1)x + \sqrt{10 + 12\sqrt{5}}y \geq 0 \right\}$$

then the diameter of  $S$  is :

A.  $\frac{3}{2}(\sqrt{5} - 1)$

B.  $3(\sqrt{5} - 1)$

C.  $3\sqrt{2}$

D. 3

**Answer: D**



**Watch Video Solution**

9. Let  $L_1$ ,  $L_2$  and  $L_3$  be the lengths of tangents drawn from a point P to circles  $x^2 + y^2 = 4$ ,  $x^2 + y^2 - 4x = 0$  and  $x^2 + y^2 - 4y = 0$  respectively. If  $L_1^4 = L_2^2 L_3^2 + 16$  then the locus of P are the curves,  $C_1$  (a straight line) and  $C_2$  (a circle).

Q. Circum centre of the triangle formed by  $C_1$  and two other lines which are at angle of  $45^\circ$  with  $C_1$  and tangent of  $C_2$  is :

A. (1, 1)

B. (0, 0)

C. (-1, -1)

D. (2, 2)

**Answer: B**



**View Text Solution**

10. Let  $L_1$ ,  $L_2$  and  $L_3$  be the lengths of tangents drawn from a point P to circles  $x^2 + y^2 = 4$ ,  $x^2 + y^2 - 4x = 0$  and  $x^2 + y^2 - 4y = 0$  respectively. If  $L_1^4 = L_2^2 L_3^2 + 16$  then the locus of P are the curves,  $C_1$  (a straight line) and  $C_2$  (a circle).

Q. If  $S_1$ ,  $S_2$  and  $S_3$  are three circles congruent to  $C_2$  and touch both  $C_1$  and  $C_2$ , then the area of triangle formed by joining centres of the circles  $S_1$ ,  $S_2$  and  $S_3$  is (in square units)

- A. 2
- B. 4
- C. 8
- D. 16

**Answer: C**



**View Text Solution**

1. Tangents are drawn to circle  $x^2 + y^2 = 1$  at its intersection points (distinct) with the circle  $x^2 + y^2 + (\lambda - 3)x + (2\lambda + 2)y + 2 = 0$ . The locus of intersection of tangents is a straight line, then the slope of that straight line is .



Watch Video Solution

2. The radical centre of the three circles is at the origin. The equations of the two of the circles are  $x^2 + y^2 = 1$  and  $x^2 + y^2 + 4x + 4y - 1 = 0$ . If the third circle passes through the points  $(1, 1)$  and  $(-2, 1)$ , and its radius can be expressed in the form of  $\frac{p}{q}$ , where  $p$  and  $q$  are relatively prime positive integers. Find the value of  $(p + q)$ .



Watch Video Solution

3. Let  $S = \{(x, y) \mid x, y \in R, x^2 + y^2 - 10x + 16 = 0\}$ . The largest value of  $\frac{y}{x}$  can be put in the form of  $\frac{m}{n}$  where  $m, n$  are relatively prime

natural numbers, then  $m^2 + n^2 =$



**Watch Video Solution**

4. In the above problem, the complete range of the expression  $x^2 + y^2 - 26x + 12y + 210$  is  $[a, b]$ , then  $b - 2a =$



**View Text Solution**

5. If the line  $y=2-x$  is tangent to the circle  $S$  at the point  $P(1,1)$  and circle  $S$  is orthogonal to the circle  $x^2 + y^2 + 2x + 2y - 2 = 0$  then find the length of tangent drawn from the point  $(2,2)$  to the circle  $S$



**Watch Video Solution**

6. Two circles having radii  $r_1$  and  $r_2$  passing through vertex  $A$  of triangle  $ABC$ . One of the circle touches the side  $BC$  at  $B$  and the other circle touches the side  $BC$  at  $C$ . If  $a = 5\text{cm}$  and  $A = 30^\circ$  find  $\sqrt{r_1 r_2}$

[Watch Video Solution](#)

7. A circle  $S$  of radius ' $a$ ' is the director circle of another circle  $S_1$ .  $S_2$  is the director circle of  $S_1$  and so on. If the sum of radii of  $S, S_1, S_2, S_3, \dots$  circles is '2' and  $a = (k - \sqrt{k})$ , then the value of  $k$  is .....

[View Text Solution](#)

8. If  $r_1$  and  $r_2$  be the maximum and minimum radius of the circle which pass through the point  $(4, 3)$  and touch the circle  $x^2 + y^2 = 49$ , then  $\frac{r_1}{r_2}$  is .....

[Watch Video Solution](#)

9. Let  $C$  be the circle  $x^2 + y^2 - 4x - 4y - 1 = 0$ . The number of points common to  $C$  and the sides of the rectangle determined by the lines  $x = 2, x = 5, y = -1$  and  $y = 5$  equal to

[Watch Video Solution](#)

10. Two congruent circles with centered at (2, 3) and (5, 6) which intersect at right angles, have radius equal to  $2\sqrt{3}$  (b) 3 (c) 4 (d) none of these

[Watch Video Solution](#)

11. The sum of abscissa and ordinate of a point on the circle  $x^2 + y^2 - 4x + 2y - 20 = 0$  which is nearest to  $\left(2, \frac{3}{2}\right)$  is :

[Watch Video Solution](#)

12. AB is any chord of the circle  $x^2 + y^2 - 6x - 8y - 11 = 0$  which subtends an angle  $\frac{\pi}{2}$  at (1, 2). If locus of midpoint of AB is a circle  $x^2 + y^2 - 2ax - 2by - c = 0$ ; then find the value of  $(a + b + c)$ .

[Watch Video Solution](#)



13. If circles  $x^2 + y^2 = c$  with radius  $\sqrt{3}$  and  $x^2 + y^2 + ax + by + c = 0$  with radius  $\sqrt{6}$  intersect at two points A and B. If length of  $AB = \sqrt{l}$ . Find  $l$ .



**Watch Video Solution**