



MATHS

BOOKS - VIKAS GUPTA MATHS (HINGLISH)

COMPOUND ANGLES

Exercise 1 Single Choice Problems

1. $\left(\cos^4 \frac{\pi}{24} - \sin^4 \frac{\pi}{24}\right)$ equals :

A. $\frac{1}{\sqrt{2}}$

B. $\frac{\sqrt{6} - \sqrt{2}}{4}$

C. $\frac{\sqrt{6} + \sqrt{2}}{4}$

D. $\frac{\sqrt{3} + 1}{2}$

Answer: C



Watch Video Solution

2. If $a \sin x + b \cos(c + x) + b \cos(c - x) = \alpha$, $\alpha > a$, then the minimum value of $|\cos c|$ is :

A. $\sqrt{\frac{\alpha^2 - a^2}{b^2}}$

B. $\sqrt{\frac{\alpha^2 - a^2}{2b^2}}$

C. $\sqrt{\frac{\alpha^2 - a^2}{3b^2}}$

D. $\sqrt{\frac{\alpha^2 - a^2}{4b^2}}$

Answer: D



Watch Video Solution

3. If all values of $x \in (a, b)$ satisfy the inequality $\tan x \tan 3x < -1$, $x \in \left(0, \frac{\pi}{2}\right)$, then the maximum value (b, -a) is :

A. $\frac{\pi}{12}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{6}$

D. $\frac{\pi}{4}$

Answer: A



Watch Video Solution

4. $\sum_{r=1}^8 \tan(rA)\tan((r+1)A)$ where $A = 36^\circ$ is :

A. $-10 - \tan A$

B. $-10 + \tan A$

C. -10

D. -9

Answer: C



Watch Video Solution

5. Let $f(x) = 2 \operatorname{cosec} 2x + \sec x + \operatorname{cosec} x$, then the minimum value of $f(x)$ for $x \in \left(0, \frac{\pi}{2}\right)$ is

A. $\frac{1}{\sqrt{2} - 1}$

B. $\frac{2}{\sqrt{2} - 1}$

C. $\frac{1}{\sqrt{2} + 1}$

D. $\frac{2}{\sqrt{2} + 1}$

Answer: B



Watch Video Solution

6. Find the exact value of $\operatorname{cosec} 10^\circ + \operatorname{cosec} 50^\circ - \operatorname{cosec} 70^\circ$

A. 4

B. 5

C. 6

D. 8

Answer: C



Watch Video Solution

7. If $u = \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} + \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta}$, then the difference between maximum and minimum values of u^2 is

A. $2(a^2 + b^2)$

B. $2\sqrt{a^2 + b^2}$

C. $(a + b)^2$

D. $(a - b)^2$

Answer: D



Watch Video Solution

8. If $u_n = \sin(n\theta)\sec^n \theta$, $v_n = \cos(n\theta)\sec^n \theta$, $n \in N$, $n \neq 1$, then

$$\frac{v_n - v_{n-1}}{v_{n-1}} + \frac{1}{n} \left(\frac{u_n}{v_n} \right) =$$

A. $-\cos \theta + \frac{1}{n} \tan(n\theta)$

B. $\cot \theta + \frac{1}{n} \tan(n\theta)$

C. $\tan \theta + \frac{1}{n} \tan(n\theta)$

D. $-\tan \theta + \frac{\tan(n\theta)}{n}$

Answer: D

 [Watch Video Solution](#)

9. If $\alpha \cos^3 \theta + \beta \cos^4 \theta = 16 \cos^6 \theta + 9 \cos^2 \theta$ is an identity then-

A. $a = 1, b = 24$

B. $a = 3, b = 24$

C. $a = 4, b = 2$

D. $a = 7, b = 18$

Answer: A

 [Watch Video Solution](#)

10. Maximum value of $\cos x(\sin x + \cos x)$ is equal to :

A. $\sqrt{2}$

B. 2

C. $\frac{\sqrt{2} + 1}{2}$

D. $\sqrt{2} + 1$

Answer: C



Watch Video Solution

11. If $\frac{\sin A}{\sin B} = \frac{\sqrt{3}}{2}$ and $\frac{\cos A}{\cos B} = \frac{\sqrt{5}}{2}$, $0 < A, B < \frac{\pi}{2}$ then

$\tan A + \tan B$ is equal to :

A. $\sqrt{\frac{3}{5}}$

B. $\sqrt{\frac{5}{3}}$

C. $\frac{\sqrt{3} + \sqrt{5}}{\sqrt{5}}$

D. $\frac{\sqrt{3} + \sqrt{5}}{\sqrt{3}}$

Answer: C



Watch Video Solution

12. Single Choice Questions: 1. Let $0 \leq \alpha, \beta, \gamma, \delta, \leq \pi$ where β and γ are not complementary such that $2 \cos \alpha + 6 \cos \beta + 7 \cos \gamma + 9 \cos \delta = 0$ and $2 \sin \alpha - 6 \sin \beta + 7 \sin \gamma - 9 \sin \delta = 0$ if $\frac{\cos(\alpha + \delta)}{\cos(\beta + \gamma)} = \frac{m}{n}$ when m and n are relatively prime positive numbers, then the value of $(m + n)$ is equal to (A)11 (B)10 (C)9 (D)7

A. 11

B. 10

C. 9

D. 2

Answer: B



Watch Video Solution

13. If $-\pi < \theta < -\frac{\pi}{2}$, then $\left| \sqrt{\frac{1 - \sin \theta}{1 + \sin \theta}} + \sqrt{\frac{1 + \sin \theta}{1 - \sin \theta}} \right|$ is equal to :

- A. $2 \sec \theta$
- B. $-2 \sec \theta$
- C. $2 \sec \frac{\theta}{2}$
- D. $-\sec \frac{\theta}{2}$

Answer: B



Watch Video Solution

14. If $A = \sum_{r=1}^3 \cos \frac{2r\pi}{7}$ and $B = \sum_{r=1}^3 \cos \frac{2^r\pi}{7}$, then :

- A. $A + B = 0$
- B. $2A + B = 0$
- C. $A + 2B = 0$


D. $A = B$

Answer: D



Watch Video Solution

15. In a $\triangle PQR$ (as shown in figure) if $x : y : z = 2 : 3 : 6$, then the value of $\angle QPR$ is :



A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{2}$

Answer: B



Watch Video Solution

16. If $A = \sum_{r=1}^3 \cos \frac{2r\pi}{7}$ and $B = \sum_{r=1}^3 \cos \frac{2^r\pi}{7}$ then :

A. $A + B = 0$

B. $2A + B = 0$

C. $A + 2B = 0$

D. $A - B = 0$

Answer: D



Watch Video Solution

17. Let $f(x) = \sin x + 2 \cos^2 x$, $x \in \left[\frac{\pi}{6}, \frac{2\pi}{3} \right]$, then maximum value of $f(x)$ is

A. 1

B. $\frac{3}{2}$

C. 2

D. $\frac{5}{2}$

Answer: C



[Watch Video Solution](#)

18. In triangle ABC , $\angle C = \frac{2\pi}{3}$ then the value of $\cos^2 A + \cos^2 B - \cos A \cdot \cos B$ is equal

A. $\frac{3}{4}$

B. $\frac{3}{2}$

C. $\frac{1}{2}$

D. $\frac{1}{4}$

Answer: A



Watch Video Solution

19. The number of solutions of the equation

$$4 \sin^2 x + \tan^2 x + \cot^2 x + \operatorname{cosec}^2 x = 6 \text{ in } [0, 2\pi] :$$

A. 1

B. 2

C. 3

D. 4

Answer: D



Watch Video Solution

20. If $\sin A$, $\cos A$ and $\tan A$ are in G.P, the $\cos^3 A + \cos^2 A$ is equal to :

- A. 1
- B. 2
- C. 4
- D. none

Answer: A



[Watch Video Solution](#)

21. find the range of function $f(x) = \sin\left(x + \frac{\pi}{6}\right) + \cos\left(x - \frac{\pi}{6}\right)$

- A. $[-\sqrt{2}, \sqrt{2}]$
- B. $[-\sqrt{2}(\sqrt{3} + 1), \sqrt{2}(\sqrt{3} + 1)]$
- C. $\left[-\frac{\sqrt{3} + 1}{\sqrt{2}}, \frac{\sqrt{3} + 1}{\sqrt{2}}\right]$
- D. $\left[-\frac{\sqrt{3} - 1}{\sqrt{2}}, \frac{\sqrt{3} - 1}{\sqrt{2}}\right]$

Answer: C



Watch Video Solution

22. The value of

$\tan(\log_2 6) \cdot \tan(\log_2 3) \cdot \tan 1$ is always equal to :

A. $\tan(\log_2 6) + \tan(\log_2 3) + \tan 1$

B. $\tan(\log_2 6) - \tan(\log_2 3) - \tan 1$

C. $\tan(\log_2 6) - \tan(\log_2 3) + \tan 1$

D. $\tan(\log_2 6) + \tan(\log_2 3) - \tan 1$

Answer: B



Watch Video Solution

23. In $\triangle ABC$, $a = 3$, $b = 4$ and $c = 5$, then value of $\sin A + \sin 2B + \sin 3C$ is

A. $\frac{24}{25}$

B. $\frac{14}{25}$

C. $\frac{64}{25}$

D. none

Answer: B



Watch Video Solution

24.

If

$$A + B + C = 180^\circ, \text{ then } \frac{\cos A \cos C + \cos(A + B)\cos(B + C)}{\cos A \sin C - \sin(A + B)\cos(B + C)}$$

simplifies to :

A. $-\cot C$

B. 0

C. $\tan C$

D. $\cot C$

Answer: D



Watch Video Solution

25. If $\alpha + \gamma = 2\beta$ then the expression $\frac{\sin \alpha - \sin \gamma}{\cos \gamma - \cos \alpha}$ simplifies to:

- A. $\tan \beta$
- B. $-\tan \beta$
- C. $\cot \beta$
- D. $-\cot \beta$

Answer: C



Watch Video Solution

26. The product $\left(\cos \frac{x}{2}\right) \cdot \left(\cos \frac{x}{4}\right) \cdot \left(\cos \frac{x}{8}\right) \cdot \dots \cdot \left(\cos \frac{x}{256}\right)$

is equal to :

A. $\frac{\sin x}{128\sin\frac{x}{256}}$

B. $\frac{\sin x}{256\sin\frac{x}{256}}$

C. $\frac{\sin x}{128\sin\frac{x}{128}}$

D. $\frac{\sin x}{512\sin\frac{x}{512}}$

Answer: B



Watch Video Solution

27. The value of the expression

$$\frac{\sin 7\alpha + 6 \sin 5\alpha + 17 \sin 3\alpha + 12 \sin \alpha}{\sin 6\alpha + 5 \sin 4\alpha + 12 \sin 2\alpha}, \text{ where } \alpha = \frac{\pi}{5} \text{ is equal to :}$$

A. $\frac{\sqrt{5} - 1}{4}$

B. $\frac{\sqrt{5} + 1}{4}$

C. $\frac{\sqrt{5} + 1}{2}$

D. $\frac{\sqrt{5} - 1}{2}$

Answer: C



Watch Video Solution

28. In a triangle ABC if $\sum \tan^2 A = \sum \tan A \tan B$, then largest angle of the triangle in radian will be :

A. $\frac{2\pi}{3}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{2}$

D. $\frac{3\pi}{4}$

Answer: B



Watch Video Solution

29. Which one of the following values is not the solution of the equation

$$\log_{|\sin x|} (|\cos x|) + \log_{|\cos x|} (|\sin x|) = 2$$

A. $\frac{7\pi}{4}$

B. $\frac{11\pi}{4}$

C. $\frac{3\pi}{4}$

D. $\frac{3\pi}{8}$

Answer: D

 [Watch Video Solution](#)

30. If $f(x) = \sin^6 x + \cos^6 x$, then range of $f(x)$ is $\left[\frac{1}{4}, 1\right]$ (b) $\left[\frac{1}{4}, \frac{3}{4}\right]$

(c) $\left[\frac{3}{4}, 1\right]$ (d) none of these

A. $\left[\frac{1}{4}, 1\right]$

B. $\left[\frac{1}{4}, \frac{3}{4}\right]$

C. $\left[\frac{3}{4}, 1\right]$

D. $[1, 2]$

Answer: A

 [Watch Video Solution](#)

31. If $\frac{2 \sin \alpha}{\{1 + \cos \alpha + \sin \alpha\}} = y$, then $\frac{\{1 - \cos \alpha + \sin \alpha\}}{1 + \sin \alpha} =$

A. $\frac{1}{y}$

B. y

C. $1 - y$

D. $1 + y$

Answer: B



Watch Video Solution

32. If $\frac{\tan^3 A}{1 + \tan^2 A} + \frac{\cot^3 A}{1 + \cot^2 A} = p \sec A \operatorname{cosec} A + q \sin A \cos A$, then :

A. $p = 2, q = 1$

B. $p = 1, q = 2$

C. $p = 1, q = -2$

D. $p = 2, q = -1$

Answer: C



Watch Video Solution

33. If θ lies in the second quadrant. Then the value of

$\sqrt{\frac{1 - \sin \theta}{1 + \sin \theta}} + \sqrt{\frac{1 + \sin \theta}{1 - \sin \theta}}$ is equal to :

- A. $2 \sec \theta$
- B. $-2 \sec \theta$
- C. $2 \operatorname{cosec} \theta$
- D. 2

Answer: B



Watch Video Solution

34. The minimum value of $(\sin \theta + \operatorname{cosec} \theta)^2 + (\cos \theta + \sec \theta)^2 =$

A. 7

B. 8

C. 9

D. none of these

Answer: C



[Watch Video Solution](#)

35. If $\log_3 \sin x - \log_3 \cos x - \log_3(1 - \tan x) - \log_3(1 + \tan x) = -1$,

then $\tan 2x$ is equal to (wherever defined)

A. -2

B. $\frac{3}{2}$

C. $\frac{2}{3}$

D. 6

Answer: C



Watch Video Solution

36. If $\sin \theta + \cos \theta = 2$, then the value of $\sin^8 \theta + \cos^8 \theta$ is equal to

A. 2

B. 2^4

C. 2^8

D. more than 2^8

Answer: A



Watch Video Solution

37. If $\tan^3 \theta + \cot^3 \theta = 52$, then the value of $\tan^2 \theta + \cot^2 \theta$ is equal to :

A. 14

B. 15

C. 16

D. 17

Answer: A



[Watch Video Solution](#)

38. The maximum value of $(\log)_{20}(3s \in x - 4 \cos x + 15)$ – a. 1 b. 2 c. 3

d. 4

A. 1

B. 2

C. 3

D. 4

Answer: A



[Watch Video Solution](#)

39. $x^2 + y^2 = 9$ & $4a^2 + 9b^2 = 16$, then maximum value of $4a^2x^2 + 9b^2y^2 - 12abxy$ is -

- A. 81
- B. 100
- C. 121
- D. 144

Answer: D

 [Watch Video Solution](#)

40. If $A = \sqrt{\sin 2 - \sin \sqrt{3}}$, $B = \sqrt{\cos 2 - \cos \sqrt{3}}$, then which of the following statement is true ?

- A. A and B both are real numbers and $A > B$
- B. A and B both are real numbers and $A < B$

C. Exactly one of A and B is not real number

D. Both A and B are not real numbers

Answer: D



[Watch Video Solution](#)

41. The number of real values of x such that

$$(2^x + 2^{-x} - 2 \cos x)(3^{x+\pi} + 3^{-x-\pi} + 2 \cos x)(5^{\pi-x} + 5^{x-\pi} - 2 \cos x) =$$

is :

A. 1

B. 2

C. 3

D. infinite

Answer: B



[Watch Video Solution](#)

42. The equation $e^{\sin x} - e^{-\sin x} - 4 = 0$ has (A) infinite number of real roots (B) no real roots (C) exactly one real root (D) exactly four real roots

A. infinite number of real roots

B. no real roots

C. exactly one real root

D. exactly four real roots

Answer: B



[Watch Video Solution](#)

43. If $\pi < \alpha < \frac{3\pi}{2}$ then the expression

$\sqrt{4\sin^4 \alpha + \sin^2 2\alpha} + 4\cos^2\left(\frac{\pi}{4} - \frac{\alpha}{2}\right)$ is equal to (A) $2 + 4\sin \alpha$ (B) $2 - 4\cos \alpha$ (C) 2 (D) $2 - 4\sin \alpha$

A. $2 + 4\sin \alpha$

B. $2 - 4 \cos \alpha$

C. 2

D. $2 - 4 \sin \alpha$

Answer: C

 [Watch Video Solution](#)

44. $\left(\cos\left(\frac{\pi}{12}\right) - \sin\left(\frac{\pi}{12}\right) \right) \left(\tan\left(\frac{\pi}{12}\right) + \cot\left(\frac{\pi}{12}\right) \right) =$

A. $\frac{1}{\sqrt{2}}$

B. $4\sqrt{2}$

C. $\sqrt{2}$

D. $2\sqrt{2}$

Answer: D

 [Watch Video Solution](#)

45. $\tan(100^\circ) + \tan(125^\circ) + \tan(100^\circ)\tan(125^\circ) =$

A. 0

B. $\frac{1}{2}$

C. -1

D. 1

Answer: D



Watch Video Solution

46. If $\sin x + \sin^2 = 1$, then $\cos^8 x + 2\cos^6 x + \cos^4 x =$

A. 2

B. 1

C. 3

D. $\frac{1}{2}$

Answer: B



Watch Video Solution

47. maximum value of $\log_5(3x + 4y)$, if $x^2 + y^2 = 25$ is

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

48. The number of values of θ between $-\pi$ and $\frac{3\pi}{2}$ that satisfies the equation $5 \cos 2\theta + 2 \cos^2 \frac{\theta}{2} + 1 = 0$ is :

A. 3

B. 4

C. 5

D. 6

Answer: C



Watch Video Solution

49.

Given

that

$\sin \beta = \frac{4}{5}, 0 < \beta < \pi$ and $\tan \beta > 0$, then $((3 \sin(\alpha + \beta) - 4 \cos(\alpha +$

is equal to :

A. 2

B. 3

C. 4

D. 5

Answer: D



Watch Video Solution

50. The maximum value of the function $f(x) = \sin\left(x + \frac{\pi}{6}\right) + \cos\left(x + \frac{\pi}{6}\right)$ in the interval $\left(0, \frac{\pi}{2}\right)$ occurs at $\frac{\pi}{12}$ (b) $\frac{\pi}{6}$ (c) $\frac{\pi}{4}$ (d) $\frac{\pi}{3}$

A. $\frac{\pi}{12}$

B. $\frac{\pi}{6}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{2}$

Answer: A



Watch Video Solution

51. The equation $\sin x(\sin x + \cos x) = k$ has real solutions, where k is a real number. Then

A. $1 - \sqrt{2} \leq a \leq 1 + \sqrt{2}$

B. $2 - \sqrt{3} \leq a \leq 2 + \sqrt{3}$

C. $0 \leq a \leq 2 + \sqrt{3}$

D. $\frac{1 - \sqrt{2}}{2} \leq a \leq \frac{1 + \sqrt{2}}{2}$

Answer: D



[Watch Video Solution](#)

52. The value of $\cos 12^\circ \cos 24^\circ \cos 36^\circ \cos 48^\circ \cos 72^\circ \cos 84^\circ$, is

A. $\frac{1}{64}$

B. $\frac{1}{128}$

C. $\frac{1}{256}$

D. $\frac{1}{512}$

Answer: B



[Watch Video Solution](#)

53. The ratio of the maximum value to minimum value of $2 \cos^2 \theta + \cos \theta + 1$ is :

A. 32 : 7

B. 32 : 9

C. 4 : 1

D. 2 : 1

Answer: A



[Watch Video Solution](#)

54. If all values of $x \in (a, b)$ satisfy the inequality $\tan x \tan 3x < -1$, $x \in \left(0, \frac{\pi}{2}\right)$, then the maximum value of $(b - a)$ is :

A. $\frac{\pi}{12}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{6}$

D. $\frac{\pi}{4}$

Answer: A



Watch Video Solution

55. If a regular polygon of n sides has circum radius R and inradius r then each side of polygon is:

A. $(R + r)\tan\left(\frac{\pi}{2n}\right)$

B. $2(R + r)\tan\left(\frac{\pi}{2n}\right)$

C. $(R + r)\sin\left(\frac{\pi}{2n}\right)$

D. $2(R + r)\cot\left(\frac{\pi}{2n}\right)$

Answer: B

 Watch Video Solution

56. The value of $\cos 12^\circ + \cos 84^\circ + \cos 156^\circ + \cos 132^\circ$ is $\frac{1}{2} 1 - \frac{1}{2} \frac{1}{8}$

A. $\frac{1}{8}$

B. $-\frac{1}{2}$

C. 1

D. $\frac{1}{2}$

Answer: B

 Watch Video Solution

57. $\frac{\sin \theta}{\cos(3\theta)} + \frac{\sin(3\theta)}{\cos(9\theta)} + \frac{\sin(9\theta)}{\cos(27\theta)} + \frac{\sin(27\theta)}{\cos(81\theta)} =$

A. $\frac{\sin(81\theta)}{2 \cos(80\theta) \cos \theta}$

B. $\frac{\sin(80\theta)}{2 \cos(81\theta) \cos \theta}$

C. $\frac{\sin(81\theta)}{\cos(80\theta) \cos \theta}$

D. $\frac{\sin(80\theta)}{2 \cos(81\theta) \cos \theta}$

Answer: B



Watch Video Solution

58. The value of $\left(\sin \frac{\pi}{9}\right) \left(4 + \sec \frac{\pi}{9}\right)$ is :

A. $\frac{1}{2}$

B. $\sqrt{2}$

C. 1

D. $\sqrt{3}$

Answer: D



Watch Video Solution

59. If $\frac{dy}{dx} = \sin\left(\frac{x\pi}{2}\right) \cos(x\pi)$, then y is strictly increasing in :

A. (3, 4)

B. $\left(\frac{5}{2}, \frac{7}{2}\right)$

C. (2, 3)

D. $\left(\frac{1}{2}, \frac{3}{2}\right)$

Answer: B



Watch Video Solution

60. Smallest positive value of θ satisfying

$$8 \sin \theta \cos 2\theta \sin 3\theta \cos 4\theta = \cos 6\theta$$
 is

A. $\frac{\pi}{18}$

B. $\frac{\pi}{22}$

C. $\frac{\pi}{24}$

D. none of these

Answer: A

 [Watch Video Solution](#)

61. If an angle A of a triangle ABC is given by $3 \tan A + 1 = 0$, then $\sin A$ and $\cos A$ are the roots of the equation

A. $10x^2 - 2\sqrt{10}x + 3 = 0$

B. $10x^2 - 2\sqrt{10}x - 3 = 0$

C. $10x^2 + 2\sqrt{10}x + 3 = 0$

D. $10x^2 + 2\sqrt{10}x - 3 = 0$

Answer: D

 [Watch Video Solution](#)

62. If θ is an acute angle and $\tan \theta = \frac{1}{\sqrt{7}}$, then the value of $\frac{\cos^2 \theta - \sec^2 \theta}{\cos^2 \theta + \sec^2 \theta}$ is

A. $3/4$

B. $1/2$

C. 2

D. $5/4$

Answer: A

 [Watch Video Solution](#)

63. If $2 \cos \theta + \sin \theta = 1$ ($\theta \neq \frac{\pi}{2}$) then $7 \cos \theta + 6 \sin \theta$ is equal to

A. 1 or 2

B. 2 or 3

C. 2 or 4

D. 2 or 6

Answer: D

 [Watch Video Solution](#)

64. If $\sin \theta + \cos \theta = 2$, then the value of $\sin^8 \theta + \cos^8 \theta$ is equal to

A. 2

B. 2^4

C. 2^8

D. more than 2^8

Answer: A



[Watch Video Solution](#)

65. If $\tan^3 \theta + \cot^3 \theta = 52$, then the value of $\tan^2 \theta + \cot^2 \theta$ is equal to :

A. 14

B. 15

C. 16

D. 17

Answer: A



Watch Video Solution

66. If ABCD is a cyclic quadrilateral such that $12 \tan A - 5 = 0$ and $5 \cos B + 3 = 0$ then $\tan C + \tan D$ is equal to :

A. $\frac{21}{12}$

B. $\frac{11}{12}$

C. $-\frac{11}{12}$

D. $-\frac{21}{12}$

Answer: B



Watch Video Solution

67. If $\frac{\pi}{2} < \theta < \frac{3\pi}{2}$ then $\sqrt{\tan^2 \theta - \sin^2 \theta}$ is equal to :

A. $\tan \theta \sin \theta$

B. $-\tan \theta \sin \theta$

C. $\tan \theta - \sin \theta$

D. $\sin \theta - \tan \theta$

Answer: B

 [Watch Video Solution](#)

68. The value of $\frac{\sin 10^\circ + \sin 20^\circ}{\cos 10^\circ + \cos 20^\circ}$ equals

A. $2 + \sqrt{3}$

B. $\sqrt{2} - 1$

C. $2 - \sqrt{3}$

D. $\sqrt{2} + 1$

Answer: C

 [Watch Video Solution](#)

69. The value of the expression $\sin^6 \theta + \cos^6 \theta + 3 \sin^2 \theta \cdot \cos^2 \theta$ equals

A. 0

B. 1

C. 2

D. 3

Answer: B



[Watch Video Solution](#)

70. $\frac{\sin x + \cos x}{\sin x - \cos x} - \frac{\sec^2 x + 2}{\tan^2 x - 1} =$, where $x \in \left(0, \frac{\pi}{2}\right)$

A. $\frac{1}{\tan x + 1}$

B. $\frac{2}{1 + \tan x}$

C. $\frac{2}{1 + \cot x}$

D. $\frac{2}{1 - \tan x}$

Answer: B



Watch Video Solution

71. If $\frac{\cot \alpha + \cot(270^\circ + \alpha)}{\cot \alpha - \cot(270^\circ + \alpha)} - 2 \cos(135^\circ + \alpha) \cos(315^\circ - \alpha) = \lambda$,
where $\alpha \in \left(0, \frac{\pi}{2}\right)$, then $\lambda =$

A. 0

B. 1

C. 2

D. 4

Answer: C



Watch Video Solution

72. The expression $\frac{\sin \alpha + \cos \alpha}{\cos \alpha - \sin \alpha} \tan\left(\frac{\pi}{4} + \alpha\right) + 1, \alpha \in \left(-\frac{\pi}{4}, \frac{\pi}{4}\right)$ simplifies to :

A. $\operatorname{cosec}^2\left(\frac{\pi}{4} - \alpha\right)$

B. $\sec^2\left(\frac{\pi}{4} - \alpha\right)$

C. $\tan^2\left(\frac{\pi}{4} - \alpha\right)$

D. $\cot^2\left(\frac{\pi}{4} - \alpha\right)$

Answer: A



Watch Video Solution

73. The value of expression $\frac{\tan \alpha + \sin \alpha}{2\cos^2 \frac{\alpha}{2}}$ for $\alpha = \frac{\pi}{4}$ is :

A. 4

B. 3

C. 2

D. 1

Answer: D



Watch Video Solution

74. $\cos 2\alpha - \cos 3\alpha - \cos 4\alpha + \cos 5\alpha$ simplifies to :

A. $-4\sin \frac{\alpha}{2} \sin \alpha \cos \frac{7\alpha}{2}$

B. $4\sin \frac{\alpha}{2} \sin \alpha \cos \frac{7\alpha}{2}$

C. $-4\sin \frac{\alpha}{2} \sin \frac{7\alpha}{2} \cos \alpha$

D. $-4\sin \alpha \cos \frac{\alpha}{2} \sin \frac{7\alpha}{2}$

Answer: A



Watch Video Solution

75. If $\tan \gamma = \sec \alpha \sec \beta + \tan \alpha \tan \beta$, then $\cos 2\gamma$ is necessarily (A) ≥ 0 (B) ≤ 0 (C) < 0 (D) > 0

A. -1

B. $\frac{1}{2}$

C. $-\frac{1}{2}$

D. 0

Answer: D



[Watch Video Solution](#)

76. If $\cos ecx = \frac{2}{\sqrt{3}}$ and $\cot x = -\frac{1}{\sqrt{3}}$ for $x \in [0, 2\pi]$ then

$\cos x + \cos 2x + \cos 3x + \dots + \cos 100x$ is

A. $\frac{1}{2}$

B. $-\frac{1}{2}$

C. $-\frac{\sqrt{3}}{2}$

D. $\frac{\sqrt{3}}{2}$

Answer: B



Watch Video Solution

77. The value of $\sum_{r=0}^{10} \frac{\cos^3(r\pi)}{3}$ is equal to $\frac{1}{4}$ (b) $\frac{1}{8}$ (c) $-\frac{1}{4}$ (d) $-\frac{1}{8}$

A. $-\frac{7}{8}$

B. $-\frac{9}{8}$

C. $-\frac{3}{8}$

D. $-\frac{1}{8}$

Answer: D



Watch Video Solution

78. Write the value of the expression $\frac{1 - 4 \sin 10^\circ \sin 70^\circ}{2 \sin 10^\circ}$

A. 1

B. 2

C. $\sqrt{3}$

D. $\frac{\sqrt{3}}{2}$

Answer: A



Watch Video Solution

79. If $x, y \in \mathbb{R}$ and satisfy $(x + 5)^2 + (y - 12)^2 = 14^2$ then the minimum value of $x^2 + y^2$ is

A. 2

B. 1

C. $\sqrt{3}$

D. $\sqrt{2}$

Answer: B

 [Watch Video Solution](#)

80. If $\theta_1, \theta_2, \theta_3$ are the three values of $\theta \in [0, 2\pi]$ for which $\tan \theta = \lambda$

then the value of

$$\frac{\tan(\theta_1)}{3} \frac{\tan(\theta_2)}{3} + \frac{\tan(\theta_2)}{3} \frac{\tan(\theta_3)}{3} + \frac{\tan(\theta_3)}{3} \frac{\tan(\theta_1)}{3}$$
 is equal to

A. -3

B. -2

C. 2

D. 3

Answer: A

 [Watch Video Solution](#)

81. If

$$\tan \alpha = \frac{b}{a}, a > b > 0 \text{ and if } 0 < \alpha < \frac{\pi}{4}, \text{ then } \sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}}$$

is equal to :

A. $\frac{2 \sin \alpha}{\sqrt{2\alpha}}$

B. $\frac{2 \cos \alpha}{\sqrt{\cos 2\alpha}}$

C. $\frac{2 \sin \alpha}{\sqrt{\sin 2\alpha}}$

D. $\frac{2 \cos \alpha}{\sqrt{\sin 2\alpha}}$

Answer: B



Watch Video Solution

82. Minimum value of $3 \sin \theta + 4 \cos \theta$ in the interval $\left[0, \frac{\pi}{2}\right]$ is :

A. -5

B. 3

C. 4

D. $\frac{7}{\sqrt{2}}$

Answer: B



Watch Video Solution

83. If $f(n) = \prod_{r=1}^n \cos r$, $n \in N$, then

A. $|f(n)| > |f(n + 1)|$

B. $f(5) > 0$

C. $f(4) > 0$

D. $|f(n)| < |f(n + 1)|$

Answer: A



Watch Video Solution

84. If $\tan \theta + \sin \theta = m$ and $\tan \theta - \sin \theta = n$, then

A. 16

B. 22

C. 18

Answer: AA rectangular button with a red border and a red play button icon on the left, followed by the text "Watch Video Solution" in red.

85.

Let

$$t_1 = (\sin \alpha)^{\cos \alpha}, t_2 = (\sin \alpha)^{\sin \alpha}, t_3 = (\cos \alpha)^{\cos \alpha}, t_4 = (\cos \alpha)^{\sin \alpha},$$

where $\alpha \in \left(0, \frac{\pi}{4}\right)$, then which of the following is correct

A. $t_3 > t_1 > t_2$

B. $t_4 > t_2 > t_1$

C. $t_4 > t_1 > t_2$

D. $t_1 > t_3 > t_2$

Answer: BA rectangular button with a red border and a red play button icon on the left, followed by the text "Watch Video Solution" in red.

86. If $\cos A = \frac{3}{4}$ then the value of $\sin\left(\frac{A}{2}\right)\sin\left(\frac{5A}{2}\right)$ is

A. 11

B. -11

C. 12

D. 4

Answer: A



Watch Video Solution

87. $\cos(\alpha + \beta) + \sin(\alpha - \beta) = 0$ and $\tan \beta = \frac{1}{2009}$

A. 2

B. 1

C. 3

D. 4

Answer: B



Watch Video Solution

88. If $2^x = 3^y = 6^{-z}$ find the value of $\left(\frac{1}{x} + \frac{1}{y} + \frac{1}{z}\right)$

A. 0

B. 1

C. 2

D. 3

Answer: A



Watch Video Solution

89. Let α, β be such that it $\pi < \alpha - \beta < 3\pi$. if $\sin \alpha + \sin \beta = -\frac{21}{65}$
and $\cos \alpha + \cos \beta = -\frac{27}{65}$ then the value of $\cos\left(\frac{\alpha - \beta}{2}\right)$ is

A. $\frac{-3}{\sqrt{130}}$

B. $\frac{3}{\sqrt{130}}$

C. $\frac{6}{65}$

D. $-\frac{6}{65}$

Answer: A



Watch Video Solution

90. If $u = \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} + \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta}$, then the difference between maximum and minimum values of u^2 is

A. $2(a^2 + b^2)$

B. $(a + b)^2$

C. $2\sqrt{a^2 + b^2}$

D. $(a - b)^2$

Answer: D

 [Watch Video Solution](#)

91. If $P = (\tan(3^{n+1}\theta) - \tan\theta)$

A. $P = 2Q$

B. $P = 3Q$

C. $2P = Q$

D. $3P = Q$

Answer: A

 [Watch Video Solution](#)

92. If $270^\circ < \theta < 360^\circ$, then $\sqrt{2 + \sqrt{2 + 2\cos\theta}}$ is equal to

A. $-2\sin\left(\frac{\theta}{4}\right)$

B. $2\sin\left(\frac{\theta}{4}\right)$

C. $\pm 2\sin\frac{\theta}{4}$

D. $2\cos\frac{\theta}{4}$

Answer: B



Watch Video Solution

93. If $y = (\sin x + \cos x) + (\sin 4x + \cos 4x)^2$, then :

A. $y > 0 \forall x \in R$

B. $y \geq 0 \forall x \in R$

C. $y < 2 + \sqrt{2} \forall x \in R$

D. $y = 2 + \sqrt{2}$ for some $x \in R$

Answer: C



View Text Solution

94.

If

$\cos x + \cos y + \cos z = \sin x + \sin y + \sin z = 0$ then $\cos(x - y) =$

A. 0

B. $-\frac{1}{2}$

C. 2

D. 1

Answer: B



[Watch Video Solution](#)

95. Find the exact value of $\operatorname{cosec} 10^\circ + \operatorname{cosec} 50^\circ - \operatorname{cosec} 70^\circ$

A. 4

B. 5

C. 6

D. 8

Answer: C

 [Watch Video Solution](#)

96. If $270^\circ < \theta < 360^\circ$, then $\sqrt{2 + \sqrt{2 + 2 \cos \theta}}$ is equal to

A. $-2 \sin\left(\frac{\theta}{4}\right)$

B. $2 \sin\left(\frac{\theta}{4}\right)$

C. $\pm 2 \sin \frac{\theta}{4}$

D. $2 \cos \frac{\theta}{4}$

Answer: B

 [Watch Video Solution](#)

Exercise 2 One Or More Than One Answer Is Are Correct

1. $\cot 12^\circ \cdot \cot 24^\circ \cdot \cot 28^\circ \cdot \cot 32^\circ \cdot \cot 48^\circ \cdot \cot 88^\circ = \dots\dots\dots$

A. $\tan 45^\circ$

B. 2

C. $2 \tan 15^\circ \cdot \tan 45^\circ \cdot \tan 75^\circ$

D. $\tan 15^\circ \cdot \tan 45^\circ \cdot \tan 75^\circ$

Answer: A:D



Watch Video Solution

2. If the equation $\operatorname{cosec}^4 x - 2 \operatorname{cosec}^2 x + a^2 = 0$ has at least one solution, then the sum of all possible integral values of a is equal to a. 4 b. 3 c. 2 d.

0

A. -1

B. 0

C. 1

D. 2

Answer: A::B::C



Watch Video Solution

3. Which of the following is/are true ?

A. $\tan 1 > \tan^{-1} 1$

B. $\sin 1 > \cos 1$

C. $\tan 1 < \sin 1$

D. $\cos(\cos 1) > \frac{1}{\sqrt{2}}$

Answer: A::B::D



Watch Video Solution

4. Which of the following is/are +ve ?

A. $\log_{\sin 1} \tan 1$

B. $\log_{\cos 1}(1 + \tan 3)$

C. $\log_{\log_{10} 5}(\cos \theta + \sec \theta)$

D. $\log_{\tan 15^\circ}(2 \sin 18^\circ)$

Answer: B::D



Watch Video Solution

5. If $\sin \alpha + \cos \alpha = \frac{\sqrt{3} + 1}{2}$, $0 < \alpha < 2\pi$, then possible values $\tan \frac{\alpha}{2}$

can take is/are :

A. $2 - \sqrt{3}$

B. $\frac{1}{\sqrt{3}}$

C. 1

D. $\sqrt{3}$

Answer: A::B



Watch Video Solution

6. If $3 \sin \beta = \sin(2\alpha + \beta)$ then

A. $(\cot \alpha + \cot(\alpha + \beta))(\cot \beta - 3 \cot(2\alpha - \beta)) = 6$

B. $\sin \beta = \cos(\alpha + \beta) \sin \alpha$

C. $\tan(\alpha + \beta) = 2 \tan \alpha$

D. $2 \sin \beta = \sin(\alpha + \beta) \cos \alpha$

Answer: A::B::C::D



Watch Video Solution

7. If $\sin(x + 20^\circ) = 2 \sin x \cos 40^\circ$, where $x \in (0, \frac{\pi}{2})$, then which of the following hold(s) good? $\cos 2x = \frac{1}{2}$ (b) $\cos ec 4x = 2 \frac{\sec x}{2} = \sqrt{6} - \sqrt{2}$

(d) $\frac{\tan x}{2} = (2 - \sqrt{3})$

A. $\sec \frac{x}{2} = \sqrt{6} - \sqrt{2}$

B. $\cot \frac{x}{2} = 2 + \sqrt{3}$

C. $\tan 4x = \sqrt{3}$

D. $\operatorname{cosec} 4x = 2$

Answer: A::B



Watch Video Solution

8. If $2(\cos(x - y) + \cos(y - z) + \cos(z - x)) = -3$, then :

A. $\cos x \cos y \cos z = 1$

B. $\cos x + \cos y + \cos z = 0$

C. $\sin x + \sin y + \sin z = 1$

D. $\cos 3x + \cos 3y + \cos 3z = 12 \cos x \cos y \cos z$

Answer: B::D



View Text Solution

9. If $0 < x < \frac{\pi}{2}$ and $\sin^n x + \cos^n x \geq 1$, then 'n' may belong to interval :

A. $[1, 2]$

B. $[3, 4]$

C. $[-\infty, 2]$

D. $[-1, 1]$

Answer: A::C::D



Watch Video Solution

10.

If

$$x = \sin(\alpha - \beta) \cdot \sin(\gamma - \delta), y = \sin(\beta - \gamma) \cdot \sin(\alpha - \delta), z = \sin(\gamma - \alpha) \cdot \sin(\beta - \delta)$$

, then :

A. $x + y + z = 0$

B. $x^3 + y^3 + z^3 = 3xyz$

C. $x + y - z = 0$

$$D. x^3 + y^3 - z^3 = 3xyz$$

Answer: A::B



Watch Video Solution

11.

If

$$x = X \cos \theta - Y \sin \theta, y = X \sin \theta + Y \cos \theta \text{ and } x^2 + 4xy + y^2 = AX^2 +$$

then

A. $\theta = \frac{\pi}{6}$

B. $\theta = \frac{\pi}{4}$

C. $A = 3$

D. $B = -1$

Answer: B::C::D



Watch Video Solution

12. If $2a = 2\tan 10^\circ + \tan 50^\circ$, $2b = \tan 20^\circ + \tan 50^\circ$

$2c = 2\tan 10^\circ + \tan 70^\circ$, $2d = \tan 20^\circ + \tan 70^\circ$

Then which of the following is/are correct ?

A. $a + d = b + c$

B. $a + b = c$

C. $a > b < c > d$

D. $a < b < c < d$

Answer: A::B::D



[Watch Video Solution](#)

13. Which of the following real numbers when simplified are neither terminating nor repeating decimal ?

A. $\sin 75^\circ \cdot \cos 75^\circ$

B. $\log_2 28$

C. $\log_3 5 \cdot \log_5 6$

D. $8^{-(\log_{27} 3)}$

Answer: B::C



Watch Video Solution

14. If $a = \sin x \cos^3 x$ and $b = \cos x \sin^3 x$ then

A. $\alpha - \beta > 0$, for all x in $\left(0, \frac{\pi}{4}\right)$

B. $\alpha - \beta < 0$, for all x in $\left(0, \frac{\pi}{4}\right)$

C. $\alpha + \beta > 0$, for all x in $\left(0, \frac{\pi}{2}\right)$

D. $\alpha + \beta < 0$, for all x in $\left(0, \frac{\pi}{2}\right)$

Answer: A::C



Watch Video Solution

15. If $\frac{\pi}{2} < \theta < \pi$, then possible answers of $\sqrt{2 + \sqrt{2 + 2 \cos 4\theta}}$ is/are :

- A. $2 \cos \theta$
- B. $2 \sin \theta$
- C. $-2 \sin \theta$
- D. $-2 \cos \theta$

Answer: B::D



[Watch Video Solution](#)

16. If $\cot^3 \alpha + \cot^2 \alpha + \cot \alpha = 1$ then which of the following is/are correct :

- A. $\cos 2\alpha \tan \alpha = 1$
- B. $\cos 2\alpha \cdot \tan \alpha = -1$
- C. $\cos 2\alpha - \tan 2\alpha = -1$
- D. $\cos 2\alpha - \tan 2\alpha = 1$

Answer: B::D

 Watch Video Solution

17. The value of x in $\left(0, \frac{\pi}{2}\right)$ satisfying

$$\frac{\sqrt{3}-1}{\sin x} + \frac{\sqrt{3}+1}{\cos x} = 4\sqrt{2}$$

is/are $\frac{\pi}{12}$ (b) $\frac{5\pi}{12}$ (c) $\frac{7\pi}{24}$ (d) $\frac{11\pi}{36}$

A. $\frac{\pi}{15}$

B. $\frac{\pi}{12}$

C. $\frac{11\pi}{36}$

D. $\frac{3\pi}{10}$

Answer: B::C

 Watch Video Solution

18. If $\alpha > \frac{1}{\sin^6 x + \cos^6 x} \forall x \in R$, then α can be

A. 3

B. 4

C. 5

D. 6

Answer: B::D



Watch Video Solution

19. If $x \in \left(0, \frac{\pi}{2}\right)$ and $\sin x = \frac{3}{\sqrt{10}}$,

Let $k = \log_{10} \sin x + \log_{10} \cos x + 2 \log_{10} \cot x + \log_{10} \tan x$ then the value of k satisfies

A. $k = 0$

B. $k + 1 = 0$

C. $k - 1 = 0$

D. $k^2 - 1 = 0$

Answer: B::D

 [Watch Video Solution](#)

20. If A, B, C are angles of $\triangle ABC$ and $\tan A \tan C = 3, \tan B \tan C = 6$, then

A. $A = \frac{\pi}{4}$

B. $\tan A \tan B = 2$

C. $\frac{\tan A}{\tan C} = 3$

D. $\tan B = 2 \tan A$

Answer: A::B::D

 [Watch Video Solution](#)

21. The value of $\frac{\sin x - \cos x}{\sin^3 x}$ is equal to :

A. $\operatorname{cosec}^2 x (1 - \cot x)$

B. $1 - \cot x + \cot^2 x - \cot^3 x$

C. $\operatorname{cosec}^2 x - \cot x - \cot^3 x$

D. $\frac{1 - \cot x}{\sin^2 x}$

Answer: A::B::C::D

 [Watch Video Solution](#)

22. If $f(x) = \sin^2 x + \sin^2\left(x + \frac{2\pi}{3}\right) + \sin^2\left(x + \frac{4\pi}{3}\right)$ then :

A. $f\left(\frac{\pi}{15}\right) = \frac{3}{2}$

B. $f\left(\frac{15}{\pi}\right) = \frac{2}{3}$

C. $f\left(\frac{\pi}{10}\right) = \frac{3}{2}$

D. $f\left(\frac{10}{\pi}\right) = \frac{2}{3}$

Answer: A::C

 [Watch Video Solution](#)

23. The range of $y = \frac{\sin 4x - \sin 2x}{\sin 4x + \sin 2x}$ satisfies

A. $y \in \left(-\infty, \frac{1}{3}\right)$

B. $y \in \left(\frac{1}{3}, 1\right)$

C. $y \in (1, 3)$

D. $y \in (3, \infty)$

Answer: A::D



Watch Video Solution

24. If $\sqrt{2} \cos A = \cos B + \cos^3 B$, and $\sqrt{2} \sin A = \sin B - \sin^3 B$ then $\sin(A - B) =$

A. $\frac{1}{2}$

B. $\frac{1}{3}$

C. $-\frac{1}{2}$

D. $-\frac{1}{3}$

Answer: B::D

 [Watch Video Solution](#)

25. If $\alpha > \frac{1}{\sin^6 x + \cos^6 x} \forall x \in R$, then α can be

A. 3

B. 4

C. 5

D. 6

Answer: C::D

 [Watch Video Solution](#)

26. If $\cot^3 \alpha + \cot^2 \alpha + \cot \alpha = 1$ then which of the following is/are correct :

A. $\cos 2\alpha \tan \alpha = 1$

B. $\cos 2\alpha \cdot \tan \alpha = -1$

C. $\cos 2\alpha - \tan 2\alpha = -1$

D. $\cos 2\alpha - \tan 2\alpha = 1$

Answer: B::D



[Watch Video Solution](#)

Exercise 3 Comprehension Type Problems

1. Let $l = \sin \theta$, $m = \cos \theta$ and $n = \tan \theta$.

Q. If $\theta = 5$ radian, then :

A. $l > m$

B. $l < m$

C. $l = m$

D. none of these

Answer: B

 [Watch Video Solution](#)

2. Let $l = \sin \theta$, $m = \cos \theta$ and $n = \tan \theta$.

Q. If $\theta = -1042^\circ$, then :

A. $n > 1$

B. $n < 1$

C. $n = 1$

D. nothing can be said

Answer: B

 [Watch Video Solution](#)

3. Let $l = \sin \theta$, $m = \cos \theta$ and $n = \tan \theta$.

Q. If $\theta = 7$ radian, then :

A. $l + m > 0$

B. $l + m < 0$

C. $l + m = 0$

D. nothing can be said

Answer: A



Watch Video Solution

4. Let a, b, c are respectively the sines and p, q, r are respectively the consines of α , $\alpha + \frac{2\pi}{3}$ and $\alpha + \frac{4\pi}{3}$, then :

Q. The value of $(a + b + c)$ is :

A. 0

B. $\frac{3}{4}$

C. 1

D. none of these

Answer: A



Watch Video Solution

5. Let a, b, c are respectively the sines and p, q, r are respectively the consines of $\alpha, \alpha + \frac{2\pi}{3}$ and $\alpha + \frac{4\pi}{3}$, then :

Q. The value of $(ab + bc + ca)$ is :

A. 0

B. $-\frac{3}{4}$

C. $-\frac{1}{2}$

D. -1

Answer: B

[Watch Video Solution](#)

6. Let a, b, c are respectively the sines and p, q, r are respectively the cosines of $\alpha, \alpha + \frac{2\pi}{3}$ and $\alpha + \frac{4\pi}{3}$, then :

Q. The value of $(qc - rb)$ is :

A. 0

B. $-\frac{\sqrt{3}}{2}$

C. $\frac{\sqrt{3}}{2}$

D. depends on α

Answer: C

[Watch Video Solution](#)

7. Consider a right angle triangle ABC right angle at B such that $AC = \sqrt{8 + 4\sqrt{3}}$ and $AB = 1$. A line through vertex A meet BC at D such that $AB = BD$. An arc DE of radius AD is drawn from vertex A to meet

AC at E and another arc DF of radius CD is drawn from vertex C to meet AC at F. On the basis of above information, answer the following questions.

Q. $\sqrt{\tan A + \cot C}$ is equal to :

A. $\sqrt{3}$

B. 1

C. $2 + \sqrt{3}$

D. $\sqrt{3} + 1$

Answer: D



Watch Video Solution

8. Consider a right angle triangle ABC right angle at B such that $AC = \sqrt{8 + 4\sqrt{3}}$ and $AB = 1$. A line through vertex A meet BC at D such that $AB = BD$. An arc DE of radius AD is drawn from vertex A to meet AC at E and another arc DF of radius CD is drawn from vertex C to meet AC at F. On the basis of above information, answer the following

questions.

Q. $\log\left(\frac{AE + CF}{CD}\right)$ is equal to :

A. $\sqrt{2}$

B. 1

C. 0

D. -1

Answer: B



Watch Video Solution

9. Consider a triangle ABC such that $\cot A + \cot B + \cot C = \cot \theta$. Now answer the following :

Q. The possible value of θ is :

A. 60°

B. 25°

C. 35°

D. 45°

Answer: B



[View Text Solution](#)

10. Consider a triangle ABC such that $\cot A + \cot B + \cot C = \cot \theta$.

Now answer the following :

Q. $\sin(A - \theta)\sin(B - \theta)\sin(C - \theta) = :$

A. $\tan^3 \theta$

B. $\cot^3 \theta$

C. $\sin^3 \theta$

D. $\cos^3 \theta$

Answer: C



[Watch Video Solution](#)

11. Consider the function $f(x) = \frac{\sqrt{1 + \cos x} + \sqrt{1 - \cos x}}{\sqrt{1 + \cos x} - \sqrt{1 - \cos x}}$ then

Q. If $x \in (\pi, 2\pi)$ then $f(x)$ is

A. $\cot\left(\frac{\pi}{2} + \frac{x}{2}\right)$

B. $\tan\left(\frac{\pi}{4} + \frac{x}{2}\right)$

C. $\cot\left(\frac{\pi}{4} - \frac{x}{2}\right)$

D. $\tan\left(\frac{\pi}{4} - \frac{x}{2}\right)$

Answer: D



Watch Video Solution

12. If the function $f(x) = \frac{\sqrt{1 + \cos x} + \sqrt{1 - \cos x}}{\sqrt{1 + \cos x} - \sqrt{1 - \cos x}}$ if the value of

$f\left(\frac{\pi}{3}\right) = a + b\sqrt{c}$ then $a + b + c =$

A. 4

B. 5

C. 6

D. 7

Answer: C



Watch Video Solution

Exercise 4 Matching Type Problems

Column-I		Column-II
(A) If $(1 + \tan 5^\circ)(1 + \tan 10^\circ) \dots (1 + \tan 45^\circ) = 2^{k+1}$ then 'k' equals	(P)	0
(B) Sum of positive integral values of 'a' for which $a^2 - 6 \sin x - 5a \leq 0 \forall x \in R$ is	(Q)	2
(C) The minimum value of $\frac{\left(a + \frac{1}{a}\right)^4 - \left(a^4 + \frac{1}{a^4}\right) - 2}{\left(a + \frac{1}{a}\right)^2 + a^2 + \frac{1}{a^2}}$ is	(R)	5
(D) Number of real roots of the equation $\sum_{k=1}^3 (x-k)^2 = 0$ is	(S)	4
	(T)	5

1.



Watch Video Solution

Column-I		Column-II	
(A)	Maximum value of $y = \frac{1 - \tan^2(\pi/4 - x)}{1 + \tan^2(\pi/4 - x)}$	(P)	1
(B)	Minimum value of $\log_3 \left(\frac{5 \sin x - 12 \cos x + 26}{13} \right)$	(Q)	0
(C)	Minimum value of $y = -2 \sin^2 x + \cos x + 3$	(R)	$\frac{7}{8}$
(D)	Maximum value of $y = 4 \sin^2 \theta + 4 \sin \theta \cos \theta + \cos^2 \theta$	(S)	5
		(T)	6

2.

 [Watch Video Solution](#)

Exercise 5 Subjective Type Problems

1. Let $P = \frac{\sin 80^\circ \sin 65^\circ \sin 35^\circ}{\sin 20^\circ + \sin 50^\circ + \sin 110^\circ}$, then the value of $24P$ is :

 [Watch Video Solution](#)

2. The value of expression $(1 - \cot 23^\circ)(1 - \cot 22^\circ)$ is equal to :

 [Watch Video Solution](#)

3. If $\tan A$ and $\tan B$ the roots of the quadratic equation,

$$4x^2 - 7x + 1 = 0$$

then

evaluate

$$4 \sin^2(A + B) - 7 \sin(A + B) \cdot \cos(A + B) + \cos^2(A + B).$$



[Watch Video Solution](#)

4. $A_1A_2A_3 \dots A_{18}$ is a regular 18 sided polygon. B is an external point such that A_1A_2B is an equilateral triangle. If $A_{18}A_1$ and A_1B are adjacent sides of a regular n sided polygon, then $n =$



[Watch Video Solution](#)

5. If $10 \sin^4 \alpha + 15 \cos^4 \alpha = 6$ then the value of $9 \cos^4 \alpha + 8 \sec^4 \alpha - 75$ is



[Watch Video Solution](#)

6. The value of $\left(1 + \tan \frac{3\pi}{8} \cdot \tan \frac{\pi}{8}\right) + \left(1 + \tan \frac{5\pi}{8} \cdot \tan \frac{3\pi}{8}\right) + \left(1 + \tan \frac{7\pi}{8} \cdot \tan \frac{5\pi}{8}\right)$

 [Watch Video Solution](#)

7. If $\alpha = \frac{\pi}{7}$ then find the value of $\left(\frac{1}{\cos \alpha} + \frac{2 \cos \alpha}{\cos 2\alpha}\right)$

 [Watch Video Solution](#)

8. Given the for $a, b, c, d \in R$, if a

$\sec(200^\circ) - c \tan(200^\circ) = d$ and $b \sec(200^\circ) + d \tan(200^\circ) = c$,

then find the value of $\left(\frac{a^2 + b^2 + c^2 + d^2}{bd - ac}\right) \sin 20^\circ$.

 [Watch Video Solution](#)

9. The expression $2 \cos \frac{\pi}{17} \cdot \cos \frac{9\pi}{17} + \cos \frac{7\pi}{17} + \cos \frac{9\pi}{17}$ simplifies to an integer P. Find the value of P.



Watch Video Solution

10. If the expression $\frac{\sin \theta \sin 2\theta + \sin 3\theta \sin 6\theta + \sin 4\theta \sin 13\theta}{\sin \theta \cos 2\theta + \sin 3\theta \cos 6\theta + \sin 4\theta \cos 13\theta} = \tan k\theta$, where $k \in \mathbb{N}$.

Find the value of k .



Watch Video Solution

11. Let $a = \sin 10^\circ$, $b = \sin 50^\circ$, $c = \sin 70^\circ$, then $8abc \left(\frac{a+b}{c} \right) \left(\frac{1}{a} + \frac{1}{b} - \frac{1}{c} \right)$

is equal to



Watch Video Solution

12. If $\sin^3 \theta + \sin^3 \left(\theta + \frac{2\pi}{3} \right) + \sin^3 \left(\theta + \frac{4\pi}{3} \right) = a \sin b\theta$. Find the value of $\left| \frac{b}{a} \right|$.



Watch Video Solution

13. If $\sum_{r=1}^n \left(\frac{\tan 2^{r-1}}{\cos 2^r} \right) = \tan p^n - \tan q$, then find the value of $(p + q)$.

 [Watch Video Solution](#)

14. If $x = \sec \theta - \tan \theta$ and $y = \operatorname{cosec} \theta + \cot \theta$, then $y - x - xy =$

 [Watch Video Solution](#)

15. If $\cos 18^\circ - \sin 18^\circ = \sqrt{n} \sin 27^\circ$, then $n =$

 [Watch Video Solution](#)

16. The value of $3(\sin 1 - \cos 1)^4 + 6(\sin 1 + \cos 1)^2 + 4(\sin^6 1 + \cos^6 1)$ is equal to

 [Watch Video Solution](#)

17. Q. $x = a$ satisfy the equation

$$3^{\sin 2x + 2 \cos^2 x} + 3^{1 - \sin 2x + 2 \sin^2 x} = 28(\sin 2a - \cos 2a)^2 + 8 \sin 4a$$
 is

equal to:

 [Watch Video Solution](#)

18. The minimum value of $(\sin \theta + \operatorname{cosec} \theta)^2 + (\cos \theta + \sec \theta)^2 =$

 [Watch Video Solution](#)

19. If $\tan 20^\circ + \tan 40^\circ + \tan 80^\circ - \tan 60^\circ = \lambda \sin 40^\circ$,

 [Watch Video Solution](#)

20. If K° lies between 360° and 540° and K° satisfies the equation

$$1 + \cos 10x \cos 6x = 2 \cos^2 8x + \sin^2 8x, \quad \text{then } \frac{K}{10} =$$

 [Watch Video Solution](#)

21. If $\cos 20^\circ + 2 \sin^2 55^\circ = 1 + \sqrt{2} \sin K^\circ$, $K \in (0, 90)$, then $K =$

 [Watch Video Solution](#)

22. If the value of $\cos\left(\frac{2\pi}{7}\right) + \cos\left(\frac{4\pi}{7}\right) + \cos\left(\frac{6\pi}{7}\right) + \cos\left(\frac{l\pi}{7}\right) = -\frac{l}{2}$ Find the value of l

 [Watch Video Solution](#)

23. Let α be the smallest integral value of x , $x > 0$ such that $\tan 19x = \frac{\cos 96^\circ + \sin 96^\circ}{\cos 96^\circ - \sin 96^\circ}$. The last digit of α is

 [Watch Video Solution](#)

24. Find the value of the expression $\frac{\sin 20^\circ (4 \cos 20^\circ + 1)}{\cos 20^\circ \cos 30^\circ}$



 Watch Video Solution

25. If the value of $\cos\left(\frac{2\pi}{7}\right) + \cos\left(\frac{4\pi}{7}\right) + \cos\left(\frac{6\pi}{7}\right) + \cos\left(\frac{l\pi}{7}\right) = -\frac{l}{2}$ Find the value of l

 Watch Video Solution

26. If $\cos A = \frac{3}{4}$ and $k \sin\left(\frac{A}{2}\right) \sin\left(\frac{5A}{2}\right) = \frac{11}{8}$. Find k .

 Watch Video Solution

27. Find the least value of the expression $3 \sin^2 x + 4 \cos^2 x$.

 Watch Video Solution

28. Q. If $\tan \alpha$ and $\tan \beta$ are the roots of equation $x^2 - 12x - 3 = 0$, then the value of $\sin^2(\alpha + \beta) + 2\sin(\alpha + \beta)\cos(\alpha + \beta) + 5\cos^2(\alpha + \beta)$ is



[Watch Video Solution](#)

29. The value of $\frac{\cos 24^\circ}{2\tan 33^\circ \sin^2(57^\circ)} + \frac{\sin 162^\circ}{\sin 18^\circ - \cos 18^\circ \tan 9^\circ} + \cos 162^\circ$ is equal to



[Watch Video Solution](#)

30. Find the value of $\tan \theta(1 + \sec 2\theta)(1 + \sec 4\theta)(1 + \sec 8\theta)$, when $\theta = \frac{\pi}{32}$



[Watch Video Solution](#)

31. If λ be the minimum value of $y = (\sin x + \operatorname{cosec} x)^2 + (\cos x + \sec x)^2 + (\tan x + \cot x)^2$ where $x \in R$. Find $\lambda - 6$.



[Watch Video Solution](#)