



India's Number 1 Education App

## MATHS

### BOOKS - VIKAS GUPTA MATHS (HINGLISH)

#### DIFFERENTIAL EQUATIONS

##### Exercise Single Choice Problems

1.  $\frac{dy}{dx} \left( \frac{1 + \cos x}{y} \right) = -\sin x$  and  $f\left(\frac{\pi}{2}\right) = -1$ , then  $f(0)$  is:

A. -2

B. -1

C. -3

D. -4

**Answer: A**



doubtnut | doubt, not out



Watch Video Solution

2. The differential equation satisfied by family of curves

$y = Ae^x + Be^{3x} + Ce^{5x}$  where A,B,C are arbitrary constants is:

A.  $\frac{d^3y}{dx^3} - 9\frac{d^2y}{dx^2} + 23\frac{dy}{dx} + 15y = 0$

B.  $\frac{d^3y}{dx^3} - 9\frac{d^2y}{dx^2} - 23\frac{dy}{dx} + 15y = 0$

C.  $\frac{d^3y}{dx^3} + 9\frac{d^2y}{dx^2} - 23\frac{dy}{dx} + 15y = 0$

D.  $\frac{d^3y}{dx^3} + 9\frac{d^2y}{dx^2} + 23\frac{dy}{dx} - 15y = 0$

Answer: A



Watch Video Solution

3. If  $y = y(x)$  and it follows the relation  $e^{xy^2} + y \cos(x^2) = 5$  then  $y'(0)$

is equal to

A. 4

B. - 16

C. - 4

D. 16

**Answer: B**



**Watch Video Solution**

4. If  $(x^2 + y^2)dy = xydx$  and  $y(1)=1$  and  $y(x_o) = e$ , then  $x_o =$

A.  $\sqrt{3}e$

B.  $\sqrt{e^2 - \frac{1}{2}}$

C.  $\sqrt{\frac{e^2 - 1}{2}}$

D.  $\sqrt{e^2 + \frac{1}{2}}$

**Answer: A**



**Watch Video Solution**

5. The differential equation  $\frac{dy}{dx} = \frac{\sqrt{1 - y^2}}{y}$  determines a family of circles with

- A. Variable radii and fixed centre at  $(0, 1)$
- B. Variable radii and fixed centre at  $(0, -1)$
- C. Fixed radius 1 and variable centres along y-axis
- D. Fixed radius 1 and variable centres along x-axis

**Answer: C**



**Watch Video Solution**

6. Interval contained in the domain of definition of non-zero solution of the differential equation  $(x - 3)^2 y' + y = 0$  is:

A.  $R - \{3\}$

B.  $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$

C.  $\left(\frac{\pi}{8}, \frac{5\pi}{4}\right)$

D.  $(-\pi, \pi)$

**Answer: A**



**Watch Video Solution**

7. A function  $y = f(x)$  satisfies

$$(x+1)f'(x) - 2(x^2+x)f(x) = \frac{e^x}{(x+1)^2}, \forall x > 1. \text{ If } f(0) = 5, \text{ then}$$

$f(x)$  is (a)

$$(b)(c) \left( (d)(e)(f) \frac{(g)3x+5}{h} ((i)x+1)(j)(k)(l) \right) (m)e^n(o)(p)x^{((q)2(r))(s)}$$

(w) (b) [Math Processing Error] (ss) (c)

$$(d)(e) \left( (f)(g)(h) \frac{(i)6x+5}{j} ((k)x+1)(l)(m)(n) \right) (o)e^p(q)(r)x^{((s)2(t))(u)}$$

(y) (d) [Math Processing Error] (uu)

A.  $\left( \frac{3x+5}{x+1} \right) \cdot e^{x^2}$

B.  $\left( \frac{6x+5}{x+1} \right) \cdot e^{x^2}$

C.  $\left( \frac{6x+5}{(x+1)^2} \right) \cdot e^{x^2}$

D.  $\left( \frac{5x+6x}{x+1} \right) \cdot e^{x^2}$

**Answer: B**



**Watch Video Solution**

**8.** The solution of the differential equation

$2x^2y \frac{dy}{dx} = \tan(x^2y^2) - 2xy^2$ , given  $y(1) = \sqrt{\frac{\pi}{2}}$ , is

A.  $\sin(x^2y^2) - 1 = 0$

B.  $\cos\left(\frac{\pi}{2} + x^2y^2\right) + x = 0$

C.  $\sin(x^2y^2) = e^{x-1}$

D.  $\sin(x^2y^2) = e^{(x-1)}$

**Answer: C**



**Watch Video Solution**

**9.** The differential equation whose general solution is given by

$$y = \left(c_1 \cos(x + c_2) - \left(c_3 e^{(-x+c_4)} + (c_5 \sin x)\right)\right)$$

are arbitrary constants, is (a)

$$(b)(c)(d) \frac{(e)(f)d^{(g)4(h)}(i)y}{j} \left( (k)d(l)x^{(m)4(n)}(o) \right) (p)(q) - (r) \frac{(s)(t)d^{(u)}}{x}$$

(gg) (hh)

$$(ii)(jj)(kk) \frac{(ll)(mm)d^{(nn)3(o)}(pp)y}{qq} \left( (rr)d(ss)x^{(tt)3(u)}(vv) \right) (ww) (\times)$$

$$+ (mmm) \frac{(nnn)dy}{ooo} ((ppp)dx)(qqq)(rrr) + y = 0(sss)$$

(ttt)

(uuu)

$$(vvv)(www)(\times x) \frac{(yyy)(zzz)d^{(aaaa)5(bbbb)}(cccc)}{dddd} \left( (eeee)d(ffff)x^{(gggg)5} \right)$$

(mmmm) (nnnn)

$$(oooo)(pppp)(qqqq) \frac{(rrrr)(ssss)d^{(ttt)3(uuu)}(vvvv)y}{wwww} \left( (xxxx)d(yyyy)x^{(zzz)} \right)$$

$$- (eeeeee) \frac{(fffff)(ggggg)d^{(hhhhh)2(iiii)}(jjjjj)y}{kkkkk} \left( (llll)d(mmmmm)m \right) x^{(nr)}$$

$$= 0(yyyyy)$$

(zzzzz)

A.  $\frac{d^4y}{dx^4} + \frac{d^2y}{dx^2} + y = 0$

B.  $\frac{d^3y}{dx^3} + \frac{d^2y}{dx^2} + \frac{dy}{dx} - y = 0$

C.  $\frac{d^3y}{dx^3} + \frac{d^2y}{dx^2} + \frac{dy}{dx} + y = 0$

D.  $\frac{d^3y}{dx^3} + \frac{d^2y}{dx^2} + \frac{dy}{dx} = 0$

**Answer: C**



Watch Video Solution

10. If  $y = e^{(\alpha+1)x}$  be solution of differential equation  $\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 0$ , then  $\alpha$  is:

A. 0

B. 1

C. -1

D. 2

**Answer: B**



Watch Video Solution

11. The order and degree of the differential equation  $\left(\frac{dy}{dx}\right)^{1/3} - 4\frac{d^2y}{dx^2} - 7x = 0$  are  $\alpha$  and  $\beta$ , then the value of  $(\alpha + \beta)$  is:

A. 3

B. 4

C. 2

D. 5

**Answer: D**



**Watch Video Solution**

12. General solution of differential equation of

$f(x) \frac{dy}{dx} = f^2(x) + yf(x) + f'(x)y$  is:

(c being arbitrary constant.)

A.  $y = f(x) + ce^x$

B.  $y = -f(x) + ce^x$

C.  $y = -f(x) + ce^x f(x)$

D.  $y = cf(x) + e^x$

**Answer: C**



**Watch Video Solution**

**13.** The order and degree respectively of the differential equation of all tangent lines to parabola  $x^2 = 2y$  is:

A. 1, 2

B. 2, 1

C. 1, 1

D. 1, 3

**Answer: A**



**Watch Video Solution**

**14.** The general solution of the differential equation  $\frac{dy}{dx} + x(x+y) = x(x+y)^3 - 1$  is:

- A.  $\ln \left| \frac{(x+y+1)(x+y-1)}{(x+y)^4} \right| = x^2 + C$
- B.  $\ln \left| \frac{(x+y+1)(x+y-1)}{(x+y)^2} \right| = x^2 + C$
- C.  $2 \ln \left| \frac{(x+y+1)(x+y-1)}{(x+y)^2} \right| = x^2 + C$
- D.  $\ln \left| \frac{(x+y+1)(x+y-1)}{(x+y)^2} \right| = x + C$

**Answer: B**



**Watch Video Solution**

15. The general solution of  $\frac{dy}{dx} = 2y \tan x + \tan^2 x$  is:

A.  $y \cos^2 x = \frac{x}{2} - \frac{\sin 2x}{4} + C$

B.  $y \sec^2 x = \frac{x}{2} - \frac{\sin 2x}{4} + C$

C.  $y \cos^2 x = \frac{x}{2} - \frac{\cos 2x}{4} + C$

D.  $y \cos^2 x = \frac{x}{2} - \frac{\sin 2x}{4} + C$

**Answer: A**



**Watch Video Solution**

16. The solution of differential equation  $\frac{d^2y}{dx^2} = \frac{dy}{dx}$ ,  $y(0) = 3$  and  $y'(0) = 2$ :

- A. is a periodic function
- B. approaches to zero as  $x \rightarrow -\infty$
- C. has an asymptote parallel to x-axis
- D. has an asymptote parallel to y-axis

**Answer: C**



17. The solution of the differential equation  $(x^2 + 1) \frac{d^2y}{dx^2} = 2x \frac{dy}{dx}$  under the conditions  $y(0)=1$  and  $y'(0)=3$ , is

- A.  $y = x^2 + 3x + 1$

B.  $y = x^3 + 3x + 1$

C.  $y = x^4 + 3x + 1$

D.  $y = 3 \tan^{-1} x + x^2 + 1$

**Answer: B**



**Watch Video Solution**

**18.** The differential equation of the family of curves  $cy^2 = 2x + c$  (where c is an arbitrary constant.) is:

A.  $\frac{x dy}{dx} = 1$

B.  $\left( \frac{dy}{dx} \right) = \frac{2xly}{dx} + 1$

C.  $y^2 = 2xy \frac{dy}{dx} + 1$

D.  $y^2 = \frac{2ydy}{dx} + 1$

**Answer: C**



**Watch Video Solution**

19. The solution of the equation  $\frac{dy}{dx} + \frac{1}{x} = \frac{1}{x^2} \tan y \sin y$  is:

A.  $2y = \sin y(1 - 2cx^2)$

B.  $2x = \cot y(1 + 2cx^2)$

C.  $2x = \sin y(1 - 2cx^2)$

D.  $2x \sin y = 1 - 2cx^2$

**Answer: C**



[View Text Solution](#)

20. Solution of the differential equation  $xdy - \sqrt{x^2 + y^2}dx = 0$  is :

A.  $y - \sqrt{x^2 + y^2} = cx^2$

B.  $y + \sqrt{x^2 + y^2} = cx$

C.  $x = \sqrt{x^2 + y^2} = cx^2$

D.  $y + \sqrt{x^2 + y^2} = cx^2$

**Answer: D**



**Watch Video Solution**

21. Let  $f(x)$  be differentiable on the interval  $(0, \infty)$  such that  $f(1) = 1$  and  $\lim_{t \rightarrow x} \frac{t^2 f(x) - x^2 f(t)}{t - x} = 1$  for each  $x > 0$ . Then  $f(x) =$

A.  $\frac{1}{4x} + \frac{3x^2}{4}$

B.  $\frac{3}{4x} + \frac{x^2}{4}$

C.  $\frac{x^2}{2} + \frac{1}{2}$

D.  $\frac{1}{4x^3} + \frac{3x}{4}$

**Answer: C**



**Watch Video Solution**

22. The population  $p(t)$  at time  $t$  of a certain mouse species satisfies the differential equation  $\left( dp \frac{t}{dt} = 0.5p(t) - 450 \right)$  If  $p(0) = 850$ , then the

time at which the population becomes zero is (1)  $2 \ln 18$  (2)  $\ln 9$  (3)  $\frac{1}{2} \ln 18$  (4)  $\ln 18$

A.  $\frac{1}{2} \ln 18$

B.  $\ln 18$

C.  $2 \ln 18$

D.  $\ln 9$

**Answer: C**



[Watch Video Solution](#)

23. The solution of the differential equation  
 $\sin 2y \frac{dy}{dx} + 2 \tan x \cos^2 y = 2 \sec x \cos^3 y$  is: (where  $C$  is arbitrary constant)

A.  $\cos y \sec x = \tan x + C$

B.  $\sec y \cos x = \tan x + C$

C.  $\sec y \sec x = \tan x + C$

D.  $\tan y \sec x = \sec x + C$

**Answer: C**



**Watch Video Solution**

24. The solution of the differential equation  $\frac{dy}{dx} = (4x + y + 1)^2$ , is

A.  $4x + y + 1 = 2 \tan(2x + y + C)$

B.  $4x + y + 1 = 2 \tan(x + 2y + C)$

C.  $4x + y + 1 = 2 \tan(2y + c)$

D.  $4x + y + 1 = 2 \tan(2x + c)$

**Answer: D**



**Watch Video Solution**

25. If a curve is such that line joining origin to any point  $P(x, y)$  on the curve and the line parallel to y-axis through P are equally inclined to tangent to curve at P, then the differential equation of the curve is:

A.  $x \left( \frac{dy}{dx} \right)^2 - 2y \frac{dy}{dx} = x$

B.  $x \left( \frac{dy}{dx} \right)^2 + 2y \frac{dy}{dx} = x$

C.  $y \left( \frac{dy}{dx} \right)^2 - 2x \frac{dy}{dx} = x$

D.  $y \left( \frac{dy}{dx} \right)^2 - 2y \frac{dy}{dx} = x$

**Answer: A**



**Watch Video Solution**

26. If  $y = f(x)$  satisfy the differential equation  $\frac{dy}{dx} + \frac{y}{x} = x^2$ ,  $f(1) = 1$ , then value of  $f(3)$  equals:

A. 7

B. 5

C. 9

D. 27

**Answer: A**



**Watch Video Solution**

27. Let  $y = f(x)$  and  $\frac{x}{y} \frac{dy}{dx} = \frac{3x^2 - y}{2y - x^2}$ ,  $f(1) = 1$  then the possible value of  $\frac{1}{3}f(3)$  equals :

A. 9

B. 4

C. 3

D. 2

**Answer: C**



**Watch Video Solution**

## Exercise One Or More Than One Answer Is Are Correct

1. Let  $y=f(x)$  be a real valued function satisfying  $x \frac{dy}{dx} = x^2 + y - 2$ ,  $f(1)=1$

then  $f(3)$  equal

A.  $f(x)$  is minimum at  $x = 1$

B.  $f(x)$  is maximum at  $x = 1$

C.  $f(3) = 5$

D.  $f(2) = 3$

**Answer: A::C**



Watch Video Solution

2. solution of differential equation  $x \cos x \frac{dy}{dx} + y(x \sin x + \cos x) = 1$

is

A.  $xy = \sin x + c \cos x$

B.  $xy \sec x = \tan x + c$

C.  $xy + \sin x + c \cos x = 0$

D. None of these

**Answer: A::B**



**Watch Video Solution**

3. If a function satisfies  $(x - y)f(x + y) - (x + y)f(x - y) = 2(x^2y - y^3)$   $\forall x, y \in R$  and  $f(1) = 0$ , then

A.  $f(x)$  must be polynomial function

B.  $f(3) = 12$

C.  $f(0) = 0$

D.  $f(3) = 13$

**Answer: A::B::C**



4. A function  $y=f(x)$  satisfies the differential equation  $f(x)\sin 2x - \cos x + (1 + \sin^2 x)f'(x) = 0$  with initial condition  $y(0) = 0$ . The value of  $f\left(\frac{\pi}{6}\right)$  is equal to

A.  $\frac{2}{5}$

B.  $\frac{3}{5}$

C.  $\frac{1}{5}$

D.  $\frac{4}{5}$

**Answer: A**



**Watch Video Solution**

5. Solution of the differential equation  $(2 + 2x^2\sqrt{y})ydx + (x^2\sqrt{y} + 2)xdy = 0$  is/are:

A.  $xy\left(x^2\sqrt{y} + 5\right) = c$

B.  $xy\left(x^2\sqrt{y} + 3\right) = c$

C.  $xy(y^2\sqrt{x} + 3) = c$

D.  $xy(y^2\sqrt{x} + 5) = c$

**Answer: B**



**Watch Video Solution**

6. If  $y(x)$  satisfies the differential equation  $\frac{dy}{dx} = \sin 2x + 3y \cot x$  and  $y\left(\frac{\pi}{2}\right) = 2$  then which of the following statement(s) is/are correct ?

A.  $y\left(\frac{\pi}{6}\right) = 0$

B.  $y'\left(\frac{\pi}{3}\right) = \frac{9 - 3\sqrt{2}}{2}$

C.  $y(x)$  increases in the interval

D.  $\int_{-\pi/2}^{\pi/2} y(x) dx = x$

**Answer: A::C**



**Watch Video Solution**

### Exercise Comprehension Type Problem

1. A differentiable function  $y = g(x)$  satisfies  $\int_0^x (x - t + 1)g(t)dt = x^4 + x^2$  for all  $x \geq 0$  then  $y=g(x)$  satisfies the differential equation

A.  $\frac{dy}{dx} - y = 12x^2 + 2$

B.  $\frac{dy}{dx} + 2y = 12x^2 + 2$

C.  $\frac{dy}{dx} + y = 12x^2 + 2$

D.  $\frac{dy}{dx} + y = 12x + 2$

**Answer: C**



**Watch Video Solution**

2. A differentiable function  $y = g(x)$  satisfies  
 $\int_0^x (x-t+1)g(t)dt = x^4 + x^2$  for all  $x \geq 0$  then  $y=g(x)$  satisfies the differential equation

A. 0

B. 1

C.  $e^2$

D. Data insufficient

**Answer: A**



**Watch Video Solution**

3. Suppose  $f$  and  $g$  are differentiable functions such that  $xg(f(x))f'(g(x))g'(x) = (g(x))g'(f(x))f'(x) \forall x \in R$  and  $f$  is positive  $\forall n \in R$ . Also

$$\int_0^x f(g(t))dt = \frac{1}{2}(1 - e^{-2x}) \quad \forall x \in R, g(f(0)) = 1 \text{ and } h(x) = \frac{g(f(x))}{f(g(x))}$$

The graph of  $y = h(x)$  is symmetric with respect to line:

A.  $x = -1$

B.  $x = 0$

C.  $x = 1$

D.  $x = 2$

**Answer: C**



**Watch Video Solution**

4. Let  $f$  and  $g$  be differentiable functions such that:

$$xg(f(x))f'(g(x))g'(x) = f(g(x))g'(f(x))f'(x) \quad \forall x \in R \quad \text{Also,}$$

$$f(x) > 0 \text{ and } g(x) > 0 \quad \forall x \in R \quad \int_0^x f(g(t))dt = 1 - \frac{e^{-2x}}{2}, \quad \forall x \in R$$

and  $g(f(0)) = 1$ ,  $h(x) = \frac{g(f(x))}{f(g(x))} \quad \forall x \in R$  Now answer the question:

$$f(g(0)) + g(f(0)) = \text{(A) 1 (B) 2 (C) 3 (D) 4}$$

A. 1

B. 2

C. 3

**Answer: B****Watch Video Solution**

5. Suppose  $f$  and  $g$  are differentiable functions such that  $xg(f(x))f'(g(x))g'(x) = (g(x))g'(f(x))f'(x) \forall x \in R$  and  $f$  is positive  $\forall n \in R$ . Also

$$\int_0^x f(g(t))dt = \frac{1}{2}(1 - e^{-2x}) \quad \forall x \in R, g(f(0)) = 1 \text{ and } h(x) = \frac{g(f(x))}{f(g(x))}$$

The largest possible value of  $h(x) \forall x \in R$  is:

A. 1

B.  $e^{1/3}$ 

C. e

D.  $e^2$ **Answer: C****Watch Video Solution**

6. Given a function 'g' which has a derivative  $g'(x)$  for every real  $x$  and satisfies  $g'(0) = 2$  and  $g(x + y) = e^y g(x) + e^y g(y)$  for all  $x$  and  $y$  then:

A.  $x(2 + xe^x)$

B.  $e(e^x + 1)$

C.  $2xe^x$

D.  $x + \ln(x + 1)$

**Answer: C**



7. Given a function 'g' which has a derivative  $g'(x)$  for every real  $x$  and satisfies  $g'(0) = 2$  and  $g(x + y) = e^y g(x) + e^y g(y)$  for all  $x$  and  $y$  then:

A. R

B.  $\left[ -\frac{2}{e}, \infty \right)$

C.  $\left[ -\frac{1}{e}, \infty \right)$

D.  $[0, \infty)$

**Answer: B**



**Watch Video Solution**

8. Given a function 'g' which has a derivative  $g'(x)$  for every real  $x$  and satisfies  $g'(0) = 2$  and  $g(x + y) = e^y g(x) + e^y g(y)$  for all  $x$  and  $y$  then:

A. 0

B. 1

C. 2

D. Does not exist

**Answer: A**



**Watch Video Solution**

## Exercise Matching Type Problems

| Column-I (Differential equation) |  | Column-II Solution (Integral curves) |                             |
|----------------------------------|--|--------------------------------------|-----------------------------|
| (A)                              | $y - x \frac{dy}{dx} = y^2 + \frac{dy}{dx}$  | (P)                                  | $y = A_1 x^2 + A_2 x + A_3$ |
| (B)                              | $(2x - 10y^3) \frac{dy}{dx} + y = 0$   | (Q)                                  | $x^2 y^2 + 1 = cy$          |
| (C)                              | $\left( \frac{dy}{dx} \right) \left( \frac{d^3 y}{dx^3} \right) - 3 \left( \frac{d^2 y}{dx^2} \right)^2 = 0$ | (R)                                  | $(x+1)(1-y) = cy$           |
| (D)                              | $(x^2 y^2 - 1) dy + 2xy^3 dx = 0$  | (S)                                  | $x = A_1 y^2 + A_2 y + A_3$ |
|                                  |  | (T)                                  | $xy^2 = 2y^5 + c$           |

1.



Watch Video Solution

## Exercise Subjective Type Problems

1. Find the value of  $|a|$  for which the area of triangle included between the coordinate axes and any tangent to the curve  $x^a y = \lambda^a$  is constant (where  $\lambda$  is constnat.),



Watch Video Solution

2. Let  $y = f(x)$  satisfies the differential equation  $xy(1 + y)dx = dy$ . If  $f(0) = 1$  and  $f(2) = \frac{e^2}{k - e^2}$ , then find the value of  $k$ .



Watch Video Solution

3. If  $y^2 = 3\cos^2 x + 2\sin^2 x$ , then the value of  $y^4 + y^3 \frac{d^2y}{dx^2}$  is



Watch Video Solution

4. Let  $f(x)$  be a differentiable function in  $[-1, \infty)$  and  $f(0) = 1$  such

that  $\lim_{t \rightarrow x+1} \frac{t^2 f(x+1) - (x+1)^2(t)}{f(t) - f(x+1)} = 1$ . Find the value of  $\lim_{x \rightarrow 1} \frac{\ln(f(x)) - \ln 2}{x - 1}$



Watch Video Solution

5. Let  $y = (a \sin x + (b + c)\cos x)e^{x+d}$ , where a,b,c and d are parameters represent a family of curves, then differential equation for the given family of curves is given by  $y'' - \alpha y' + \beta y = 0$ , then  $\alpha + \beta =$



**Watch Video Solution**

6. Let  $y = f(x)$  satisfies the differential equation  $xy(1+y)dx = dy$ . If  $f(0) = 1$  and  $f(2) = \frac{e^2}{k - e^2}$ , then find the value of k.



**Watch Video Solution**