



MATHS

BOOKS - VIKAS GUPTA MATHS (HINGLISH)

VECTOR & 3D DIMENSIONAL GEOMETRY

Exercise 1 Single Choice Problems

1. If $ax + by + cz = p$, then minimum value of $x^2 + y^2 + z^2$ is

A. $\left(\frac{p}{a + b + c}\right)^2$

B. $\frac{p^2}{a^2 + b^2 + c^2}$

C. $\frac{a^2 + b^2 + c^2}{p^2}$

D. 0

Answer: B



2. If the angle between the vectors \vec{a} and \vec{b} is $\frac{\pi}{3}$ and the area of the triangle with adjacent sides parallel to \vec{a} and \vec{b} is 3 is

A. $\sqrt{3}$

B. $2\sqrt{3}$

C. $4\sqrt{3}$

D. $\frac{\sqrt{3}}{2}$

Answer: B



Watch Video Solution

3. Let B_1, C_1 and D_1 are points on AB, AC and AD of the parallelogram $ABCD$, such that $\overrightarrow{AB_1} = k_1 \overrightarrow{AC}, \overrightarrow{AC_1} = k_2 \overrightarrow{AC}$ and $\overrightarrow{AD_1} = k_3 \overrightarrow{AD}$, where k_1, k_2 and k_3 are scalar.

A. λ_1, λ_3 and λ_2 are in AP

B. λ_1, λ_3 and λ_2 are in GP

C. λ_1, λ_3 and λ_2 are in HP

D. $\lambda_1 + \lambda_2 + \lambda_3 = 0$

Answer: C



Watch Video Solution

4. Let $\vec{a} = 2\hat{i} + \hat{j} - 2\hat{k}$ and $\vec{b} = \hat{i} + \hat{j}$. If \vec{c} is a vector such that $\vec{a} \cdot \vec{c} = |\vec{c}|$, $|\vec{c} - \vec{a}| = 2\sqrt{2}$ and the angle between $\vec{a} \times \vec{b}$ and \vec{c} is 30° then $\left| \left(\vec{a} \times \vec{b} \right) \times \vec{c} \right|$ is equal to :

A. $\frac{2}{3}$

B. $\frac{3}{2}$

C. 2

D. 3

Answer: B



Watch Video Solution

5. If acute angle between the line $\vec{r} = \hat{i} + 2\hat{j} + \lambda(4\hat{i} - 3\hat{k})$ and xy-plane is θ_1 and acute angle between planes $x + 2y = 0$ and $2x + y = 0$ is θ_2 , then $(\cos^2 \theta_1 + \sin^2 \theta_2)$ equals to :

A. 1

B. $\frac{1}{4}$

C. $\frac{2}{3}$

D. $\frac{3}{4}$

Answer: A



Watch Video Solution

6. If a, b, c, x, y, z are real and $a^2 + b^2 + c^2 = 25, x^2 + y^2 + z^2 = 36$ and $ax + by + cz = 30$, then $\frac{a + b + c}{x + y + z}$ is equal to :

A. 1

B. $\frac{6}{5}$

C. $\frac{5}{6}$

D. $\frac{3}{4}$

Answer: C



Watch Video Solution

7. If \vec{a} and \vec{b} are non-zero, non-collinear vectors such that $|\vec{a}| = 2, \vec{a} \cdot \vec{b} = 1$ and angle between \vec{a} and \vec{b} is $\frac{\pi}{3}$. If \vec{r} is any vector such that $\vec{r} \cdot \vec{a} = 2, \vec{r} \cdot \vec{b} = 8, \left(\vec{r} + 2\vec{a} - 10\vec{b} \right) \cdot \left(\vec{a} \times \vec{b} \right) = 4\sqrt{3}$ and satisfy to $\vec{r} + 2\vec{a} - 10\vec{b} = \lambda \left(\vec{a} \times \vec{b} \right)$, then λ is equal to :

A. $\frac{1}{2}$

B. 2

C. $\frac{1}{4}$

D. None of these

Answer: D



Watch Video Solution

8. Let $\vec{a} = 3\hat{i} + 2\hat{j} + 4\hat{k}$, $\vec{b} = 2(\hat{i} + \hat{k})$ and $\vec{c} = 4\hat{i} + 2\hat{j} + 3\hat{k}$. Sum of the values of α for which the equation $x\vec{a} + y\vec{b} + z\vec{c} = \alpha(x\hat{i} + y\hat{j} + z\hat{k})$ has non-trivial solution is:

A. -1

B. 4

C. 7

D. 8

Answer: C



Watch Video Solution

9. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = \hat{i} - \hat{j} + \hat{k}$, $\vec{c} = \hat{i} + 2\hat{j} - \hat{k}$, then the value of

$$\begin{vmatrix} \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{b} \cdot \vec{a} & \vec{b} \cdot \vec{b} & \vec{b} \cdot \vec{c} \\ \vec{c} \cdot \vec{a} & \vec{c} \cdot \vec{b} & \vec{c} \cdot \vec{c} \end{vmatrix} \text{ is equal to :}$$

A. 2

B. 4

C. 16

D. 64

Answer: C



Watch Video Solution

10. \vec{a} and \vec{b} are two vectors such that $|\vec{a}| = 1$, $|\vec{b}| = 4$ and $\vec{a} \cdot \text{Vecb} = 2$. If $\vec{c} = (2\vec{a} \times \vec{b}) - 3\vec{b}$ then find angle between \vec{b} and \vec{c} .

A. $\frac{\pi}{6}$

B. $\frac{\pi}{3}$

C. $\frac{2\pi}{3}$

D. $\frac{5\pi}{3}$

Answer: D



Watch Video Solution

11. If $\vec{a}, \vec{b}, \vec{c}$ are unit vectors, then the value of $|\vec{a} - 2\vec{b}|^2 + |\vec{b} - 2\vec{c}|^2 + |\vec{c} - 2\vec{a}|^2$ does not exceed to :

A. 9

B. 12

C. 18

D. 21

Answer: D



Watch Video Solution

12. Two adjacent sides OA and OB of a rectangle $OACB$ are represented by \vec{a} and \vec{b} respectively, where o is origin. If $16\left|\vec{a} \times \vec{b}\right| = 3\left|\vec{a} + \vec{b}\right|^2$ and θ is the angle between the diagonals OC and AB, then the value(s) of $\tan\left(\frac{\theta}{2}\right)$

A. $\frac{1}{\sqrt{2}}$

B. $\frac{1}{2}$

C. $\frac{1}{\sqrt{3}}$

D. $\frac{1}{3}$

Answer: D

[Watch Video Solution](#)

13. The vectors $\overrightarrow{AB} = 3\hat{i} + 4\hat{k}$ and $\overrightarrow{AC} = 5\hat{i} - 2\hat{j} + 4\hat{k}$ are the sides of a triangle ABC. The length of the median through A is (A) $\sqrt{72}$ (B) $\sqrt{33}$ (C) $\sqrt{2880}$ (D) $\sqrt{18}$

A. $\sqrt{288}$

B. $\sqrt{72}$

C. $\sqrt{33}$

D. $\sqrt{18}$

Answer: C

[Watch Video Solution](#)

14. If $\vec{a} = 2\hat{i} + \lambda\hat{j} + 3\hat{k}$, $\vec{b} = 3\hat{i} + 3\hat{j} + 5\hat{k}$, $\vec{c} = \lambda\hat{i} + 2\hat{j} + 2\hat{k}$ are linearly dependent vectors, then the number of possible values of λ is :

A. 0

B. 1

C. 2

D. More than 2

Answer: C



Watch Video Solution

15. The scalar triple product

$\left[\vec{a} + \vec{b} - \vec{c} \quad \vec{b} + \vec{c} - \vec{a} \quad \vec{c} + \vec{a} - \vec{b} \right]$ is equal to

A. 0

B. $\left[\vec{a} \vec{b} \vec{c} \right]$

C. $2 \left[\vec{a} \vec{b} \vec{c} \right]$

D. $4 \left[\vec{a} \vec{b} \vec{c} \right]$

Answer: D



Watch Video Solution

16. If \hat{a} and \hat{b} are unit vectors then the vector defined as $\vec{V} = (\hat{a} \times \hat{b}) \times (\hat{a} + \hat{b})$ is collinear to the vector :

A. $\hat{a} + \hat{b}$

B. $\hat{b} - \hat{a}$

C. $2\hat{a} - \hat{b}$

D. $\hat{a} + 2\hat{b}$

Answer: B



Watch Video Solution

17. The sine of angle formed by the lateral face ADC and plane of the base ABC of the tetrahedron ABCD, where $A = (3, -2, 1)$, $B = (3, 1, 5)$, $C = (4, 0, 3)$ and $D = (1, 0, 0)$, is :

A. $\frac{2}{\sqrt{29}}$

- B. $\frac{5}{\sqrt{29}}$
- C. $\frac{3\sqrt{3}}{\sqrt{29}}$
- D. $\frac{-2}{\sqrt{29}}$

Answer: B



Watch Video Solution

18. Let $\vec{a}_r = x_r \hat{i} + y_r \hat{j} + z_r \hat{k}$, $r = 1, 2, 3$ three mutually perpendicular

unit vectors then the value of $\begin{vmatrix} x_1 & -x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{vmatrix}$ is equal to

- A. 0
- B. ± 1
- C. ± 2
- D. ± 4

Answer: B



Watch Video Solution

19. If \vec{a} , \vec{b} and \vec{c} are three non coplanar vectors and \vec{r} is any vector in space, then

$$\left(\vec{a} \times \vec{b} \right), \left(\vec{r} \times \vec{c} \right) + \left(\vec{b} \times \vec{c} \right) \times \left(\vec{r} \times \vec{a} \right) + \left(\vec{c} \times \vec{a} \right) \times \left(\vec{r} \times \vec{b} \right)$$

(A) $\left[\vec{a} \vec{b} \vec{c} \right] \vec{r}$ (B) $2 \left[\vec{a} \vec{b} \vec{c} \right] \vec{r}$ (C) $3 \left[\vec{a} \vec{b} \vec{c} \right] \vec{r}$ (D) $4 \left[\vec{a} \vec{b} \vec{c} \right] \vec{r}$

A. $\left[\vec{a} \vec{b} \vec{c} \right] \vec{r}$

B. $2 \left[\vec{a} \vec{b} \vec{c} \right] \vec{r}$

C. $4 \left[\vec{a} \vec{b} \vec{c} \right] \vec{r}$

D. 0

Answer: B



Watch Video Solution

20. E and F are the interior points on the sides BC and CD of a parallelogram ABCD. Let $\vec{BE} = 4\vec{EC}$ and $\vec{CF} = 4\vec{FD}$. If the line EF

meets the diagonal AC in G, then $\overrightarrow{AG} = \lambda \overrightarrow{AC}$, where λ is equal to :

A. $\frac{1}{3}$

B. $\frac{21}{25}$

C. $\frac{7}{13}$

D. $\frac{21}{5}$

Answer: B



Watch Video Solution

21. If \vec{a} and \vec{b} are unit vectors and \vec{c} is such that \vec{c} is such that $\vec{c} = \vec{c} = \vec{a} \times \vec{c} + \vec{b}$ then maximum value of $\left[\vec{a} \vec{b} \vec{c} \right]$ is

A. 1

B. $\frac{1}{2}$

C. 2

D. $\frac{3}{2}$

Answer: B



Watch Video Solution

22. Consider the matrices $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 1 & 2 \\ 1 & -1 & 1 \end{bmatrix}$ $B = \begin{bmatrix} 2 & 1 & 3 \\ 4 & 1 & -1 \\ 2 & 2 & 3 \end{bmatrix}$
 $C = \begin{bmatrix} 14 \\ 12 \\ 2 \end{bmatrix}$ $D = \begin{bmatrix} 13 \\ 11 \\ 14 \end{bmatrix}$. Now $x = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ is such that solutions of
equation $AX = C$ and $BX = D$ represent two points L and M
respectively in 3 dimensional space. If L' and M' are reflections of L
and M in the plane $x+y+z=9$ then find coordinates of L, M, L', M'

A. (3, 4, 2)

B. (5, 3, 4)

C. (7, 2, 3)

D. (1, 5, 6)

Answer: A



Watch Video Solution

23. The value of α for which point $M(\alpha\hat{i} + 2\hat{j} + \hat{k})$, lie in the plane containing three points $A(\hat{i} + \hat{j} + \hat{k})$ and $C(3\hat{i} - \hat{k})$ is :

A. 1

B. 2

C. $\frac{1}{2}$

D. $-\frac{1}{2}$

Answer: B



Watch Video Solution

24. Q is the image of point P(1, -2, 3) with respect to the plane $x - y + z = 7$. The distance of Q from the origin is.

A. $\sqrt{\frac{70}{3}}$

B. $\frac{1}{2}\sqrt{\frac{70}{3}}$

C. $\sqrt{\frac{35}{3}}$

D. $\sqrt{\frac{15}{2}}$

Answer: A



Watch Video Solution

25. \hat{a} , \hat{b} and $\hat{a} - \hat{b}$ are unit vectors. The volume of the parallelopiped, formed with \hat{a} , \hat{b} and $\hat{a} \times \hat{b}$ as coterminous edges is :

A. 1

B. $\frac{1}{4}$

C. $\frac{2}{3}$

D. $\frac{3}{4}$

Answer: D



Watch Video Solution

26. A line passing through $P(3, 7, 1)$ and $R(2, 5, 7)$ meet the plane $3x + 2y + 11z - 9 = 0$ at Q . Then PQ is equal to :

- A. $\frac{5\sqrt{41}}{59}$
- B. $\frac{\sqrt{41}}{59}$
- C. $\frac{50\sqrt{41}}{59}$
- D. $\frac{25\sqrt{41}}{59}$

Answer: D



Watch Video Solution

27. Let $\vec{a}, \vec{b}, \vec{c}$ be three non-zero non coplanar vectors and \vec{p}, \vec{q} and \vec{r} be three vectors given by $\vec{p} = \vec{a} + \vec{b} - 2\vec{c}, \vec{q} = 3\vec{a} - 2\vec{b} + \vec{c}$ and $\vec{r} = \vec{a} - 4\vec{b} + 2\vec{c}$

If the volume of the parallelopiped determined by \vec{a}, \vec{b} and \vec{c} is V_1 and that of the parallelopiped determined by \vec{a}, \vec{q} and \vec{r} is V_2 , then $V_2 : V_1 =$

A. 10

B. 15

C. 20

D. None of these

Answer: B



Watch Video Solution

28. If the two lines represented by $x + ay = b$, $z + cy = d$ and $x = a'y + b'$, $z = c'y + d'$ be perpendicular to each other, then the value of $aa' + cc'$ is :

A. 1

B. 2

C. 3

D. 4

Answer: A



Watch Video Solution

29. The distance between the line

$\vec{r} = 2\hat{i} - 2\hat{j} + 3\hat{k} + \lambda(\hat{i} - \hat{j} + 4\hat{k})$ and the plane

$\vec{r} \cdot (\hat{i} + 5\hat{j} + \hat{k}) = 5$ is

A. $\frac{10}{9}$

B. $\frac{10}{3\sqrt{3}}$

C. $\frac{3}{10}$

D. $\frac{10}{3}$

Answer: B



Watch Video Solution

30. If $(\vec{a} \times \vec{b}) \times \vec{c} = \vec{a} \times (\vec{b} \times \vec{c})$, where \vec{a} , \vec{b} and \vec{c} are any three vectors such that $\vec{a} \cdot \vec{b} \neq 0$, $\vec{b} \cdot \vec{c} \neq 0$, then \vec{a} and \vec{c} are :

A. Inclined at an angle of $\frac{\pi}{3}$

B. Inclined at an angle of $\frac{\pi}{6}$

C. Perpendicular

D. Parallel

Answer: D



Watch Video Solution

31. Let \vec{r} be position vector of variable point in cartesian plane OXY such that $\vec{r} \cdot (\vec{r} + 6\hat{j}) = 7$ cuts the co-ordinate axes at four distinct points, then the area of the quadrilateral formed by joining these points is :

A. $4\sqrt{7}$

B. $6\sqrt{7}$

C. $7\sqrt{7}$

D. $8\sqrt{7}$

Answer: D



Watch Video Solution

32. If $|\vec{a}| = 2$, $|\vec{b}| = 5$ and $\vec{a} \cdot \vec{b} = 0$, then $\vec{a} \times \left(\vec{a} \times \left(\vec{a} \times \left(\vec{a} \times \left(\vec{a} \times \left(\vec{a} \times \left(\vec{a} \times \vec{b} \right) \right) \right) \right) \right) \right) \right)$ is equal to :

A. $64\vec{a}$

B. $64\vec{b}$

C. $-64\vec{a}$

D. $-64\vec{b}$

Answer: D



Watch Video Solution

33. If O (origin) is a point inside the triangle PQR such that $\vec{OP} + k_1 \vec{OQ} + k_2 \vec{OR} = 0$, where k_1, k_2 are constants such that $\frac{\text{Area}(\Delta PQR)}{\text{Area}(\Delta OQR)} = 4$, then the value of $k_1 + k_2$ is :

A. 2

B. 3

C. 4

D. 5

Answer: B



Watch Video Solution

34. Let PQ and QR be diagonals of adjacent faces of a rectangular box, with its centre at O. If $\angle QOR, \angle ROP$ and $\angle POQ$ are θ, ϕ and Ψ respectively then the value of ' $\cos \theta + \cos \phi + \cos \Psi$ ' is :

A. -2

B. $-\sqrt{3}$

C. -1

D. 0

Answer: C



Watch Video Solution

35. The value of $\begin{vmatrix} \vec{a} & \vec{b} & \vec{c} \\ \vec{a} \cdot \vec{p} & \vec{b} \cdot \vec{p} & \vec{c} \cdot \vec{p} \\ \vec{a} \cdot \vec{q} & \vec{b} \cdot \vec{q} & \vec{c} \cdot \vec{q} \end{vmatrix}$ is equal to :

A. $(\vec{p} \times \vec{q}) \begin{bmatrix} \vec{a} \times \vec{b} & \vec{b} \times \vec{c} & \vec{c} \times \vec{a} \end{bmatrix}$

B. $2(\vec{p} \times \vec{q}) \begin{bmatrix} \vec{a} \times \vec{b} & \vec{b} \times \vec{c} & \vec{c} \times \vec{a} \end{bmatrix}$

C. $4(\vec{p} \times \vec{q}) \begin{bmatrix} \vec{a} \times \vec{b} & \vec{b} \times \vec{c} & \vec{c} \times \vec{a} \end{bmatrix}$

D. $(\vec{p} \times \vec{q}) \sqrt{\begin{bmatrix} \vec{a} \times \vec{b} & \vec{b} \times \vec{c} & \vec{c} \times \vec{a} \end{bmatrix}}$

Answer: D



View Text Solution

36.

If

$$\vec{r} = a(\vec{m} \times \vec{n}) + b(\vec{n} \times \vec{l}) + c(\vec{l} \times \vec{m}) \text{ and } [\vec{l} \vec{m} \vec{n}] = 4, \text{ find}$$

:

A. $\frac{1}{4}$

B. $\frac{1}{2}$

C. 1

D. 2

Answer: A**Watch Video Solution**

37. The volume of tetrahedron, for which three co-terminus edges are

$$\vec{a} - \vec{b}, \vec{b} + 2\vec{c} \text{ and } 3\vec{a} - \vec{c} \text{ is :}$$

A. 6k

B. 7k

C. 30k

D. 42k

Answer: D



Watch Video Solution

38. The equation of a plane passing through the line of intersection of the planes :

$x + 2y + z - 10 = 0$ and $3x + y - z = 5$ and passing through the origin is :

A. $5x + 3z = 0$

B. $5x - 3z = 0$

C. $5x + 4y + 3z = 0$

D. $5x - 4y + 3z = 0$

Answer: B



Watch Video Solution

39. Find the locus of a point whose distance from x-axis is equal the distance from the point $(1, -1, 2)$:

A. $y^2 + 2x - 2y - 4z + 6 = 0$

B. $x^2 + 2x - 2y - 4z + 6 = 0$

C. $x^2 - 2x + 2y - 4z + 6 = 0$

D. $z^2 - 2x + 2y - 4z + 6 = 0$

Answer: C



Watch Video Solution

Exercise 2 One Or More Than One Answer Is Are Correct

1. If equation of three lines are :

$$\frac{x}{1} = \frac{y}{2} = \frac{z}{3}, \frac{x}{2} = \frac{y}{1} = \frac{z}{3} \text{ and } \frac{x-1}{1} = \frac{2-y}{1} = \frac{z-3}{0}, \text{ then}$$

which of the following statement(s) is/are correct ?

- A. Triangle formed by the line is equilateral
- B. Triangle formed by the lines is isosceles
- C. Equation of the plane containing the lines is $x + y = z$
- D. Area of the triangle formed by the lines is $\frac{3\sqrt{3}}{2}$

Answer: B::C::D



Watch Video Solution

2. If $\vec{a} = \hat{i} + 6\hat{j} + 3\hat{k}; \vec{b} = 3\hat{i} + 2\hat{j} + \hat{k}$ and $\vec{c} = (\alpha + 1)\hat{i} + (\beta - 1)\hat{j} + \hat{k}$ are linearly dependent vectors and $|\vec{c}| = \sqrt{6}$; then the possible value(s) of $(\alpha + \beta)$ can be:

A. 1

B. 2

C. 3

D. 4

Answer: A::C



Watch Video Solution

3. Consider the lines:
 $L_1: \frac{x-2}{1} = \frac{y-1}{7} = \frac{z+2}{-5}, L_2: x-4 = y+3 = -z$ Then which of the following is/are correct ? (A) Point of intersection of L_1 and L_2 is $(1, -6, 3)$

A. Point of intersection of L_1 and L_2 is $(1, -6, 3)$

B. Equation of plane containing L_1 and L_2 is $x + 2y + 3z + 2 = 0$

C. Acute angle between L_1 and L_2 is $\cot^{-1}\left(\frac{13}{15}\right)$

D. Equation of plane containing L_1 and L_2 is $x + 2y + 2z + 3 = 0$

Answer: A::B::C



Watch Video Solution

4. Let \hat{a} , \hat{b} and \hat{c} be three unit vectors such that $\hat{a} = \hat{b} + (\hat{b} \times \hat{c})$, then the possible value(s) of $|\hat{a} + \hat{b} + \hat{c}|^2$ can be :

A. 1

B. 4

C. 16

D. 9

Answer: A::D



Watch Video Solution

5. The value(s) of μ for which the straight lines $\vec{r} = 3\hat{i} - 2\hat{j} - 4\hat{k} + \lambda_1(\hat{i} - \hat{j} + \mu\hat{k})$ and

$\vec{r} = 5\hat{i} - 2\hat{j} + \hat{k} + \lambda_2(\hat{i} + \mu\hat{j} + 2\hat{k})$ are coplanar is/are :

A. $\frac{5 + \sqrt{33}}{4}$

B. $\frac{-5 + \sqrt{33}}{4}$

C. $\frac{5 - \sqrt{33}}{4}$

D. $\frac{-5 - \sqrt{33}}{4}$

Answer: A::C



Watch Video Solution

6.

If

$$\hat{i} \times \left[\left(\vec{a} - \hat{j} \right) \times \hat{i} \right] \times \left[\left(\vec{a} - \hat{k} \right) \times \hat{j} \right] + \vec{k} \times \left[\left(\vec{a} - \vec{i} \right) \times \hat{k} \right] = 0,$$

then find vector \vec{a} .

A. $x + y = 1$

B. $y + z = \frac{1}{2}$

C. $x + z = 1$

D. None of these

Answer: A::C



Watch Video Solution

7. The value of expression $\left[\vec{a} \times \vec{b} \quad \vec{c} \times \vec{d} \quad \vec{e} \times \vec{f} \right]$ is equal to :

A. $\left[\vec{a} \vec{b} \vec{d} \right] \left[\vec{c} \vec{e} \vec{f} \right] - \left[\vec{a} \vec{b} \vec{c} \right] \left[\vec{d} \vec{e} \vec{f} \right]$

B. $\left[\vec{a} \vec{b} \vec{e} \right] \left[\vec{f} \vec{c} \vec{d} \right] - \left[\vec{a} \vec{b} \vec{f} \right] \left[\vec{e} \vec{c} \vec{d} \right]$

C. $\left[\vec{c} \vec{d} \vec{a} \right] \left[\vec{b} \vec{e} \vec{f} \right] - \left[\vec{c} \vec{d} \vec{b} \right] \left[\vec{a} \vec{e} \vec{f} \right]$

D. $\left[\vec{b} \vec{c} \vec{d} \right] \left[\vec{a} \vec{e} \vec{f} \right] - \left[\vec{b} \vec{c} \vec{f} \right] \left[\vec{a} \vec{e} \vec{d} \right]$

Answer: A::B::C



Watch Video Solution

8. If \vec{a} , \vec{b} , \vec{c} and \vec{d} are the position vectors of the points A, B, C and D respectively in three dimensional space no three of A, B, C, D are collinear and satisfy the relation $3\vec{a} - 2\vec{b} + \vec{c} - 2\vec{d} = 0$, then

A. A, B, C and D are coplanar

B. The line joining the points B and D divides the line joining the point A and C in the ratio of 2:1

C. The line joining the points A and C divides the line joining the points B and D in the ratio of 1:1

D. The four vectors \vec{a} , \vec{b} , \vec{c} and \vec{d} are linearly dependent .

Answer: A::C::D



Watch Video Solution

9. If OAB is a tetrahedron with edges and \hat{p} , \hat{q} , \hat{r} are unit vectors along bisectors of

$$\overrightarrow{OA}, \overrightarrow{OB}, \overrightarrow{OC}, \overrightarrow{OA}$$

respectively

and

$$\hat{a} = \frac{\overrightarrow{OA}}{|\overrightarrow{OA}|}, \hat{b} = \frac{\overrightarrow{OB}}{|\overrightarrow{OB}|}, \hat{c} = \frac{\overrightarrow{OC}}{|\overrightarrow{OC}|}, \text{ then :}$$

$$\text{A. } \frac{[\hat{a}\hat{b}\hat{c}]}{[\hat{p}\hat{q}\hat{r}]} = \frac{3\sqrt{3}}{2}$$

$$\text{B. } \frac{\begin{bmatrix} \hat{a} + \hat{b} & \hat{b} + \hat{c} & \hat{c} + \hat{a} \end{bmatrix}}{\begin{bmatrix} \hat{p} + \hat{q} & \hat{q} + \hat{r} & \hat{r} + \hat{p} \end{bmatrix}} = \frac{3\sqrt{3}}{4}$$

$$\text{C. } \frac{\begin{bmatrix} \hat{a} + \hat{b} & \hat{b} + \hat{c} & \hat{c} + \hat{a} \end{bmatrix}}{[\hat{p}\hat{q}\hat{r}]} = \frac{3\sqrt{3}}{2}$$

$$\text{D. } \frac{[\hat{a}\hat{b}\hat{c}]}{\begin{bmatrix} \hat{p} + \hat{q} & \hat{q} + \hat{r} & \hat{r} + \hat{p} \end{bmatrix}} = \frac{3\sqrt{3}}{4}$$

Answer: A::D



Watch Video Solution

10. Let \hat{a} and \hat{c} are unit vectors and $|\vec{b}| = 4$. If the angle between \hat{a} and \hat{c} is $\cos^{-1}\left(\frac{1}{4}\right)$, and $\hat{b} - 2\hat{c} = \lambda\hat{a}$, then the value of λ can be :

A. 2

B. 3

C. -3

D. 4

Answer: C::D



Watch Video Solution

11. Consider the lines $x = y = z$ and line $2x + y + z - 1 = 0 = 3x + y + 2z - 2$, then

A. The shortest distance between the two lines is $\frac{1}{\sqrt{2}}$

B. The shortest distance between the two lines is $\sqrt{2}$

C. Plane containing the line L_2 and parallel to line L_1 is

$$z - x + 1 = 0$$

D. Perpendicular distance of origin from plane containing line L_2 and

parallel to line L_1 is $\frac{1}{\sqrt{2}}$

Answer: A::D

[Watch Video Solution](#)

12. Let $\vec{r} = \sin x (\vec{a} \times \vec{b}) + \cos y (\vec{b} \times \vec{c}) + 2(\vec{c} \times \vec{a})$, where \vec{a}, \vec{b} are non-collinear and \vec{c}, \vec{d} are also non-collinear then :

A. π^2

B. $\frac{5\pi^2}{4}$

C. $\frac{35\pi^2}{4}$

D. $\frac{37\pi^2}{4}$

Answer: B::D

[Watch Video Solution](#)

13. If $(\vec{a} \times \vec{b}) \times (\vec{c} \times \vec{d}) = h\vec{a} + k\vec{b} = r\vec{c} + s\vec{d}$, where \vec{a}, \vec{b} are non-collinear and \vec{c}, \vec{d} are also non-collinear then :

A. $h = \begin{bmatrix} \vec{b} & \vec{c} & \vec{d} \end{bmatrix}$

$$\text{B. } k = \begin{bmatrix} \vec{a} & \vec{c} & \vec{d} \end{bmatrix}$$

$$\text{C. } r = \begin{bmatrix} \vec{a} & \vec{b} & \vec{d} \end{bmatrix}$$

$$\text{D. } s = - \begin{bmatrix} \vec{a} & \vec{b} & \vec{c} \end{bmatrix}$$

Answer: B::C::D



Watch Video Solution

14. Let a be a real number and $\vec{\alpha} = \hat{i} + 2\hat{j}$, $\vec{\beta} = 2\hat{i} + a\hat{j} + 10\hat{k}$, $\vec{\gamma} = 12\hat{i} + 20\hat{j} + a\hat{k}$ be three vectors, then $\vec{\alpha}$, $\vec{\beta}$ and $\vec{\gamma}$ are linearly independent for :

A. $a > 0$

B. $a < 0$

C. $a = 0$

D. No value of a

Answer: A::B::C

[Watch Video Solution](#)

15. The volume of a tetrahedron prism $ABCA_1B_1C_1$ is equal to 3. Find the coordinates of the vertex A_1 , if the coordinate of the base vertices of the prism are $A(1, 0, 1)$, $B(2, 0, 0)$ and $C(0, 1, 0)$.

A. (2, 2, 2)

B. (0, 2, 0)

C. (0, -2, 2)

D. (0, -2, 0)

Answer: A::D

[Watch Video Solution](#)

16. If $\vec{a} = x\hat{i} + y\hat{j} + z\hat{k}$, $\vec{b} = y\hat{i} + z\hat{j} + x\hat{k}$, and $\vec{c} = z\hat{i} + x\hat{j} + y\hat{k}$, then $\vec{a} \times (\vec{b} \times \vec{c})$ is :

A. Parallel to $(y - z)\hat{i} + (z - x)\hat{j} + (x - y)\hat{k}$

B. Orthogonal to $\hat{i} + \hat{j} + \hat{k}$

C. Orthogonal to $(y + z)\hat{i} + (z + x)\hat{j} + (x + y)\hat{k}$,

D. Orthogonal to $x\hat{i} + y\hat{j} + z\hat{k}$

Answer: A::B::C::D



Watch Video Solution

17. If a line has a vector equation, $\vec{r} = 2\hat{i} + 6\hat{j} + \lambda(\hat{i} - 3\hat{j})$ then which of the following statements holds good ?

A. the line is parallel to $2\hat{i} + 6\hat{j}$

B. the line passes through the point $3\hat{i} + 3\hat{j}$

C. the line passes through the point $\hat{i} + 9\hat{j}$

D. the line is parallel to xy plane

Answer: B::C::D

 [Watch Video Solution](#)

18. Let M, N, P and Q be the mid points of the edges AB, CD, AC and BD respectively of the tetrahedron ABCD. Further, MN is perpendicular to both AB and CD and PQ is perpendicular to both AC and BD. Then which of the following is/are correct:

A. $AB = CD$

B. $BC = DA$

C. $AC = BD$

D. $AN = BN$

Answer: A::B::C::D

 [Watch Video Solution](#)

19. The solution vectors \vec{r} of the equation $\vec{r} \times \hat{i} = \hat{j} + \hat{k}$ and $\vec{r} \times \hat{j} = \hat{k} + \hat{i}$ represent two straight lines which

are :

- A. Intersecting
- B. Non coplanar
- C. Coplanar
- D. Non intersecting

Answer: B::D



Watch Video Solution

20. Which of the following statement(s) is/are incorrect ?

A. The lines

$$\frac{x-4}{-3} = \frac{y+6}{-1} = \frac{z+6}{-1} \text{ and } \frac{x-1}{-1} = \frac{y-2}{-2} = \frac{z-3}{2} \text{ are}$$

orthogonal

B. The planes $3x - 2y - 4z = 3$ and the plane $x - y - z = 3$ are

orthogonal

C. The function $f(x) = \ln(e^{-2} + e^x)$ is monotonic increasing

$$\forall x \in \mathbb{R}$$

D. If g is the inverse of the function,

$$f(x) = \ln(e^{-2} + e^x) \text{ then } g(x) = \ln(e^x - e^{-2})$$

Answer: A::B



Watch Video Solution

21. The lines with vector equations are,

$$\vec{r}_1 = 3\hat{i} + 6\hat{j} + \lambda(-4\hat{i} + 3\hat{j} + 2\hat{k}) \text{ and } \vec{r}_2 = -2\hat{i} + 7\hat{j} + \mu(-4\hat{i} +$$

are such that :

A. they are coplanar

B. they do not intersect

C. they are skew

D. the angle between them is $\tan^{-1}(3/7)$

Answer: B::C::D



View Text Solution

Exercise 3 Comprehension Type Problems

1. The vertices of $\triangle ABC$ are $(2, 0, 0)$, $B(0, 1, 0)$, $C(0, 0, 2)$. Its orthocentre is H and circumcentre is S. P is a point equidistant from A, B, C and the origin O.

Q. The z-coordinate of H is :

A. 1

B. $1/2$

C. $1/6$

D. $1/3$

Answer: D



View Text Solution

2. The vertices of $\triangle ABC$ are $(2, 0, 0)$, $B(0, 1, 0)$, $C(0, 0, 2)$. Its orthocentre is H and circumcentre is S. P is a point equidistant from A, B, C and the origin O.

Q. The y-coordinate of S is :

A. $5/6$

B. $1/3$

C. $1/6$

D. $1/2$

Answer: C



View Text Solution

3. The vertices of $\triangle ABC$ are $(2, 0, 0)$, $B(0, 1, 0)$, $C(0, 0, 2)$. Its orthocentre is H and circumcentre is S. P is a point equidistant from A, B, C and the

origin O.

Q. PA is equal to :

A. 1

B. $\sqrt{2}$

C. $\sqrt{\frac{3}{2}}$

D. $\frac{3}{2}$

Answer: D



View Text Solution

4. Consider a plane $\pi: \vec{r} \cdot \vec{n} = d$ (where \vec{n} is not a unit vector). There are two points $A(\vec{a})$ and $B(\vec{b})$ lying on the same side of the plane.

Q. If foot of perpendicular from A and B to the plane π are P and Q respectively, then length of PQ be :

A. $\frac{\left| \left(\vec{b} - \vec{a} \right) \cdot \vec{n} \right|}{\left| \vec{n} \right|}$

B. $\left| \left(\vec{b} - \vec{a} \right) \cdot \vec{n} \right|$

C. $\frac{\left| \left(\vec{b} - \vec{a} \right) \times \vec{n} \right|}{\left| \vec{n} \right|}$

D. $\left| \left(\vec{b} - \vec{a} \right) \times \vec{n} \right|$

Answer: C



Watch Video Solution

5. Consider a plane $\pi: \vec{r} \cdot \vec{n} = d$ (where \vec{n} is not a unit vector). There are two points $A(\vec{a})$ and $B(\vec{b})$ lying on the same side of the plane.

Q. Reflection of $A(\vec{a})$ in the plane π has the position vector :

A. $\vec{a} + \frac{2}{(\vec{n})^2} (d - \vec{a} \cdot \vec{n}) \vec{n}$

B. $\vec{a} - \frac{1}{(\vec{n})^2} (d - \vec{a} \cdot \vec{n}) \vec{n}$

C. $\vec{a} + \frac{2}{(\vec{n})^2} (d + \vec{a} \cdot \vec{n}) \vec{n}$

D. $\vec{a} + \frac{2}{(\vec{n})^2} \vec{n}$

Answer: A



View Text Solution

6. Consider a plane $\pi: \vec{r} \cdot \vec{n} = d$ (where \vec{n} is not a unit vector). There are two points $A(\vec{a})$ and $B(\vec{b})$ lying on the same side of the plane.

Q. If a plane π_1 is drawn from the point $A(\vec{a})$ and another plane π_2 is drawn from point $B(\vec{b})$ parallel to π , then the distance between the planes π_1 and π_2 is :

A. $\frac{\left| \left(\vec{a} - \vec{b} \right) \cdot \vec{n} \right|}{\left| \vec{n} \right|}$

B. $\left| \left(\vec{a} - \vec{b} \right) \cdot \vec{n} \right|$

C. $\left| \left(\vec{a} - \vec{b} \right) \times \vec{n} \right|$

D. $\frac{\left| \left(\vec{a} - \vec{b} \right) \times \vec{n} \right|}{\left| \vec{n} \right|}$

Answer: A



View Text Solution

7. Consider a plane $\Pi: \vec{r} \cdot (2\hat{i} + \hat{j} - \hat{k}) = 5$, a line $L_1: \vec{r} = (3\hat{i} - \hat{j} + 2\hat{k}) + \lambda(2\hat{i} - 3\hat{j} - \hat{k})$ and a point $a(3, -4, 1) \cdot L_2$ is a line passing through A intersecting L_1 and parallel to plane Π .

Q. Equation of L_2 is :

A. $\vec{r} = (1 + \lambda)\hat{i} + (2 - 3\lambda)\hat{j} + (1 - \lambda)\hat{k}; \lambda \in R$

B. $\vec{r} = (3 + \lambda)\hat{i} - (4 - 2\lambda)\hat{j} + (1 + 3\lambda)\hat{k}, \lambda \in R$

C. $\vec{r} = (3 + \lambda)\hat{i} - (4 + 3\lambda)\hat{j} + (1 - \lambda)\hat{k}, \lambda \in R$

D. None of the above

Answer: C



View Text Solution

8. Consider a plane $\Pi: \vec{r} \cdot (2\hat{i} + \hat{j} - \hat{k}) = 5$, a line $L_1: \vec{r} = (3\hat{i} - \hat{j} + 2\hat{k}) + \lambda(2\hat{i} - 3\hat{j} - \hat{k})$ and a point

$a(3, -4, 1) \cdot L_2$ is a line passing through A intersecting L_1 and parallel to plane Π .

Q. Plane containing L_1 and L_2 is :

- A. parallel to yz-plane
- B. parallel to x-axis
- C. parallel to y-axis
- D. passing through origin

Answer: B



View Text Solution

9. Consider a plane $\Pi : \vec{r} \cdot (2\hat{i} + \hat{j} - \hat{k}) = 5$, a line $L_1 : \vec{r} = (3\hat{i} - \hat{j} + 2\hat{k}) + \lambda(2\hat{i} - 3\hat{j} - \hat{k})$ and a point $a(3, -4, 1) \cdot L_2$ is a line passing through A intersecting L_1 and parallel to plane Π .

Q. Line L_1 intersects plane Π at Q and xy-plane at R the volume of

tetrahedron OAQR is :

(where 'O' is origin)

A. 0

B. $\frac{14}{3}$

C. $\frac{3}{7}$

D. $\frac{7}{3}$

Answer: D



View Text Solution

10. Consider three planes :

$$2x + py + 6z = 8, x + 2y + qz = 5 \text{ and } x + y + 3z = 4$$

Q. Three planes intersect at a point if :

A. $p = 2, q \neq 3$

B. $p \neq 2, q \neq 3$

C. $p \neq 2, q = 3$

D. $p = 2, q = 3$

Answer: B



Watch Video Solution

11. Consider three planes :

$$2x + py + 6z = 8, x + 2y + qz = 5 \text{ and } x + y + 3z = 4$$

Q. Three planes do not have any common point of intersection if :

A. $p = 2, q \neq 3$

B. $p \neq 2, q \neq 3$

C. $p \neq 2, q = 3$

D. $p = 2, q = 3$

Answer: C



Watch Video Solution

12. The points A, B and C with position vectors \vec{a} , \vec{b} and \vec{c} respectively lie on a circle centered at origin O. Let G and E be the centroid of $\triangle ABC$ and $\triangle ACD$ respectively where D is mid point of AB.

Q. If OE and CD are mutually perpendicular, then which of the following will be necessarily true ?

A. $\left| \vec{b} - \vec{a} \right| = \left| \vec{c} - \vec{a} \right|$

B. $\left| \vec{b} - \vec{a} \right| = \left| \vec{b} - \vec{c} \right|$

C. $\left| \vec{c} - \vec{a} \right| = \left| \vec{c} - \vec{b} \right|$

D. $\left| \vec{b} - \vec{a} \right| = \left| \vec{c} - \vec{a} \right| = \left| \vec{b} - \vec{c} \right|$

Answer: A



View Text Solution

13. The points A, B and C with position vectors \vec{a} , \vec{b} and \vec{c} respectively lie on a circle centered at origin O. Let G and E be the centroid of $\triangle ABC$ and $\triangle ACD$ respectively where D is mid point of AB.

Q. If GE and CD are mutually perpendicular, then orthocenter of ΔABC must lie on :

- A. median through A
- B. median through C
- C. angle bisector through A
- D. angle bisector through B

Answer: B



View Text Solution

14. The points A, B and C with position vectors \vec{a} , \vec{b} and \vec{c} respectively lie on a circle centered at origin O. Let G and E be the centroid of ΔABC and ΔACD respectively where D is mid point of AB.

Q. If $\left[\vec{AB} \vec{AC} \vec{AB} \times \vec{AC} \right] = \lambda \left[\vec{AE} \vec{AG} \vec{AE} \times \vec{AG} \right]$, then the value of λ is :

A. -18

B. 18

C. -324

D. 324

Answer: D



View Text Solution

15. Consider a tetrahedron $D - ABC$ with position vectors of its angular points as

$A(1, 1, 1), B(1, 2, 3), C(1, 1, 2)$

and centre of tetrahedron $\left(\frac{3}{2}, \frac{3}{4}, 2\right)$.

Q. Shortest distance between the skew lines AB and CD :

A. $\frac{1}{2}$

B. $\frac{1}{3}$

C. $\frac{1}{4}$

D. $\frac{1}{5}$

Answer: B



[View Text Solution](#)

16. Consider a tetrahedron $D - ABC$ with position vectors of its angular points as

$A(1, 1, 1), B(1, 2, 3), C(1, 1, 2)$

and centre of tetrahedron $\left(\frac{3}{2}, \frac{3}{4}, 2\right)$.

Q. If N be the foot of the perpendicular from point D on the plane face ABC then the position vector of N are :

A. $(-1, 1, 2)$

B. $(1, -1, 2)$

C. $(1, 1, -2)$

D. $(-1, -1, 2)$

Answer: B



[View Text Solution](#)

17. In a triangle AOB, R and Q are the points on the side OB and AB respectively such that $3OR = 2RB$ and $2AQ = 3QB$. Let OQ and AR intersect at the point P (where O is origin).

Q. If the point P divides OQ in the ratio of $\mu : 1$, then μ is :

A. $\frac{2}{19}$

B. $\frac{2}{17}$

C. $\frac{2}{15}$

D. $\frac{10}{9}$

Answer: D



View Text Solution

18. In a triangle AOB, R and Q are the points on the side OB and AB respectively such that $3OR = 2RB$ and $2AQ = 3QB$. Let OQ and AR intersect at the point P (where O is origin).

Q. If the ratio of area of quadrilateral PQBR and area of $\triangle OPA$ is $\frac{\alpha}{\beta}$ then $(\beta - \alpha)$ is (where α and β are coprime numbers) :

A. 1

B. 9

C. 7

D. 0

Answer: D



View Text Solution

Exercise 4 Matching Type Problems

| Column-I | | Column-II | |
|----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|-----|
| (A) | If \vec{a} , \vec{b} and \vec{c} are three mutually perpendicular vectors where $ \vec{a} = \vec{b} = 2$, $ \vec{c} = 1$, then $\{\vec{a} \times \vec{b} \quad \vec{b} \times \vec{c} \quad \vec{c} \times \vec{a}\}$ is | (P) | -12 |
| (B) | If \vec{a} and \vec{b} are two unit vectors inclined at $\frac{\pi}{3}$, then $16[\vec{a} \quad \vec{b} + (\vec{a} \times \vec{b}) \quad \vec{b}]$ is | (Q) | 0 |
| (C) | If \vec{b} and \vec{c} are orthogonal unit vectors and $\vec{b} \times \vec{c} = \vec{a}$ then $[\vec{a} + \vec{b} + \vec{c} \quad \vec{a} + \vec{b} \quad \vec{b} + \vec{c}]$ is | (R) | 16 |
| (D) | If $[\vec{x} \quad \vec{y} \quad \vec{a}] = [\vec{x} \quad \vec{y} \quad \vec{b}] = [\vec{a} \quad \vec{b} \quad \vec{c}] = 0$, each vector being a non-zero vector, then $[\vec{x} \quad \vec{y} \quad \vec{c}]$ is | (S) | 1 |
| | | (T) | 4 |

1.



Watch Video Solution

Exercise 5 Subjective Type Problems

1. A straight line L intersects perpendicularly both the lines :

$$\frac{x+2}{2} = \frac{y+6}{3} = \frac{z-34}{-10} \quad \text{and} \quad \frac{x+6}{4} = \frac{y-7}{-3} = \frac{z-7}{-2},$$

then the square of perpendicular distance of origin from L is



View Text Solution

2. If \hat{a} , \hat{b} and \hat{c} are non-coplanar unit vectors such that $[\hat{a}\hat{b}\hat{c}] = [\hat{b} \times \hat{c} \quad \hat{c} \times \hat{a} \quad \hat{a} \times \hat{b}]$, then find the projection of $\hat{b} + \hat{c}$ on $\hat{a} \times \hat{b}$.



Watch Video Solution

3. Let OA, OB, OC be coterminal edges of a cuboid. If l, m, n be the shortest distances between the sides OA, OB, OC and their respective skew body diagonals to them, respectively, then find

$$\frac{\left(\frac{1}{l^2} + \frac{1}{m^2} + \frac{1}{n^2}\right)}{\left(\frac{1}{OA^2} + \frac{1}{OB^2} + \frac{1}{OC^2}\right)}.$$



View Text Solution

4. Let OABC be a tetrahedron whose edges are of unit length. If $\vec{OA} = \vec{a}$, $\vec{OB} = \vec{b}$, and $\vec{OC} = \alpha(\vec{a} + \vec{b}) + \beta(\vec{a} \times \vec{b})$, then $(\alpha\beta)^2 = \frac{p}{q}$ (where p & q are relatively prime to each other). then the value of $\left[\frac{q}{2}p\right]$ is



Watch Video Solution

5. Let \vec{v}_0 be a fixed vector and $\vec{v}_0 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$. Then for $n \geq 0$ a sequence is defined $\vec{v}_{n+1} = \vec{v}_n + \left(\frac{1}{2}\right)^{n+1} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}^{n+1} \vec{v}_0$ then $\lim_{n \rightarrow \infty} \vec{v}_n = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}$. Find $\frac{\alpha}{\beta}$.



View Text Solution

6. If A is the matrix $\begin{bmatrix} 1 & -3 \\ -1 & 1 \end{bmatrix}$, then $A - \frac{1}{3}A^2 + \frac{1}{9}A^3 - \dots + \left(-\frac{1}{3}\right)^n A^{n+1} + \dots \infty = \frac{3}{13} \begin{bmatrix} 1 & a \\ b & 1 \end{bmatrix}$. Find $\left|\frac{a}{b}\right|$.



View Text Solution

7. A sequence of 2×2 matrices $\{M_n\}$ is defined as follows

$$M_n = \begin{bmatrix} \frac{1}{(2n+1)!} & \frac{1}{(2n+2)!} \\ \sum_{k=0}^n \frac{(2n+2)!}{(2k+2)!} & \sum_{k=0}^n \frac{(2n+1)!}{(2k+1)!} \end{bmatrix} \quad \text{then}$$

$\lim_{n \rightarrow \infty} \det. (M_n) = \lambda - e^{-1}$. Find λ .



Watch Video Solution

8. Let $|\vec{a}| = 1$, $|\vec{b}| = 1$ and $|\vec{a} + \vec{b}| = \sqrt{3}$. If \vec{c} be a vector such that $\vec{c} = \vec{a} + 2\vec{b} - 3(\vec{a} \times \vec{b})$ and $p|\left(\vec{a} \times \vec{b}\right) \times \vec{c}|$, then find $[p^2]$.
(where $[]$ represents greatest integer function).



Watch Video Solution

9. Let $\vec{r} = (\vec{a} \times \vec{b})\sin x + (\vec{b} \times \vec{c})\cos y + 2(\vec{c} \times \vec{a})$, where $\vec{a}, \vec{b}, \vec{c}$ are non-zero and non-coplanar vectors. If \vec{r} is orthogonal to $\vec{a} + \vec{b} + \vec{c}$, then find the minimum value of $\frac{4}{\pi^2}(x^2 + y^2)$.



Watch Video Solution

10. The plane denoted by $P_1: 4x + 7y + 4z + 81 = 0$ is rotated through a right angle about its line of intersection with the plane

$P_2: 5x + 3y + 10z = 25$. If the plane in its new position is denoted by P , and the distance of this plane from the origin is d , then find the value of $[k/2]$ (where $[\cdot]$ represents greatest integer less than or equal to k).



Watch Video Solution

11. ABCD is a regular tetrahedron, A is the origin and B lies on x-axis. ABC lies in the xy-plane $|\overrightarrow{AB}| = 2$ Under these conditions, the number of possible tetrahedrons is :



Watch Video Solution

12. A, B, C, D are four points in the space and satisfy $|\overrightarrow{AB}| = 3$, $|\overrightarrow{BC}| = 7$, $|\overrightarrow{CD}| = 11$ and $|\overrightarrow{DA}| = 9$. Then find the value of $\overrightarrow{AC} \cdot \overrightarrow{BD}$.



Watch Video Solution

13. Let OABC be a regular tetrahedron of edge length unity. Its volume be V and $6V = \sqrt{\frac{p}{q}}$ where p and q are relatively prime. The find the value of $(p+q)$



Watch Video Solution

14. If \vec{a} and \vec{b} are non zero, non collinear vectors and $\vec{a}_1 = \lambda \vec{a} + 3 \vec{b}$, $\vec{b}_1 = 2 \vec{a} + \lambda \vec{b}$, $\vec{c}_1 = \vec{a} + \vec{b}$. Find the sum of all possible real values of λ so that points A_1, B_1, C_1 whose position vectors are $\vec{a}_1, \vec{b}_1, \vec{c}_1$ respectively are collinear is equal to.



Watch Video Solution

15. Let P and Q are two points on curve $y = \log_{\frac{1}{2}}\left(x - \frac{1}{2}\right) + \log_2 \sqrt{4x^2 - 4x + 1}$ and P is also on $x^2 + y^2 = 10$. Q lies inside the given circle such that its abscissa is

integer. Find the smallest possible value of $\vec{OP} \cdot \vec{OQ}$ where 'O' being origin.



Watch Video Solution

16. Let P and Q are two points on curve $y = \log_{\frac{1}{2}}\left(x - \frac{1}{2}\right) + \log_2 \sqrt{4x^2 - 4x + 1}$ and P is also on $x^2 + y^2 = 10$. Q lies inside the given circle such that its abscissa is integer. find the largest possible value of $|\vec{PQ}|$.



Watch Video Solution

17. If $a, b, c, l, m, n \in \mathbb{R} - \{0\}$ such that $al + bm + cn = 0, bl + cm + an = 0, cl + am + bn = 0$. If a, b, c are distinct and $f(x) = ax^3 + bx^2 + cx + 2$. Find $f(1)$:



Watch Video Solution

18. Let \vec{u} and \vec{v} be unit vectors such that $\vec{u} \times \vec{v} + \vec{u} = \vec{w}$ and $\vec{w} \times \vec{u} = \vec{v}$. Find the value of $\left[\vec{u} \vec{v} \vec{w} \right]$.



Watch Video Solution