



PHYSICS

BOOKS - NIKITA PHYSICS (HINGLISH)

QUESTION PAPER - MH-CET 2018

Mcq

1. The path length of oscillation of simple pendulum of length 1 m is 16 cm. Its maximum velocity is (take, $g = \pi^2 m / s^2$)

A. $2\pi cm / s$

B. $4\pi cm / s$

C. $8\pi cm / s$

D. $16\pi cm / s$

Answer: C



[Watch Video Solution](#)

2. A vessel completely filled with water has holes 'A' and 'B' at depths 'h' and '2h' from the top, respectively. Hole 'A' is a square of side 'L' and 'B' is a circle of radius 'r'. The water flowing out per second from both the holes is same. Then, 'L' is equal to

A. $r^{\frac{1}{2}} (\pi)^{\frac{1}{2}} (3)^{\frac{1}{2}}$

B. $r \cdot (\pi)^{\frac{1}{2}} (3)^{\frac{1}{4}}$

C. $r \cdot (\pi)^{\frac{1}{4}} (3)^{\frac{1}{4}}$

D. $r^{\frac{1}{2}} (\pi)^{\frac{1}{3}} (3)^{\frac{1}{2}}$

Answer: C



[Watch Video Solution](#)

3. A transistor is used as a common emitter amplifier with a load resistance $2\text{K}\Omega$. The input resistance is 150Ω . Base current is changed by

20 μA which results in a change in collector current by 1.5 mA. The voltage gain of the amplifier is

- A. 900
- B. 1000
- C. 1100
- D. 1200

Answer: B



[Watch Video Solution](#)

4. A disc has mass 'M' and radius 'R'. How much tangential force should be applied to the rim of the disc, so as to rotate with angular velocity ' ω ' in time 't' ?

- A. $\frac{MR\omega}{4t}$
- B. $\frac{MR\omega}{2t}$
- C. $\frac{MR\omega}{t}$

D. $MR\omega t$

Answer: C



[Watch Video Solution](#)

5. A circular coil of radius R carries a current i . The magnetic field at its centre is B . The distance from the centre on the axis of the coil where the magnetic field will be $B/8$ is

A. $R\sqrt{2}$

B. $R\sqrt{3}$

C. $2R$

D. $3R$

Answer: B



[Watch Video Solution](#)

6. Two periodic waves of intensities I_1 and I_2 pass through a region at the same time in the same direction. The sum of the maximum and minimum intensities is:

A. $(I_1 + I_2)$

B. $2(I_1 + I_2)$

C. $(\sqrt{I_1} + \sqrt{I_2})$

D. $(\sqrt{I_1} - \sqrt{I_2})$

Answer: B



[Watch Video Solution](#)

7. An alternating voltage $E = 200\sqrt{2}\sin(100t)$ is connected to a 1 microfarad capacitor through an AC ammeter. The reading of the ammeter shall be

A. $5mA$

B. $10mA$


C. 15mA

D. 20mA

Answer: D

 Watch Video Solution

8. In the following network, the current flowing through 15Ω resistance is



A. 0.8A

B. 1.0A

C. 1.2A

D. 1.4A

Answer: C



Watch Video Solution

9. The angle made by incident ray of light with the reflecting surface is called

- A. glancing angle
- B. angle of incidence
- C. angle of deviation
- D. angle of refraction

Answer: A



Watch Video Solution

10. In a non - uniform circular motaion the ratio of tangential to radial acceleration is (where, r = radius of circle, v = speed of the particle, α = angular acceleration)

A. $\frac{\alpha^2 r^2}{v}$

B. $\frac{\alpha^2 r}{v^2}$

C. $\frac{\alpha r^2}{v^2}$

D. $\frac{v^2}{r^2 \alpha}$

Answer: C



[Watch Video Solution](#)

11. If numerical aperture of a microscope is increased, then its

A. resolving power remains constant

B. resolving power becomes zero

C. limit of resolution is decreased

D. limit of resolution is increased

Answer: B



[Watch Video Solution](#)

12. In amplitude modulation

A. amplitude remains constant but frequency changes

B. both amplitude and frequency do not change

C. both amplitude and frequency change

D. amplitude of the carrier wave changes according to information
signal

Answer: D



[Watch Video Solution](#)

13. If M_z = magnetisation of paramagnetic sample B= external magnetic field T= absolute temperature, C= curie constant then according to Curie's law in magnetism, the correct relation is

A. $M_x = \frac{T}{CB}$

B. $M_z = \frac{CB}{T}$

C. $C = \frac{M_z B}{T}$

D. $C = \frac{T^2}{M_z B}$

Answer: B



Watch Video Solution

14. An electron of stationary hydrogen atom jumps from 4th energy level to ground level. The velocity that the photon acquired as a result of electron transition will be (where, h = Planck's constant, R =Rydberg's constant m = mass of photon)

A. $\frac{9Rh}{16m}$

B. $\frac{11hR}{16m}$

C. $\frac{13hR}{16m}$

D. $\frac{15hR}{16m}$

Answer: D



Watch Video Solution

15. A metal wire of density ρ floats on water surface horizontally. If is NOT to sink in water, then maximum radius of wire is proportional to (where, T =surface tension of water, g =gravitational acceleration)

A. $\sqrt{\frac{T}{\log}}$

B. $\sqrt{\frac{\pi\rho g}{T}}$

C. $\frac{T}{\pi\rho g}$

D. $\frac{\pi\rho g}{T}$

Answer: D



Watch Video Solution

16. A sphere of mass m moving with velocity v collides head-on on another sphere of same mass which is at rest. The ratio of final velocity of second sphere to the initial velocity of the first sphere is (where, e is coefficient of restitution and collision is inelastic)

A. $\frac{e - 1}{2}$

B. $\frac{e}{2}$

C. $\frac{e + 1}{2}$

D. e

Answer: C



Watch Video Solution

17. For a particle performing linear SHM, its average speed over one oscillation is (where, a = amplitude of SHM, n = frequency of oscillation)

A. $2an$

B. $4an$

C. $6an$

D. $8an$

Answer: B



[Watch Video Solution](#)

18. An ideal transformer converts 220 V AC to 3.3 kV AC to transmit a power of 4.4 kW. If primary coil has 600 turns, then alternating current in secondary coil is

A. $\frac{1}{3} A$

B. $\frac{4}{3} A$

C. $\frac{5}{3}A$

D. $\frac{7}{3}A$

Answer: B



[Watch Video Solution](#)

19. A conducting wire has length L_1 and diameter d_1 . After stretching the same wire length becomes L_2 and diameter d_2 , The ratio of resistances before and after stretching is

A. $d_2^4 : d_1^4$

B. $d_1^4 : d_2^4$

C. $d_2^2 : d_1^2$

D. $d_1^2 : d_2^2$

Answer: B



[Watch Video Solution](#)

20. The molar specific heat of an ideal gas at constant pressure and constant volume is C_p and C_v respectively. If R is the universal gas constant and the ratio of C_p to C_v is γ , then C_v .

A. $\frac{1 - \gamma}{1 + \gamma}$

B. $\frac{1 + \gamma}{1 - \gamma}$

C. $\frac{\gamma - 1}{R}$

D. $\frac{R}{\gamma - 1}$

Answer: D



Watch Video Solution

21. In a capillary tube having area of cross - section A , water rises to a height h . If cross-sectional area is reduced to $\frac{A}{9}$, the rise of water in the capillary tube is

A. $4h$

B. 3h

C. 2h

D. h

Answer: B



[Watch Video Solution](#)

22. On adjusting the $P - N$ junction diode in forward biased

A. is one in which width of depletion layer increases

B. is one in which potential barrier increases

C. acts as closed switch

D. acts as open switch

Answer: C



[Watch Video Solution](#)

23. An alternating electric field, of frequency ν , is applied across the dees (radius= R) of a cyclotron that is being used to accelerate protons (mass= m) the operating magnetic field (B) used in the cyclotron and the kinetic energy (K) of the proton beam, produced by it, are given by:

A. $\frac{2\pi m\nu}{e}, 2\pi^2 m\nu^2 R^2$

B. $\frac{2\pi^2 m\nu}{e^2}, 4\pi^2 m\nu^2 \pi^2$

C. $\frac{\pi m\nu}{e}, \pi^2 m\nu^2 R^2$

D. $\frac{2\pi^2 m^2 \nu^2}{e}, 2\pi^2 m^2 \nu^2 R^2$

Answer: A



Watch Video Solution

24. A ray of light is incident normally on a glass slab of thickness 5 cm and refractive index 1.6. The time taken to travel by a ray from source to surface of slab is same as to travel through glass slab. The distance of source from the surface is

A. 4 cm

B. 8 cm

C. 12 cm

D. 16 cm

Answer: B



Watch Video Solution

25. A string is vibrating in its fifth overtone between two rigid supports 2.4 m apart. The distance between successive node and antinode is

A. $0.1m$

B. $0.2m$

C. $0.6m$

D. $0.8m$

Answer: B

26.

If

$$\vec{A} = 3\hat{i} - 2\hat{j} + \hat{k}, \vec{B} = \hat{i} - 3\hat{j} + 5\hat{k} \text{ and } \vec{C} = 2\hat{i} + \hat{j} - \hat{k}$$

from a right angled triangle then out of the following which one is satisfied ?

A. $\vec{A} = \vec{B} + \vec{C}$ and $A^2 = B^2 + C^2$

B. $\vec{A} = \vec{B} + \vec{C}$ and $B^2 = A^2 + C^2$

C. $\vec{B} = \vec{A} + \vec{C}$ and $B^2 = A^2 + C^2$

D. $\vec{B} = \vec{A} + \vec{C}$ and $A^2 = B^2 + C^2$

Answer: B

27. A square frame ABCD is formed by four identical rods each of mass 'm' and length 'l'. This frame is in X Y - plane such that side AB coincides with

X-axis and side AD along Y-axis. The moment of inertia of the frame about X-axis is

A. $\frac{5ml^2}{3}$

B. $\frac{2ml^2}{3}$

C. $\frac{4ml^2}{3}$

D. $\frac{ml^2}{12}$

Answer: A



Watch Video Solution

28. A unit vector is represented as $(0.8\hat{i} + b\hat{j} + 04\hat{k})$. Hence, the value of b must be

A. 0.4

B. $\sqrt{0.6}$

C. 0.2

D. $\sqrt{0.2}$

Answer: D



[Watch Video Solution](#)

29. Magnetic susceptibility for a paramagnetic and diamagnetic materials is respectively,

- A. small, positive and small, positive
- B. large, positive and small, negative
- C. small, positive and small, negative
- D. large, negative and large, positive

Answer: C



[Watch Video Solution](#)

30. Frequency of a particle executing SHM is 10 Hz. The particle is suspended from a vertical spring. At the highest point of its oscillation the spring is unstretched. Maximum speed of the particle is $\left(g = 10\frac{m}{s}\right)$

A. $2\pi m / s$

B. $\pi m / s$

C. $\frac{1}{2\pi} m / s$

D. $\frac{1}{\pi} m / s$

Answer: D



[Watch Video Solution](#)

31. The moment of inertia of a ring about an axis passing through the centre and perpendicular to its plane is I . It is rotating with angular velocity ω . Another identical ring is gently placed on it, so that their centres coincide. If both the rings are rotating about the same axis, then loss in kinetic energy is

A. $\frac{I\omega^2}{2}$

B. $\frac{I\omega^2}{4}$

C. $\frac{I\omega^2}{6}$

D. $\frac{I\omega^2}{8}$

Answer: B



Watch Video Solution

32. A bomb at rest explodes into 3 parts of same mass. The momentum of two parts is $-3p\hat{i}$ and $2p\hat{j}$. Respectively. The magnitude of momentum of the third part is

A. P

B. $\sqrt{5}P$

C. $\sqrt{11}P$

D. $\sqrt{13}P$

Answer: D



[Watch Video Solution](#)

33. In a photocell frequency of incident radiation is increased by keeping other factors constant ($\nu > \nu_0$), the stopping potential

- A. decreases
- B. increases
- C. becomes zero
- D. first decreases and then increases

Answer: B



[Watch Video Solution](#)

34. A body crosses the topmost point of a vertical circle with a critical speed. Its centripetal acceleration, when the string is horizontal will be

A. g

B. 3g

C. 4g

D. 6g

Answer: B

 [Watch Video Solution](#)

35. The expression for electric field intensity at a points outside uniformly charged thin plane sheet is (where, d is the distance of point from plane sheet)

A. independent of d

B. directly proportional to \sqrt{d}

C. directly proportional to d

D. directly proportional to $\frac{1}{\sqrt{d}}$

Answer: A



[Watch Video Solution](#)

36. When source of sound moves towards a stationary observer, the wavelength of sound received by him

- A. decreases while frequency increases
- B. remains the same whereas frequency increases
- C. increases and frequency also increases
- D. decreases while frequency remains the same

Answer: A



[Watch Video Solution](#)

37. When a shunt of 4Ω is attached to a galvanometer, the deflection reduces to $\frac{1}{5}$ th. If an additional shunt of 2Ω is attached. What will be the

deflection?

A. $\frac{1}{2}$

B. $\left(\frac{1}{3}\right)^{rd}$

C. $\left(\frac{1}{4}\right)^{th}$

D. $\left(\frac{1}{8.5}\right)^{th}$

Answer: D



Watch Video Solution

38. A body is thrown from the surface of the earth with velocity u m/s. The maximum height in metre above the surface of the earth upto which it will reach is (where, R = radius of earth, g =acceleration due to gravity)

A. $\frac{u^2 R}{2gR - u^2}$

B. $\frac{2u^2 R}{gR - u^2}$

C. $\frac{u^2 R^2}{2gR^2 - u^2}$

D. $\frac{u^2 R}{9R - u^2}$

Answer: A



Watch Video Solution

39. A series combination of N_1 capacitors (each of capacity C_1) is charged to potential difference $3V$. Another parallel combination of N_2 capacitors (each of capacity C_2) is charged to potential difference V . The total energy stored in both the combinations is same, The value of C_1 in terms of C_2 is

A. $\frac{C_2 N_1 N_2}{9}$

B. $\frac{C_2 N_1^2 N_2^2}{9}$

C. $\frac{C_2 N_1}{9 N_2}$

D. $\frac{C_2 N_2}{9 N_1}$

Answer: A



Watch Video Solution

40. Heat energy is incident on the surface at the rate of 1000 J/min. If coefficient of absorption is 0.8 and coefficient of reflection is 0.1 then heat energy transmitted by the surface in 5 minute is

A. 100 J

B. 500 J

C. 710 J

D. 900 J

Answer: B

 [Watch Video Solution](#)

41. Two metal wires P and Q of same length and material are stretched by same load. Their masses are in the ratio $m_1 : m_2$. The ratio of elongation of wire P to that of Q is

A. $m_1^2 : m_2^2$

B. $m_2^2 : m_1^2$

C. $m_1 : m_1$

D. $m_1 : m_2$

Answer: D

 [Watch Video Solution](#)

42. Let $x = \left[\frac{a^2 b^2}{c} \right]$ be the physical quantity. If the percentage error in the measurement of physical quantities a, b, and c is 2, 3 and 4 per cent respectively, then percentage error in the measurement of x is

A. 7 %

B. 14 %

C. 21 %

D. 28 %

Answer: B



[Watch Video Solution](#)

43. Draw a graph showing the variation of stopping potential with frequency of the incident radiation. What does the slope of the line with frequency axis indicate? What information can be obtained from the values of intercept on the potential axis?

A. (1)

B. (2)

C. (2)

D. (4)

Answer: A



[Watch Video Solution](#)

44. In compound microscope, the focal length and aperture of the objective used is respectively

- A. large and large
- B. large and small
- C. short and large
- D. short and small

Answer: B



[Watch Video Solution](#)

45. The energy of an electron having de-Broglie wavelength λ is

(where, h =Plank's constant, m = mass of electron)

- A. $\frac{h}{2mA}$
- B. $\frac{h^2}{2mA^2}$
- C. $\frac{h^2}{2m^2\lambda^2}$

D. $\frac{h^2}{2m^2\lambda}$

Answer: B



Watch Video Solution

46. n number of waves are produced on a string in 0.5 s. Now, the tension in the string is doubled (Assume length and radius constant), the number of waves produced in 0.5s for the same harmonic will be

A. n

B. $\sqrt{2}n$

C. $\frac{n}{\sqrt{2}}$

D. $\frac{n}{\sqrt{5}}$

Answer: C



Watch Video Solution

47. The increase in energy of a metal bar of length L and cross-sectional area A when compressed with a load M along its length is (where, Y = Young's modulus of the material of metal bar)

A. $\frac{FL}{2AY}$

B. $\frac{F^2L}{2AY}$

C. $\frac{FL}{AY}$

D. $\frac{F^2L^2}{2AY}$

Answer: C



[Watch Video Solution](#)

48. The ratio of magnetic fields due to a bar magnet at the two axial points P_1 and P_2 which are separated from each other by 10 cm is 25 : 2 . Point P_1 is situated at 10 cm from the centre of the magnet. Magnetic length of the bar magnet is (Points P_1 and P_2 are on the same side of magnet and distance of P_2 from the centre is greater than distance of P_1 from the centre of magnet)

A. 5 cm

B. 10 cm

C. 15 cm

D. 20 cm

Answer: B



Watch Video Solution

49. A satellite is revolving in a circular orbit at a height h above the surface of the earth of radius R . The speed of the satellite in its orbit is one-fourth the escape velocity from the surface of the earth. The relation between h and R is

A. $h = 2R$

B. $h = 3R$

C. $h = 5R$

D. $h = 7R$

Answer: D



[Watch Video Solution](#)

50. A pipe closed at one end has length 83 cm. The number of possible natural oscillations of air column whose frequencies lie below 1000 Hz are (take, velocity of sound in air 332 m/s)

A. 3

B. 4

C. 5

D. 6

Answer: C



[Watch Video Solution](#)