



## MATHS

### BOOKS - MARVEL MATHS (HINGLISH)

### QUESTION PAPER 2017

#### Question

1. The number of principal solutions of  $\tan 2\theta = 1$  is

- A. one
- B. two
- C. three
- D. four

**Answer: B**



Watch Video Solution

2. The objective function  $Z = 4x_1 + 5x_2$ , Subject to  $2x_1 + x_2 \geq 7$ ,  $2x_1 + 3x_2 \leq 15$ ,  $x_2 \leq 3$ ,  $x_1, x_2 \geq 0$  has minimum value at the point

- A. On x-axis
- B. On y-axis
- C. At the origin
- D. On the line parallel to x-axis

**Answer: A**



Watch Video Solution

3. If  $z_1$  and  $z_2$  are z-coordinates of the points of trisection of the segment joining the points A(2,1,4), B(-1,3,6), then  $z_1 + z_2 +$

A. 1

B. 4

C. 5

D. 10

**Answer: D**



[Watch Video Solution](#)

4. The maximum value of  $f(x) = \frac{\log x}{x}$  ( $x \neq 0, x \neq 1$ ) is

A. e

B.  $\frac{1}{e}$

C.  $e^2$

D.  $\frac{1}{e^2}$

**Answer: B**



[Watch Video Solution](#)

$$5. \int_0^1 x \tan^{-1} x dx =$$

A.  $\frac{\pi}{4} + \frac{1}{2}$

B.  $\frac{\pi}{4} - \frac{1}{2}$

C.  $\frac{1}{2} - \frac{\pi}{4}$

D.  $-\frac{\pi}{4} - \frac{1}{2}$

**Answer: B**



**Watch Video Solution**

6. The statement pattern  $(\sim p \wedge q)$  is logically equivalent to

A.  $(p \vee q) \vee \sim P$

B.  $(p \vee q) \wedge \sim P$

C.  $(p \wedge q) \rightarrow P$

D.  $(p \vee q) \rightarrow p$

**Answer: B**

 [Watch Video Solution](#)

7. If  $g(x)$  is the inverse function of  $f(x)$  and  $f'(x) = \frac{1}{1+x^4}$ , then  $g'(x)$  is

A.  $1 + [g(x)]^4$

B.  $1 - [g(x)]^4$

C.  $1 + [f(x)]^4$

D.  $\frac{1}{1 + [g(x)]^4}$

**Answer: A**

 [Watch Video Solution](#)

8. The inverse of the matrix  $\begin{bmatrix} 1 & 0 & 0 \\ 3 & 3 & 0 \\ 5 & 2 & -1 \end{bmatrix}$  is

A.  $-\frac{1}{3} \begin{bmatrix} -3 & 0 & 0 \\ 3 & 1 & 0 \\ 9 & 2 & -3 \end{bmatrix}$

B.  $-\frac{1}{3} \begin{bmatrix} -3 & 0 & 0 \\ 3 & -1 & 0 \\ -9 & -2 & 3 \end{bmatrix}$

C.  $-\frac{1}{3} \begin{bmatrix} 3 & 0 & 0 \\ 3 & -1 & 0 \\ -9 & -2 & 3 \end{bmatrix}$

D.  $-\frac{1}{3} \begin{bmatrix} -3 & 0 & 0 \\ -3 & -1 & 0 \\ -9 & -2 & 3 \end{bmatrix}$

Answer: B



Watch Video Solution

9. if  $\int \frac{1}{\sqrt{9-16x^2}} dx = \alpha \sin^{-1}(\beta x) + c$ . then  $\alpha + \frac{1}{\beta} =$

A. 1

B.  $\frac{7}{12}$

C.  $\frac{19}{12}$

D.  $\frac{9}{12}$

**Answer: A**



**Watch Video Solution**

10.  $O(0,0)$ ,  $A(1,2)$ ,  $B(3,4)$  are the vertices of  $\triangle OAB$ . The joint equation of the altitude and median drawn from  $O$  is

A.  $x^2 + 7xy - y^2 = 0$

B.  $x^2 + 7xy + y^2 = 0$

C.  $3x^2 - xy - 2y^2 = 0$

D.  $3x^2 + xy - 2y^2 = 0$

**Answer: D**



**Watch Video Solution**

11.  $f(x) = \begin{cases} \left(\tan\frac{\pi}{4} + x\right)^{1/x}, & x \neq 0 \\ k, & x = 0 \end{cases}$  for what value of  $k$ ,  $f(x)$  is

continuous at  $x = 0$  ?

A.  $e$

B.  $e^{-1}$

C.  $e^2$

D.  $e^{-2}$

**Answer: B**



[Watch Video Solution](#)

12. For a invertible matrix  $A$  if  $A(adjA) = \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix}$  then  $|A| =$

A. 100

B.  $-100$

C. 10



D.  $-10$

**Answer: C**



**Watch Video Solution**

13. The Solution of the differential equation  $\frac{dy}{dx} = \tan\left(\frac{y}{x}\right) + \frac{y}{x}$  is

A.  $\cos\left(\frac{y}{x}\right) = cx$

B.  $\sin\left(\frac{y}{x}\right) = cx$

C.  $\cos\left(\frac{y}{x}\right) = cy$

D.  $\sin\left(\frac{y}{x}\right) = cy$

**Answer: B**



**Watch Video Solution**

14. In  $\Delta ABC$  if  $\sin^2 A + \sin^2 B = \sin^2 B = \sin^2 C$  and  $1(AB) = 10$ , then the maximum value of the area of  $\Delta ABC$  is

A. 50

B.  $10\sqrt{2}$

C. 25

D.  $25\sqrt{2}$

Answer: C



[View Text Solution](#)

15. If  $x = f(t)$  and  $y = g(t)$  are differentiable functions of  $t$  then  $\frac{d^2y}{dx^2}$  is

A.  $\frac{f'(t) \cdot g''(t) - g'(t) \cdot f''(t)}{[f'(t)]^3}$

B.  $\frac{f'(t) \cdot g''(t) - g'(t) \cdot f''(t)}{[f'(t)]^2}$

C.  $\frac{g'(t) \cdot f''(t) - f'(t) \cdot g''(t)}{[f'(t)]^3}$

D.  $\frac{g'(t) \cdot f''(t) + f'(t) \cdot g''(t)}{[f'(t)]^3}$

Answer: A



Watch Video Solution

16. The equation of line equally inclined to co -ordinate axes and passing through (3,2,-5) is

A.  $\frac{x + 3}{1} = \frac{y - 2}{1} = \frac{z + 5}{1}$

B.  $\frac{x - 3}{1} = \frac{y - 2}{1} = \frac{5 + z}{-1}$

C.  $\frac{x + 3}{-1} = \frac{y - 2}{1} = \frac{z + 5}{-1}$

D.  $\frac{x + 3}{-1} = \frac{2 - y}{1} = \frac{z + 5}{-1}$

Answer: B



Watch Video Solution

17. If  $\int_0^{\pi/2} \log \cos x dx = \frac{\pi}{2} \log \left( \frac{1}{2} \right)$ , then  $\int_0^{\pi/2} \log \sec x dx =$

A.  $\frac{\pi}{2} \log \left( \frac{1}{2} \right)$

B.  $1 - \frac{\pi}{2} \log \left( \frac{1}{2} \right)$

C.  $1 + \frac{\pi}{2} \log \left( \frac{1}{2} \right)$

D.  $\frac{\pi}{2} \log 2$

**Answer: D**



[Watch Video Solution](#)

18. A boy tosses fair coin 3 times. If he gets Rs 2X for X heads, then his expected gain equals to Rs.....

A. 1

B.  $\frac{3}{2}$

C. 3

D. 4

**Answer: B**



[Watch Video Solution](#)

**19.** Which of the following statement pattern is a tautology?

A.  $p \vee (q \rightarrow p)$

B.  $\sim q \rightarrow \sim P$

C.  $(1 \rightarrow p) \vee (\sim p \leftrightarrow q)$

D.  $p \wedge \sim p$

**Answer: C**



[Watch Video Solution](#)

20. If the angle between the planes  $\bar{r} \cdot (m\hat{i} - \hat{j} + 2\hat{k}) + 3 = 0$  and  $\bar{r} \cdot (2\hat{i} - m\hat{j} - \hat{k}) - 5 = 0$  is  $\frac{\pi}{3}$ , then  $m =$

- A. 2
- B.  $\pm 3$
- C. 3
- D. -2

**Answer: C**



[Watch Video Solution](#)

21. If the origin and the point  $p(2, 3, 4), q(1, 2, 3)R(x, y, z)$  are coplanar then

- A.  $x - 2y - z = 0$
- B.  $x + 2y + z = 0$

C.  $x - 2y + z = 0$

D.  $2x - 2y + z = 0$

**Answer: C**



**Watch Video Solution**

22. if lines represented by equation  $px^2 - qy^2 = 0$  are distinct, then

A.  $pq > 0$

B.  $pq < 0$

C.  $pq = 0$

D.  $p + q = 0$

**Answer: A**



**Watch Video Solution**

23. Let  $\square PQRS$  be a quadrilateral. If M and N are the mid-points of the sides PQ and RS respectively, then  $PS+QR=$

A.  $3\overline{MN}$

B.  $4\overline{MN}$

C.  $2\overline{MN}$

D.  $2\overline{NM}$

**Answer: C**



[Watch Video Solution](#)

24. If slopes of lines represented by  $kx^2 + 5xy + y^2 = 0$  differ by 1, then k=

A. 2

B. 3



C. 6

D. 8

**Answer: C**



[Watch Video Solution](#)

25. If vector  $r$  with direction cosines  $l, m, n$  is equally inclined to the coordinate axes, then the total number of such vectors is

A. 4

B. 6

C. 8

D. 2

**Answer: C**



[Watch Video Solution](#)

26. If  $\int \frac{1}{(x^2 + 4)(x^2 + 9)} dx = A \tan^{-1} \frac{x}{2} + B \tan^{-1} \left( \frac{x}{3} \right) + C$ , then

A-B=

A.  $\frac{1}{6}$

B.  $\frac{1}{30}$

C.  $-\frac{1}{30}$

D.  $-\frac{1}{6}$

**Answer: A**



**Watch Video Solution**

27. If  $\alpha$  and  $\beta$  are roots of the equation  $x^2 + 5|x| - 6 = 0$ , then the value of  $|\tan^{-1} \alpha - \tan^{-1} \beta|$  is

A.  $\frac{\pi}{2}$

B. 0

C.  $\pi$

D.  $\frac{\pi}{4}$

**Answer: A**



**Watch Video Solution**

28. If  $x = a\left(t - \frac{1}{t}\right)$ ,  $y = a\left(t + \frac{1}{t}\right)$ , where  $t$  be the parameter, then

$$\frac{dy}{dx} = ?$$

A.  $\frac{y}{x}$

B.  $\frac{-x}{y}$

C.  $\frac{x}{y}$

D.  $\frac{-y}{x}$

**Answer: C**



**Watch Video Solution**

29. The point on the curve  $y = \sqrt{x-1}$  where the tangent is perpendicular to the line  $2x + y - 5 = 0$  is

A.  $(2, -1)$

B.  $(10, 3)$

C.  $(2, 1)$

D.  $(5, -2)$

**Answer: C**



**Watch Video Solution**

30. If

$$\int \sqrt{\frac{x-5}{x-7}} dx = A\sqrt{x^2 - 12x + 35} + \log|x - 6 + \sqrt{x^2 - 12x + 35}| + C$$

, then

A.  $-1$

B.  $\frac{1}{2}$

C.  $-\frac{1}{2}$

D. 1

**Answer: D**



[Watch Video Solution](#)

31. A r.v.  $X \sim B(n, p)$ . If values of mean and variance of  $X$  are 18 and 12 respectively, then total number of possible values of  $X$  are

A. 54

B. 55

C. 12

D. 18

**Answer: B**



[Watch Video Solution](#)

32. The area of the region bounded by the lines  $y = 2x + 1$ ,  $y = 3x + 1$  and  $x = 4$  is

A.  $16\text{sq unit}$

B.  $\frac{121}{3}$  sq, unit

C.  $\frac{121}{6}$ sq,unit

D.  $8$  sq,unit

**Answer: D**



[Watch Video Solution](#)

33. A box contains 6 pens , 2 of which are defective Two pens are taken randomly from the box .If r.v.  $X$ , : Number of defective pens obtained , then standard deviation of  $x$ =

A.  $\pm \frac{4}{3\sqrt{5}}$

B.  $\frac{2}{3}$

C.  $\frac{16}{45}$

D.  $\frac{4}{3\sqrt{5}}$

**Answer: B**



**Watch Video Solution**

**34.** If the volume of spherical ball is increasing at the rate of  $4\pi$  cc/s, then the rate of change of its surface area when the volume is  $288\pi$  cc is

A.  $\frac{4}{3}\pi x m^2 / \text{sec}$

B.  $\frac{2}{3}\pi cm^2 / \text{sec}$

C.  $4\pi cm^2 / \text{sec}$

D.  $2\pi c \frac{m^2}{\text{sec}}$

**Answer: A**



Watch Video Solution

35. If  $f(x) = \log(\sec^2 x)^{\cot^2 x}$  for  $x \neq 0$  for  $x=0$  is continuous at  $x=0$ , then K is

A.  $e^{-1}$

B. 1

C.  $e$

D. 0

Answer: B



Watch Video Solution

36. If  $c$  denotes the contradiction, then dual of the compound statement  $\sim p \wedge (q \vee c)$  is

A.  $\sim P \vee (q \wedge t)$



B.  $\sim P \wedge (q \vee t)$

C.  $p \vee (\sim q \vee t)$

D.  $\sim p \vee (q \wedge c)$

**Answer: A**



**Watch Video Solution**

37. All parabolas whose axis is the Y-axis.

A.  $x \frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$

B.  $x \frac{d^2y}{dx^2} + \frac{dy}{dx} = 0$

C.  $\frac{d^2y}{dx^2} - y = 0$

D.  $\frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$

**Answer: A**



**Watch Video Solution**

38.  $\int_0^3 [x] dx = \underline{\hspace{2cm}}$  where  $[x]$  is greatest integer function

A. 3

B. 0

C. 2

D. 1

**Answer: A**



[Watch Video Solution](#)

39. The objective function of LPP defined over the convex set attains its optimum value at

A. at least two of the corner points

B. all the corner points

C. at least one of the corner points

D. none of the corner points

**Answer: C**



**Watch Video Solution**

40. IF the inverse of the matrix  $\begin{bmatrix} \alpha & 14 & -1 \\ 2 & 3 & 1 \\ 6 & 2 & 3 \end{bmatrix}$  does not exist then the value of  $\alpha$  is

A. 1

B. -1

C. 0

D. -2

**Answer: D**



**Watch Video Solution**

41. If  $f(x) = x$  for  $x \leq 0$

= 0 for  $x > 0$  then  $f(x)$  at  $x=0$  is

- A. continuous but not differentiable
- B. Not continuous but differentiable
- C. Continuous and differentiable
- D. Not continuous and not differentiable

**Answer: A**



[Watch Video Solution](#)

42. The equation of the plane through  $(-1,1,2)$ , whose normal makes equal acute angles with coordinate axes is

A.  $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 2$

B.  $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 6$

C.  $\vec{r} \cdot (3\hat{i} - 3\hat{j} + 3\hat{k}) = 2$

$$D. \bar{r} \cdot (\hat{i} - \hat{j} + \hat{k}) = 3$$

**Answer: A**



**Watch Video Solution**

**43.** Probability that a person will develop immunity after vaccination is 0.8 If 8 people are given the vaccine then probability that all develop immunity is

A.  $(0.2)^8$

B.  $(0.8)^8$

C. 1

D.  ${}^8C_6(0.2)^6(0.8)^2$

**Answer: B**



**Watch Video Solution**

44. If the distance of points  $2\hat{i} + 3\hat{j} + \lambda\hat{k}$  from the plane  $\vec{r} \cdot (3\hat{i} + 2\hat{j} + 6\hat{k}) = 13$  is 5 units then  $\lambda =$

A.  $6, -\frac{17}{3}$

B.  $6, \frac{17}{3}$

C.  $-6, -\frac{17}{3}$

D.  $-6, \frac{17}{3}$

**Answer: A**



**Watch Video Solution**

45. The value of  $\cos^{-1} \left[ \cot \left( \frac{\pi}{2} \right) \right] + \cos^{-1} \left[ \sin \left( \frac{2\pi}{3} \right) \right]$  is

A.  $\frac{2\pi}{3}$

B.  $\frac{\pi}{3}$

C.  $\frac{\pi}{2}$

D.  $\pi$

**Answer: A**



[Watch Video Solution](#)

**46.** The particular solution of the differential equation  $xdy + 2ydx = 0$ ,  
when  $x = 2, y = 1$  is

A.  $xy = 4$

B.  $x^2y = 4$

C.  $xy^2 = 4$

D.  $x^2y^2 = 4$

**Answer: B**



[Watch Video Solution](#)

47.  $ABC$  is a triangle and  $A = (2, 3, 5)$ ,  $B = (-1, 3, 2)$  and  $C = (\lambda, 5, \mu)$ . If the median through  $A$  is equally inclined to the axes, then find the value of  $\lambda$  and  $\mu$ .

A. 10, 7

B. 9, 10

C. 7, 9

D. 7, 10

**Answer: D**



[Watch Video Solution](#)

48. For the following distribution function  $F(x)$  of a r.v  $X$

$x$	1	2	3	4	5	6
$F(x)$	0.2	0.37	0.48	0.62	0.85	1

$$p(3 < x \leq 5) =$$

A. 0.48



B. 0.37

C. 0.27

D. 1.47

**Answer: B**



**Watch Video Solution**

49. The lines  $\frac{x-1}{2} = \frac{y+1}{2} = \frac{z-1}{4}$  and  $\frac{x-3}{1} = \frac{y-k}{2} = \frac{z}{1}$

intersect each other at point

A.  $(-2, -4, 5)$

B.  $(-2, -4, -5)$

C.  $(2, 4, -5)$

D.  $(2, -4, -5)$

**Answer: B**



**Watch Video Solution**

50.  $\int \frac{\sec^8 x}{\operatorname{cosec} x} dx =$

A.  $\frac{\sec^8 x}{8} + c$

B.  $\frac{\sec^7 x}{7} + c$

C.  $\frac{\sec^6 x}{6} + c$

D.  $\frac{\sec^9 x}{9} + c$

**Answer: B**



**Watch Video Solution**