



PHYSICS

BOOKS - TARGET PHYSICS (HINGLISH)

MEASUREMENTS

Classical Thinking Introduction

1. The atomic molecular and nuclear phenomes are the part of _____ domian.

A. macroscopic

B. microscopic

C. megascopic

D. electroscopic

Answer: b



Watch Video Solution

2. Nano size of gold has _____ colour .

A. yellow

B. red

C. pink

D. orange

Answer: c



Watch Video Solution

3. Maxwell's equal relate to _____

A. law of gravitation

B. basic laws of electromagnetism

C. laws of electrostatics

D. nuclear model of an atom

Answer: b



Watch Video Solution

Classical Thinking Need Of Measurements

1. _____ is needed for the experimental verification of various theories .

A. unit

B. Symbol

C. Instrument

D. Measurement

Answer: d



Watch Video Solution

Classical Thinking Units Of Measurements

1. The reference standard used for the measurement of physical quantity is called _____.

A. standard quantity

B. dimension

C. constant

D. unit

Answer: d



Watch Video Solution

2. Which of the following is NOT a characteristic of a good unit ?

A. It is invariable

B. It is reproducible

C. It is perishable

D. It is easily available

Answer: c



Watch Video Solution

Classical Thinking System Of Units

1. Unit are classified into _____group.

A. 2

B. 4

C. 5

D. 6

Answer: a



Watch Video Solution

2. A set of fundamental and derived units is known as _____ .

A. supplementary units

B. system of units

C. complementary units

D. metric units

Answer: b



Watch Video Solution

3. The physical quantity having the same unit in all the systems of unit is _____ .

A. length

B. time

C. mass

D. foot

Answer: b



Watch Video Solution

Classical Thinking S I Units

1. S.I system of unit contains _____ supplementary unit.

A. 7

B. 2

C. 4

D. many

Answer: b



Watch Video Solution

2. In which of following system, scientific data can be exchanged between different parts of the world?

A. M.K.S

B. C.G.S

C. F.P.S

D. S.I

Answer: d



Watch Video Solution

Classical Thinking Fundamental And Derived Units

1. Out of the following units, which is NOT a fundamental unit?

A. newton

B. second

C. punod

D. kg

Answer: a



Watch Video Solution

2. Temperature can be expressed as a derived quantity in terms of any of the following

- A. length and mass
- B. mass and time
- C. length, mass and time
- D. none of these

Answer: d



Watch Video Solution

3. Which of the following is NOT a derived unit?

A. joule

B. erg

C. dyne

D. mole

Answer: d



Watch Video Solution

4. Which of the following is the CORRECT way of writing units?

A. 25 ms length

B. 30kg

C. 5 Newton

D. 10 N

Answer: d



Watch Video Solution

5. Which one of the following methods is used to measure distance of a planet or a star from the earth?

A. echo

B. direct

C. parallax

D. paradox

Answer: c



Watch Video Solution

6. The mass of the body depends only on

A. temperature

B. pressure

C. quantity of matter contained in the
body.

D. location of the body from the observer.

Answer: c



Watch Video Solution

7. Which of the following represents a unified atomic mass unit (u)?

A. $\frac{1}{12}$ of the mass of an atom of ^{12}C in kg

B. $\frac{1}{12}$ of the mass of an atom of ^{12}C in g

C. 8.333 of the mass of an atoms of ^{12}C in
g

D. 0.833 of the mass of an atom of ^{12}C in
kg

Answer: d



Watch Video Solution

8. In cesium atomic clock_____ is used .

A. cesium-122 atom

B. cesium-132 atom

C. cesium-133 atom

D. cesium-134 atom

Answer: c



Watch Video Solution

9. A _____ is the interval from one noon to the next noon.

A. mean solar day

B. solar day

C. lunar day

D. day

Answer: b



Watch Video Solution

10. Light year is a unit of

A. distance

B. time

C. temperature

D. luminous intensity

Answer: a



Watch Video Solution

11. Which of the following quantity is expressed as force per unit area

A. work

B. pressure

C. volume

D. density

Answer: b



Watch Video Solution

12. The physical quantity having the unit dyne

g^{-1} is _____ .

A. velocity

B. mass

C. force

D. acceleration

Answer: d



Watch Video Solution

13. Write the SI unit of luminous intensity and the amount of substance.

A. watt

B. lux

C. lumen

D. candela

Answer: d



Watch Video Solution

14. Which of the following is a supplementary unit ?

A. Steradian

B. candela

C. kelvin

D. pascal

Answer: a



Watch Video Solution

15. The pressure of 10^3 dyne/cm^2 is equivalent to

A. 10 N/m^2

B. 10^2 N/m^2

C. $10^{-2} N / m^2$

D. $10^{-1} N / m^2$

Answer: b



Watch Video Solution

Classical Thinking Dimensional Analysis

1. $[L^{-1} M^1 T^{-2}]$ are the dimensions of

A. joule constant

B. gravitational constant

C. pressure

D. force

Answer: c



Watch Video Solution

2. Checking the correctness of equations using the method of dimensions is based on

A. equality of inertial frame of reference.

B. the types of system of units

C. the method of measurement

D. principal of homogeneity of dimension.

Answer: d



Watch Video Solution

3. A unitless quantity

A. always has a non - zero dimension.

B. may have a non - zero dimension .

C. never has a zero dimension.

D. has no dimensions.

Answer: d



Watch Video Solution

4. Which of the following is NOT a dimensionless quantity?

A. angle

B. strain

C. specific gravity

D. density

Answer: d



Watch Video Solution

5. The unit of plane angle is radian , hence its dimension are

A. $[M^0 L^0 T^0]$

B. $[M^1 L^{-1} T^{-2}]$

C. $[M^0 L^1 T^{-1}]$

D. $[M^1 L^0 T^{-1}]$

Answer: a



Watch Video Solution

6. Dimensional equation CANNOT be used

A. to check the correctness

B. to drive the relation between different
physical quantities.

C. to find out constant of proportionality

which may be pure number

D. to change from one system of units to

another system

Answer: c



Watch Video Solution

7. If the dimensional formula for the physical quantity is $[M^1 L^2 T^{-2}]$ then the physical quantity is _____ .

A. torque

B. impulse

C. force

D. force per unit area

Answer: a



Watch Video Solution

8. If the dimensions of a physical quantity are given by $[L^a, M^b T^c]$, then the physical quantity will be

A. force ,if $a = -1$, $b=0,c=- 2$

B. pressure, if $a=-1,b=1,c=-2$

C. velocity , if $a =1$, $b=0,c=1$

D. acceleration , if $a=1,b=1,c=2$

Answer: b



Watch Video Solution

Classical Thinking Order Of Magnitude And Significant Figure

1. The value of the magnitude rounded off to the nearest integral power of 10 is called _____.

- A. significant figure
- B. uncertain number
- C. significant number
- D. order of magnitude

Answer: d



Watch Video Solution

2. Order of magnitude of $(10^6 + 10^3)$ is

A. 10^8

B. 10^9

C. 10^6

D. 10^3

Answer: c



Watch Video Solution

3. The length of a rod is 5×10^2 m , the order of magnitude of the length of the rod is

A. $10^3 m$

B. $10^2 m$

C. $10^1 m$

D. $10^{-1} m$

Answer: b



Watch Video Solution

4. The charge on the electron is $1.6 \times 10^{-19} \text{ C}$

. The order of magnitude is

A. 10^{19} C

B. 10^{18} m

C. 10^{-18} C

D. 10^{-19} C

Answer: d



Watch Video Solution

5. Significant figures depends upon the _____
of the measuring instrument .

A. length

B. readings

C. number

D. accuracy

Answer: d



Watch Video Solution

6. The number of significant figures in 0.0009 is

A. 4

B. 3

C. 2

D. 1

Answer: d



Watch Video Solution

7. The number of significant figures in 0.400 is

A. 1

B. 2

C. 3

D. 4

Answer: c



Watch Video Solution

8. The number of significant figures 0.0500 is

A. 4

B. 3

C. 2

D. 1

Answer: b



Watch Video Solution

9. State the number of significant figures in 6.032 J .

A. 4

B. 3

C. 2

D. 1

Answer: a



Watch Video Solution

Classical Thinking Accuracy And Errors In Measurements

1. The different between the true value and measured value is called _____.

A. mistake

B. error

C. significant figure

D. fault

Answer: b





2. If the pointer of the voltmeter is not exactly at the zero of the scale the the error is called _____.

- A. instrumental error
- B. systematic error
- C. personal error
- D. random error

Answer: b



[View Text Solution](#)

3. Zero error of an instrument introduces

- A. systematic error
- B. random error
- C. instrumental error
- D. none of these

Answer: a



[Watch Video Solution](#)

4. Accidental error can be minimised by

A. taking only one reading

B. taking small magnitude of the quantity.

C. selecting instrument with greater least
count

D. selecting instrument with small least
count.

Answer: d



Watch Video Solution

5. Zero error of an instrument introduces

A. faulty construction of instrument

B. wrong setting of instrument

C. lack of concentration of observer

D. wrong procedure of handling the instrument .

Answer: a



Watch Video Solution

6. Error due to non-removal of parallax between pointer and its image in case of magnetic compass needle causes _____.

A. instrumental error

B. persistent error

C. personal error

D. random error

Answer: c



Watch Video Solution

7. Instrumental error can be minimised by

A. taking large number of readings

B. using different accurate instrument for
the same reading

C. adjusting zero of the instrument

D. mainataining the temperature of the
surrounding.

Answer: b



Watch Video Solution

8. The magnitude of the difference between mean value and each individual value is called _____ .

- A. absolute error
- B. error in reading
- C. most probable error
- D. true error

Answer: a



Watch Video Solution

9. The formula for percentage error is

A. Percentage error $\frac{|\Delta a_m|}{a_m} \times 100 \%$

B. Percentage error

$$= \frac{1}{n} \sum_{i=1}^n |\Delta a_i| \times 100 \%$$

C. Percentage error $= \frac{a_m}{|\Delta a_m|} \times 100 \%$

D. Percentage error $\frac{1}{n} \sum_{i=1}^n a_i \times 100 \%$

Answer: a



Watch Video Solution

10. If $Y = a/b$, the maximum percentage error in the measurement of Y will be

A. $\frac{\Delta a/a}{\Delta b/b}$

B. $\frac{\Delta a}{a} + \frac{\Delta b}{b}$

C. $\frac{\Delta a}{a} - \frac{\Delta b}{b}$

D. $\frac{\Delta b/b}{\Delta a/a}$

Answer: b



Watch Video Solution

11. Given : $l_1 = 44.2 \pm 0.1$ and

$l_2 = 23.1 \pm 0.1$ the uncertainty in $l_1 + l_2$ is

A. 0

B. 0.1

C. 0.2

D. 0.4

Answer: c



Watch Video Solution

12. Two resistance $R_1 = 50 \pm 2$ ohm and $R_2 = 60$ ohm connected in series, the equivalent resistance of the series combination is

A. (110 ± 4) ohm

B. (110 ± 2) ohm

C. (110 ± 5) ohm

D. (110 ± 6) ohm

Answer: c



Watch Video Solution

13. If $x = a^n$ then relative error is (when n is power of a).



Watch Video Solution

14. Thickness of the paper measured by micrometer screw gauge of least count 0.01 mm is 1.03 mm, the percentage error in the measurement of thickness of paper is

A. 1.1 %

B. 1 %

C. 0.97 %

D. 0.8 %

Answer: c



Watch Video Solution

Classical Thinking Miscellaneous

1. One micron is related to centimetre as

A. 1 micron = 10^{-8} cm

B. 1 micron = 10^{-5} cm

C. 1 micron = 10^{-5} cm

D. 1 micron = 10^{-4} cm

Answer: d



Watch Video Solution

Critical Thinking Units Of Measurements

1. If u_1 and u_2 are the units selected in two systems of measurement and n_1 and n_2 their numerical values, then

A. $n_1 u_1 = n_2 u_2$

B. $n_1 u_1 + n_2 u_2 = 0$

C. $n_1 n_2 = u_1 u_2$

D. $(n_1 + u_3) = (n_2 + u_2)$

Answer: a



Watch Video Solution

Critical Thinking System Of Units

1. Which of the following system of units is not based on units of mass, length and time alone

A. S.I

B. M.K.S

C. F.P.S

D. C.G.S

Answer: a



Watch Video Solution

Critical Thinking Fundamental And Derived Units

1. The physical denote by

$$\frac{\text{mass} \times \text{perssuer}}{\text{density}}$$

A. force

B. momentum

C. angular momentum

D. wrok

Answer: d



Watch Video Solution

2. Universal time is based on

- A. rotation of the earth on it axis
- B. Earth's orbital motion around the sun .
- C. vibration of cesium atom
- D. oscillation of quartz crystal

Answer: c



Watch Video Solution

3. 1 a.m.u is equivalent to

A. $1.6 \times 10^{-27} \text{ kg}$

B. 934 Me

C. $1.6 \times 10^{-24} \text{ kg}$

D. All of the above

Answer: d



Watch Video Solution

4. The S.I. unit of momentum is

A. $\frac{kg}{m}$

B. $\frac{kgm}{s}$

C. $\frac{kgm^2}{s}$

D. $(kg \times \text{newton})$

Answer: b



Watch Video Solution

5. Curie is a unit of

A. energy of γ rays

B. half life

C. radioactivity

D. intensity of γ rays

Answer: c



Watch Video Solution

6. $S = A(1 - e^{Bxt})$, where S is speed and x is displacement. The units of B is

A. $m^{-1}s^{-1}$

B. $m^{-2}s$

C. s^{-2}

D. s^{-1}

Answer: a



Watch Video Solution

7. To determine the young's modulus of a wire , the formula is $Y = \frac{F}{A} \cdot \frac{L}{\Delta l}$, where $L = l$ length , $A =$ area of cross - section of the wire , $\Delta L =$ change in the length of the wire when stretched with a force F . Find the conversion factor to change it from CGS to MKS system.

A. 1

B. 10

C. 0.1

D. 0.01

Answer: c



Watch Video Solution

8. The moon subtends an angle of 57 minutes at the base line equal to radius of earth. What is the distance of moon from earth. Given radius of earth is 6400 km.

A. $11.22 \times 10^9 \text{ m}$

B. $3.86 \times 10^8 \text{ m}$

C. $3.68 \times 10^9 \text{ m}$

D. $3.68 \times 10^8 \text{ m}$

Answer: b



Watch Video Solution

9. The angular diameter of the sun is $1920''$. If the distance of the sun from the earth is $1.5 \times 10^{11} \text{ m}$, what is the linear diameter of the sun ?

A. $2.6 \times 10^9 \text{ m}$

B. $0.7 \times 10^9 m$

C. $5.2 \times 10^9 m$

D. $1.4 \times 10^9 m$

Answer: d



Watch Video Solution

Critical Thinking Dimensional Analysis

1. The fundamental physical quantities that have same dimension in the

dimensional formula of Torque and Angular Momentum are

A. mass, time

B. time , length

C. mass, length

D. time,mole

Answer: c



Watch Video Solution

2. Which one of the following represents the correct dimensions of the coefficient of viscosity?

A. $[M^1 L^{-1} T^2]$

B. $[M^1 L^{-1} T^{-1}]$

C. $[M^1 L^{-1} T^{-1}]$

D. $[M^1 L^{-2} T^{-2}]$

Answer: b



Watch Video Solution

3. Dimensions of electrical resistance are

A. $[M^1 L^2 T^2 I^{-1}]$

B. $[M^1 L^2 T^2 I^{-1}]$

C. $[M^1 L^1 T^{-3} I^{-2}]$

D. $[M^1 L^2 T^{-3} I^{-2}]$

Answer: d



Watch Video Solution

4. The dimensions of length in electric field, electric flux and electric dipole moment are respectively

A. L, L^2, L^3

B. L^3, L^2, L

C. L^{-1}, L^3, L^3

D. L, L^3, L

Answer: d



Watch Video Solution

5. If L denotes the inductance of an inductor through which a current i is flowing, the dimensions of Li^2 are

A. $[L^2 M^1 T^{-2}]$

B. Not expressible in LMT

C. $[L^1 M^1 T^{-2}]$

D. $[L^2 M^2 T^{-2}]$

Answer: a



Watch Video Solution

6. If the magnitude of length is halved and that of mass is double then dimension of force is

A. $[M^2 L^{-2} T^{-2}]$

B. $[M^2 L^{-1/2} T^{-2}]$

C. $[M^2 L^{1/2} T^{-2}]$

D. $[M^1 L^1 T^{-2}]$

Answer: d



Watch Video Solution

7. Out of the following pairs, which one does NOT have identical dimensions?

A. Energy and moment of force

B. Work and torque

C. Density and surface energy

D. Pressure and stress

Answer: c



Watch Video Solution

8. Which of the following equations is dimensionally correct?

A. pressure = Energy per units volume

B. pressure = Energy per units area

C. pressure = Momentum \times Volume \times
time

D. pressure = Force \times area

Answer: a



Watch Video Solution

9. The dimensional formul for impulse is same as the dimensional formula for

A. acceleration

B. force

C. momentum

D. rate of change in momentum

Answer: c



Watch Video Solution

10. The dimensions of is that of

A. Velocity

B. Time

C. Capacitance

D. Distance

Answer: a



Watch Video Solution

11. Which of the following pairs has the same dimensions?

- A. Current density and charge density
- B. Angular momentum and momentum
- C. Spring constant and surface energy
- D. Force and torque

Answer: c



Watch Video Solution

12. A small steel ball of mass m and radius r is falling under gravity through a viscous liquid of coefficient of viscosity η . If g is the value of acceleration due to gravity. Then the terminal velocity of the ball is proportional to (ignore buoyancy)

A. $v \propto \frac{m g r}{\eta}$

B. $v \propto m g \eta r$

C. $v \propto \frac{m g}{\eta r}$

D. $v \propto \frac{\eta m g}{r}$

Answer: c



Watch Video Solution

13. A force F is given by $F = at + bt^2$, where t is time . What are the dimensions of a and b ?

- A. $[M^1 L^2 T^{-1}]$ and $[M^1 L^1 T^0]$
- B. $[M^1 L^1 T^{-3}]$ and $[M^1 L^1 T^{-4}]$
- C. $[M^1 L^1 T^{-4}]$ and $[M^1 L^1 T^1]$
- D. $[M^1 L^{-3} T^1]$ and $[M^1 L^1 T^{-4}]$

Answer: b



Watch Video Solution

14. Write the dimensions of a/b in the relation

$F = a\sqrt{x} + bt^2$ where F is force x is distance

and t is time.

A. $[M^0 L^0 T^{-2}]$

B. $[M^0 L^{1/2} T^2]$

C. $[M^0 L^{1/2} T^2]$

D. $[M^0 L^{-1/2} T^2]$

Answer: c



Watch Video Solution

15. For the equation $F = A^a v^b d^c$ where F is force, A is area, v is velocity and d is density with the dimensional analysis gives the following values for the exponents.

A. 1,2,1

B. 2,1,1

C. 1,1,2

D. 0,1,1

Answer: a



Watch Video Solution

16. Using the principle of homogeneity of dimensions, find which of the following relation is correct? [T is the period, a is the radius of the orbit and M is the mass of the sun.

$$\text{A. } T^2 = \frac{4\pi^2 a^3}{G}$$

$$\text{B. } T^2 = \frac{4\pi^2 a^3}{GM}$$

$$\text{C. } T^2 = 4\pi^2 a^3$$

$$\text{D. } T^2 = \frac{4\pi^2 a^3}{GM^2}$$

Answer: b



Watch Video Solution

17. The period T of a soap bubble under *SHM* is given by $T = P^a D^b S^c$, where P is pressure, D , is density and S is surface tension. Then the values of a , b and c are

A. $-\frac{3}{2}, \frac{1}{2}, 1$

B. $-1, -2, 3$

C. $\frac{1}{2}, -\frac{3}{2} - \frac{1}{2}$

D. $1, 2, \frac{1}{3}$

Answer: a



Watch Video Solution

18. The equation of a wave is given by

$$Y = A \sin \omega \left(\frac{x}{v} - k \right), \text{ where } \omega \text{ is the angular}$$

velocity and v is the linear velocity. Find the dimension of k .

A. LT

B. T

C. T^{-1}

D. T^2

Answer: b



Watch Video Solution

19. The dimensions of $\frac{a}{b}$ in the equation

$$P = \frac{a - t^2}{bx}$$

where P is pressure, x is

distance and t is time are

A. $[M^1 L^2 T^{-2}]$

B. $[M^1 L^0 T^{-2}]$

C. $[M^{-1} L^{-2} T^2]$

D. $[M^1 L^{-2} T^2]$

Answer: b



Watch Video Solution

20. The equation of the state of some gases

can be expressed as $\left(P + \frac{a}{V^2} (V - b) = RT \right)$

. Here P is the pressure, V is the volume, T is

the absolute temperature and a, b, R are

constants. The dimensions of $\frac{a}{b}$ are

A. $[M^1 L^2 T^{-2}]$

B. $[ML^{-1}T^{-2}]$

C. $[M^0 L^3 T^0]$

D. $[M^0 L^6 T^0]$

Answer: a



Watch Video Solution

21. If the speed of light c , acceleration due to gravity (g) and pressure (p) are taken as the fundamental quantities then the dimension of gravitational constant is

A. $[c^2 g^0 p^{-2}]$

B. $[c^0 g^2 p^{-1}]$

C. $[c g^3 p^{-2}]$

D. $[c^1 g^0 p^{-1}]$

Answer: b



Watch Video Solution

22. The value of acceleration due to gravity is 980cm s^{-2} . What will be its value if the unit of length is kilometer and that of time is minute?

A. 980 km min^3

B. 98 km min^3

C. 35.28 km min^{-2}

D. 28.35 km min^{-2}

Answer: c



Watch Video Solution

Critical Thinking Order Of Magnitude And Significant Figure

1. The magnitude of any physical quantity can be expressed as $A \times 10^n$ where n is a number called order of magnitude and A is

$$A. 0.1 \leq A \leq 1$$

B. $0.5 \leq A < 5$

C. $5 \leq A < 9$

D. $1 \leq A > 9$

Answer: b



Watch Video Solution

2. The radius of the earth is 6400 km, the order of magnitude is

A. 10^7m

B. $10^4 m$

C. $10^3 m$

D. $10^2 m$

Answer: a



Watch Video Solution

3. The radius of the earth of 49 and the order of magnitude of 51

A. is same

B. differs by 1

C. is 1

D. is 2

Answer: b



Watch Video Solution

4. Find the number of seconds in 1 year.

Express them in order magnitude.

A. $8.64 \times 10^4 s, 10^5 s$

B. $6.84 \times 10^4 s, 10^4 s$

C. $8.64 \times 10^5 s, 10^5 s$

D. $6.85 \times 10^4 s, 10^5 s$

Answer: a



Watch Video Solution

5. Figure which is of some significance but it does not necessarily denote certainty is defined as _____

A. special figures

B. characteristic figures

C. unknown figures

D. significant figure

Answer: d



Watch Video Solution

6. The number of significant figures in all the given numbers 25.12, 2009, 4.156 and 1.217×10^{-4} is

A. 1

B. 2

C. 3

D. 4

Answer: d



Watch Video Solution

7. The value of $\pi/53.2$ with due regard to significant figures is,

A. 13.000

B. 13.00

C. 13.0

D. 13

Answer: c



Watch Video Solution

8. In the reading 2.614 cm of measurement with a vernier calliper, only uncertain figure is

A.

B.

C.

D.

Answer: c



Watch Video Solution

9. The side of a rectangle are 6.01 m and 12 m

Taking the significant figures into account , the

area of the rectangle is

A. 72.00cm^2

B. 72.1cm^2

C. 72m^2

D. 72.12cm^2

Answer: c



Watch Video Solution

Critical Thinking Accuracy And Errors In Measurements

1. Estimate the mean absolute error from the following data .

20.17,21.23,20.79,22.07,21.78

A. 0.85

B. 0.58

C. 0.03

D. 0.01

Answer: b



View Text Solution

2. In the expression $A = \frac{xy^3}{z^2}$ the percentage

error is given by

A. $\left(\frac{\Delta x}{x} + 3\frac{\Delta y}{y} - 2\frac{\Delta z}{z} \right) \times 100\%$

B. $\left(\frac{\Delta x}{x} + \frac{3\Delta y}{y} + \frac{2\Delta z}{z} \right) \times 100\%$

C. $\left(\frac{\Delta x}{x} - \frac{3\Delta y}{y} - \frac{2\Delta z}{y} \right) \times 100\%$

D. $\left(\frac{\Delta x}{x} - 3\frac{\Delta y}{y} + 2\frac{\Delta z}{z} \right) \times 100$

Answer: b



Watch Video Solution

3. The least count of a screw gauge is 0.005 cm . The diameter of wire is 0.002 cm as measured by it. The percentage error in measurement is

A. 0.25

B. 0.2

C. 0.15

D. 0.05

Answer: a



Watch Video Solution

4. The error in measurement of radius of a sphere is 0.1% then error in its volume is -

A. 0.1 %

B. 0.2 %

C. 0.25 %

D. 0.3 %

Answer: d



Watch Video Solution

5. The pressure on a square plate is measured by measuring the force on the plate and the length of the sides of the plate. If the maximum error in measurement of force and length are respectively 4% and 2%, the maximum error in the measurement of pressure is

A. 0.01

B. 0.02

C. 0.06

D. 0.08

Answer: d



Watch Video Solution

6. The percentage error in the measurement of mass of a body is 0.75% and the percentage error in the measurement of its speed is 1.85% . Then the percentage error in the measurement of its kinetic energy is

A. 7.05%

B. 4.45 %

C. 2.6 %

D. 1.1 %

Answer: b



Watch Video Solution

7. The error in the measurement of length (L) of time simple pendulum is 0.1% and the error in the period (T) is 3%. The maximum possible error in the measurement of $\frac{L}{T^2}$ is

A. 2.9 %

B. 3.1 %

C. 5.9 %

D. 6.1 %

Answer: d



Watch Video Solution

8. The period of oscillation of a simple

pendulum is given by $T = 2\pi\sqrt{\frac{l}{g}}$ where l is

about 100 cm and is known to have 1 mm accuracy. The period is about 2 s. The time of 100 oscillation is measured by a stop watch of least count 0.1 s. The percentage error is g is

A. 0.1 %

B. 1 %

C. 0.3 %

D. 0.8 %

Answer: c



Watch Video Solution

9. The length , breadth and height of a rectangular block of wood were measured to be $l = 13.12 \pm 0.02\text{cm}$, $b = 7.18 \pm 0.01 \text{ cm}$, $h = 4.16 \pm 0.02 \text{ cm}$

The percentage error in the volume of the block will be .

A. 0.07

B. 0.77 %

C. 0.72 %

D. 0.27 %

Answer: b



Watch Video Solution

10. The heat dissipated in a resistance can be determined from the relation : $H = \frac{I^2 R t}{4.2}$ cal

If the maximum errors in the measurement of current , resistance , and time are 2%, 1%, and 1%, what would be the maximum error in the dissipated heat ?

A. 0.05

B. 0.04

C. 0.06

D. 0.5 %

Answer: c



View Text Solution

Critical Thinking Miscellaneous

1. If momentum (p), area (A) and time(t) are taken to be fundamental quantities then energy has the dimensional formula

A. $[P^1 A^{-1/2} T^{-1}]$

B. $[P^1 A^{1/2} T^{-1}]$

C. $[P^2 A^{-1} T^{-1}]$

D. $[P^1 A^{-1} T^{-1}]$

Answer: b



Watch Video Solution

2. Assertion : Avogadro number is the number of atoms in one gram mole.

Reason : Avogadro number is a dimensionless constant.

A. Assertion is True , Reason is True , Reason is a correct explanation for Assertion.

B. Assertion is True , Reason is True , Reason is not a correct explanation for Assertion

C. Assertion is True , Reason is False

D. Assertion is False , Reason is False

Answer: c



Watch Video Solution

3. Assertion : Avogadro number is the number of atoms in one gram mole.

Reason : Avogadro number is a dimensionless constant.

A. Assertion is True , Reason is True ,
Reason is a correct explanation for
Assertion

B. Assertion is True , Reason is True ,
Reason is not a correct explanation for
Assertion

C. Assertion is True, Reason is False

D. Assertion is False, Reason is False .

Answer: a



Watch Video Solution

Competitive Thinking Fundamental And Derived Units

1. S.I. Unit of surface tension is:

A. $\frac{N}{m^2}$

B. $\frac{J}{m}$

C. $J - m$

D. $\frac{J}{m^2}$

Answer: d



Watch Video Solution

2. Which of the following is not the unit of energy ?

A. watt - hour

B. electron volt

C. N m

D. kgm^2s^{-2}

Answer: c



3. The units of constants a in van der Waals' equations is

A. Nm^2 / mole

B. Nm^2 / mole

C. N^2m / mole

D. m^2

Answer: d



[Watch Video Solution](#)

4. Units of a and b in van der Waal's equation

$$\left(P + \frac{an^2}{V^2}\right)(V - nb) = nRT \text{ are}$$

A. m^3 / mole

B. m^2 / mole

C. m/mole

D. m^3

Answer: a



Watch Video Solution

5. The S. I units of the constant in Wein's displacement law are

A. cmK^{-1}

B. mk

C. cm^2k^{-1}

D. cmk^{-2}

Answer: b



Watch Video Solution

6. Which of the following is the unit of specific heat

A. kcal / gm k

B. cal/gm k

C. j/kg k

D. erg / kg k

Answer: c



Watch Video Solution

7. A-s is unit of

A. capacitance

B. charge

C. energy

D. power

Answer: b



Watch Video Solution

8. S.I unit of specific resistance is

A. Ωcm

B. Ωm

C. Ω / cm

D. mho-cm

Answer: b



Watch Video Solution

9. The unit of permeability of vacuum (μ_0) is

_____ .

A. $\frac{N}{A}$

B. $\frac{N}{A^2}$

C. NA

D. $\frac{J}{A^2}$

Answer: b



Watch Video Solution

10. 1 Tesla =

A. 1 Wb/m

B. 1 J/Am

C. 1 N/Am

D. 1Am/N

Answer: c



Watch Video Solution

11. Units of λ in radiocativity is

A. m

B. (Unit of half life)⁻¹

C. (year)⁻¹

D. sec

Answer: b



Watch Video Solution

12. If $x = at + bt^2$ where x is in meter (m) and t is in hour (hr) then unit of b will be .

A. m^2 / hr

B. m

C. m / hr

D. m / hr^2

Answer: d



Watch Video Solution

13. If the unit of length and force be increased four times, then the unit of energy is

- A. increased 4 times
- B. increased 8 times
- C. increased 16 times
- D. decreased 16 times

Answer: c



Watch Video Solution

14. The surface tension of a liquid is 10^8 dyne cm^{-1} . It is equivalent to

A. $10^7 N / m$

B. $10^6 N / m$

C. $10^5 N / m$

D. $10^4 N / m$

Answer: c



Watch Video Solution

Competitive Thinking Dimensional Analysis

1. If R and L represent respectively resistance and self inductance, which of the following combinations has the dimensions of frequency

A. $\sqrt{\frac{L}{R}}$

B. $\frac{L}{R}$

C. $\sqrt{\frac{R}{L}}$

D. $\frac{R}{L}$

Answer: b



Watch Video Solution

2. A quantity X is given by $\epsilon_0 L \frac{\Delta V}{\Delta t}$, where ϵ_0 is the permittivity of free space, L is a length, ΔV is a potential difference and Δt is a time interval. The dimensional formula for X is the same as that of -

A. Resistance

B. Charge

C. Voltage

D. Current

Answer: d



[Watch Video Solution](#)

3. Planck constant has the same dimensions as

A. energy

B. angular momentum

C. mass

D. force

Answer: b



[Watch Video Solution](#)

4. The dimensions of Planck's constant is the same as the product of

A. Force and time

B. Force, displacement and time

C. Force and velocity

D. Force and displacement

Answer: b



Watch Video Solution

5. Which of the following set have different dimensions?

A. Pressure, Young's modulus , stress

B. e.m.f potential difference , electric potential

C. Heat , work done energy

D. dipole moment , electric flux, electric field

Answer: d





Watch Video Solution

6. The dimension of universal gravitational constant are

A. $[ML^2T^{-1}]$

B. $[M^{-1}L^3T^{-2}]$

C. $[M^{-1}L^2T^{-2}]$

D. $[ML^3T^{-2}]$

Answer: b



Watch Video Solution

7. Dimensions of angular momentum is

A. $[M^1 L^2 T^{-1}]$

B. $[M^1 L^{-2} T^{-2}]$

C. $[M^1 L^2 T^{-1}]$

D. $[M^1 L^0 T^{-1}]$

Answer: c



Watch Video Solution

8. The dimensions of $\epsilon_0\mu_0$ are

A. $[M^0 L^{-2} T^2]$

B. $[M^0 L^2 T^{-2}]$

C. $[M^0 L^1 T^{-1}]$

D. $[M^0 L^{-1} T^1]$

Answer: c



Watch Video Solution

9. The surface tension is $T = \frac{F}{l}$, then the dimensions of surface tension is

- A. $[M^1 L^2 T^2]$
- B. $[M^1 L^0 T^{-2}]$
- C. $[M^1 L^2 T^{-2}]$
- D. $[M^0 L^0 T^{-2}]$

Answer: b



Watch Video Solution

10. Dimension of force constant is given by

A. $[M^1 L^1 T^{-2}]$

B. $[M^0 L^1 T^{-1}]$

C. $[M^1 L^0 T^{-2}]$

D. $[M^1 L^0 T^{-1}]$

Answer: c



Watch Video Solution

11. The dimensions of k in equation $F = kx$ are

A. $[M^1 L^0 T^{-2}]$

B. $[M^0 L^1 T^{-1}]$

C. $[M^1 L^1 T^{-2}]$

D. $[M^1 L^0 T^{-1}]$

Answer: a



Watch Video Solution

12. An object is moving through the liquid. The viscous damping force acting on it is

proportional to the velocity. Then dimensions of constant of proportionality are

A. $[M^1 L^{-1} T^{-1}]$

B. $[M^1 L^1 T^{-1}]$

C. $[M^0 L^1 T^{-1}]$

D. $[M^1 L^0 T^{-1}]$

Answer: d



Watch Video Solution

13. The dimensional formula for Reynold's number is

A. $[L^0 M^0 T^0]$

B. $[L^1 M^1 T^1]$

C. $[L^{-1} M^1 T^1]$

D. $[L^1 M^1 T^{-1}]$

Answer: a



Watch Video Solution

14. The unit of universal gas constant is

A. $[ML^2T^{-2}\theta^{-}]$

B. $[ML^2T^{-2}\theta]$

C. $[ML^3T^{3/2}T^{-2}]$

D. None of these

Answer: a



Watch Video Solution

15. The relation between force F and density d

is $F = \frac{x}{\sqrt{d}}$.

The dimension of x is

A. $[L^{-1/2} M^{3/2} T^{-2}]$

B. $[L^{1/2} M^{1/2} T^{-2}]$

C. $[L^{-1} M^{1/2} T^{-2}]$

D. $[L^1 M^{1/2} T^{-2}]$

Answer: a



Watch Video Solution

16. In the following dimensionally correct equation $F = \frac{X}{\text{Linear Density}}$, where F is the force. What is the dimensional formula for X?

A. $[M^2 L^0 T^{-2}]$

B. $[M^0 L^0 T^{-1}]$

C. $[L^2 T^{-2}]$

D. $[M^0 L^2 T^{-2}]$

Answer: a



Watch Video Solution

17. The units of constants a in van der Waal's equation is

A. $[M^1 L^{-1} T^{-2} mol^{-2}]$

B. $[M^1 L^3 T^{-2} mol^{-2}]$

C. $[M^1 L^5 T^{-2} mol^{-2}]$

D. $[M^1 L^3 T^{-2} mol^{-1}]$

Answer: c



Watch Video Solution

18. Let $[\epsilon_0]$ denote the dimensional formula of the permittivity of vacuum. If M= mass, L=length, T=Time and A= electric current, then:

A. $[\epsilon_0] = [M^1 L^{-3} T^2 A]$

B. $[\epsilon_0] = [M^{-1} L^{-3} T^4 A^2]$

C. $[\epsilon_0] = [M^{-1} L^{-3} T^4 A^2]$

D. $[M^0 L^0 T^0 A^0]$

Answer: b



Watch Video Solution

19. The dimensional formula for electric field is

A. $[M^1 L^{-2} T^{-3} A^{-2}]$

B. $[M^1 L^2 T^{-3} A^{-1}]$

C. $[M^1 L^1 T^3 A^1]$

D. $[M^0 L^0 T^0 A^0]$

Answer: c



Watch Video Solution

20. The dimensional formula of capacitance is _____. Take Q as the dimension formula of charge.

A. $[M^1 L^{-2} T^{-2} Q^{-2}]$

B. $[M^1 L^2 T^{-2} Q^{-2}]$

C. $[M^1 L^1 T^2 Q^1]$

D. $[M^{-1} L^{-2} T^2 Q^2]$

Answer: d



Watch Video Solution

21. The dimensions of $\frac{1}{2} \epsilon_0 E^2$, where ϵ_0 is permittivity of free space and E is electric field, is :-

A. $[L^{-2} M^1 T^2]$

B. $[L^1 M^1 T^{-2}]$

C. $[L^2 M^1 T^2]$

D. $[L^1 M^1 T^1]$

Answer: b



Watch Video Solution

22. Dimension of magnetic flux is

A. $[M^0 L^1 T^0 A^1]$

B. $[M^0 L^{-3} T^0 A^1]$

C. $[M^0 L^1 T^1 A^{-1}]$

D. $[M^0 L^2 T^0 A^1]$

Answer: a



Watch Video Solution

23. Select the dimensional formula of $B^2 / 2\mu_0$

.

A. $[M^1 L^1 T^0 A^1]$

B. $[M^0 L^{-3} T^{-2} A^{-1}]$

C. $[M^0 L^1 T^1 A^{-1}]$

D. $[M^0 L^2 T^0 A^1]$

Answer: d



View Text Solution

24. Dimensional formula of magnetic flux is

A. $[M^1 L^1 T^2]$

B. $[M^{-1} L^1 T^{-2}]$

C. $[M^{-1} L^{-1} T^{-1} A^1]$

D. $[M^1 L^{-1} T^{-2}]$

Answer: b



Watch Video Solution

25. The dimensional formula of mobility is

_____ .

A. $[M^{-1}L^2T^2A^1]$

B. $[M^1L^2T^{-2}A^{-1}]$

C. $[M^1L^{-2}T^2A^1]$

D. $[M^1L^3T^{-2}A^{-1}]$

Answer: d



Watch Video Solution

26. Dimensions of solar constant are

A. $[MLT^{-2}]$

B. $[M^0L^0T^0]$

C. $[ML^0T^{-3}]$

D. $[M^0LT^{-3}]$

Answer: c



Watch Video Solution

27. The velocity v of a particle at time t is given by $v = at + \frac{b}{t + c}$, where a , b and c are constants. The dimensions of a , b , c are respectively :-

- A. L, LT and T^2
- B. LT^2, L and T
- C. L^2, T and LT^3
- D. LT^3, LT and L

Answer: b



Watch Video Solution

28. If E, M, J and G respectively denote energy , mass angular momentum and gravitational constant then $\frac{EJ^2}{M^2G^2}$ has the dimensions of .

A. length

B. angle

C. mass

D. time

Answer: b



29. The function f is given by $f = A \sin \alpha x + B \cos \beta t$, where x is displacement and t is the time. The dimensions of α / β is

- A. velocity
- B. velocity gradient
- C. angular velocity
- D. angular momentum

Answer: a



Watch Video Solution

30. $X = 3YZ^2$ find dimension of Y in (MKSA) system, if X and Z are the dimension of capacity and magnetic field respectively

A. $[M^{-3}L^{-2}T^{-4}A^{-1}]$

B. $[M^1L^{-2}]$

C. $[M^{-3}L^{-2}T^4A^4]$

D. $[M^{-3}L^{-2}T^8A^4]$

Answer: d



Watch Video Solution

31. Plank 's constant (h) speed of length in vacium (C) and newton 's gravitational constant (G) are three fundamental constant .Which of the following combinations of these has the dimension of length?

A. $\sqrt{\frac{Ge}{h^{\frac{1}{2}}}}$

B. $\frac{\sqrt{hG}}{c^{3/2}}$

C. $\frac{\sqrt{hG}}{c^{3/2}}$

D. $\sqrt{\frac{hc}{G}}$

Answer: b



Watch Video Solution

32. A spherical liquid drop is placed on a horizontal plane . A small disturbance cause the volume of the drop to oscillate . The time period oscillation (T) of the liquid drop depends on radius (r) of the drop , density (ρ)

and surface tension (S) of the liquid.

Which amount the following will be be a possible expression for T (where k is a dimensionless constant)?

A. $k\sqrt{\frac{\rho r}{S}}$

B. $k\sqrt{\frac{\rho^2 r}{S}}$

C. $k\sqrt{\frac{\rho^2 r}{S}}$

D. $k\sqrt{\frac{\rho r^2}{S^2}}$

Answer: c



Watch Video Solution

33. A physical energy of the dimension of length that can be formula cut of c , G and $\frac{e^2}{4\pi\epsilon_0}$ is [c is velocity of light G is universal constant of gravitation e is charge

A. $\frac{1}{c^2} \left[G \frac{e^2}{4\pi \epsilon_0} \right]^{1/2}$

B. $c^2 \left[G \frac{e^2}{4\pi \epsilon_0} \right]^{1/2}$

C. $\frac{1}{c^2} \left[\frac{e^2}{G4\pi \epsilon_0} \right]^{1/2}$

D. $\frac{1}{e} G \frac{e^2}{4\pi \epsilon_0}$

Answer: a



Watch Video Solution

34. In the relation, $P = \frac{\alpha}{\beta} e^{\frac{\alpha Z}{k\theta}}$ P is pressure, Z is distance, k is Boltzmann constant and θ is the temperature. The dimensional formula of β will be-

A. $[M^0 L^2 T^0]$

B. $[M^1 L^2 T^{-1}]$

C. $[M^1 L^0 T^{-1}]$

D. $[M^0 L^2 T^{-1}]$

Answer: a



Watch Video Solution

35. The number of significant figure in
0.002305 is

A. 6

B. 4

C. 7

D. 2

Answer: b



Watch Video Solution

36. The respective number of significant figures for the numbers 23.023, 0.0003 and 2.1×10^{-3} are

A. 4,4,2

B. 5,1,2

C. 5,1,5

D. 5,5,2

Answer: b



Watch Video Solution

37. In a experiment the angle are required to be measured using an instrument. 26 divisions of the main scale exactly coincide with the 30 divisions of the vernier scale . If the smallest division of the main scale is half -a-degree ($= 0.5^\circ$) then the least count of the instrument is .

A. one minute

B. half minute

C. one degree

D. half degree

Answer: a



Watch Video Solution

38. A vernier calipers has 1mm marks on the main scale. It has 20 equal divisions on the Verier scale which match with 16 main scale

divisions. For this Vernier calipers, the least count is

A. 0.02 cm

B. 0.05 cm

C. 0.1 mm

D. 0.2 mm

Answer: d



Watch Video Solution

39. The diameter of a cylinder is measured using a Vernier callipers with no zero error. It is found that the zero of the Vernier scale lies between 5.10cm and 5.15cm of the main scale. The Vernier scale has 50 divisions equivalent to 2.45cm . The 24^{th} division of the Vernier scale exactly coincides with one of the main scale divisions. the diameter of the cylinder is

A. 5.112 cm

B. 5.124 cm

C. 5.136 cm

D. 5.148 cm

Answer: b



Watch Video Solution

40. A student measured the length of a rod and wrote it as 3.50 cm. Which instrument did he use to measure it ?

A. A meter scale

B. A vernier calliper where the 10 divisions in vernier scale matches with 9 division in main scale and main scale has 10 divisions in 1 cm

C. A screw gauge having 100 divisions in the circular scale and pitch as 1 mm

D. A screw gauge having 50 divisions in the circular scale and pitch as 1mm

Answer: b



Watch Video Solution

41. In a vernier callipers, one main scale division is x cm and n divisions of the vernier scale coincide with $(n-1)$ divisions of the main scale. The least count (in cm) of the callipers is

∴

A. $\left(\frac{n-1}{n}\right) \times x$

B. $\frac{nx}{(n-1)}$

C. $\frac{x}{n}$

D. $\frac{x}{(n-1)}$

Answer: c



Watch Video Solution

42. A screw gauge gives the following reading when used to measure the diameter of a wire.

Main scale reading : 0mm

Circular scale reading : $52 \div \text{isions}$

Given that 1mm on main scale corresponds to 100 divisions of the circular scale. the diameter of wire from the above data is :

A. 0.52 cm

B. 0.052 cm

C. 0.026 cm

D. 0.005 cm

Answer: b



Watch Video Solution

43. A spectrometer gives the following reading when used to measure the angle of a prism.

Main scale reading = 58.5 degree

Vernier scale reading = 0.9 divisions

Given that 1 divisions on main scale corresponds to 0.5 degree. Total divisions on the vernier scale is 30 and match with 29 divisions of the main scale. the angle of the prism from the above data is :

A. 58.59 degree

B. 58.77 degree

C. 58.65 degree

D. 59 degree

Answer: a



Watch Video Solution

Competitive Thinking Accuracy And Errors In Measurements

1. Choose the INCORRECT statement out of the following .

A. Every measurement by any measuring instrument has some error.

- B. Every calculated physical quantity that is based on measured values has some error
- C. A measurement can have more accuracy but less precision and vice versa
- D. The percentage error is different form relative error

Answer: d



Watch Video Solution

2. Assertion: The error in the measurement of radius of sphere is 0.3% . The permissible error in its surface area is 0.6% .

Reason: The permissible error is calculated by

the formula $\frac{\Delta A}{A} = \frac{4\Delta r}{r}$.

A. Assertion is True , Reason is True ,

Reason is a correct explanation for

Assertion .

B. Assertion is True , Reason is True ,

Reason is not a correct explanation for

Assertion

C. Assertion is True , Reason is False

D. Assertion is False, Reason is False

Answer: c



Watch Video Solution

3. If $Y = a - b$, the maximum percentage error in the measurement of Y will be

A. $\left[\frac{\Delta a}{a - b} - \frac{\Delta b}{a - b} \right] \times 100$

$$\text{B. } \left[\frac{\Delta a}{a - b} + \frac{\Delta b}{b - a} \right] \times 100$$

$$\text{C. } \left[\frac{\Delta a}{a} - \frac{\Delta b}{b} \right] \times 100$$

$$\text{D. } \left[\frac{\Delta a}{a} - \frac{\Delta b}{b} \right] \times 100$$

Answer: b



Watch Video Solution

4. The radius of a sphere is $(5.3 \pm 0.1)\text{cm}$. The percentage error in its volume is

$$\text{A. } 3 + 6.01 \times \frac{100}{5.3}$$

B. $\frac{1}{3} \times 0.01 \times \frac{100}{5.3}$

C. $\left(\frac{3 \times 0.1}{5.3} \right) \times 100$

D. $\frac{0.1}{5.3} \times 100$

Answer: c



Watch Video Solution

5. Resistance of a given wire is obtained by measuring the current flowing in it and the voltage difference applied across it. If the percentage errors in the measurement of the

current and the voltage difference are 3% each, then error in the value of resistance of the wire is

A. 0.06

B. Zero

C. 0.01

D. 0.03

Answer: a



Watch Video Solution

6. In an experiment four quantities a, b, c and d are measured with percentage error 1%, 2%, 3%, and 4% respectively. Quantity P is calculated as follows:

$$P = \frac{a^3 b^2}{cd} \quad \text{\% error in } P \text{ is}$$

A. 0.14

B. 0.1

C. 0.07

D. 0.04

Answer: a



Watch Video Solution

7. Let $x = \left[\frac{a^2 b^2}{c} \right]$ be the physical quantity. If the percentage error in the measurement of physical quantities $a, b,$ and c is 2, 3 and 4 per cent respectively, then percentage error in the measurement of x is

A. 0.07

B. 0.14

C. 0.21

D. 0.28

Answer: c



Watch Video Solution

8. Two full turns of the circular scale of a screw gauge cover a distance of 1 mm on its main scale . The total number of divisions on the circular scale is 50. Further more it is found that the screw gauge has a zero error of - 0.02 mm . While measuring the diameter of a thin

wire , a student notes the main scale reading of 4 mm and the number of circular scale divisions in line with the main scale as 37. The diameter of the wire is

A. 3.73 cm

B. 3.67 cm

C. 3.38 cm

D. 3.32 cm

Answer: c



Watch Video Solution

9. A screw gauge with a pitch of 0.5mm and a circular scale with 50 divisions is used to measure the thickness of a thin sheet of Aluminium. Before starting the measurement, it is found that when the jaws of the screw gauge are brought in contact, the 45^{th} division coincides with the main scale line and the zero of the main scale is barely visible. What is the thickness of the sheet if the main scale reading is 0.5mm and the 25^{th} division coincides with the main scale line?

A. 0.80 cm

B. 0.70 cm

C. 0.50 cm

D. 0.75 cm

Answer: a



Watch Video Solution

10. A student measured the diameter of a small steel ball using a screw gauge of least count 1.001cm . The main scale reading is 5mm and

zero of circular scale division coincides with 25 divisions above the reference level. If screw gauge has a zero error of -0.004cm , the correct diameter of the ball is

A. 0.521cm

B. 0.525cm

C. 0.053 cm

D. 0.526 cm

Answer: d



Watch Video Solution

11. The density of a solid ball is to be determined in an experiment. The diameter of the ball is measured with a screw gauge, whose pitch is 0.5mm and there are 50 divisions on the circular scale. The reading on the main scale is 2.5mm and that on circular scale is 20 divisions. If the measured mass of the ball has a relative error of 2% , the relative percentage error in the density is

A. 0.9%

B. 2.4 %

C. 3.1 %

D. 4.2 %

Answer: c



Watch Video Solution

12. A force \vec{F} applied on a square plate of length L . If the percentage error in the determination of L is 3% and in F is 4% , the

permissible error in the calculation of pressure is

A. 0.13

B. 0.1

C. 0.07

D. 0.12

Answer: b



Watch Video Solution

13. The density of a material in the shape of a cube is determined by measuring three sides of the cube and its mass. If the relative errors in measuring the mass and length are respectively 1.5 % and 1 %, the maximum error in determining the density is:

A. 4.5 %

B. 6 %

C. 2.5 %

D. 3.5 %

Answer: a



Watch Video Solution

14. In the determination of Young's modulus

$\left(\left(Y = \frac{4MLg}{\pi ld^2} \right) \right)$ by using Searle's method, a

wire of length $L = 2m$ and diameter

$d = 0.5mm$ is used. For a load $M = 2.5kg$, an

extension $l = 0.25mm$ in the length of the

wire is observed. Quantities D and l are

measured using a screw gauge and a

micrometer, respectively. They have the same

pitch of 0.5mm . the number of divisions on their circular scale is 100. the contribution to the maximum probable error of the Y measurement

A. due to the errors in the measurements of d and l are the same.

B. due to the error in the measurement of d is twice that due to the error in the measurement of l .

C. due to the error in the measurement of l is twice that due to the error in the

measurement of d .

D. due to the error in the measurement of d is four times that due to the error in the measurement of l .

Answer: a



Watch Video Solution

15. If the time period of a simple pendulum is

$T = 2\pi\sqrt{l/g}$, then the fractional error in

acceleration due to gravity is

A. $\frac{4\pi^2 \Delta l}{\Delta T^2}$

B. $\frac{\Delta l}{l} - 2\frac{\Delta l}{T}$

C. $\frac{\Delta l}{l} + 2\frac{\Delta T}{T}$

D. None of these

Answer: c



Watch Video Solution

16. A student measures the value of g with the help of a simple pendulum using the formula

$g = \frac{4\pi^2 L}{T^2}$. He measures length L with a meter scale having least count 1 mm and finds it 98.0 cm . The time period is measured with the help of a watch of least count 0.1s . The time of 20 oscillations is found to be 40.4 s. The error Δg in the measurement of g is (in m / s^2).

A. $9.68 \left[\frac{0.1}{98} + 0.1 \right]$

B. $9.86 \left[\frac{1}{98} + 0.1 \right]$

C. $9.68 \left[\frac{0.1}{98} + \frac{0.1}{20} \right]$

D. $9.68 \left[\frac{1}{98} + \frac{1}{20} \right]$

Answer: a



Watch Video Solution

17. The period of oscillation of a simple

pendulum is $T = 2\pi \sqrt{\frac{L}{g}}$. Measured value of L

is 20.0cm known to 1mm accuracy and time for

100 oscillation of the pendulum is found to be

90s using a wrist watch of 1s resolution. The

accuracy in the determination of g is :

A. 0.02

B. 0.03

C. 0.01

D. 0.05

Answer: b



Watch Video Solution

18. The following observations were taken for determining the surface tension of water by capillary tube method: diameter of capillary , $D = 1.25 \times 10^{-2}m$ and rise of water in

capillary , $h = 1.45 \times 10^{-2}m$. Taking $g = 9.80ms^{-2}$ and using the relation $T = (rgh/2) \times 10^3Nm^{-1}$, what is the possible error in measurement of surface tension T ?

(a) 2.4 % (b) 15 % (c) 1.6 % (d) 0.15%

A. 0.15 %

B. 1.5 %

C. 2.4 %

D. 10 %

Answer: b



Watch Video Solution

Competitive Thinking Miscellaneous

1. One femtometer is equivalent to

A. 10^{15} m

B. 10^{-15} m

C. 10^{-12} m

D. 10^{12} m

Answer: b



Watch Video Solution

2. Which of the following units denotes the dimensions ML^2/Q^2 , where Q denotes the electric charge ?

A. henry (H)

B. H/m^2

C. weber (Wb)

D. Wb/m^2

Answer: a



Watch Video Solution

3. Which of the following physical quantities represent the dimensions of $\frac{b}{a}$ in the relations and t time ?

- A. Power
- B. Surface tension
- C. Torsional constant
- D. Force

Answer: c



Watch Video Solution

4. If force (F), velocity (V) and time (T) are taken as fundamental units, then the dimensions of mass are

A. $[FVT^{-1}]$

B. $[FVT^{-2}]$

C. $[FV^{-1}T^{-1}]$

D. $[FV^{-1}T]$

Answer: d



Watch Video Solution

5. If energy (E), velocity (V) and time (T) are chosen as the fundamental quantities, the dimensions formula of surface tension will be

A. $[EV^{-2}T^{-1}]$

B. $[EV^{-1}T^{-2}]$

C. $[EV^{-2}T^{-2}]$

D. $[E^{-2}V^{-1}T^{-3}]$

Answer: c



Watch Video Solution

6. If the velocity of surface wave (v) depends upon surface tension (T), coefficient of viscosity (η) and density (ρ) then the expression for v will be .

A. $\frac{T^2}{\rho\eta}$

B. $\frac{T}{\eta}$

C. $\frac{\eta\rho}{T^3}$

D. $\frac{\rho}{\eta}$

Answer: b



Watch Video Solution

7. The ratio of the dimension of Planck's constant and that of moment of inertia is the dimension of

A. angular

B. time

C. velocity

D. frequency

Answer: d



Watch Video Solution

Evaluation Test

1. When dimensions of a given physical quantity are given, the physical quantity is unique .

A. The statement and its converse both are true .

B. The statement is false but its converse is true.

C. The statement is true but its converse is true .

D. The statement is true but its converse is false .

Answer: c



Watch Video Solution

2. The quantities A and B are related by the relation $A/B = m$, where m is the linear mass density and A is the force, the dimensions of B will be

A. latent heat

B. pressure

C. work

D. momentum

Answer: a



Watch Video Solution

3. The 'rad' is the correct unit used to report the measurement of :

A. the rate of decay of radioactive source

B. the ability of a beam of gamma ray photons to produce ions in a target .

C. the energy delivered by radiation to a target

D. the biological effect of radiation .

Answer: d



Watch Video Solution

4. The dimension of magnetic field in M , L , T and C (coulomb) is given as

A. $[M^1 L^2 T^{-2} C^{-2}]$

B. $[L^{-2} T^2 C^{+2}]$

C. $[M^1 L^{-2} T^2 C^2]$

D. $[M^1 L^{-2} T^2 C^2]$

Answer: c



Watch Video Solution

5. Two full turns of the circular scale of a screw gauge cover a distance of 1 mm on its main scale. The total number of divisions on the circular scale is 50. Further it is found that the screw gauge has a zero error of - 0.02 mm. While measuring the diameter of a thin wire, a student notes the main scale reading of 4 mm and the number of circular scale divisions

in line with the main scale as 37. The diameter of the wire is

A. 4.37 mm

B. 4.39 mm

C. 4.74 mm

D. 4.76 mm

Answer: b



Watch Video Solution

6. The potential energy of a particle varies with distance x from a fixed origin as $U = \frac{A\sqrt{x}}{x^2 + B}$, where A and B are dimensional constants, then find the dimensional formula for AB .

A. $[M^1 L^{7/2} T^{-2}]$

B. $[M^1 L^{11/2} T^{-2}]$

C. $[M^1 L^{5/2} T^{-2}]$

D. $[M^1 L^{9/2} T^{-2}]$

Answer: b



Watch Video Solution

7. Assertion : The number 37800 has three significant digits .

Reason : All non - zero digits are significant .

A. Assertion is True , Reason is True , Reason is a correct explanation for Assertion.

B. Assertion is True , Reason is True, Reason is not a correct explanation for Assertion .

C. Assertion is True , Reason in False .

D. Assertion is False but , Reason is False .

Answer: b



Watch Video Solution

8. If $A = B + \frac{C}{D + E}$ the dimensions of B and C are $[M^0LT^{-1}]$ and $[M^0LT^0]$, respectively . Find the dimensions of A, D and E.

A. $A = [M^0 L^0 T^{-1}]$, $D = [T]$, $E = [LT]$

B. $A = [MLT^0]$, $D = [T^2]$, $E = [T^2]$

C.

$$A = [MLT^{-1}], D = [MT], E = [MT]$$

D. $A = [M^0 LT^{-1}]$, $D[T]$, $E = [T]$

Answer: d



Watch Video Solution

9. In the measurement of a physical quantity

$$X = \frac{A^2 B}{C^{1/3} D^3} .$$
 The percentage errors

introduced in the measurement of the quantities A,B,C and D are 1%, 3% , 4% and 5% respectively . Then the minimum amount of percentage of error in the measurement of X is contributed by

A. A

B. B

C. C

D. D

Answer: c



Watch Video Solution

10. If $E =$ energy , $G =$ gravitational constant , $I =$ impulse and $M =$ mass the dimension $\frac{GI^2M}{E^2}$

is same as that of

A. spring constant

B. wavelenght

C. energy gradient

D. Rydberg constant

Answer: b



Watch Video Solution

11. Choose the incorrect statement

A. A dimensionally correct equation may be correct .

B. A dimensionally correct equation may

C. A dimensionally incorrect equations
must be incorrect

D. A dimensionally incorrect equation be
correct

Answer: d



Watch Video Solution

12.5

A. 0.11

B. 0.04

C. 0.19

D. 0.09

Answer: c



View Text Solution

13. Number of particles is given by

$$n = -D \frac{n_2 - n_1}{x_2 - x_1} \text{ crossing a unit area}$$

perpendicular to X -axis in unit time, where n_1

and n_2 are number of particles per unit

volume for the value of x meant to x_2 and x_1 .

Find dimensions of D called as diffusion constant

A. $[M^0 L^1 T^2]$

B. $[M^0 L^2 T^4]$

C. $[M^0 L^1 T^{-3}]$

D. $[M^0 L^2 T^{-1}]$

Answer: d



Watch Video Solution

14. When a screw gauge is completely closed, zero of circular scale is 4 divisions above the reference line of graduation. If L.C of screw gauge is 10^{-3} cm, the zero error is

A. -4×10^{-3} cm

B. $+4 \times 10^{-3}$ cm

C. -0.004 mm

D. $+0.004$ mm

Answer: a



Watch Video Solution

15. Which of the following is not dimensionless?

A. Relative refractive index

B. Relative permittivity

C. Relative density

D. Relative velocity

Answer: d



Watch Video Solution

16. The jaws of a vernier callipers touch the inner wall of calorimeter without any undue pressure. The position of zero of vernier scale on the main scale reads 3.48. The 6th of vernier scale division is coinciding with any main scale division. Vernier constant of callipers is 0.01 cm. Find actual internal diameter of calorimeter, when it is observed that the vernier scale has a zero error of -0.03cm.

A. 3.37 cm

B. 3.57 cm

C. 3.42 cm

D. 3.54 cm

Answer: b



Watch Video Solution

17. The thin metallic strip of vernier callipers moves downward from top to bottom in such a way that it just touches the surface of beaker. Main scale reading of calliper is 6.4 cm whereas its vernier constant is 0.1 mm. The 4th

of vernier scale division is coinciding with main scale division. The actual depth of berker in mm is (when zero of vernier coincides with zero of main scale)

A. 6.64 cm

B. 6.42 cm

C. 6.44 cm

D. 6.13 cm

Answer: c



Watch Video Solution

