


CHEMISTRY

NCERT - NCERT CHEMISTRY(ENGLISH)

EQUILIBRIUM

Solved Example

1. The following concentration were obtained for the formation of NH_3 from N_2 and H_2 at equilibrium for the reaction



$$[N_2] = 1.5 \times 10^{-2} M$$


$$[H_2] = 3.0 \times 10^{-2} M$$

$$[NH_3] = 1.2 \times 10^{-2} M$$

Calculate equilibrium constant.

 Watch Video Solution

2. At equilibrium, the concentrations of $N_2 = 3.0 \times 10^{-3} M$, $O_2 = 4.2 \times 10^{-3} M$, and $NO = 2.8 \times 10^{-3} M$ in a sealed vessel at $800K$. What will be K_c for the reaction




 Watch Video Solution

3. PCl_5 , PCl_3 and Cl_2 are at equilibrium at $500K$ and having concentration $1.59M PCl_3$, $1.59M Cl_2$ and $1.41M PCl_5$. Calculate K_c for the reaction,



 Watch Video Solution


4. The value of $K_c = 4.24$ at $800K$ for the reaction.



Calculate equilibrium concentration of CO_2 , H_2 , CO and H_2O at $800K$. If only CO and H_2O are present initially at concentrations of $0.10M$ each.

 [Watch Video Solution](#)

5. For the equilibrium




the value of the equilibrium constant, K_c is 3.75×10^{-6} at $1069K$.

Calculate the K_p for the reaction at this temperature?

 [Watch Video Solution](#)

6. The value of K_p for the reaction



is 3.0 bar at 1000K. If initially $P_{CO_2} = 0.48$ bar, $P_{CO} = 0$ bar and pure graphite is present then determine equilibrium partial pressure of CO and CO_2 .

 [Watch Video Solution](#)

7. The value of K_c for the reaction $2A \rightleftharpoons B + C$ is 2.0×10^{-3} . At a given time, the composition of reaction mixture is $[A] = [B] = [C] = 3 \times 10^{-4}M$. In which direction the reaction will proceed?

 [Watch Video Solution](#)

8. 13.8g of N_2O_4 was placed in a 1L reaction vessel at 400K and allowed to attain equilibrium



The total pressure at equilibrium was found to be 9.15 bar.

Calculate K_c , K_p and partial pressure at equilibrium.

 [View Text Solution](#)

9. 3.00 mol of PCl_5 kept in 1L closed reaction vessel was allowed to attain equilibrium at 3.80K. Calculate composition of the mixture at equilibrium $K_c = 1.80$

 [Watch Video Solution](#)

10. The value of ΔG^\ominus for the phosphorylation of glucose in glycolysis is $13.8kJmol^{-1}$. Find the value of K_c at 298K

 [Watch Video Solution](#)

11. Hydrolysis of sucrose gives



Equilibrium constant K_c for the reaction is 2×10^{13} at $300K$.

Calculate ΔG^\ominus at $300K$.

 [Watch Video Solution](#)

12. Write the conjugate bases for the following Brddotoned acids

(a) HF (b) H_2SO_4 (c) HCO_3^\ominus

 [Watch Video Solution](#)

13. Wirte the conjugate acids for the following Brdddotted bases:

a. $\overset{\ominus}{N}H_2$ b. NH_3 c. $HCOO^{\ominus}$

 [Watch Video Solution](#)

14. The species: H_2O , HCO_3^{\ominus} , HSO_4^{\ominus} and NH_3 can act both as Bronsted acids and bases. For each case give the corresponding conjugate acid and base.

 [Watch Video Solution](#)

15. Classify the following species into Lewis acids and Lewis bases and show how these act as Lewis acid / base:

a. $\overset{\ominus}{O}H$, b. F^{\ominus} , c. H^{\oplus} , d. BCl_3

 [Watch Video Solution](#)

16. The concentration of hydrogen ion in a sample of soft drink is $3.8 \times 10^{-3} M$. What is its pH ?

 [Watch Video Solution](#)

17. The pH of $10^{-8} M$ solution of HCl in water is

 [Watch Video Solution](#)

18. The ionization constant of HF is 3.2×10^{-4} . Calculate the degree of ionization of HF in its $0.02 M$ solution. Calculate the concentration of all species present in the solution and its pH .

 [View Text Solution](#)

19. The pH of $0.1M$ monobasic acid is 4.50. Calculate the concentration of species, H^{\oplus} , A^{\ominus} , and HA at equilibrium. Also determine the value of K_a and pK_a of the monobasic acid.

 [View Text Solution](#)

20. Calculate the pH of 0.08 solution of $HOCl$ (hydrochlorous acid). The ionisation constant of the acid is 2.5×10^{-5} . Determine the percent dissociation of $HOCl$.

 [Watch Video Solution](#)

21. The pH of $0.004M$ hydrazine ($NH_2.NH_2$) solution is 9.7. Calculate its ionisation constant K_b and pK_b .

 [Watch Video Solution](#)

22. Calculate the pH of the solution in which $0.2M NH_4Cl$ and $0.1M NH_3$ are present. The pK_b of ammonia solution is 4.75.

 [View Text Solution](#)

23. Determine the degree of ionization and pH of $0.05M$ of ammonia solution. The ionization constant of ammonia can be taken from Table 7.7. Also calculate the ionization constant of the conjugate acid of ammonia.

 [Watch Video Solution](#)

24. 50.0 mL of 0.10 M ammonia solution is treated with 25.0 mL of $0.10M HCl$. If $K_b(NH_3) = 1.77 \times 10^{-5}$, the pH of the resulting solution will be

 [Watch Video Solution](#)

25. The pK_a of acetic acid and pK_b of ammonium hydroxide are 4.76 and 4.75 respectively. Calculate the pH of ammonium acetate solution.

 [Watch Video Solution](#)

26. Calculate the solubility of M_2X_3 in pure water, assuming that neither kind of ion reacts with H_2O . The solubility product of M_2X_3 , $K_{sp} = 1.1 \times 10^{-23}$.

 [Watch Video Solution](#)

27. The values of K_{sp} of two sparingly soluble salts, $Ni(OH)_2$ and $AgCN$ are 2.0×10^{-15} and 6×10^{-7} respectively, which salt is more soluble? Explain



Watch Video Solution

28. The solubility product of $Ni(OH)_2$ is 2.0×10^{-15} . The molar solubility of $Ni(OH)_2$ in $0.1MNaOH$ solution is



Watch Video Solution

Exercise


1. A liquid is in equilibrium with its vapour in a sealed container at a fixed temperature. The volume of the container is suddenly increased.

- what is the initial effect of the change on vapour pressure?
- How do rates of evaporation and condensation change initially?
- What happens when equilibrium is restored finally and what will be the final vapour pressure?

 [Watch Video Solution](#)


2. What is K_c for the following equilibrium concentration of each substance is:

$$[SO_2] = 0.60M, [O_2] = 0.82M \text{ and } [SO_3] = 1.90M?$$





 [Watch Video Solution](#)

3. At a certain temperature and a total pressure of $10^5 Pa$, iodine vapour contains 40% by volume of I atoms, Calculate K_p for the equilibrium.






 [Watch Video Solution](#)

4. Write the expression for the equilibrium constant K_c for each of the following reactions:





c.



 [Watch Video Solution](#)

5. Find out the value of K_c for each of the following equilibrium from the value of K_p :



 [Watch Video Solution](#)

6. For the following equilibrium, $K_c = 6.3 \times 10^{14}$ at 1000K




Both the forward and reverse reactions in the equilibrium are elementary bimolecular reactions. What is K_c , for the reverse reaction?

 [Watch Video Solution](#)

7. Explain why pure liquids and solids can be ignored while writing the equilibrium constant expression?

 [Watch Video Solution](#)

8. Reaction between nitrogen and oxygen takes place as following:




If a mixture of 0.482mole N_2 and 0.933mole of O_2 is placed in a

reaction vessel of volume 10litre and allowed to form N_2O at a temperature for which $K_c = 2.0 \times 10^{-37} \text{litremol}^{-1}$. Determine the composition of equilibrium mixture.

 [Watch Video Solution](#)


9. Nitric oxide reacts with bromine and gives nitrosyl-bromide as per reaction given below:



When 0.087mole of NO and 0.0437mole of Br_2 are mixed in a closed container at constant temperature, 0.0518mole of $NOBr$ is obtained at equilibrium. Calculate equilibrium amount of nitric oxide and bromine.

 [Watch Video Solution](#)


10. At $450K$, $K_p = 2.0 \times 10^{10}$ / bar for the given reaction at equilibrium.



What is K_c at this temperature?

 [Watch Video Solution](#)

11. A sample of $HI(g)$ is placed in flask at a pressure of $0.2atm$. At equilibrium. The partial pressure of $HI(g)$ is $0.04atm$. What is K_p for the given equilibrium?



 [Watch Video Solution](#)

12. A mixture of $1.57mol$ of N_2 , $1.92mol$ of H_2 and $8.13mol$ of NH_3 is introduced into a $20L$ reaction vessel at $500K$. At this

temperature, the equilibrium constant K_c for the reaction $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ is 1.7×10^2 . Is the reaction mixture at equilibrium? If not, what is the direction of the net reaction?

 [Watch Video Solution](#)


13. The equilibrium constant expression for a gas reaction is :

$$K_c = \frac{[NH_3]^4 [O_2]^5}{[NO]^4 [H_2O]^6}$$

Write the balanced chemical equation corresponding to this expression.

 [Watch Video Solution](#)


14. One mole of H_2O and one mole of CO are taken in a 10litre vessel and heated to 725K. At equilibrium, 40percent of water (by mass) reacts with carbon monoxide according to the equation,



Calculate the equilibrium constant for the reaction.

 [Watch Video Solution](#)

15. At $700K$ equilibrium constant for the reaction,



is 54.8 . If 0.5mollitre^{-1} of $HI_{(g)}$ is present at equilibrium at

$700K$, what are the concentrations of $H_{2(g)}$ and $I_{2(g)}$, assuming

that we initially started with $HI_{(g)}$ and allowed it to reach


equilibrium at $700K$.

 [Watch Video Solution](#)

16. What is the equilibrium concentration of each of the substance


in the equilibrium when the initial concentration of ICl was

0.78M?




 [Watch Video Solution](#)

17. $K_p = 0.04\text{atm}$ at 899K for the equilibrium shown below. What is the equilibrium concentration of C_2H_6 when it is placed in a flask at 4.0atm pressure and allowed to come to equilibrium?



 [Watch Video Solution](#)

18. The ester, ethyl acetate is formed by the reaction between ethanol and acetic acid and equilibrium is represented as:



(a) Write the concentration ratio (reaction quotient), Q_e , for this reaction. Note that water is not in excess and is not a solvent in

this reaction.

(b) At $293K$, if one starts with 1.00mole of acetic acid and 0.180 of ethanol, there is 0.171mole of ethyl acetate in the final equilibrium mixture. Calculate the equilibrium constant.

(c) Starting with 0.500mole of ethanol and 1.000mole of acetic acid and maintaining it at $293K$, 0.214mole of ethyl acetate is found after some time. Has equilibrium been reached?




[View Text Solution](#)

19. A sample of pure PCl_5 was introduced into an evacuated vessel at $473K$. After equilibrium was attained, concentration of PCl_5 was found to be $0.5 \times 10^{-1} \text{mollitre}^{-1}$. If value of K_c is $8.3 \times 10^{-3} \text{mollitre}^{-1}$. What are the concentrations of PCl_3 and Cl_2 at equilibrium ?



[Watch Video Solution](#)


20. One of the reaction that takes place in producing steel from iron ore is the reduction of iron(II) oxide by carbon monoxide to give iron metal and CO_2 .



What are the equilibrium partial pressure of CO and CO_2 at $1050K$ if the partial pressure are: $p_{CO} = 1.4\text{atm}$ and $p_{CO_2} = 0.80\text{atm}$?

 [Watch Video Solution](#)

21. Equilibrium constant, K_c for the reaction,



at $500K$ is $0.061\text{litre}^2\text{mole}^{-2}$. At a particular time, the analysis shows that composition of the reaction mixture is $3.00\text{mollitre}^{-1}N_2$, $2.00\text{mollitre}^{-1}H_2$, and $0.500\text{mollitre}^{-1}NH_3$.

Is the reaction at equilibrium? If not, in which direction does the reaction tend to proceed to reach equilibrium?

 [Watch Video Solution](#)


22. Bromine monochloride, ($BrCl$) decomposes into bromine and chlorine and reaches the equilibrium.



For which $K_c = 32$ at $500K$. If initially pure $BrCl$ is present at a concentration of $3.30 \times 10^{-3} \text{mollitre}^{-1}$, what is its molar concentration in the mixture at equilibrium?

 [Watch Video Solution](#)


23. At $1127K$ and $1atm$ pressure, a gaseous mixture of CO and CO_2 in equilibrium with solid carbon has 90.55% CO by mass:



Calculate K_c for the reaction at the above temperature.

 [Watch Video Solution](#)

24. Calculate (a) ΔG^\ominus and (b) the equilibrium constant for the formation of NO and O_2 at $298K$



where

$$\Delta_f G^\ominus(NO_2) = 52.0kJmol^{-1}$$



$$\Delta_f G^\ominus(NO) = 87.0kJmol^{-1}$$

$$\Delta_f G^\ominus(O_2) = 0kJmol^{-1}$$

 [Watch Video Solution](#)






25. Does the number of moles of reaction products increase, decrease, or remain same when each of the following equilibrium

is subjected to a decrease in pressure by increasing the volume?




Watch Video Solution

26. Which of the following reactions will get affected by increasing the pressure? Also, mention whether change will cause the reaction the reaction to go into forward of backward direction.



Watch Video Solution


27. The equilibrium constant for the following reaction is 1.6×10^5 at $1024K$



find the equilibrium pressure of all gases if 10.0 bar of HBr is introduced into a sealed container at $1024K$.

 [Watch Video Solution](#)

28. Dihydrogen gas is obtained from natural gas by partial oxidation with steam as per following endothermic reaction:



- Write an expression for K_p for the above reaction.
- How will the value of K_p and composition of equilibrium mixture be affected by
 - Increasing the pressure

ii. Increasing the temperature

iii. Using a catalyst?

 [Watch Video Solution](#)

29. Describe the effect of:


a. Addition of H_2

b. Addition of CH_3OH

c. Removal of CO

d. Removal of CH_3OH


on the equilibrium of the reaction:



 [Watch Video Solution](#)

30. At $473K$, equilibrium constant K_c for decomposition of phosphorus pentachloride, PCl_5 is 8.3×10^{-3} . If decomposition

is depicted as,




- a. Write an expression for K_c for the reaction.
- b. What is the value of K_c for the reverse reaction at the same temperature?
- c. What would be the effect on K_c if
 - i. More PCl_5 is added
 - ii. Pressure is increased
 - iii. The temperature is increased?



[Watch Video Solution](#)




31. Dihydrogen gas used in Haber's process is produced by reacting methane from natural gas with high temperature steam. The first stage of the two 2 stage reaction involves the formation of CO and H_2 . In second stage, CO formed in first stage is reacted with more steam in water gas shift reaction,



If a reaction vessel at $400^\circ C$ is charged with an equimolar mixture of CO and steam such that $p_{CO} = p_{H_2O} = 4.0$ bar, what will be the partial pressure of H_2 at equilibrium? $K_p = 0.1$ at $400^\circ C$.

 [Watch Video Solution](#)

32. Predict which of the following reactions will have appreciable concentration of reactants and products:



 [Watch Video Solution](#)

33. The value of K_c for the reaction $3O_2(g) \rightleftharpoons 2O_3(g)$ is 2.0×10^{-50} at $25^\circ C$. If the equilibrium concentration of O_2 in air

at 25°C is 1.6×10^{-2} , what is the concentration of O_3 ?

 [Watch Video Solution](#)

34. The reaction, $\text{CO}(g) + 3\text{H}_2(g) \rightleftharpoons \text{CH}_4(g) + \text{H}_2\text{O}(g)$ is at equilibrium at 1300K in a 1L flask. It also contains 0.30mol of CO , 0.10mol of H_2 and 0.02mol of H_2O and an unknown amount of CH_4 in the flask. Determine the concentration of CH_4 in the mixture. The equilibrium constant K_c for the reaction at the given temperature is 3.90 .

 [Watch Video Solution](#)

35. What is meant by the conjugate acid-base pair? Find the conjugate acid / base for the following species:

HNO_2 , CN^{\ominus} , HClO_4 , F^{\ominus} , OH^{\ominus} , CO_3^{2-} , and S^{2-}

 [Watch Video Solution](#)

36. Which of the followings are Lewis acids: H_2O , BF_3 , H^{\oplus} and NH_4 ?

 [Watch Video Solution](#)

37. Write the conjugate bases for the following Brddotonsted acids
(a) HF (b) H_2SO_4 (c) HCO_3^{\ominus}

 [Watch Video Solution](#)

38. Wirte the conjugate acids for the following Brdddotosted bases:

a. $\overset{\ominus}{N}H_2$ b. NH_3 c. $HCOO^{\ominus}$

 [Watch Video Solution](#)

39. The species: H_2O , HCO_3^\ominus , HSO_4^\ominus and NH_3 can act both as Bronsted acids and bases. For each case give the corresponding conjugate acid and base.

 [Watch Video Solution](#)

40. Classify the following species into Lewis acids and Lewis bases and show how these act as Lewis acid / base:

a. OH^\ominus , b. F^\ominus , c. H^\oplus , d. BCl_3

 [Watch Video Solution](#)

41. The concentration of hydrogen ion in a sample of soft drink is $3.8 \times 10^{-3} M$. What is its pH ?

 [Watch Video Solution](#)

42. The pH of a sample of vinegar is 3.76, Calculate the concentration of hydrogen ion in it.

 [Watch Video Solution](#)

43. The ionization constant of HF , $HCOOH$ and HCN at $298K$ are 6.8×10^{-4} , 1.8×10^{-4} and 4.8×10^{-9} respectively. Calculate the ionization constant of the corresponding conjugate base.

 [Watch Video Solution](#)

44. The ionization constant of phenol is 1.0×10^{-10} . What is the concentration of phenolate ion in $0.05M$ solution of phenol? What will be its degree of ionization if the solution is also $0.01M$ in sodium phenolate?

 [Watch Video Solution](#)

45. The first ionization constant of H_2S is 9.1×10^{-8} . Calculate the concentration of HS^{\ominus} ion in its $0.1M$ solution. How will this concentration be affected if the solution is $0.1M$ in HCl also? If the second dissociation constant if H_2S is 1.2×10^{-13} , calculate the concentration of S^{2-} under both conditions.

 [Watch Video Solution](#)

46. The ionization constant of acetic acid 1.74×10^{-5} . Calculate the degree of dissociation of acetic acid in its $0.05M$ solution. Calculate the concentration of acetate ion in the solution and its pH .

 [Watch Video Solution](#)

47. It has been found that the pH of a $0.01M$ solution of an organic acid is 4.15. Calculate the concentration of the anion, the ionization constant of the acid and its pK_a .

 [Watch Video Solution](#)

48. Assuming complete dissociation, calculate the pH of the following solutions,

a. $0.003M HCl$, b. $0.005M NaOH$,

c. $0.002M HBr$, d. $0.002M KOH$

 [Watch Video Solution](#)

49. Calculate the pH of the following solutions:

a. $2g$ of $TlOH$ dissolved in water to give 2 litre of solution.

b. $0.3g$ of $Ca(OH)_2$ dissolved in water to give $500mL$ of solution.

c. $0.3g$ of $NaOH$ dissolved in water to give $200mL$ of solution.

d. $1mL$ of $13.6MHCl$ is diluted with water to give 1 litre of solution.



[View Text Solution](#)

50. The degree of ionization of a $0.1M$ bromoacetic acid solution is 0.132 . Calculate the pH of the solution and the pK_a of bromoacetic acid.



[View Text Solution](#)

51. The pH of $0.005M$ codenine ($C_{18}H_{21}NO_3$) solution is 9.95 . Calculate its ionisation constant and pK_b .



[View Text Solution](#)

52. What is the pH of $0.001M$ aniline solution? The ionization constant of aniline 4.27×10^{-10} . Calculate the degree of ionization of aniline in the solution. Also calculate the ionization constant of the conjugate acid of aniline.



[View Text Solution](#)

53. Calculate the degree of ionisation of $0.05M$ acetic acid if its pK_a value is 4.74. How is the degree of dissociation affected when its solution also contains

a. $0.01M$, b. $0.1M$ in HCl ?



[View Text Solution](#)

54. The ionisation constant of dimethylamine is 5.4×10^{-4} . Calculate its degree of ionization in its $0.02M$ solution. What

percentage of dimethylamine is ionized if the solution is also $0.1M$ in $NaOH$?

 [View Text Solution](#)

55. Calculate the hydrogen ion concentration in the following biological fluids whose pH are given below:

a. Human muscle-fluid, 6.83

b. Human stomach fluid, 1.2

c. Human blood, 7.38

d. Human saliva, 6.4.

 [Watch Video Solution](#)

56. The pH of milk, black coffee, tomato juice, lemon juice and egg white are 6.8, 5.0, 4.2, 2.2 and 7.8 respectively. Calculate corresponding hydrogen ion concentration in each.



[Watch Video Solution](#)

57. If 0.561g of (KOH) is dissolved in water to give. 200mL of solution at 298K . Calculate the concentration of potassium, hydrogen and hydroxyl ions. What is its pH ?



[Watch Video Solution](#)

58. The solubility of $\text{Sr}(\text{OH})_2$ at 298K is 19.23gL^{-1} of solution. Calculate the concentrations of strontium and hydroxyl ions and the pH of the solution.



[View Text Solution](#)

59. The ionization constant of propionic acid is 1.32×10^{-5} . Calculate the degree of ionization of the acid in its 0.05M solution

and also its pH. What will be its degree of ionization in the solution of $0.01N HCl$?

 [Watch Video Solution](#)

60. The pH of $0.1M$ solution of cyanic acid ($HCNO$) is 2.34. Calculate the ionization constant of the acid and its degree of ionisation in the solution.

 [Watch Video Solution](#)

61. The ionization constant of nitrous acid is 4.5×10^{-4} . Calculate the pH of $0.04M$ sodium nitrite solution and also its degree of hydrolysis.

 [Watch Video Solution](#)

62. A $0.02M$ solution of pyridinium hydrochloride has $pH = 3.44$.

Calculate the ionization constant of pyridine.

 [Watch Video Solution](#)

63. Predict if the solutions of the following salts are neutral, acidic or basic: $NaCl$, KBr , $NaCN$, NH_4NO_3 , $NaNO_2$ and KF

 [Watch Video Solution](#)

64. The ionization constant of chloroacetic acid is 1.35×10^{-3} .

What will be the pH of $0.1M$ acid and its $0.1M$ sodium salt solution?

 [Watch Video Solution](#)

65. Ionic product of water at $310K$ is 2.7×10^{-14} . What is the pH of natural water at this temperature?

 [Watch Video Solution](#)

66. Calculate the pH of the resultant mixture:

a. $10mL$ of $0.2M Ca(OH)_2$ + $25mL$ of $0.1M HCl$

b. $10mL$ of $0.01M H_2SO_4$ + $10mL$ of $0.01M Ca(OH)_2$.

c. $10mL$ of $0.1M H_2SO_4$ + $10mL$ of $0.1M KOH$.

 [View Text Solution](#)

67. Determine the solubilities of silver chromate, barium chromate, ferric hydroxide, lead chloride and mercurous iodide at $298K$ from their solubility product constants given in Table 7.9. Determine also the molarities of individual ions.

 [View Text Solution](#)

68. The solubility product constant of Ag_2CrO_4 and $AgBr$ are 1.1×10^{-12} and 5.0×10^{-13} respectively. Calculate the ratio of the molarities of their saturated solutions.

 [Watch Video Solution](#)

69. Equal volumes of 0.002 M solution of sodium iodate and cupric chlorate are mixed together. Will it lead to precipitation of copper iodate?

(for cupric iodate $K = 7.4 \times 10^{-8}$).

 [Watch Video Solution](#)

70. The ionisation constant of benzoic acid ($PhCOOH$) is 6.46×10^{-5} and K_{sp} for silver benzoate is 2.5×10^{-3} . How many times is silver benzoate more soluble in a buffer of $pH 3.19$ compared to its solubility in pure water?

 [Watch Video Solution](#)

71. What is the maximum concentration of equimolar solutions of ferrous sulphate and sodium sulphide so that when mixed in equal volumes, there is no precipitation of iron sulphide? (For iron sulphide, $K_{sp} = 6.3 \times 10^{-18}$).

 [Watch Video Solution](#)

72. What is the minimum volume of water required to dissolve 1.0g of calcium sulphate at 298K?

(For calcium sulphate, K_{sp} is 9.1×10^{-6}).

 [Watch Video Solution](#)

73. The concentration of sulphide ion in $0.1M HCl$ solution saturated with hydrogen sulphide is $1.0 \times 10^{-19}M$. If $10mL$ of this is added to $5mL$ of $0.04M$ solution of the following: $FeSO_4$, $MnCl_2$, $ZnCl_2$ and $CdCl_2$. In which of these solutions precipitation will take place?

 [Watch Video Solution](#)