



MATHS

BOOKS - ARIHANT MATHS (HINGLISH)

COORDINATE SYSTEM AND COORDINATES

Example

1. Draw the polar coordinates

$$\left(2, \frac{\pi}{3}\right), \left(-2, \frac{\pi}{3}\right), \left(-2, -\frac{\pi}{3}\right) \text{ and } \left(2, -\frac{\pi}{3}\right)$$

on the plane.



Watch Video Solution

2. Draw the polar coordinate $\left(3, \frac{5\pi}{4}\right)$ on the plane.



[Watch Video Solution](#)

3. Find the cartesian coordinates of the points whose polar coordinates are

$$\left(5, \pi - \tan^{-1}\left(\frac{4}{3}\right)\right)$$



[Watch Video Solution](#)

4. Find the cartesian coordinates of the points whose polar coordinates are

$$\left(5\sqrt{2}, \frac{\pi}{4}\right)$$



[Watch Video Solution](#)

5. Find the polar coordinates of the points whose cartesian coordinates are

$$(-2, -2)$$



[Watch Video Solution](#)

6. Find the polar coordinates of the points whose cartesian coordinates are

$$(-3, 4)$$



[Watch Video Solution](#)

 Watch Video Solution

7. Transform to Cartesian coordinates the equations:

$$r^2 = a^2 \cos 2\theta$$



Watch Video Solution

8. Transform the equation $x^2 + y^2 = ax$ into polar form.



Watch Video Solution

9. Prove that the distance of the point $(a \cos \alpha, a \sin \alpha)$ from the origin is independent of α

 [Watch Video Solution](#)

10. The distance between the points $(a \cos \alpha, a \sin \alpha)$ and $(a \cos \beta, a \sin \beta)$ where $a > 0$

 [Watch Video Solution](#)

11. If $P(x, y)$ is a point equidistant from the points $A(6, -1)$ and $B(2, 3)$, show that $x - y = 3$.

 [Watch Video Solution](#)

12. Using distance formula, show that the points $(1, 5)$, $(2, 4)$ and $(3, 3)$ are collinear.



Watch Video Solution

13. An equilateral triangle has one vertex at $(0, 0)$ and another at $(3, \sqrt{3})$. What are the coordinates of the third vertex ?



Watch Video Solution

14. Show that four points $(1, -2)$, $(3, 6)$, $(5, 10)$ and $(3, 2)$ are the vertices of a parallelogram.



Watch Video Solution

15. Let the opposite angular points of a square be $(3, 4)$ and $(1, -1)$. Find the coordinates of the remaining angular points.



Watch Video Solution

16. Find the circumcentre of the triangle whose vertices are $(-2, -3)$, $(-1, 0)$ and $(7, -6)$. Also find the

radius of the circumcircle.



[Watch Video Solution](#)

17. If the line segment joining the point $A(a, b)$ and $B(c, d)$ subtends an angle θ at the

origin. Prove that $\cos \theta = \frac{ac + bd}{\sqrt{(a^2 + b^2) \cdot (c^2 + d^2)}}$



[Watch Video Solution](#)

18. Show that the triangle, the coordinates of whose vertices are given by integers, can never be an equilateral triangle.



[Watch Video Solution](#)

 [Watch Video Solution](#)

19. In any triangle ABC , prove that $AB^2 + AC^2 = 2(AD^2 + BD^2)$, where D is the midpoint of BC .

 [Watch Video Solution](#)

20. Let $ABCD$ be a rectangle and P be any point in its plane. Show that $AP^2 + PC^2 = PB^2 + PD^2$.

 [Watch Video Solution](#)

21. Prove that the points $(0, 0)$, $\left(3, \frac{\pi}{2}\right)$ and $\left(3, \frac{\pi}{6}\right)$ are the vertices of an equilateral triangle.



Watch Video Solution

22. Find the coordinates of the point which divides the line segment joining the points $A(5,-2)$ and $B(9,6)$ in the ratio 3:1

A. $(8, 4)$

B. $(8, -4)$

C. $(-8, 4)$

D. $(8, 4)$

Answer: A



Watch Video Solution

23. Find the lengths of the medians of a triangle whose vertices are $A(-1, 3)$, $B(1, -1)$ and $C(5, 1)$.



Watch Video Solution

24. In what ratio does the line $y - x + 2 = 0$ cut the line joining $(3, -1)$ and $(8, 9)$?



Watch Video Solution

25. The coordinates of three consecutive vertices of a parallelogram are $(1, 3)$, $(-1, 2)$ and $(2, 5)$. Then find the coordinates of the fourth vertex.



Watch Video Solution

26. In what ratio does the x -axis divide the line segment joining the points $(2, -3)$ and $(5, 6)$?



Watch Video Solution

27. The mid-points of the sides of a triangle are $(1, 2)$, $(0, -1)$ and $(2, -1)$. Find the coordinates of the vertices of a triangle with the help of two unknowns.



Watch Video Solution

28. Prove that the mid point of the hypotenuse of a right triangle is equidistant from its vertices.



Watch Video Solution

29. The line segment joining the mid-points of any two sides of a triangle is parallel to the third side and

equal to half of it.



[Watch Video Solution](#)

30. Find the coordinates of a point which divides externally the line joining $(1, -3)$ and $(-3, 9)$ in the ratio $1 : 3$.



[Watch Video Solution](#)

31. The line segment joining $A(6, 3)$ to $B(-1, -4)$ is doubled in length by having its length added to each end, then the ordinates of new ends are



[Watch Video Solution](#)

 [Watch Video Solution](#)

32. Using section formula show that the points $(1,-1)$, $(2, 1)$ and $(4, 5)$ are collinear.



[Watch Video Solution](#)

33. Find the ratio in which the point $(2, y)$ divides the line segment $(4,3)$ and $(6,3)$. hence find the value of y



[Watch Video Solution](#)

34. Find the harmonic conjugates of the point $R(5, 1)$ with respect to the points $P(2, 10)$ and $Q(6, -2)$



Watch Video Solution

35. Two vertices of a triangle are $(-1, 4)$ and $(5, 2)$. If its centroid is $(0, -3)$, find the third vertex.



Watch Video Solution

36. The vertices of a triangle are $(1, 2)$, $(h, -3)$ and $(-4, k)$. Find the value of $\sqrt{\{(h + k)^2 + (h + 3k)^2\}}$. If the centroid of the triangle be at point $(5, -1)$.



Watch Video Solution

37. If $D(-2, 3)$, $E(4, -3)$ and $F(4, 5)$ are the mid-points of the sides BC , CA and AB of the sides BC , CA and AB of triangle ABC , then find $\sqrt{(|AG|^2 + |BG|^2 - |CG|^2)}$ where, G is the centroid of $\triangle ABC$.



Watch Video Solution

38. If G be the centroid of a triangle ABC and P be any other point in the plane prove that $PA^2 + PB^2 + PC^2 = GA^2 + GB^2 + GC^2 + 3GP^2$



Watch Video Solution

39. If G be the centroid of the ΔABC , then prove that $AB^2 + BC^2 + CA^2 = 3(GA^2 + GB^2 + GC^2)$



[Watch Video Solution](#)

40. The vertices of a triangle are $(1, a)$, $(2, b)$ and $(c^2, -3)$. (1) Prove that its centroid can not lie on the y-axis. (2) Find the condition that the centroid may lie on the x-axis for any value of $a, b, c \in R$



[Watch Video Solution](#)

41. The vertices of a triangle are $(1, a)$, $(2, b)$ and $(c^2, -3)$. (i) Prove that its centroid can not lie on the y-axis. (ii) Find the condition that the centroid may lie on the x-axis for any value of $a, b, c \in \mathbb{R}$.

 [Watch Video Solution](#)

42. Find the coordinates of incentre of the triangle whose vertices are $(4, -2)$, $(-2, 4)$ and $(5, 5)$.

 [Watch Video Solution](#)

43. If $\left(\frac{3}{2}, 0\right)$, $\left(\frac{3}{2}, 6\right)$ and $(-1, 6)$ are mid-points of the sides of a triangle, then find

Centroid of the triangle



[Watch Video Solution](#)

44. If $\left(\frac{3}{2}, 0\right)$, $\left(\frac{3}{2}, 6\right)$ and $(-1, 6)$ are mid-points of the sides of a triangle, then find

Centroid of the triangle



[Watch Video Solution](#)

45. If a vertex of a triangle be $(1, 1)$ and the middle points of the sides through it be $(-2, 3)$ and $(5, 2)$, find the other vertices.



[Watch Video Solution](#)

46. If G is the centroid and I the in-centre of the triangle, with vertices $A(-36, 7)$, $B(20, 7)$ and $C(0, -8)$, then, find the value of GI



[Watch Video Solution](#)

47. In a ΔABC with vertices $A(1,2)$, $B(2,3)$ and $C(3, 1)$

and

$$\angle A = \angle B = \cos^{-1} \left(\frac{1}{\sqrt{10}} \right), \angle C = \cos^{-1} \left(\frac{4}{5} \right),$$

then find the circumcentre of ΔABC .



[Watch Video Solution](#)

48. Find the circumcentre of the triangle whose vertices are $(2, 2)$, $(4, 2)$ and $(0, 4)$.



[Watch Video Solution](#)

49. Find the circumcentre of triangle ABC if $A(7, 4)$, $B(3, -2)$ and $\angle C = \frac{\pi}{3}$.

 [Watch Video Solution](#)

50. Find the orthocentre of ΔABC if $A \equiv (0, 0)$, $B \equiv (3, 5)$ and $C \equiv (4, 7)$.

 [Watch Video Solution](#)

51. If a triangle has its orthocenter at $(1, 1)$ and circumcentre $(\frac{3}{2}, \frac{3}{4})$ then centroid is:

 [Watch Video Solution](#)

52. The vertices of a triangle are $A(x_1, x_1 \tan \theta_1)$, $B(x_2, x_2 \tan \theta_2)$ and $C(x_3, x_3 \tan \theta_3)$.

if the circumcentre of ΔABC coincides with the origin and $H(x, y)$ is the orthocentre, show that

$$\frac{y}{x} = \frac{\sin \theta_1 + \sin \theta_2 + \sin \theta_3}{\cos \theta_1 + \cos \theta_2 + \cos \theta_3}$$



[Watch Video Solution](#)

53. The coordinates of A , B , C are $(6, 3)$, $(-3, 5)$ and $(4, -2)$ respectively and P is any point (x, y) .

Show that the ratio of the areas of triangles PBC

and ABC is $\left| \frac{x + y - 2}{7} \right|$.



[Watch Video Solution](#)

54. Find the area of the pentagon whose vertices are $A(1, 1)$, $B(7, 21)$, $C(7, -3)$, $D(12, 2)$, and $E(0, -3)$



[Watch Video Solution](#)

55. Show that the points $(a, 0)$, $(0, b)$ and $(1, 1)$ are collinear, if $\frac{1}{a} + \frac{1}{b} = 1$



[Watch Video Solution](#)

56. Prove that the coordinates of the vertices of an equilateral triangle can not all be rational.



[Watch Video Solution](#)

57. The coordinates of two points A and B are (3, 4) and (5, -2) respectively. Find the coordinates of any point P if $PA = PB$ and area of $\triangle APB$ is 10.



[Watch Video Solution](#)

58. Find the area of the triangle formed by the straight lines $7x - 2y + 10 = 0$, $7x + 2y - 10 = 0$

and $9x + y + 2 = 0$ (without sloving the vertices of the triangle).

 [Watch Video Solution](#)

59. Find the locus of a point which moves such that its distance from the origin is three times its distance from x-axis.

 [Watch Video Solution](#)

60. The locus of the moving point P such that $2PA = 3PB$, where A is (0,0) and B is (4,-3), is

 [Watch Video Solution](#)

61. The sum of the squares of the distances of a moving point from two fixed points $(a,0)$ and $(-a, 0)$ is equal to a constant quantity $2c^2$. Find the equation to its locus.

[Watch Video Solution](#)

62. A point moves so that the sum of its distances from $(ae, 0)$ and $(-ae, 0)$ is $2a$, prove that the equation to its locus is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where $b^2 = a^2(1 - e^2)$.

[Watch Video Solution](#)

63. Find the equation of the locus of a point which moves so that the difference of its distances from the points $(3, 0)$ and $(-3, 0)$ is 4 units.

 [Watch Video Solution](#)

64. The ends of the hypotenuse of a right angled triangle are $(6, 0)$ and $(0, 6)$. Find the locus of the third vertex.

 [Watch Video Solution](#)

65. Find the equation to the locus of a point which moves so that the sum of its distances from $(3,0)$ and $(-3,0)$ is less than 9.

 [Watch Video Solution](#)

66. Find the locus of a point whose coordinate are given by $x = t + t^2$, $y = 2t + 1$, where t is variable.

 [Watch Video Solution](#)

67. A stick of length 10 units rests against the floor and a wall of a room. If the stick begins to slide on floor then the locus of its middle point is:



Watch Video Solution

68. Locus of the point of intersection of the lines
 $x \cos \alpha + y \sin \alpha = a$ and $x \sin \alpha - y \cos \alpha = b$
where α is variable.



Watch Video Solution

69. A variable line cuts X-axis at A, Y -axis at B, where
 $OA = a, OB = b$ (O as origin) such that $a^2 + b^2 = 1$.
Find the locus of
centroid of $\triangle OAB$



Watch Video Solution

70. A variable line cuts x-axis at A, y-axis at B where $OA = a, OB = b$ (O as origin) such that then the locus of circumcentre of ΔOAB is-

 [Watch Video Solution](#)

71. Two points $P(a,0)$ and $Q(-a,0)$ are given, R is a variable on one side of the line PQ such that $\angle RPQ - \angle RQP$ is a positive constant 2α . Find the locus of point R.

 [Watch Video Solution](#)

72. The equation of curve referred to the new axes, axes retaining their directions, and origin $(4, 5)$ is $X^2 + Y^2 = 36$. Find the equation referred to the original axes.



[Watch Video Solution](#)

73. Shift the origin to a suitable point so that the equation $y^2 + 4y + 8x - 2 = 0$ will not contain a term in y and the constant term.



[Watch Video Solution](#)

74. At what point the origin be shifted, if the coordinates of a point $(-1, 8)$ become $(-7, 3)$?



[Watch Video Solution](#)

75. If the axes are turned through 45° , find the transformed form of the equation

$$3x^2 + 3y^2 + 2xy = 2.$$



[View Text Solution](#)

76. Prove that if the axes be turned through $\frac{\pi}{4}$ the equation $x^2 - y^2 = a^2$ is transformed to the form

$xy = \lambda$. Find the value of λ .



[View Text Solution](#)

77. Though what angle should the axes be rotated so that the equation $9x^2 - 2\sqrt{3}xy + 7y^2 = 10$ may be changed to $3x^2 + 5y^2 = 5$?



[Watch Video Solution](#)

78. If (x, y) and (X, Y) be the coordinates of the same point referred to two sets of rectangular axes with the same origin and if $ux + vy$, when u and v are

independent of X and Y become $VX + UY$, show that

$$u^2 + v^2 = U^2 + V^2.$$



[Watch Video Solution](#)

79. What does the equation

$$2x^2 + 4xy - 5y^2 + 20x - 22y - 14 = 0$$
 become

when referred to the rectangular axes through the

point $(-2, -3)$, the new axes being inclined at an

angle at 45° with the old axes?



[Watch Video Solution](#)

80. Given the equation $4x^2 + 2\sqrt{3}xy + 2y^2 = 1$, through what angle should the axes be rotated so that the term in xy be wanting from the transformed equation.



Watch Video Solution

81. Find λ if $(\lambda, \lambda + 1)$ is an interior point of $\triangle ABC$ where, $A \equiv (0, 3)$, $B \equiv (-2, 0)$ and $C \equiv (6, 1)$.



Watch Video Solution

82. Locus of centroid of the triangle whose vertices are $(a \cos t, a \sin t)$, $(b \sin t, -b \cos t)$ and $(1, 0)$, where t is a

A. $(3x - 1)^2 + (3y)^2 = a^2 - b^2$

B. $(3x - 1)^2 + (3y)^2 = a^2 + b^2$

C. $(3x + 1)^2 + (3y)^2 = a^2 + b^2$

D. $(3x + 1)^2 + 3y^2 = a^2 - b^2$

Answer: B



Watch Video Solution

83. The incentre of triangle with vertices

$(1, \sqrt{3})$, $(0, 0)$ and $(2, 0)$ is

A. $\left(1, \frac{\sqrt{3}}{2}\right)$

B. $\left(\frac{2}{3}, \frac{1}{\sqrt{3}}\right)$

C. $\left(\frac{2}{3}, \frac{\sqrt{3}}{2}\right)$

D. $\left(1, \frac{1}{\sqrt{3}}\right)$

Answer: D



Watch Video Solution

84. Orthocentre of triangle with vertices $(0, 0)$, $(3, 4)$ and $(4, 0)$ is

A. $\left(3, \frac{5}{4}\right)$

B. $(3, 12)$

C. $\left(3, \frac{3}{4}\right)$

D. $(3, 9)$

Answer: C



Watch Video Solution

85. If x_1, x_2, x_3 as well as y_1, y_2, y_3 are in GP, with the same common ratio, then the points $(x_1, y_1), (x_2, y_2)$ and (x_3, y_3)

- A. lie on a straight line
- B. lie on an ellipse
- C. lie on a circle
- D. are vertices of a triangle

Answer: A



Watch Video Solution

86. Let A be the image of (2, -1) with respect to Y - axis

Without transforming the origin, coordinate axis are

turned at an angle 45° in the clockwise direction.

Then, the coordinates of A in the new system are

A. $\left(-\frac{1}{\sqrt{2}}, -\frac{3}{\sqrt{2}} \right)$

B. $\left(-\frac{3}{\sqrt{2}}, -\frac{1}{\sqrt{2}} \right)$

C. $\left(\frac{1}{\sqrt{2}}, \frac{3}{\sqrt{2}} \right)$

D. $\left(\frac{3}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right)$

Answer: A



Watch Video Solution

87. Let S_1, S_2, \dots , be squares such that for each $n \geq 1$, the length of a side of S_n equals the length of a diagonal of S_{n+1} . If the length of a side S_1 is 10 cm, then for which of the following value of n is the area of S_n less than 1 sq cm ?

A. 7

B. 8

C. 9

D. 10

Answer: B::C::D





[Watch Video Solution](#)

88. If each of the vertices of a triangle has integral coordinates, then the triangles may be

- A. right angled
- B. equilateral
- C. isosceles
- D. scalene

Answer: A::C::D



[Watch Video Solution](#)

89. ABC is an isosceles triangle. If the coordinates of the base are B(1, 3) and C(-2, 7). The coordinates of vertex A can be



Watch Video Solution

90. If $A\left(\alpha, \frac{1}{\alpha}\right), B\left(\beta, \frac{1}{\beta}\right), C\left(\gamma, \frac{1}{\gamma}\right)$ be the vertices of a ΔABC , where α, β are the roots of $x^2 - 6ax + 2 = 0$, β, γ are the roots of $x^2 - 6bx + 3 = 0$ and γ, α are the roots of $x^2 - 6cx + 6 = 0$, a, b, c being positive.

The value of $a + b + c$ is

A. 1

B. 2

C. 3

D. 5

Answer: B



Watch Video Solution

91. If $A\left(\alpha, \frac{1}{\alpha}\right), B\left(\beta, \frac{1}{\beta}\right), C\left(\gamma, \frac{1}{\gamma}\right)$ be the vertices of a ΔABC , where α, β are the roots of $x^2 - 6ax + 2 = 0$, β, γ are the roots of $x^2 - 6bx + 3 = 0$ and γ, α are the roots of

$$x^2 - 6cx + 6 = 0, a, b, c \text{ being positive.}$$

The coordinates of centroid of ΔABC is

A. $\left(1, \frac{11}{9}\right)$

B. $\left(\frac{1}{3}, \frac{11}{18}\right)$

C. $\left(2, \frac{11}{18}\right)$

D. $\left(\frac{2}{3}, \frac{11}{19}\right)$

Answer: C



Watch Video Solution

92. If $A\left(\alpha, \frac{1}{\alpha}\right), B\left(\beta, \frac{1}{\beta}\right), C\left(\gamma, \frac{1}{\gamma}\right)$ be the vertices of a ΔABC , where α, β are the roots of

$x^2 - 6ax + 2 = 0$, β, γ are the roots of
 $x^2 - 6bx + 3 = 0$ and γ, α are the roots of
 $x^2 - 6cx + 6 = 0$, a, b, c being positive.

The coordinates of orthocentre of ΔABC is

A. $\left(-\frac{1}{2}, -2\right)$

B. $\left(-\frac{1}{3}, -3\right)$

C. $\left(-\frac{1}{5}, -5\right)$

D. $\left(-\frac{1}{6}, -6\right)$

Answer: D



Watch Video Solution

93. If the points $(-2, 0)$, $\left(-1, \frac{1}{\sqrt{3}}\right)$ and $(\cos \theta, \sin \theta)$ are collinear, then the number of value of $\theta \in [0, 2\pi]$ is



Watch Video Solution

94. Statement I : The area of the triangle formed by the points $A(100, 102)$, $B(102, 105)$, $C(104, 107)$ is same as the area formed by $A'(0, 0)$, $B'(2, 3)$, $C'(4, 5)$.

Statement II : The area of the triangle is constant with respect to translation.

- A. Statement I is true, Statement II is true,
Statement II is a correct explanation for
Statement I.
- B. Statement I is true, Statement II is true,
Statement II is not a correct explanation for
Statement I.
- C. Statement I is true, Statement II is false.
- D. Statement I is false, Statement II is true.

Answer: A



Watch Video Solution

95. Statement I : If centroid and circumcentre of a triangle are known its orthocentre can be found

Statement II : Centroid, orthocentre and circumcentre of a triangle are collinear.

A. Statement I is true, Statement II is true,

Statement II is a correct explanation for

Statement I.

B. Statement I is true, Statement II is true,

Statement II is not a correct explanation for

Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

Answer: B



[View Text Solution](#)

96. The four points $A(\alpha, 0)$, $B(\beta, 0)$, $C(\gamma, 0)$ and $D(\delta, 0)$ are such that α, β are the roots of equation $ax^2 + 2h'x + b' = 0$. Show that the sum of the ratios in which C and D divide AB is zero, if $ab' + a'b = 2hh'$.



[View Text Solution](#)

97. If m_1 and m_2 are roots of equation $x^2 + (\sqrt{3} + 2)x + \sqrt{3} - 1 = 0$ the the area of the triangle formed by lines $y = m_1x$, $y = m_2x$, $y = c$ is:



Watch Video Solution

98. x coordinates of two points B and C are the roots of equation $x^2 + 4x + 3 = 0$ and their y coordinates are the roots of equation $x^2 - x - 6 = 0$. If x coordinate of B is less than the x coordinate of C and y coordinate of B is greater than the y coordinate of

C and coordinates of a third point A be $(3, -5)$, find the length of the bisector of the interior angle at A.



[Watch Video Solution](#)

99. The distance between the two parallel lines is 1 unit. A point A is chosen to lie between the lines at a distance 'd' from one of them. Triangle ABC is equilateral with B on one line and C on the other parallel line. The length of the side of the equilateral triangle is



[Watch Video Solution](#)

100.

In

a

ΔABC , $A \equiv (\alpha, \beta)$, $B \equiv (1, 2)$, $C \equiv (2, 3)$ and

point A lies on the line $y = 2x + 3$ where $\alpha, \beta \in I$. If

the area of ΔABC be such that $[\Delta] = 2$, where $[.]$

denotes the greatest integer function, find all

possible coordinates of A.



Watch Video Solution

101. Let S be the square of unit area. Consider any

quadrilateral which has one vertex on each side of S.

If a, b, c and d denote the lengths of the sides of the

quadrilateral, prove that $2 \leq a^2 + b^2 + c^2 + d^2 \leq 4$.

 [Watch Video Solution](#)

102. The circumcentre of a triangle having vertices $A(a, a \tan \alpha)$, $B(b, b \tan \beta)$, $C(c, c \tan \gamma)$ is at origin, where $\alpha + \beta + \gamma = \pi$. Then the orthocentre lies on

 [Watch Video Solution](#)

Exercise For Session 1

1. The polar coordinates of the point whose cartesian coordinates are $(-1, -1)$ is

A. $\left(\sqrt{2}, \frac{\pi}{4}\right)$

B. $\left(\sqrt{2}, \frac{3\pi}{4}\right)$

C. $\left(\sqrt{2}, -\frac{\pi}{4}\right)$

D. $\left(\sqrt{2}, -\frac{3\pi}{4}\right)$

Answer: D



Watch Video Solution

2. The cartesian coordinates of the point whose polar coordinates are $\left(13, \pi - \tan^{-1}\left(\frac{5}{12}\right)\right)$ is

A. (12, 5)

B. (-12, 5)

C. (-12, -5)

D. (12, -5)

Answer: B



Watch Video Solution

3. The transform equation of $r^2 \cos^2 \theta = a^2 \cos 2\theta$ to cartesian form is $(x^2 + y^2)x^2 = a^2 \lambda$, then value of λ is

A. $y^2 - x^2$

B. $x^2 - y^2$


C. xy

D. x^2y^2

Answer: B

 Watch Video Solution

4. The coordinates of P' in the figure is



A. $\left(3, \frac{\pi}{3}\right)$

B. $\left(3, -\frac{\pi}{3}\right)$

C. $\left(-3, -\frac{\pi}{3}\right)$


D. $\left(-3, \frac{\pi}{3}\right)$

Answer: B



Watch Video Solution

5. The cartesian coordinates of the point Q in the figure is



- A. $(\sqrt{3}, 1)$
- B. $(-\sqrt{3}, 1)$
- C. $(-\sqrt{3}, -1)$
- D. $(\sqrt{3}, -1)$

Answer: B



Watch Video Solution

6. A point lies on X-axis at a distance 5 units from Y-axis. What are its coordinates ?



[Watch Video Solution](#)

7. A point lies on Y-axis at a distance 4 units from X-axis. What are its coordinates ?



[Watch Video Solution](#)

8. A point lies on negative direction of X-axis at a distance 6 units from Y-axis. What are its coordinates ?



[Watch Video Solution](#)

9. Transform the equation $y = x \tan \alpha$ to polar form.



[Watch Video Solution](#)

10. Transform the equation $r = 2a \cos \theta$ to cartesian form.



[Watch Video Solution](#)

Exercise For Session 2

1. If the distance between the points $(a, 2)$ and $(3, 4)$ be 8, then a equals to

A. $2 + 3\sqrt{3}$

B. $2 - 3\sqrt{15}$

C. $2 \pm 3\sqrt{15}$

D. $3 \pm 2\sqrt{15}$

Answer: D



Watch Video Solution

2. The three points $(-2, 2)$, $(8, -2)$ and $(-4, -3)$ are the vertices of

- A. an isosceles triangle
- B. an equilateral triangle
- C. a right angled triangle
- D. None of these

Answer: C

 [Watch Video Solution](#)

3. The distance between the points $\left(3, \frac{\pi}{4}\right)$ and $\left(7, \frac{5\pi}{4}\right)$

A. 8

B. 10

C. 12

D. 14

Answer: B



Watch Video Solution

4. Let $A(6, -1)$, $B(1, 3)$ and $C(x, 8)$ be three points such that $AB = BC$ then the value of x are

A. 3, 5

B. $-3, 5$

C. $3, -5$

D. $-3, -5$

Answer: B



Watch Video Solution

5. The points $(a + 1, 1)$, $(2a + 1, 3)$ and $(2a + 2, 2a)$ are collinear if

A. $a = -1, 2$

B. $a = \frac{1}{2}, 2$

C. $a = 2, 1$

$$D. a = -\frac{1}{2}, 2$$

Answer: D

 [Watch Video Solution](#)

6. Let $A = (3, 4)$ and B is a variable point on the lines $|x| = 6$. If $AB \leq 4$, then find the number of position of B with integral coordinates.

A. 5

B. 6

C. 10

D. 12

Answer: A



Watch Video Solution

7. The number of points on X-axis which are at a distance c units ($c < 3$) from $(2, 3)$ is

A. 1

B. 2

C. 0

D. 3

Answer: C



Watch Video Solution

8. The point on the axis of y which its equidistant from $(-1, 2)$ and $(3, 4)$, is

A. $(0, 3)$

B. $(0, 4)$

C. $(0, 5)$

D. $(0, -6)$

Answer: C



9. Find the distance between the points $(at_1^2, 2at_1)$ and $(at_2^2, 2at_2)$, where t_1 and t_2 are the roots of the equation $x^2 - 2\sqrt{3}x + 2 = 0$ and $a > 0$.



Watch Video Solution

10. If P and Q are two points whose coordinates are $(at^2, 2at)$ and $\left(\frac{a}{t^2}, \frac{2a}{t}\right)$ respectively and S is the point $(a, 0)$. Show that $\frac{1}{SP} + \frac{1}{sQ}$ is independent of t.



Watch Video Solution

11. Show that the points $(3, 4)$, $(8, -6)$ and $(13, 9)$ are the vertices of a right angled triangle.

 [Watch Video Solution](#)

12. Prove that the points $(0, -1)$, $(6, 7)$, $(-2, 3)$ and $(8, 3)$ are the vertices of a rectangle.

 [Watch Video Solution](#)

13. Find the circumcentre and circumradius of the triangle whose vertices are $(-2, 3)$, $(2, -1)$ and $(4, 0)$.



[Watch Video Solution](#)

14. The vertices of a triangle are $A(1, 1)$, $B(4, 5)$ and $C(6, 13)$. Find $\cos A$.



[Watch Video Solution](#)

15. The opposite vertices of a square are $(2, 6)$ and $(0, -2)$. Find the coordinates of the other vertices.



[Watch Video Solution](#)

16. If the point (x, y) is equidistant from the points (a, b) and $(a - b, a + b)$, prove that $bx = ay$.



Watch Video Solution

17. If a and b are between 0 and 1 such that the points $(a, 1)$, $(1, b)$ and $(0, 0)$ form an equilateral triangle then the values of 'a' and 'b' respectively



Watch Video Solution

18. An equilateral triangle has one vertex at (3, 4) and another at (-2, 3). Find the coordinates of the third vertex.



[Watch Video Solution](#)

19. If P be any point in the plane of square ABCD, prove that

$$PA^2 + PC^2 = PB^2 + PD^2$$



[Watch Video Solution](#)

1. The coordinates of the middle points of the sides of a triangle are $(4, 2)$, $(3, 3)$ and $(2, 2)$, then coordinates of centroid are

A. $(3, 7/3)$

B. $(3, 3)$

C. $(4, 3)$

D. $(3, 4)$

Answer: A



Watch Video Solution

2. The incentre of the triangle whose vertices are $(-36, 7)$, $(20, 7)$ and $(0, -8)$ is

A. $(0, -1)$

B. $(-1, 0)$

C. $(1, 1)$

D. $\left(\frac{1}{2}, 1\right)$

Answer: B



Watch Video Solution

3. If the orthocentre and centroid of a triangle are $(-3, 5)$ and $(3, 3)$ then its circumcentre is

A. $(6, 2)$

B. $(3, -1)$

C. $(-3, 5)$

D. $(-3, 1)$

Answer: A



Watch Video Solution

4. An equilateral triangle has each side to a. If the coordinates of its vertices are (x_1, y_1) , (x_2, y_2) and (x_3, y_3) then the square of the determinat

$$\begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} \text{ equals}$$

A. $3a^4$

B. $\frac{3a^4}{2}$

C. $\frac{3}{4}a^4$

D. $\frac{3}{8}a^4$

Answer: C



Watch Video Solution

5. The vertices of a triangle are $A(0, 0)$, $B(0, 2)$ and $C(2, 0)$. The distance between circumcentre and orthocentre is

A. $\sqrt{2}$

B. $\frac{1}{\sqrt{2}}$

C. 2

D. $\frac{1}{2}$

Answer: A



Watch Video Solution

6. Area of the triangle with vertices (a, b) , (x_1, y_1) and (x_2, y_2) where a, x_1, x_2 which in G.P. common ratio r , b, y_1, y_2 are in G.P with common ratio s , is

A. $ab(r - 1)(s - 1)(s - r)$

B. $\frac{1}{2}ab(r + 1)(s + 1)(s - r)$

C. $\frac{1}{2}ab(r - 1)(s - 1)(s - r)$

D. $ab(r + 1)(s + 1)(r - s)$

Answer: C



Watch Video Solution

7. The points $(x + 1, 2)$, $(1, x + 2)$, $\left(\frac{1}{x + 1}, \frac{2}{x + 1}\right)$ are collinear, then x is equal to

A. -4

B. -8

C. 4

D. 8

Answer: A



Watch Video Solution

8. The vertices of a triangle are $(6, 0)$, $(0, 6)$ and $(6, 6)$.

Then distance between its circumcentre and centroid,
is

A. $2\sqrt{2}$

B. 2

C. $\sqrt{2}$

D. 1

Answer: C



View Text Solution

9. The nine point centre of the triangle with vertices

$(1, \sqrt{3})$, $(0, 0)$ and $(2, 0)$ is

A. $\left(1, \frac{\sqrt{3}}{2}\right)$

B. $\left(\frac{2}{3}, \frac{1}{\sqrt{3}}\right)$

C. $\left(\frac{2}{3}, \frac{\sqrt{3}}{2}\right)$

D. $\left(1, \frac{1}{\sqrt{3}}\right)$

Answer: D



Watch Video Solution

10. The vertices of a triangle are $(0, 0)$, $(1,0)$ and $(0,1)$.

Then excentre opposite to $(0, 0)$ is

A. $\left(1 - \frac{1}{\sqrt{2}}, 1 + \frac{1}{\sqrt{2}}\right)$

B. $\left(1 + \frac{1}{\sqrt{2}}, 1 + \frac{1}{2}\right)$

C. $\left(1 + \frac{1}{\sqrt{2}}, 1 - \frac{1}{\sqrt{2}}\right)$

D. $\left(1 - \frac{1}{\sqrt{2}}, 1 - \frac{1}{\sqrt{2}}\right)$

Answer: B



Watch Video Solution

11. If α, β, γ are the real roots of the equation $x^3 - 3px^2 + 3qx - 1 = 0$, then find the centroid of the triangle whose vertices are $\left(\alpha, \frac{1}{\alpha}\right)$, $\left(\beta, \frac{1}{\beta}\right)$ and $\left(\gamma, \frac{1}{\gamma}\right)$.



[Watch Video Solution](#)

12. If centroid of a triangle be $(1, 4)$ and the coordinates of its any two vertices are $(4, -8)$ and $(-9, 7)$, find the area of the triangle.



[Watch Video Solution](#)

13. Find the centroid and incentre of the triangle whose vertices are $(1, 2)$, $(2, 3)$ and $(3, 4)$.



Watch Video Solution

14. Show that the area of the triangle with vertices $(\lambda, \lambda - 2)$, $(\lambda + 3, \lambda)$ and $(\lambda + 2, \lambda + 2)$ is independent of λ .



Watch Video Solution

15. Prove that the $(a, b + c)$, $(b, c + a)$ and $(c, a + b)$ are collinear.



[Watch Video Solution](#)

16. Prove that the points (a, b) , (c, d) and $(a-c, b-d)$ are collinear, if $ad = bc$.



[Watch Video Solution](#)

17. If the points (x_1, y_1) , (x_2, y_2) , and (x_3, y_3) are collinear show that

$$\frac{y_2 - y_3}{x_2 x_3} + \frac{y_3 - y_1}{x_3 x_1} + \frac{y_1 - y_2}{x_1 x_2} = 0$$



[Watch Video Solution](#)

18. The coordinates of points A,B,C and D are $(-3, 5)$, $(4, -2)$, $(x, 3x)$ and $(6, 3)$ respectively and Area of $\frac{\Delta ABC}{\Delta BCD} = \frac{2}{3}$, find x.



Watch Video Solution

19. Find the area of the hexagon whose consecutive vertices are $(5, 0)$, $(4, 2)$, $(1, 3)$, $(-2, 2)$, $(-3, -1)$ and $(0, -4)$



Watch Video Solution

1. The equation of the locus of points equidistant from $(-1, -1)$ and $(4, 2)$ is

A. $3x - 5y - 7 = 0$

B. $5x + 3y - 9 = 0$

C. $4x + 3y + 2 = 0$

D. $x - 3y + 5 = 0$

Answer: B



Watch Video Solution

2. The equation of the locus of a point which moves so that its distance from the point $(ak, 0)$ is k times its distance from the point $\left(\frac{a}{k}, 0\right)$ ($k \neq 1$) is

A. $x^2 - y^2 = a^2$

B. $2x^2 - y^2 = 2a^2$

C. $xy = a^2$

D. $x^2 + y^2 = a^2$

Answer: D



Watch Video Solution

3. If the coordinates of a variable point P be $\left(t + \frac{1}{t}, t - \frac{1}{t}\right)$, where t is the variable quantity, then the locus of P is

A. $xy = 8$

B. $2x^2 - y^2 = 8$

C. $x^2 - y^2 = 4$

D. $2x^2 + 3y^2 = 5$

Answer: C



Watch Video Solution

4. If the coordinates of a variable point be $(\cos \theta + \sin \theta, \sin \theta - \cos \theta)$, where θ is the parameter, then the locus of P is

A. $x^2 - y^2 = 4$

B. $x^2 + y^2 = 2$

C. $xy = 3$

D. $x^2 + 2y^2 = 3$

Answer: B



Watch Video Solution

5. If a point moves such that twice its distance from the axis of x exceeds its distance from the axis of y by 2, then its locus is

A. $x - 2y = 2$

B. $x + 2y = 2$

C. $2y - x = 2$

D. $2y - 3x = 5$

Answer: C



Watch Video Solution

6. The equation $4xy - 3x^2 = a^2$ become when the axes are turned through an angle $\tan^{-1} 2$ is

A. $x^2 + 4y^2 = a^2$

B. $x^2 - 4y^2 = a^2$

C. $4x^2 + y^2 = a^2$

D. $4x^2 - y^2 = a^2$

Answer: B



Watch Video Solution

7. Transform the equation

$x^2 - 3xy + 11x - 12y + 36 = 0$ to parallel axes

through the point $(-4, 1)$ becomes $ax^2 + bxy + 1 = 0$

then $b^2 - a =$

A. $\frac{1}{4}$

B. $\frac{1}{16}$

C. $\frac{1}{64}$

D. $\frac{1}{256}$

Answer: C



Watch Video Solution

8. Find the locus of a point equidistant from the point $(2,4)$ and the y -axis.



Watch Video Solution

9. Find the equation of the locus of the points twice as from $(-a, 0)$ as from $(a, 0)$.



Watch Video Solution

10. OA and OB are two perpendicular straight lines. A straight line AB is drawn in such a manner that $OA + OB = 8$. Find the locus of the mid point of AB .

 [Watch Video Solution](#)

11. The ends of a rod of length l move on two mutually perpendicular lines. Find the locus of the point on the rod which divides it in the ratio $1 : 2$.

 [Watch Video Solution](#)

12. The coordinates of three points O, A, B are $(0, 0)$, $(0, 4)$ and $(6, 0)$ respectively. A point P moves so that the area of $\triangle POA$ is always twice the area of $\triangle POB$. Find the equation to both parts of the locus of P .

 [Watch Video Solution](#)

13. What does the equation

$(a - b)(x^2 + y^2) - 2abx = 0$ become if the origin is shifted to the point $\left(\frac{ab}{a - b}, 0\right)$ without rotation?

 [Watch Video Solution](#)

14. The equation $x^2 + 2xy + 4 = 0$ is transformed to the parallel axes through the point $(6, \lambda)$. For what value of λ its new form passes through the new origin ?

 [View Text Solution](#)

15. Show that if the axes be turned through $7\frac{1}{2}^\circ$, the equation $\sqrt{3}x^2 + (\sqrt{3} - 1)xy - y^2 = 0$ become free of xy in its new form.



Watch Video Solution

16. Find the angle through which the axes may be turned so that the equation $Ax + By + C = 0$ may reduce to the form $x = \text{constant}$, and determine the value of this constant.



Watch Video Solution

17.

Transform

$$12x^2 + 7xy - 12y^2 - 17x - 31y - 7 = 0 \quad \text{to}$$

rectangular through the point $(1, -1)$ inclined at an

angle $\tan^{-1}\left(\frac{4}{3}\right)$ to the original axes.



[View Text Solution](#)

Exercise Single Option Correct Type Questions

1. Vertices of a variable triangle are

$(3, 4)$, $(5 \cos \theta, 5 \sin \theta)$ and $(5 \sin \theta, -5 \cos \theta)$, where

$\theta \in R$. Locus of its orthocentre is

A. $x^2 + y^2 + 6x + 8y - 25 = 0$

$$B. x^2 + y^2 - 6x + 8y - 25 = 0$$

$$C. x^2 + y^2 + 6x - 8y - 25 = 0$$

$$D. x^2 + y^2 - 6x - 8y - 25 = 0$$

Answer: D



Watch Video Solution

2. If a rod AB of length 2 units slides on coordinate axes in the first quadrant. An equilateral triangle ABC is completed with C on the side away from O. Then, locus of C is

$$A. x^2 + y^2 - xy + 1 = 0$$

B. $x^2 + y^2 - xy\sqrt{3} + 1 = 0$

C. $x^2 + y^2 + xy\sqrt{3} - 1 = 0$

D. $x^2 + y^2 - xy\sqrt{3} - 1 = 0$

Answer: D



View Text Solution

3. The sides of a triangle are $3x + 4y$, $4x + 3y$ and $5x + 5y$ units, where $x > 0$, $y > 0$. The triangle is

A. right angled

B. acute angled

C. obtuse angled

D. isosceles

Answer: C



Watch Video Solution

4. Let P and Q be the points on the line joining $A(-2, 5)$ and $B(3, 1)$ such that $AP = PQ = QB$.

Then, the mid-point of PQ is

A. $\left(\frac{1}{2}, 3\right)$

B. $\left(-\frac{1}{4}, 4\right)$

C. $(2, 3)$

D. $(-1, 4)$

Answer: A



Watch Video Solution

5. A triangle ABC right angled at A has points A and B as $(2, 3)$ and $(0, -1)$ respectively. If $BC = 5$ units, then the point C is

A. $(4, 2)$

B. $(-4, 2)$

C. $(-4, 4)$

D. $(4, -4)$

Answer: A



Watch Video Solution

6. The locus of a point P which divides the line joining $(1, 0)$ and $(2 \cos \theta, \sin \theta)$ internally in the ratio $2 : 3$ for all θ is

- A. a straight line
- B. a circle
- C. a pair of straight lines
- D. a parabola

Answer: B



[View Text Solution](#)

7. The points with coordinates $(2a, 3a)$, $(3b, 2b)$ and (c, c) are collinear

- A. for no value of a, b, c
- B. for all values of a, b, c
- C. if $a, \frac{c}{5}, b$ are in HP
- D. if $a, \frac{2c}{5}, b$ are in HP

Answer: D



[Watch Video Solution](#)

8. The vertices of a triangle are $(0, 3)$, $(-3, 0)$ and $(3, 0)$.

The coordinates of its orthocentre are

A. $(0, -2)$

B. $(0, 2)$

C. $(0, 3)$

D. $(0, -3)$

Answer: C



Watch Video Solution

9. ABC is an equilateral triangle such that the vertices B and C lie on two parallel at a distance 6. If A lies between the parallel lines at a distance 4 from one of them then the length of a side of the equilateral triangle.

A. 8

B. $\sqrt{\frac{88}{3}}$

C. $\frac{4\sqrt{7}}{\sqrt{3}}$

D. None of these

Answer: C



Watch Video Solution

10. A, B, C are respectively the points (1,2), (4, 2), (4, 5).

If T_1, T_2 are the points of trisection of the line segment BC, the area of the quadrilateral $T_1S_1S_2T_2$ is

A. 1

B. $\frac{3}{2}$

C. 2

D. $\frac{5}{2}$

Answer: B



Watch Video Solution

11. (i) The points $(-1, 0)$, $(4, -2)$ and $(\cos 2\theta, \sin 2\theta)$ are collinear

(ii) The points $(-1, 0)$, $(4, -2)$ and $\left(\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}, \frac{2 \tan \theta}{1 + \tan^2 \theta}\right)$ are collinear

A. both statements are equivalent

B. statement (i) has more solution than statement (ii) for θ

C. statement (ii) has more solution than statement (i) for θ

D. None of the above

Answer: B



12. If $\alpha_1, \alpha_2, \alpha_3, \beta_1, \beta_2, \beta_3$ are the values of n for which $\sum_{r=0}^{n-1} x^{2r}$ is divisible by $\sum_{r=0}^{n-1} x^r$, then the triangle having vertices $(\alpha_1, \beta_1), (\alpha_2, \beta_2)$ and (α_3, β_3) cannot be

- A. an isosceles triangle
- B. a right angled isosceles triangle
- C. a right angled triangle
- D. an equilateral triangle

Answer: D



13. A triangle ABC with vertices $A(-1, 0)$, $B\left(-2, \frac{3}{4}\right)$, and $C\left(-3, -\frac{7}{6}\right)$ has its orthocentre at H . Then, the orthocentre of triangle BCH will be (a) $(-3, -2)$ (b) $(1, 3)$ (c) $(-1, 2)$ (d) none of these

A. $(-3, -2)$

B. $(1, 3)$

C. $(-1, 2)$

D. None of these

Answer: D



14.

If

$$\sum_{i=1}^4 (x_i^2 + y_i^2) \leq 2x_1x_3 + 2x_2x_4 + 2y_2y_3 + 2y_1y_4,$$

the points $(x_1, y_1), (x_2, y_2), (x_3, y_3), (x_4, y_4)$ are the vertices of a rectangle collinear the vertices of a trapezium none of these

- A. the vertices of a rectangle
- B. collinear
- C. the vertices of a trapezium
- D. None of these

Answer: A



Watch Video Solution

15. Without change of axes the origin is shifted to (h, k) , then from the equation $x^2 + y^2 - 4x + 6y - 7 = 0$, then term containing linear powers are missing, then point (h, k) is

A. $(3, 2)$

B. $(-3, 2)$

C. $(2, -3)$

D. $(-2, -3)$

Answer: C



Watch Video Solution

Exercise More Than One Correct Option Type Questions

1. If $(-6, -4)$, $(3, 5)$, $(-2, 1)$ are the vertices of a parallelogram, then remaining vertex can be

A. $(0, -1)$

B. $(-1, 0)$

C. $(-11, -8)$

D. $(7, 10)$

Answer: B::C::D



Watch Video Solution

2. If the point $P(x, y)$ be equidistant from the points $A(a + b, a - b)$ and $B(a - b, a + b)$ then

A. $ax = by$

B. $bx = ay$

C. $x^2 - y^2 = 2(ax + by)$

D. P can be (a, b)

Answer: B::D



Watch Video Solution

3. If the vertices P, Q, R of a triangle PQR are rational points, which of the following points of the triangle POR is (are) always rational point(s) ?

- A. centroid
- B. incentre
- C. circumcentre
- D. orthocentre

Answer: A::C::D



4. Show that the following points are the vertices of a rectangle.

(i) A(-4, -1), B(-2, -4), C(4, 0) and D(2, 3)

(ii) A(2, -2), B(14, 10), C(11, 13) and D(-1, 1)

(iii) A(0, -4), B(6, 2), C(3, 5) and D(-3, -1)

A. parallelogram

B. rectangle

C. rhombus

D. square

Answer: A:B



Watch Video Solution

5. The medians AD and BE of the triangle with vertices A(0, b), B(0, 0) and C(a, 0) are mutually perpendicular if

A. $b = a\sqrt{2}$

B. $a = b\sqrt{2}$

C. $b = -a\sqrt{2}$

D. $a = -b\sqrt{2}$

Answer: B::D



Watch Video Solution

6. The points $A(x, y)$, $B(y, z)$ and $C(z, x)$ represents the vertices of a right angled triangle, if

A. $x = y$

B. $y = z$

C. $z = x$

D. $x = y = z$

Answer: A::B::C



Watch Video Solution

7. Let the base of a triangle lie along the line $x = a$ and be of length $2a$. The area of this triangles is a^2 , if the

vertex lies on the line

A. $x = -a$

B. $x = 0$

C. $x = \frac{a}{2}$

D. $x = 2a$

Answer: B::D



Watch Video Solution

Exercise Passage Based Questions

1. ABC is a triangle right angled at A, $AB = 2AC$, $A = (1, 2)$, $B(-3, 1)$. The vertices of the triangles are in anticlockwise sense. BCEF is a square with vertices in clockwise sense.

- A. 42
- B. 51
- C. 62
- D. 102

Answer: B



Watch Video Solution

2. ABC is a triangle right angled at A, $AB = 2AC$, $A = (1, 2)$, $B(-3, 1)$. The vertices of the triangles are in anticlockwise sense. BCEF is a square with vertices in clockwise sense.

A. $\sqrt{17}\sqrt{(4 - \sqrt{3})}$

B. $\frac{\sqrt{17}}{2}\sqrt{(8 + \sqrt{3})}$

C. $\frac{\sqrt{17}}{2}\sqrt{(4 + \sqrt{3})}$

D. $\sqrt{15}\sqrt{(4 + \sqrt{3})}$

Answer: B



Watch Video Solution

3. ABC is a triangle right angled at A, $AB = 2AC$, $A = (1, 2)$, $B(-3, 1)$. The vertices of the triangles are in anticlockwise sense. BCEF is a square with vertices in clockwise sense.

A. $-\frac{1}{4}$

B. $-\frac{3}{4}$

C. $-\frac{5}{4}$

D. $-\frac{7}{4}$

Answer: D



Watch Video Solution

4. Let $O(0, 0)$ and $B\left(1, \frac{1}{\sqrt{3}}\right)$ be the vertices of a triangle. Let R be the region consisting of all those points P inside $\triangle OAB$ satisfying $d(P, OA) \leq \min\{d(P, OB), d(P, AB)\}$, where d denotes the distance from the point P to the corresponding line. Let M be peak of region R .

The perimeter of region R is equal to

A. $\sqrt{3}$

B. $\frac{1}{\sqrt{3}}$

C. 3

D. $2 - \sqrt{3}$

Answer: D



Watch Video Solution

5. Let $O(0, 0)$, $A(2, 0)$ and $B\left(1, \frac{1}{\sqrt{3}}\right)$ be the vertices of a triangle. Let R be the region consisting of all those points P inside $\triangle OAB$ satisfying $d(P, OA) \leq \min \{d(P, OB), d(P, AB)\}$, where d denotes the distance from the point P to the corresponding line. Let M be peak of region R . The perimeter of region R is equal to

A. $4 - \sqrt{3}$

B. $4 + \sqrt{3}$

C. $4 + 3\sqrt{3}$

D. $2 + 4\sqrt{(2 - \sqrt{3})}$

Answer: D



Watch Video Solution

6. Let $O(0, 0)$, $A(2, 0)$, and $B\left(1, \frac{1}{\sqrt{3}}\right)$ be the vertices of a triangle. Let R be the region consisting of all those points P inside OAB which satisfy $d(P, OA) \leq \min [d(p, OB), d(P, AB)]$, where d denotes the distance from the point to the

corresponding line. Sketch the region R and find its area.

A. $2 - \sqrt{3}$

B. $2 + \sqrt{3}$

C. $2\sqrt{3}$

D. $4 + \sqrt{3}$

Answer: A



Watch Video Solution

Exercise Single Integer Answer Type Questions

1. If the area of the triangle formed by the points $(2a, b)$, $(a + b, 2b + a)$ and $(2b, 2a)$ be Δ_1 and the area of the triangle whose vertices are $(a + b, a - b)$, $(3b - a, b + 3a)$ and $(3a - b, 3b - a)$ be Δ_2 , then the value of Δ_2 / Δ_1 is

 [Watch Video Solution](#)

2. The diameter of the nine point circle of the triangle with vertices $(3, 4)$, $(5 \cos \theta, 5 \sin \theta)$ and $(5 \sin \theta, -5 \cos \theta)$, where $\theta \in R$, is

 [Watch Video Solution](#)

3. The ends of the base of an isosceles triangle are $(2\sqrt{2}, 0)$ and $(0, \sqrt{2})$. One side is of length $2\sqrt{2}$. If Δ be the area of triangle, then the value of $[\Delta]$ is (where $[.]$ denotes the greatest integer function)



[Watch Video Solution](#)

4. If (x, y) is the incentre of the triangle formed by the points $(3, 4)$, $(4, 3)$ and $(1, 2)$, then the value of x^2 is



[Watch Video Solution](#)

5. Let P and Q be points on the line joining A(-2, 5) and B(3, 1) such that $AP = PQ = QB$. If mid-point of PQ is (a, b), then the value of $\frac{b}{a}$ is



[Watch Video Solution](#)

Exercise Statement I And Ii Type Questions

1. The vertices of a triangle are $A(1, 2)$, $B(-1, 3)$ and $C(3, 4)$. Let D, E, F divide BC, CA, AB respectively in the same ratio.

Statement I : The centroid of triangle DEF is (1, 3).

Statement II : The triangle ABC and DEF have the same centroid.

A. Statement I is true, Statement II is true,

Statement II is a correct explanation for

Statement I.

B. Statement I is true, Statement II is true,

Statement II is not a correct explanation for

Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

Answer: A



 [View Text Solution](#)

2. Statement 1 : Let the vertices of a ABC be $A(-5, -2)$, $B(7, 6)$, and $C(5, -4)$. Then the coordinates of the circumcenter are $(1, 2)$. Statement 2 : In a right-angled triangle, the midpoint of the hypotenuse is the circumcenter of the triangle.

 [Watch Video Solution](#)

3. A line segment AB is divided internally and externally in the same ratio at P and Q respectively

and M is the mid-point of AB. Statement I : MP, MB,

MQ are in G.P. Statement II : AP, AB and AQ are in H.P.



[Watch Video Solution](#)

4. Transform the equation

$x^2 - 3xy + 11x - 12y + 36 = 0$ to parallel axes

through the point $(-4, 1)$ becomes $ax^2 + bxy + 1 = 0$

then $b^2 - a =$



[Watch Video Solution](#)

Exercise Subjective Type Questions

1. If a, b, c be the p th, q th and r th terms respectively of a HP, show that the points (bc, p) , (ca, q) and (ab, r) are collinear.

 [Watch Video Solution](#)

2. A line L intersects three sides BC, CA and AB of a triangle in P, Q, R respectively, show that

$$\frac{BP}{PC} \cdot \frac{CQ}{QA} \cdot \frac{AR}{RB} = -1$$

 [Watch Video Solution](#)

3. If the points $\left(\frac{a^3}{a-1}, \frac{a^2-3}{a-1}\right)$, $\left(\frac{b^3}{b-1}, \frac{b^2-3}{b-1}\right)$, $\left(\frac{c^3}{c-1}, \frac{c^2-3}{c-1}\right)$ are collinear for 3 distinct values a, b, c and $a \neq 1, b \neq 1, c \neq 1$, then find the value of $abc - (ab + bc + ca) + 3(a + b + c)$.



[Watch Video Solution](#)

4. If $A_1, A_2, A_3, \dots, A_n$ are n points in a plane whose coordinates are $(x_1, y_1), (x_2, y_2), (x_3, y_3), \dots, (x_n, y_n)$ respectively. A_1A_2 is bisected in the point G_1 ; G_1A_3 is divided at G_2 in the ratio 1 : 2, G_2A_4 at G_3 in the ratio 1 : 3 and so on until all the points are exhausted. Show that the

coordinates of the final point so obtained are

$$\frac{x_1 + x_2 + \dots + x_n}{n} \text{ and } \frac{y_1 + y_2 + \dots + y_n}{n}$$

 [Watch Video Solution](#)

5. If by change of axes without change of origin, the

expression $ax^2 + 2hxy + by^2$ becomes

$a_1x_1^2 + 2h_1x_1y_1 + b_1y_1^2$, prove that

$$(a - b)^2 + 4h^2 = (a_1 - b_1)^2 + 4h_1^2$$

 [Watch Video Solution](#)

Exercise 7

1. If by change of axes without change of origin, the expression $ax^2 + 2hxy + by^2$ becomes

$a_1x_1^2 + 2h_1x_1y_1 + b_1y_1^2$, prove that

$$(a - b)^2 + 4h^2 = (a_1 - b_1)^2 + 4h_1^2$$



[Watch Video Solution](#)

Exercise Questions Asked In Previous 13 Years Exam

1. If a vertex of a triangle is $(1, 1)$ and the mid-points of two side through this vertex are $(-1, 2)$ and $(3, 2)$, then centroid of the triangle is

A. $\left(\frac{1}{3}, \frac{7}{3}\right)$

B. $\left(1, \frac{7}{3}\right)$

C. $\left(-\frac{1}{3}, \frac{7}{3}\right)$

D. $\left(-1, \frac{7}{3}\right)$

Answer: B



Watch Video Solution

2. Let $O(0, 0)$, $P(3, 4)$, $Q(6, 0)$ be the vertices of the triangle OPQ . The point R inside the triangle OPQ is such that the triangles OPR , PQR are of equal area.

The coordinates of R are

A. $\left(\frac{4}{3}, 3\right)$

B. $\left(3, \frac{2}{3}\right)$

C. $\left(3, \frac{4}{3}\right)$

D. $\left(\frac{4}{3}, \frac{2}{3}\right)$

Answer: C



Watch Video Solution

3. Let $A(h, k)$, $B(1, 1)$ and $C(2, 1)$ be the vertices of a right angled triangle with AC as its hypotenuse. If the area of the triangle is 1, then the set of values which k can take is given by (1) $\{1, 3\}$ (2) $\{0, 2\}$ (3) $\{-1, 3\}$ (4) $\{-3, -2\}$

A. $\{1, 3\}$

B. $\{0, 2\}$

C. $\{-1, 3\}$

D. $\{-3, -2\}$

Answer: C



Watch Video Solution

4. Three distinct point A, B and C are given in the 2-dimensional coordinates plane such that the ratio of the distance of any one of them from the point $(1, 0)$ to the distance from the point $(-1, 0)$ is equal to $\frac{1}{3}$.

Then, the circumcentre of the triangle ABC is at the point

A. $\left(\frac{5}{4}, 0\right)$

B. $\left(\frac{5}{2}, 0\right)$

C. $\left(\frac{5}{3}, 0\right)$

D. $(0, 0)$

Answer: A



Watch Video Solution

5. The x -coordinate of the incentre of the triangle that has the coordinates of mid-points its sides are

(0,1), (1,1) and (1, 0) is

A. $2 + \sqrt{2}$

B. $2 - \sqrt{2}$

C. $1 + \sqrt{2}$

D. $1 - \sqrt{2}$

Answer: B



Watch Video Solution

6. The number of points, having both coordinates are integers, that lie in the interior of the triangle with vertices (0,0), (0,41) and (41, 0) is

A. 820

B. 780

C. 901

D. 861

Answer: B



Watch Video Solution

7. Let k be an integer such that the triangle with vertices $(k, -3k)$, $(5, k)$ and $(-k, 2)$ has area 28 sq units. Then, the orthocentre of this triangle is at the point

A. $\left(2, \frac{1}{2}\right)$

B. $\left(2, -\frac{1}{2}\right)$

C. $\left(1, \frac{3}{4}\right)$

D. $\left(1, -\frac{3}{4}\right)$

Answer: A



Watch Video Solution