



## **CHEMISTRY**

## NCERT - NCERT CHEMISTRY(ENGLISH)

# ALDEHYDES, KETONES AND CARBOXYLIC ACIDS

**Solved Examples** 

**1.** Give names of the reagents to bring about the following transformations:

- (i) Hexan-1-ol to hexanal
- (ii) Cyclohexanol to cyclohexanone
- (iii) p-Fluorotoluene to
- (iv) Ethanenitrile to ethanal p-fluorobenzaldehyde
- (v) Allyl alcohol to propenal
- (vi) But-2-ene to ethanal



**2.** Arrange the following compounds in the increasing order of their boiling points:

 $CH_3CH_2CH_2CHO, CH_3CH_2CH_2CH_2OH, H_5C_2-O-C_2H_5, CH_3CH_2OH, CH_3CH_2OH$ 

Watch Video Solution

3. Would you expect benzaldehyde to be more reactive or less reactive in

nucleophilic addition reactions than propanal? Explain your answer.



**4.** An organic compound containing (x) with molecular formula  $C_8H_8O$  forms an oragne-red precipitate with 2, - -DNP reagent and gives yellow precipitate on heating with ioxdine in the presence of sodium hydroxide. It neither reduces Tollen's Fehlling's reagent, nor does it decolouries bromine water or Baeyer's reagent.

On drastic oxidation with chromic acid, it gives a carboxylic acid (Y) having molcular formula  $C_7H_6O_2$ . Identify the compounds X and Y.

View Text Solution

- 5. Write chemical reactions to affect the following transformations:
- (i) Butan-1-ol to butanoic acid
- (ii) Benzyl alcohol to phenylethanoic acid
- (iii) 3 Nitrobromobenzene to 3-nitrobenzoic acid
- (iv) 4-Methylacetophenone to benzene-1,4-dicarboxylic acid
- (v) Cyclohexene to hexane-1,6-dioic acid
- (vi) Butanal to butanoic acid.

Watch Video Solution



- **1.** Write the structures of the following compounds.
- (i)  $\alpha$ -Methoxypropionaldehyde
- (ii) 3-Hydroxybutanal
- (iii) 2-Hydroxycyclopentane carbaldehyde
- (iv) 4-Oxopentanal
- (v) Di-sec. butyl ketone
- (vi) 4-Fluoroacetophenone

Watch Video Solution

2. Write the structures of products of the following reactions,

(i) 
$$C_{g}H_{g} - C - CI \xrightarrow{\text{Anhyd. AlCl}_{3}}$$
  
(ii)  $(C_{g}H_{g}CH_{2})_{2}Cd + 2CH_{3}COCI \longrightarrow$   
(iii)  $H_{3}C - C = C - H \xrightarrow{-Hg^{2+},H_{2}SO_{4}}$   
(iv)  $O_{2}N \longrightarrow CH_{3} \xrightarrow{1.CrO_{2}Cl_{2}}$ 

**3.** Arrange the following compounds in increasing order of their boiling points.

 $CH_3CHO, CH_3CH_2OH, CH_3OCH_3, CH_3CH_2CH_3$ 

Watch Video Solution

4. Arrange the following compounds in increasing order of their reactivity

in nucleophilic addition reactions.

(i) Ethanal, Propanal, Propanone, Butanone.

(ii) Benzaldehyde, p-Tolualdehyde, p-Nitrobenzaldehyde, Acetophenone.

Hint: Consider steric effect and electronic effect.



5. Predict the products of the following reactions:



6. Give the IUPAC names of the following compounds:

(i) Ph  $CH_2CH_2COOH$ 

(ii)  $(CH_3)_2 C = CHOOH$ 





**7.** Show how each of the following compounds can be converted to benzoic acid.

- (i) Ethylbenzene
- (ii) Acetophenone
- (iii) Bromobenzene
- (iv) Phenylethene (Styrene)



8. Which acid of each pair shown here would you expect to be stronger?

(i)  $CH_3CO_2H$  or  $CH_2FCO_2H$ 

(ii)  $CH_2FCO_2H$  or  $CH_2ClCO_2H$ 

(iii)  $CH_2FCH_2CH_2CO_2H$  or  $CH_3CHFCH_2CO_2H$ 

Watch Video Solution

9. What is meant by the following terms ? Give an example of the reaction

in each case.

(i) Cyanohydrin

(ii) Acetal

(iii) Semicarbazone

(iv) Aldol

(v) Hemiacetal

(vi) Oxime

(vii) Ketal

(vii) Imine

(ix) 2,4-DNP-derivative

(x) Schiff's base

View Text Solution

**10.** Name the following compounds according to the IUPAC system of nomenclature:

i.  $CH_3CH(CH_3)CH_2CH_2CHO$ 

ii.  $CH_3CH_2COCH(C_2H_5)CH_2CH_2Cl$ 

iii.  $CH_3CH = CHCHO$ 

iv.  $CH_3COCH_2COCH_3$ 

v.  $CH_3CH(CH_3)CH_2C(CH_3)_2COCH_3$ 

vi.  $(CH_3)_3CCH_2COOH$ 

vii.  $OHCC_6H_4CHO-p$ 

Watch Video Solution

**11.** Draw the structures of following compound:

- i. 3-Methylbutanal
- ii. p-Nitropropiopehnone
- iii. P-Methylbenzaldehyde
- iv. 4-Methylpent-3-en-2-one
- v. 4-Chloropentan-2-one
- vi. 3-Bromo-4-phenylpentanoic acid
- vii. p-p'-Dihydroxybenzophenone
- viii. Hex-2-en-4-ynoic acid



**12.** Write the IUPAC names of following ketones and aldehydes. Wherever possible, give common nemes also.

i.  $CH_3CO(CH_2)_4CH_3$ 

ii.  $CH_3CH_2CHBrCH_2CH(CH_3)CHO$ 

iii.  $CH_3(CH_2)_5CHO$ 

iv. Ph - CH = CH - CHO



v.

### vi. PhCOPh

Watch Video Solution

13. Draw the structure of following derivatives:

i. 2,4-Dinitrophylhydrazone of benzaldehyde.

- ii. Cyclopropanone oxime
- iii. Actaldehyde dimethyl acetal
- iv. Semicarbazone of cyclobutanone
- v. Ethylene ketal of hexan-3-one
- vi. Methyl hemiacetal of formaldehyde

Watch Video Solution

**14.** Predict the products formed when cyclohexane carbaldehyde reacts with the following reagents:

- i. PhMgBr and then  $H_3O^+$
- ii. Tollens reagent
- iii. Semicarbazine and weak acid
- iv. Excess ethanol and acid
- v. Zinc amalgam and dilute hydrochloric acid

Watch Video Solution

**15.** Which of the following compounds would undergo aldol condensation

or the Cannizzaro reaction, or neither? Writer the structures of expected

products of aldol condensation and Cannizzaro reaction.

- i. Methanal ii. 2-Methylpentanal
- iii. Benzaldehyde iv. Benzophenone
- v. Cyclohexanone vi. 1-Phenylpropanone
- vii. Phenylacetaldehye viii. Butan-1-ol
- ix. 2,2-Dimethylbutanal



**17.** Write structure formulae and names of four possible aldol condensation products form propanal and butanal. In each case. Indicate which aldehyde acts as nucleophile and which as electrophile.

#### Watch Video Solution

**18.** Anorganic compound with molecular formula  $C_9H_{10}O$  forms 2,4-DNP derivative, reduces Tollens' reagent and undergoes Cannizzaro reaction. On vigorous oxidation, it gives 1,2-benzenedicarboxylic acid. Identify the compound.

**19.** An organic compound (A) (molecular formula  $C_8H_{16}O_2$ ) was hydrolysed with dilute sulphuric acid to give a carboxylic acid (B) and an alcohol (C). Oxidation of (C) with chromic acid produced (B). (C) on dehydration gives but-1-ene. Write equations for the reactions involved.

Watch Video Solution

**20.** Arrange the following compounds in increasing order of their property as indicated :

(a) Acetaldehyde, Acetone, Di-tert-butyl ketone, Methyl tert-butyl ketone

[Reactivity towards HCN]

(b)

 $CH_{3}CH_{2}CH(Br)COOH, CH_{3}CH(Br)CH_{2}COOH, (CH_{3})_{2}CHCOOH, (CH_{3})_{2}CHCOH, (CH_{3})$ 

[Acid strength]

(c) Benzoic acid, 4-Nitrobenzoic acid, 3,4-Dinitrobenzoic acid, 4-Methoxybenzoic acid [Acid strength] **21.** Give simple chemical test to distinguish between the following pairs of compounds.

- i. Propanal and Propanone
- ii. Acetophenone and Benzophenone
- iii. Phenol and Benzioc acid
- iv. Benzoic acid and Ethyl benzoate
- v. Pentan-2-one and Pentan-3-one
- vi. Benzaldehyde and Acetophenone
- vii. Ethanal and Propanal

View Text Solution

**22.** How will you prapare the following compounds from benzene ? You may use any inorganic reagent and any organic one having not more than one carbon atom.

i. Methyl benzonate ii. m-Nitrobenzonic acid

iii. p-Nitrobenzoic acid iv. Phenylacetic acid

v. p-Nitrobenzaldehyde

**Watch Video Solution** 

23. How will you bring about the following conversions in not more than

two steps ?

- i. Propanone to propene
- ii. Benzoic acid to Benzaldehyde
- iii. Ethanol to 3-Hydroxybutanal
- iv. Benzene to m-Nitroacetopenone
- v. Benzaldehyde to Benzophenone
- vi. Bromobenzene to 1-Phenylethonal
- vii. benzaldehyde to 3-Phenylpropan-1-ol
- viii. Benzaldehyde to  $\alpha$ -Hydrophylacetic acid
- ix. Benzoic acid to m-Nitrobenzyl alcohol

View Text Solution

- 24. Describe the following
- i. Acetylation
- ii. Cannizzaro reaction
- iii. Cross aldol condensation
- iv. Decarboxylation

Watch Video Solution

25. Complete each synthesis by giving missing starting material, reagent

or products



**View Text Solution** 

**26.** Giving plausible explanation for each of the following:

i. Cyclohexanone forms cyanohydrin good yield but 2,2,6trimethylcyclohexanone does not.

ii. There are two  $(-NH_2)$  groups in semicarbazide. However, only one is involved in the formation of semicarbazones.

iii. During the preparation of esters from a carboxylic acid and an alcohol in the easter should be removed as soon as it is formed.

#### View Text Solution

**27.** An organic compound contains 69.77 % carbon, 11.63 % hydrogen, and rest oxygen. The molecular mass of the compound is 86. It does not reduce Tollens reagent but forms an aditional compound with sodium hydrogensulphite and gives positive iodoform test. On vigorous oxidation, it gives ethanoic and propanoic acid. Write the possible structure of the compound.

**28.** Although phenoxide ion has more number of resonating structures than carboxylate ion, carboxylic acid is a stronger than phenol. Why ?

