



MATHS

BOOKS - RD SHARMA MATHS (HINGLISH)

ALGEBRA OF MATRICES

Solved Examples And Exercises

1. Express the matrix $A = \begin{bmatrix} 4 & 2 & -1 \\ 3 & 5 & 7 \\ 1 & -2 & 1 \end{bmatrix}$ as the sum of a symmetric and a skew-symmetric matrix.

 [Watch Video Solution](#)

2. A matrix which is both symmetric as well as skew-symmetric is a null matrix.

 [Watch Video Solution](#)

3. Show that positive odd integral powers of a skew-symmetric matrix are skew-symmetric and positive even integral powers of a skew-symmetric matrix are symmetric.

 [Watch Video Solution](#)

4. Show that all positive integral powers of a symmetric matrix are symmetric.

 [Watch Video Solution](#)

5. If B is a skew-symmetric matrix, write whether the matrix ABA^T is symmetric or skew-symmetric.

 [Watch Video Solution](#)

6. If matrix $A = [a_{ij}]_{2 \times 2}$, where $a_{ij} = \begin{cases} 1 & i \neq j \\ 0 & i = j \end{cases}$, then A^2 is equal to

A. $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$

B. $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$

C. $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$

D. $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

Answer: D



[Watch Video Solution](#)

7. Find the value of λ , a non-zero scalar, if

$$\lambda[102345] + 2[123 - 1 - 32] = [44104214]$$



[Watch Video Solution](#)

8. Solve the matrix equation $[x^2, y^2] - 3[x, 2y] = [-2, 9]$



[Watch Video Solution](#)

 Watch Video Solution

9. If A, B and C three matrices of the same order, then prove that

$$A = B \Rightarrow A + C = B + C$$

 Watch Video Solution

10. Find non-zero values of x satisfying the matrix equation :

$$x \begin{bmatrix} 2x & 2 \\ 3 & x \end{bmatrix} + 2 \begin{bmatrix} 8 & 5x \\ 4 & 4x \end{bmatrix} = 2 \begin{bmatrix} x^2 + 8 & 24 \\ 10 & 6x \end{bmatrix}$$

 Watch Video Solution

11. Find a matrix A , if $A + [23 \ -14] = [3 \ -6 \ -38]$

 Watch Video Solution

12. If $A = \text{diag}(1, -1, 2)$ and $B = \text{diag}(2, 3, -1)$, find $A + B, 3A + 4B$.



Watch Video Solution

13. If $A = [2, -1, 3, 1]$ and $B = [1, 4, 7, 2]$, find $3A - 2B$.



Watch Video Solution

14. Find the values of a and b if $A = B$, where

$$A = \begin{bmatrix} a + 4 & 3b \\ 8 & -6 \end{bmatrix}, B = \begin{bmatrix} 2a + 2 & b^2 + 2 \\ 8 & b^2 - 56 \end{bmatrix}$$



Watch Video Solution

15. Find the values of x and y if

$$[x + 10y^2 + 2y - 4] = [3x + 430y^2 - 5y]$$



Watch Video Solution

16. For what values of x and y are the following matrices equal ?

$$A = \begin{bmatrix} 2x + 1 & 3y \\ 0 & y^2 - 5y \end{bmatrix}, B = \begin{bmatrix} x + 3 & y^2 + 2 \\ 0 & -6 \end{bmatrix}$$

 [Watch Video Solution](#)

17. Let A, B be two matrices such that they commute. Show that for any positive integer n , $AB^n = B^n A$ (ii) $(AB)^n = A^n B^n$

 [Watch Video Solution](#)

18. Let $A = \begin{bmatrix} 2 & 3 \\ -1 & 2 \end{bmatrix}$ and $f(x) = x^2$ this rest of $f \in dA^5$

 [Watch Video Solution](#)

19. If $A = [35], B = [73]$, then find a non-zero matrix C such that $AC=AC$.

 [Watch Video Solution](#)

20. If A is a square matrix such that $A^2 = A$, show that $(I + A)^3 = 7A + I$.

 [Watch Video Solution](#)

21. Under what conditions is the matrix $A^2 - B^2 = (A - B)(A + B)$ is true?

 [Watch Video Solution](#)

22. If a is a non-zero real or complex number. Use the principle of mathematical induction to prove that:

If $A = \begin{bmatrix} a & 1 \\ 0 & a \end{bmatrix}$, then $A^n = \begin{bmatrix} a^n & na^{n-1} \\ 0 & a^n \end{bmatrix}$ for every positive integer n .

 [Watch Video Solution](#)

23. If $A = \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix}$ $B = \begin{bmatrix} \cos^2 \phi & \cos \phi \sin \phi \\ \cos \phi \sin \phi & \sin^2 \phi \end{bmatrix}$ and $\theta - \phi = (2n + 1) \frac{\pi}{2}$ Find AB .

 [Watch Video Solution](#)

24. If A and B are square matrices of order n , then prove that A and B will commute iff $A - \lambda I$ and $B - \lambda I$ commute for every scalar λ .

 [Watch Video Solution](#)

25. Show that positive odd integral powers of a skew-symmetric matrix are skew-symmetric and positive even integral powers of a skew-symmetric matrix are symmetric.

 [Watch Video Solution](#)

26. If $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$, then find the values of θ satisfying the equation $A^T + A = I_2$.

 [Watch Video Solution](#)

27. If $A = \text{diag}(abc)$, show that $A^n = \text{diag}(a^n b^n c^n)$ for all positive integer n .

 [Watch Video Solution](#)

28. If $A = \begin{bmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{bmatrix}$, then prove by principal of mathematical induction that $A^n = \begin{bmatrix} \cos n\theta & i \sin n\theta \\ i \sin n\theta & \cos n\theta \end{bmatrix}$ for all $n \in \mathbb{N}$.

 [Watch Video Solution](#)

29. Let $A = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 1 & 3 \\ 1 & 2 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \\ 0 & 1 & 1 \end{bmatrix}$. Find A^T, B^T and verify

that (i) $(A + B)^T = A^T + B^T$ (ii) $(AB)^T = B^T A^T$ (iii) $(2A)^T = 2A^T$

 [Watch Video Solution](#)

30. If $[xy, 4, z + 6, x + y] = [8, w, 0, 6]$, write the value of $(x + y + z)$.

 [Watch Video Solution](#)

31. If $A = [a_{ij}]$ is a skew-symmetric matrix, then write the value of

$$\sum_i a_{ij}$$

 [Watch Video Solution](#)

32. Express the matrix $[3 \ -2 \ -4 \ 3 \ -2 \ -5 \ -11 \ 2]$ as the sum of a symmetric and skew-symmetric matrix.

 [Watch Video Solution](#)



Watch Video Solution

33. Show that the elements on the main diagonal of a skew-symmetric matrix are all zero.



Watch Video Solution

34. If A is a skew-symmetric and $n \in \mathbb{N}$ such that $(A^n)^T = \lambda A^n$, write the value of λ .



Watch Video Solution

35. If A is a skew-symmetric matrix and n is an odd natural number, write whether A^n is symmetric or skew-symmetric or neither of the two.



Watch Video Solution

36. Show that the matrix $B^T A B$ is symmetric or skew-symmetric according as A is symmetric or skew-symmetric.

 [Watch Video Solution](#)

37. If A is a symmetric matrix and $n \in \mathbb{N}$, write whether A^n is symmetric or skew-symmetric or neither of these two.

 [Watch Video Solution](#)

38. If $S = [S_{ij}]$ is a scalar matrix such that $s_{ii} = k$ and A is a square matrix of the same order, then $AS = SA = ?$

(a) A^k

(b) $k + A$

(c) kA

(d) kS

 [Watch Video Solution](#)

39. If A is a square matrix, using mathematical induction prove that $(A^T)^n = (A^n)^T$ for all $n \in \mathbb{N}$.

 [Watch Video Solution](#)

40. If A and B are square matrices of the same order such that $AB = BA$, then show that $(A + B)^2 = A^2 + 2AB + B^2$

 [Watch Video Solution](#)

41. There are 2 families A and B. There are 4 men, 6 women and 2 children in family A, and 2 men, 2 women and 4 children in family B. The recommended daily amount of calories is 2400 for men, 1900 for women, 1800 for children and 45 grams of proteins for men, 55 grams for women and 33 grams for children. Represent the above information using matrix. Using matrix multiplication, calculate the total requirement for calories and proteins for each of the two families. What awareness can you create among people about the planned diet from this question?



[Watch Video Solution](#)

42. If $A = \begin{bmatrix} - & 1 & 2 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} - & 2 & - & 1 & - & 4 \end{bmatrix}$, verify that $(AB)^T = B^T A^T$.



[Watch Video Solution](#)

43. If: $\begin{bmatrix} xy & 5xz & + & 6y^2 & + & 1 \end{bmatrix} = \begin{bmatrix} 8w & 0 & 6 \end{bmatrix}$, then find the values of x, y, z and w



[Watch Video Solution](#)

44. The sales figure of two car dealers during January 2013 showed that dealer A sold 5 deluxe, 3 premium and 4 standard cars, while dealer B sold 7 deluxe, 2 premium and 3 standard cars. Total sales over the 2 month period of January-February revealed that dealer A sold 8 deluxe 7 premium and 6 standard cars. In the same 2 month period, dealer B sold 10 deluxe, 5 premium and 7 standard cars. Write 2×3 matrices summarizing sales data for January and 2 month period for each dealer.



Watch Video Solution

45. If $A^T = [34 - 1201]$ and $B = [-121123]$, find $A^T - B^T$



Watch Video Solution

46. If l_1, m_1, n_1 and l_2, m_2, n_2 are the direction cosines of two mutually perpendicular lines, show that the direction cosines of the line perpendicular to both of these are $m_1n_2 - m_2n_1, n_1l_2 - n_2l_1, l_1m_2 - l_2m_1$.



Watch Video Solution

47. Construct a 2×3 matrix $A = [a_{ij}]$ whose elements are given by

$$a_{ij} = \frac{1 - j}{1 + i}$$



Watch Video Solution

48. Construct a 3×2 matrix $A = [a_{ij}]$ whose elements are given by

$$a_{ij} = e^{ix} \sin jx \quad a_{ij} = e^{-ix} \cos\left(\frac{\pi}{2}i + jx\right)$$

 [Watch Video Solution](#)

49. For what values of x and y such that matrices A and B are equal ?

$$A = \begin{bmatrix} 2x + 1 & 3y \\ 0 & y^2 - 5y \end{bmatrix}, B = \begin{bmatrix} x + 3 & y^2 + 2 \\ 0 & -6 \end{bmatrix}$$

 [Watch Video Solution](#)

50. If $\begin{bmatrix} x - y & z \\ 2x - y & \omega \end{bmatrix} = \begin{bmatrix} -1 & 2 \\ 4 & 7 \end{bmatrix}$, $f \in d x, y, z, \omega$.

 [Watch Video Solution](#)

51. Construct a 3×4 matrix $A = [a_{ij}]$ whose elements are given by

$$a_{ij} = i + j \quad (\text{ii}) \quad a_{ij} = i - j$$

 [Watch Video Solution](#)

52. If $[x - y \ 2x + z \ 2x - y \ 3z + w] = [-15 \ 0 \ 13]$, $f \in d, xy, z, w$.

 [Watch Video Solution](#)

53. Find the values of x, y, z and a which satisfy the matrix equation

$$[x + 32y + xz - 14a - 6] = [0 - 732a]$$

 [Watch Video Solution](#)

54. In a certain city there are 30 colleges. Each college has 15 peons, 6 clerks, 1 typist and 1 section officer. Express the given information as a column matrix. Using scalar multiplication, find the total number of posts of each kind in all the colleges.

 [Watch Video Solution](#)

55.

If

$$A = [1 \ -123], B = [2110], \text{ prove that } (A + B)^2 \neq A^2 + 2AB + B^2$$



56. Find a matrix X such that $2A + B + X = 0$, where $A = [-1234]$, $B = [3 \ -215]$ if $A = [804 \ -236]$ and $B = [2 \ -242 \ -51]$, then find the matrix X of order 3×2 such that $2A + 3X = 5B$.



57. If X and Y are 2×2 matrices, then solve the following matrix equations for X and Y

$$2X + 3Y = \begin{bmatrix} 2 & 3 \\ 4 & 0 \end{bmatrix}, 3X + 2Y = \begin{bmatrix} -2 & 2 \\ 1 & -5 \end{bmatrix}$$


58. If $A = \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix}$, find the values of α for which $A^2 = B$

A. 1

B. 2

C. -1

D. No real values

Answer: No real values



[Watch Video Solution](#)

59. Find the value of x such that $\begin{bmatrix} 1 & x & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 2 \\ 2 & 5 & 1 \\ 15 & 3 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ x \end{bmatrix} = 0$



[Watch Video Solution](#)

60. If $A = \begin{bmatrix} 1 & -1 \\ 2 & -1 \end{bmatrix}$, $B = \begin{bmatrix} a & 1 \\ b & -1 \end{bmatrix}$ and $(A + B)^2 = A^2 + B^2$, find the values of a and b

 [Watch Video Solution](#)

61. If $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$, find x and y such that $(xI + yA)^2 = A$

 [Watch Video Solution](#)

62. If $\begin{bmatrix} 2 & -1 \\ 1 & 0 \\ -3 & 4 \end{bmatrix} A = \begin{bmatrix} -1 & -8 & -10 \\ 1 & -2 & -5 \\ 9 & 22 & 15 \end{bmatrix}$, find A .

 [Watch Video Solution](#)

63. Let $f(x) = x^2 - 5x + 6$. Find $f(A)$, if $A = \begin{bmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{bmatrix}$

 [Watch Video Solution](#)

64. If A and B are symmetric matrices of the same order, write whether $AB - BA$ is symmetric or skew-symmetric or neither of the two.

 [Watch Video Solution](#)

65. If A is a square matrix such that $A^2 = A$, then $(I + A)^3 - 7A$ is equal to

A. (a) A

B. (b) $I - A$

C. (c) I

D. (d) $3A$

Answer: (c) I

 [Watch Video Solution](#)

66. If A is a square matrix such that $A^2 = A$, then write the value of $7A - (I + A)^3$, where I is the identity matrix.

 [Watch Video Solution](#)

67. Construct a 3×4 matrix $A = [a_{ij}]$ whose elements are given by (i) $a_{ij} = i + j$ (ii) $a_{ij} = i - j$

 [Watch Video Solution](#)

68. If $[x - y \ 2x + z \ 2x - y \ 3z + w] = [-15 \ 0 \ 13]$, find x, y, z, w .

 [Watch Video Solution](#)

69. Find the values of x, y, z and a which satisfy the matrix equation

$$\begin{bmatrix} x + 3 & 2 \\ y + xz - 1 & 4a - 6 \end{bmatrix} = \begin{bmatrix} 0 & -7 \\ 3 & 2a \end{bmatrix}$$

 [Watch Video Solution](#)

70. A matrix has 12 elements. What are the possible orders it can have?

 [Watch Video Solution](#)

71. If $A = [a_{ij}]$ is a matrix given by

$A = [a_{ij}] = [4521 \quad -2715 \quad 1918 \quad 36 \quad -25]$ write the order of A and find the elements a_{24} , a_{34} . Also, show that $a_{32} = a_{23} + a_{24}$.

 [Watch Video Solution](#)

72. A matrix has 12 elements. What are the possible orders it can have?

 [Watch Video Solution](#)

73. Construct a 2×2 matrix $A = [a_{ij}]$ whose elements are given by

$$a_{ij} = \frac{(i + 2j)^2}{2}.$$





Watch Video Solution

74. Construct a 2×3 matrix $A = [a_{ij}]$ whose elements are given by

$$a_{ij} = \frac{i - j}{i + j}.$$



Watch Video Solution

75. Construct a 3×2 matrix $A = [a_{ij}]$ whose elements are given by

$$a_{ij} = e^{ix} \sin jx \text{ (ii) } a_{ij} = e^{-ix} \cos\left(\frac{\pi}{2}i + jx\right)$$



Watch Video Solution

76. Find x , y , z and w such that
$$\begin{bmatrix} x - y & 2z + w \\ 2x - y & 2x + w \end{bmatrix} = \begin{bmatrix} 5 & 3 \\ 12 & 15 \end{bmatrix}.$$



Watch Video Solution

77. Consider the following information regarding the number of men and women workers in three factories. I, II and III.

	Men workers	Women workers
I	30	25
II	25	31
III	27	26

Represent the above information in the form of 3×2 matrix. What does the entry in the third row and second column represent?



[Watch Video Solution](#)

78. Find Value of K, for which $A = \begin{bmatrix} K & 8 \\ 4 & 2K \end{bmatrix}$ is a singular matrix is: .

A. 4

B. -4

C. 4 or -4

D. 0

Answer: C



[Watch Video Solution](#)

79. For what values of x and y are the following matrices equal?

$$A = \begin{bmatrix} 2x + 1 & 3y \\ 0 & y^2 - 5y \end{bmatrix}, B = \begin{bmatrix} x + 3 & y^2 + 2 \\ 0 & -6 \end{bmatrix}$$

 [Watch Video Solution](#)

80. If a matrix has 8 elements, what are the possible orders it can have?

What if it has 5 elements?

 [Watch Video Solution](#)

81. If $A = [a_{ij}] = \begin{bmatrix} 2 & 3 & -5 \\ 1 & 4 & 9 \\ 0 & 7 & -2 \end{bmatrix}$ and $B = [b_{ij}] = \begin{bmatrix} 2 & -1 \\ -3 & 4 \\ 1 & 2 \end{bmatrix}$ then find

$$a_{22} + b_{21} \quad \text{(ii)} \quad a_{11}b_{11} + a_{22}b_{22}$$

 [Watch Video Solution](#)

82. Let A be a matrix of order 3×4 . If R_1 denotes the first row of A and C_2 denotes its second column, then determine the orders of matrices R_1 and C_2 .



[Watch Video Solution](#)

83. Construct a 2×3 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:

(i) $a_{ij} = i \times j$ (ii) $a_{ij} = 2i - j$



[Watch Video Solution](#)

84. Construct a 2×2 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:

(i) $a_{ij} = i + j$ (ii) $a_{ij} = \frac{(i + j)^2}{2}$



[Watch Video Solution](#)

85. Construct a 2×2 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:

$$(i) \frac{(i+j)^2}{2} \quad (ii) a_{ij} = \frac{(i-j)^2}{2} \quad (iii) a_{ij} = \frac{(i-2j)^2}{2}$$



Watch Video Solution

86. Construct a 2×2 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:

$$(i) a_{ij} = \frac{(2i+j)^2}{2} \quad (ii) a_{ij} = \frac{|2i-3j|}{2} \quad (iii) a_{ij} = \frac{|-3i+j|}{2}$$



Watch Video Solution

87. Construct a 2×2 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:

$$(i) a_{ij} = e^{2ix} \sin xj$$



Watch Video Solution

88. Construct a 3×4 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:

$$(i) a_{ij} = i + j \quad (ii) a_{ij} = 1 - j \quad (iii) a_{ij} = 2i$$



Watch Video Solution

89. Construct a 3×4 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:

(i) $a_{ij} = j$ (ii) $a_{ij} = \frac{1}{2} | -3i + j |$



Watch Video Solution

90. Construct a 4×3 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:

(i) $a_{ij} = 2i + \frac{i}{j}$



Watch Video Solution

91. Find x, y, a and b if $\begin{bmatrix} 3x + 4y & 2 & x - 2y \\ a + b & 2a - b & -1 \end{bmatrix} = \begin{bmatrix} 2 & 2 & 4 \\ 5 & -5 & -1 \end{bmatrix}$



Watch Video Solution

92. Find x, y, a and b if $\begin{bmatrix} 2x - 3y & a - b & 3 \\ 1 & x + 4y & 3a + 4b \end{bmatrix} = \begin{bmatrix} 1 & -2 & 3 \\ 1 & 6 & 29 \end{bmatrix}$



Watch Video Solution

93. Find the values of a , b , c and d from the following equations:

$$\begin{bmatrix} 2a + b & a - 2b \\ 5c - d & 4c + 3d \end{bmatrix} = \begin{bmatrix} 4 & -3 \\ 11 & 24 \end{bmatrix}$$



Watch Video Solution

94. Find x , y and z so that $A = B$, where $A = [x - 2 \quad 3z \quad 18zy + 26z]$,

$$B = [yz \quad 66yx \quad 2y]$$



Watch Video Solution

95. If $\begin{bmatrix} x & 3x - y \\ 2x + z & 3y - \omega \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 4 & 7 \end{bmatrix}$, find x , y , z , ω .



Watch Video Solution

96. If $\begin{bmatrix} x - y & z \\ 2x - y & \omega \end{bmatrix} = \begin{bmatrix} -1 & 4 \\ 0 & 5 \end{bmatrix}$, find x , y , z , ω .



Watch Video Solution

97. If
$$\begin{bmatrix} x + 3 & z + 4 & 2y - 7 \\ 4x + 6 & a - 1 & 0 \\ b - 3 & 3b & z + 2c \end{bmatrix} = \begin{bmatrix} 0 & 6 & 3y - 2 \\ 2x & -3 & 2c + 2 \\ 2b + 4 & -21 & 0 \end{bmatrix}$$
 Obtain the values of a , b , c , x , y and z .



Watch Video Solution

98. If $[2x + 15x0y^2 + 1] = [x + 310026]$, find the value of $(x + y)$.



Watch Video Solution

99. If $[xy4z + 6x + y] = [8w06]$, then find the values of x , y , z and w .



Watch Video Solution

100. Give an example of (i) a row matrix which is also a column matrix



[Watch Video Solution](#)

101. The sales figure of two car dealers during January 2013 showed that dealer A sold 5 deluxe, 3 premium and 4 standard cars, while dealer B sold 7 deluxe, 2 premium and 3 standard cars. Total sales over the 2 month period of January-February revealed that dealer A sold 8 deluxe 7 premium and 6 standard cars. In the same 2 month period, dealer B sold 10 deluxe, 5 premium and 7 standard cars. Write 2×3 matrices summarizing sales data for January and 2-month period for each dealer.

[Watch Video Solution](#)

102. For what values of x and y are the following matrices equal?

$$A = \begin{bmatrix} 2x + 13y \\ 0y^2 - 5y \end{bmatrix}, B = \begin{bmatrix} x + 3y^2 + 20 \\ -6 \end{bmatrix}$$

[Watch Video Solution](#)

103. Find the values of x and y if

$$\begin{bmatrix} x + 10 & y^2 + 2y \\ 0 & -4 \end{bmatrix} = \begin{bmatrix} 3x + 4 & 3 \\ 0 & y^2 - 5y \end{bmatrix}$$

 [Watch Video Solution](#)

104. Find the values of a and b if $A = B$, where $A = [a + 43b8 - 6]$,

$$B = [2a + 2b^2 + 28b^2 - 56]$$

 [Watch Video Solution](#)

105. If $A = [234046589]$ and $B = [305532047]$, find $3A - 2B$.

 [Watch Video Solution](#)

106. If $A = \begin{bmatrix} 2 & 3 & -5 \\ 1 & 2 & -1 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 5 & 1 \\ -2 & 7 & 3 \end{bmatrix}$, find $A + B$ and $A - B$

 [Watch Video Solution](#)

107. $A = \begin{bmatrix} 2 & -1 \\ 3 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 4 \\ 7 & 2 \end{bmatrix}$, find $3A - 2B$.

 [Watch Video Solution](#)

108. If $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{bmatrix}$, find $A + B$,
 $3A + 4B$.

 [Watch Video Solution](#)

109. Simplify:
 $\cos \theta [\cos \theta \sin \theta - \sin \theta \cos \theta] + \sin \theta [\sin \theta - \cos \theta \cos \theta \sin \theta]$

 [Watch Video Solution](#)

110. Find X and Y , if $X + Y = [7025]$ and $X - Y = [3003]$.

 [Watch Video Solution](#)

111. Find a matrix A , if $A + [23 - 14] = [3 - 6 - 38]$.

 [Watch Video Solution](#)

112. Find x, y, z, t if $2 \begin{bmatrix} x & z \\ y & t \end{bmatrix} + 3 \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} = 3 \begin{bmatrix} 3 & 5 \\ 4 & 6 \end{bmatrix}$.

 [Watch Video Solution](#)

113. Find non-zero values of x satisfying the matrix equation:

$$x[2x23x] + 2[85x44x] = 2[x^2 + 824106x]$$

 [Watch Video Solution](#)

114. If A, B and C are three matrices of the same order, then prove that

$$A = B \Rightarrow A + C = B + C.$$

 [Watch Video Solution](#)

115. Find a matrix X such that $2A + B + X = O$, where $A = \begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 3 & -2 \\ 1 & 5 \end{bmatrix}$.

 [Watch Video Solution](#)

116. Find a matrix A such that $2A - 3B + 5C = O$, where $B = \begin{bmatrix} -2 & 2 & 0 \\ 3 & 1 & 4 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & 0 & -2 \\ 7 & 1 & 6 \end{bmatrix}$.

 [Watch Video Solution](#)

117. Solve the matrix equation $[x^2y^2] - 3[x2y] = [-29]$.

 [Watch Video Solution](#)

118. Two farmers Ramkishan and Gurcharan Singh cultivates only three varieties of rice namely Basmati, Permal and Naura. The sale (in Rupees)

of these varieties of rice by both the farmers in the month of September and October are given by the following matrices A and B. (i) Find the combined sales in September and October for each farmer in each variety. (ii) Find the decrease in sales from September to October. (iii) if both farmers receive 2% profit on gross sales, compute the profit for each farmer and for each variety sold in October.

 [Watch Video Solution](#)

119. Compute the following sums: $\begin{bmatrix} 3 & -2 \\ 1 & 4 \end{bmatrix} + \begin{bmatrix} -2 & 4 \\ 1 & 3 \end{bmatrix}$

 [Watch Video Solution](#)

120. Let $A = [2432]$, $B = [13 - 25]$ and $C = [- 2534]$. Find each of the following: $2A - 3B$ (ii) $B - 4C$ (iii) $3A - C$ (iv) $3A - 2B + 3C$

 [Watch Video Solution](#)

121. If $A = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix}$, $B = \begin{bmatrix} -1 & 0 & 2 \\ 3 & 4 & 1 \end{bmatrix}$, $C = \begin{bmatrix} -1 & 2 & 3 \\ 2 & 1 & 0 \end{bmatrix}$, find $A + B$ and $B + C$ (ii) $2B + 3A$ and $3C - 4B$.

 [Watch Video Solution](#)

122. If $A = \begin{bmatrix} -1 & 0 & 2 \\ 3 & 1 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 0 & -2 & 5 \\ 1 & -3 & 1 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & -5 & 2 \\ 6 & 0 & -4 \end{bmatrix}$, then find $(2A - 3B + 4C)$.

 [Watch Video Solution](#)

123. If $A = \text{diag}(2 \ -5 \ 9)$, $B = \text{diag}(1 \ 1 \ -4)$ and $C = \text{diag}(-6 \ 3 \ 4)$, find $A - 2B$ (ii) $B + C - 2A$ (iii) $2A + 3B - 5C$

 [Watch Video Solution](#)

124. Given the matrices $A = [2113 \ -10024]$, $B = [97 \ -1354216]$ and $C = [2 \ -431 \ -10945]$ Verify that $(A + B) + C = A + (B + C)$.



[Watch Video Solution](#)

125. Find matrices X and Y , if $X + Y = \begin{bmatrix} 5 & 2 \\ 0 & 9 \end{bmatrix}$ and $X - Y = \begin{bmatrix} 3 & 6 \\ 0 & -1 \end{bmatrix}$.



[Watch Video Solution](#)

126. Find X if $Y = \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}$ and $2X + Y = \begin{bmatrix} 1 & 0 \\ -3 & 2 \end{bmatrix}$.



[Watch Video Solution](#)

127. Find matrices X and Y , if $2X + Y = [6 - 60 - 421]$ and $X + 2Y = [325 - 21 - 7]$.



[Watch Video Solution](#)

128. If $X - Y = [111110100]$ and $X + Y = [351 - 1141180]$, find X and Y .

 [Watch Video Solution](#)

129. Find matrix A , if $[12 - 1049] + A = [9 - 14 - 213]$.

 [Watch Video Solution](#)

130. If $A = [9178]$, $B = [15712]$, find matrix C such that $5A + 3B + 2C$ is a null matrix.

 [Watch Video Solution](#)

131. If $A = [2 - 242 - 51]$, $B = [804 - 236]$, find matrix X such that $2A + 3X = 5B$.

 [Watch Video Solution](#)

132. If $A = \begin{bmatrix} 1 & -3 & 2 \\ 2 & 0 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & -1 & -1 \\ 1 & 0 & -1 \end{bmatrix}$, find a matrix C

such that

$(A + B + C)$ is a zero matrix.

 [Watch Video Solution](#)

133. Find x, y satisfying the matrix equations:

$$[x - y \ 2 - 24x \ 6] + [3 - 22 \ 10 - 1] = [600 \ 52x + y \ 5] \quad (\text{ii})$$

$$[xy + 2z - 3] + [y \ 4 \ 5] = [49 \ 1 \ 2]$$

 [Watch Video Solution](#)

134. Find x, y satisfying the matrix equations:

$$x[2 \ 1] + y[3 \ 5] + [-8 - 11] = 0$$

 [Watch Video Solution](#)

135. If $2[345x] + [1y01] = [70105]$, find x and y .



Watch Video Solution

136. Find the value of λ , a non-zero scalar, if $\lambda[102345] + 2[123 - 1 - 32] = [44104214]$.



Watch Video Solution

137. Find a matrix X such that $2A + B + X = O$, where $A = [-1234]$,
 $B = [3 - 215]$



Watch Video Solution

138. If $A = [804 - 236]$ and $B = [2 - 242 - 51]$, then find the matrix X of order 3×2 such that $2A + 3X = 5B$.



Watch Video Solution

139. Find x , y , z and t , if $3[xyz] = [x6 - 12t] + [4x + yz + t3]$



Watch Video Solution

140. Find x , y , z and t , if $2[x57y - 3] + [3412] = [7141514]$



Watch Video Solution

141. If X and Y are 2×2 matrices, then solve the following matrix equations for X and Y . $2X + 3Y = [2340]$, $3X + 2Y = [-221 - 5]$



Watch Video Solution

142. In a certain city there are 30 colleges. Each college has 15 peons, 6 clerks, 1 typist and 1 section officer. Express the given information as a column matrix. Using scalar multiplication, find the total number of posts of each kind in all the colleges.

[Watch Video Solution](#)

143. The monthly incomes of Aryan and Babban are in the ratio 3 : 4 and their monthly expenditures are in the ratio 5 : 7. If each saves Rs. 15000 per month, find their monthly incomes using matrix method. This problem reflects which value?

[Watch Video Solution](#)

144. Let $A = \begin{bmatrix} 1 & -2 & 3 & 2 \\ -1 & 1 & -1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 3 & -1 & 2 \\ -1 & 2 & 4 & -5 \end{bmatrix}$. Find AB and BA and show that $AB \neq BA$.

[Watch Video Solution](#)

145. If A, B, C are matrices such that $A = \begin{bmatrix} x & y & z \end{bmatrix}$, $B = \begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix}$ and

$C = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ then find ABC

[Watch Video Solution](#)



Watch Video Solution

146. If $A = [1 \ -123]$, $B = [2110]$, prove that $(A + B)^2 \neq A^2 + 2AB + B^2$.



Watch Video Solution

147. If $A = [1 \ -12 \ -1]$, $B = [a1b \ -1]$ and $(A + B)^2 = A^2 + B^2$, find a and b .



Watch Video Solution

148. If $A = [01 \ -10]$, find x and y such that $(xI + yA)^2 = A$.



Watch Video Solution

149. If $A = [\alpha011]$ and $B = [1051]$, find the values of α for which $A^2 = B$.



Watch Video Solution

150. Let $A = \begin{bmatrix} 2 & -13 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 5 & 2 & 7 & 4 \end{bmatrix}$, $C = \begin{bmatrix} 2 & 5 & 3 & 8 \end{bmatrix}$. Find a matrix D such that $CD - AB = O$.



Watch Video Solution

151. Find the value of x such that $\begin{bmatrix} 1 & x & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 2 & 2 & 5 & 1 & 1 & 5 & 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 & x \end{bmatrix} = 0$



Watch Video Solution

152. If $A = \begin{bmatrix} 1 & 0 & -1 & 7 \end{bmatrix}$ and $I = \begin{bmatrix} 1 & 0 & 0 & 1 \end{bmatrix}$, then find k so that $A^2 = 8A + kI$



Watch Video Solution

153. If $\begin{bmatrix} 2 & -1 & 1 & 0 & -3 & 4 \end{bmatrix} A = \begin{bmatrix} -1 & -8 & -1 & 0 & 1 & -2 & -5 & 9 & 2 & 2 & 1 & 5 \end{bmatrix}$, find A .



Watch Video Solution

154. Let $f(x) = x^2 - 5x + 6$ then find $f(A)$ when $A = \begin{bmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{bmatrix}$

 [Watch Video Solution](#)

155. If $A = [31 \ -12]$, show that $A^2 - 5A + 7I_2 = O$.

 [Watch Video Solution](#)

156. Prove the following by principle of mathematical induction If

$$A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}, \text{ then } A^n = \begin{bmatrix} 1 + 2n & -4n \\ n & 1 - 2n \end{bmatrix} \text{ for every positive integer}$$

n

 [Watch Video Solution](#)

157. If $A_\alpha = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$, then prove that $A_\alpha A_\beta = A_{\alpha+\beta}$ for every positive integer n .

 [Watch Video Solution](#)

158. If $A_\alpha = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$, then prove that $(A_\alpha)^n = \begin{bmatrix} \cos n\alpha & \sin n\alpha \\ -\sin n\alpha & \cos n\alpha \end{bmatrix}$ for every positive integer n .

 [Watch Video Solution](#)

159. If a is a non-zero real or complex number. Use the principle of mathematical induction to prove that: If $A = \begin{bmatrix} a & 1 \\ 0 & a \end{bmatrix}$, then $A^n = \begin{bmatrix} a^n & n a^{n-1} \\ 0 & a^n \end{bmatrix}$ for every positive integer n .

 [Watch Video Solution](#)

160. If $A = [111111111]$, then prove that $A^n = [3^{n-1}3^{n-1}3^{n-1}3^{n-1}3^{n-1}3^{n-1}3^{n-1}3^{n-1}]$ for every positive integer n .

 [Watch Video Solution](#)

161. Under what conditions is the matrix equation $A^2 - B^2 = (A - B)(A + B)$ is true?

 [Watch Video Solution](#)

162. If A is any $m \times n$ such that AB and BA are both defined show that B is an $n \times m$ matrix.

 [Watch Video Solution](#)

163. A, B are two matrices such that AB and $A + B$ are both defined; show that A, B are square matrices of the same order.

 [Watch Video Solution](#)

164. If A and B are square matrices of order n , then prove that A and B will commute iff $A - \lambda I$ and $B - \lambda I$ commute for every scalar λ

 [Watch Video Solution](#)

165. If $AB = A$ and $BA = B$, then show that $A^2 = A, B^2 = B$.

 [Watch Video Solution](#)

166. Give an example of two matrices A and B such that
 $A \neq O, B \neq O, AB = O$ and $BA \neq O$ (ii)
 $A \neq O, B \neq O, AB = BA = O$.



[Watch Video Solution](#)

167. Give an example of three matrices A , B , C such that $AB = AC$ but $B \neq C$.



[Watch Video Solution](#)

168. There are two families A and B. In family A, there are 4 men, 6 women and 2 children; and in family B, there are 2 men, 2 women and 4 children. The recommended daily requirement of Calories is Men: 2400; Women: 1900; Children: 1800. Also daily requirement for protein is Men: 55 gm; Women: 45 gm and Children: 33 gm. Calculate the total requirement of calories and proteins for each of the two families. Use matrix method.



[Watch Video Solution](#)

169. Use matrix multiplication to divide Rs. 30,000 in two parts such that the total annual interest at 9% on the first part and 11% on the second

part amounts Rs. 3060



Watch Video Solution

170. Three schools A , B and C organised a mela for collecting funds for helping the rehabilitation of flood victims. They sold hand made fans, mats and plates from recycled material at a cost of Rs. 25, Rs. 100 and Rs. 50 each. The number of articles sold are given below.

School/Article	A	B	C
Hand-fans	40	25	35
Mats	50	40	50
Plates	20	30	40

Find the funds collected by each school separately by selling the above articles. Also, find the total funds collected for the purpose.



Watch Video Solution

171. Prove that the product of matrices $\begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta & \sin^2 \theta \\ \cos \theta \sin \theta & \cos^2 \theta & \sin^2 \theta \\ \sin^2 \theta & \cos \theta \sin \theta & \cos^2 \theta \end{bmatrix}$ and $\begin{bmatrix} \cos^2 \varphi & \cos \varphi \sin \varphi & \sin^2 \varphi \\ \cos \varphi \sin \varphi & \cos^2 \varphi & \sin^2 \varphi \\ \sin^2 \varphi & \cos \varphi \sin \varphi & \cos^2 \varphi \end{bmatrix}$ is the null matrix, when θ and φ differ by an odd multiple of $\frac{\pi}{2}$.



Watch Video Solution

 Watch Video Solution

172. Let $A = \begin{bmatrix} 0 & -\tan\left(\frac{\alpha}{2}\right) \\ \tan\left(\frac{\alpha}{2}\right) & 0 \end{bmatrix}$ and I be the identity matrix of order 2. Show that $I + A = (I - A)[\cos \alpha - \sin \alpha \sin \alpha \cos \alpha]$.

 Watch Video Solution

173. Let $F(x) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}$. Show that $F(x)F(y) = F(x + y)$

 Watch Video Solution

174. Let $A = \begin{bmatrix} 2 & 3 \\ -12 & 7 \end{bmatrix}$ and $f(x) = x^2 - 4x + 7$. Show that $f(A) = O$. Use this result to find A^5 .

 Watch Video Solution

175. If $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, prove that $(aI + bA)^n = a^n I + na^{n-1} bA$ where I is a unit matrix of order 2 and n is a positive integer.

 [Watch Video Solution](#)

176. Let A, B be two matrices such that they commute. Show that for any positive integer n , $AB^n = B^n A$

 [Watch Video Solution](#)

177. Let A, B be two matrices such that they commute. Show that for any positive integer n , $(AB)^n = A^n B^n$

 [Watch Video Solution](#)

178. If A is a square matrix such that $A^2 = A$, show that $(I + A)^3 = 7A + I$.



Watch Video Solution

179. If A is a square matrix such that $A^2 = I$, then find the simplified value of $(A - I)^3 + (A + I)^3 - 7A$.



Watch Video Solution

180. If $A = \begin{bmatrix} 3 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 7 & 3 \end{bmatrix}$, then find a non-zero matrix C such that $AC = BC$.



Watch Video Solution

181. Compute the indicated products: $[ab - ba][a - \mathbf{a}]$ (ii)

$[1 - 223][123 - 32 - 1]$ (iii) $[234345456][1 - 35024305]$



Watch Video Solution

182. Show that $AB \neq BA$ in each of the following cases: $A = [5 \ -167]$ and $B = [2134]$ (ii) $A = [-1100 \ -11234]$ and $B = [123010110]$ (iii) $A = [130110410]$ and $B = [010100051]$



[Watch Video Solution](#)

183. Compute the products AB and BA whichever exists in each of the following cases: $A = [1 \ -223]$ and $B = [123231]$ (ii) $A = [32 \ -10 \ -11]$ and $B = [456012]$



[Watch Video Solution](#)

184. Compute the products AB and BA whichever exists in each of the following cases: $A = [1 \ -1 \ 2 \ 3]$ and $B = [0132]$ (ii) $[a \ b][cd] + [a \ b \ c \ d][abcd]$



[Watch Video Solution](#)

185. Show that $AB \neq BA$ in each of the following cases:

$$A = [13 - 12 - 1 - 130 - 1] \text{ and } B = [-23 - 1 - 12 - 1 - 69 - 4]$$

(ii) $A = [10 - 4 - 1 - 11509 - 51]$ and $B = [121342132]$

 [Watch Video Solution](#)

186. Evaluate the following: $([13 - 1 - 4] + [3 - 2 - 11])[135246]$ (ii)

$[1 \ 2 \ 3][102201012][246]$ (iii) $[1 - 10223]([102201] - [012102])$

 [Watch Video Solution](#)

187. If $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ and $C = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, then show that

$$A^2 = B^2 = C^2 = I_2.$$

 [Watch Video Solution](#)

188. If $A = \begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 4 \\ -1 & 7 \end{bmatrix}$, find $3A^2 - 2B + I$.

 [Watch Video Solution](#)

189. If $A = \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$, prove that $(A - 2I)(A - 3I) = 0$.

 [Watch Video Solution](#)

190. If $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, show that $A^2 = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$ and $A^3 = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$.

 [Watch Video Solution](#)

191. If $A = \begin{bmatrix} ab & b^2 \\ -a^2 & -ab \end{bmatrix}$, show that $A^2 = 0$.

 [Watch Video Solution](#)

192. If $A = \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$, find A^2 .

 [Watch Video Solution](#)

193. If $A = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 3 & 5 \\ 1 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$, show that

$$AB = BA = O_3 \times 3.$$

 [Watch Video Solution](#)

194. If $A = \begin{bmatrix} 0 & c & -b \\ -c & 0 & a \\ b & -a & 0 \end{bmatrix}$ and $B = \begin{bmatrix} a^2 & ab & ac \\ ab & b^2 & bc \\ ac & bc & c^2 \end{bmatrix}$, show that

$$AB = BA = O_3 \times 3.$$

 [Watch Video Solution](#)

195. If $A = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$, show that

$$AB = A \text{ and } BA = B.$$

 [Watch Video Solution](#)

196. Let $A = \begin{bmatrix} -1 & 1 & -1 \\ 3 & -3 & 3 \\ 5 & 5 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 4 & 3 \\ 1 & -3 & -3 \\ -1 & 4 & 4 \end{bmatrix}$, compute $A^2 - B^2$.

 [Watch Video Solution](#)

197. For the following matrices verify the associativity of matrix multiplication i.e. $(AB)C = A(BC)$

(i) $A = \begin{bmatrix} 1 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ -1 & 2 \\ 0 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$

(ii) $A = \begin{bmatrix} 4 & 2 & 3 \\ 1 & 1 & 2 \\ 3 & 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 2 & -1 & 1 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$.

 [Watch Video Solution](#)

198. For the following matrices verify the distributivity of matrix multiplication over matrix addition i.e. $A(B + C) = AB + AC$

$$\cdot \quad (i) A = \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \quad (ii)$$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & 1 \\ -1 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}.$$

 [Watch Video Solution](#)

199. If $A = \begin{bmatrix} 1 & 0 & -2 \\ 3 & -1 & 0 \\ -2 & 1 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 5 & -4 \\ -2 & 1 & 3 \\ -1 & 0 & 2 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 5 & 2 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$, verify that $A(B - C) = AB - AC$

 [Watch Video Solution](#)

200. Compute the elements a_{43} and a_{22} of the matrix:

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 2 \\ 0 & 3 & 2 \\ 4 & 0 & 4 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ -3 & 2 \\ 4 & 3 \end{bmatrix} \begin{bmatrix} 0 & 1 & -1 & 2 & -2 \\ 3 & -3 & 4 & -4 & 0 \end{bmatrix}$$

 [Watch Video Solution](#)

201. If $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ p & q & r \end{bmatrix}$ and I is the identity matrix of order 3, show that $A^3 = pI + qA + rA^2$.

 [Watch Video Solution](#)

202. If ω is a complex cube root of unity, show that

$$\left(\begin{bmatrix} 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 \\ \omega^2 & 1 & \omega \end{bmatrix} + \begin{bmatrix} \omega & \omega^2 & 1 \\ \omega^2 & 1 & \omega \\ \omega & \omega^2 & 1 \end{bmatrix} \right) \begin{bmatrix} 1 \\ \omega \\ \omega^2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

 [Watch Video Solution](#)

203. If $A = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix}$, show that $A^2 = A$.

 [Watch Video Solution](#)

204. If $A = \begin{bmatrix} 4 & -1 & -4 \\ 3 & 0 & -4 \\ 3 & -1 & -3 \end{bmatrix}$, show that $A^2 = I_3$

 [Watch Video Solution](#)

205. If $[1 \ 1 \ x][102021210][111] = 0$, find x .

 [Watch Video Solution](#)

206. If $\begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix} \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} -4 & 6 \\ -9 & x \end{bmatrix}$, find x .

 [Watch Video Solution](#)

207. If $[x \ 4 \ 1] \begin{bmatrix} 2 & 1 & 2 \\ 1 & 0 & 2 \\ 0 & 2 & -4 \end{bmatrix} \begin{bmatrix} x \\ 4 \\ -1 \end{bmatrix} = 0$, find x .

 [Watch Video Solution](#)

208. If $\begin{bmatrix} 1 & -1 & x \end{bmatrix} \begin{bmatrix} 0 & 1 & -1 \\ 2 & 1 & 3 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = 0$, find x

 [Watch Video Solution](#)

209. If $A = \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix}$ and $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then prove that $A^2 - A + 2I = O$.

 [Watch Video Solution](#)

210. If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$ and $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then find λ so that $A^2 = 5A + \lambda I$

 [Watch Video Solution](#)

211. If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I_2 = O$.

 [Watch Video Solution](#)

212. If $A = \begin{bmatrix} 2 & 3 \\ -1 & 0 \end{bmatrix}$, show that $A^2 - 2A + 3I_2 = O$.

 [Watch Video Solution](#)

213. Show that the matrix $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ satisfies the equation $A^3 - 4A^2 + A = O$.

 [Watch Video Solution](#)

214. Show that the matrix $A = \begin{bmatrix} 5 & 3 \\ 12 & 7 \end{bmatrix}$ is a root of the equation $A^2 - 12A - I = 0$.

 [Watch Video Solution](#)

215. If $A = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$, find $A^2 - 5A - 14I$.

 [Watch Video Solution](#)

216. If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = O$. Use this to find A^4

 [Watch Video Solution](#)

217. If $A = \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix}$, find k such that $A^2 - kA - 2I_2 = 0$.

 [Watch Video Solution](#)

218. If $A = \begin{bmatrix} 1 & 0 \\ -1 & 7 \end{bmatrix}$, find k such that $A^2 - 8A + kI = O$.

 [Watch Video Solution](#)

219. If $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $f(x) = x^2 - 2x - 3$, show that $f(A) = O$.

 [Watch Video Solution](#)

220. If $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ and $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then find λ, μ so that $A^2 = \lambda A + \mu I$

 [Watch Video Solution](#)

221. Find the value of x for which the matrix product

$$\begin{bmatrix} 2 & 0 & 7 \\ 0 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix} \begin{bmatrix} -x & 14x & 7x \\ 0 & 1 & 0 \\ x & -4x & -2x \end{bmatrix} \text{ equal to an identity matrix.}$$

 [Watch Video Solution](#)

222. Solve the matrix equations: $\begin{bmatrix} 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 0 \\ 2 & 0 & 1 \\ 1 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 2 \\ x \end{bmatrix} = 0$

 [Watch Video Solution](#)

223. Solve the matrix equations: $\begin{bmatrix} x & -5 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix} \begin{bmatrix} x \\ 4 \\ 1 \end{bmatrix} = 0$

 [Watch Video Solution](#)

224. If $A = \begin{bmatrix} 1 & 2 & 0 \\ 3 & -4 & 5 \\ 0 & -1 & 3 \end{bmatrix}$, compute $A^2 - 4A + 3I_3$.



Watch Video Solution

225. If $f(x) = x^2 - 2x$, find $f(A)$, where $A = \begin{bmatrix} 0 & 1 & 2 \\ 4 & 5 & 0 \\ 0 & 2 & 3 \end{bmatrix}$.



Watch Video Solution

226. If $f(x) = x^3 + 4x^2 - x$, find $f(A)$, where $A = \begin{bmatrix} 0 & 1 & 2 \\ 2 & -3 & 0 \\ 1 & -1 & 0 \end{bmatrix}$.



Watch Video Solution

227. If $A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix}$, then show that A is a root of the polynomial

$$f(x) = x^3 - 6x^2 + 7x + 2.$$

 [Watch Video Solution](#)

228. If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$, then prove that $A^2 - 4A - 5I = O$.

 [Watch Video Solution](#)

229. If $A = \begin{bmatrix} 3 & 2 & 0 \\ 1 & 4 & 0 \\ 0 & 0 & 5 \end{bmatrix}$, show that $A^2 - 7A + 10I_3 = O$.

 [Watch Video Solution](#)

230. Without using the concept of inverse of a matrix, find the matrix

$$[xyzu] \text{ such that } \begin{bmatrix} 5 & -7 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} x & y \\ z & u \end{bmatrix} = \begin{bmatrix} -16 & -6 \\ 7 & 2 \end{bmatrix}.$$

 [Watch Video Solution](#)

231. Find the matrix A such that : $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} A = \begin{bmatrix} 3 & 3 & 5 \\ 1 & 0 & 1 \end{bmatrix}$

 [Watch Video Solution](#)

232. Find the matrix A such that : $[2, 1, 3] \begin{bmatrix} -1 & 0 & -1 \\ -1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = A$

 [Watch Video Solution](#)

233. Find a 2×2 matrix A such that $A \begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix} = 6I_2$.

 [Watch Video Solution](#)

234. If $A = \begin{bmatrix} 0 & 0 \\ 4 & 0 \end{bmatrix}$, find A^{16} .

 [Watch Video Solution](#)

235. If $A = \begin{bmatrix} 0 & -x \\ x & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ and $x^2 = -1$, then show that $(A + B)^2 = A^2 + B^2$.

 [Watch Video Solution](#)

236. If $A = \begin{bmatrix} 1 & 0 & -3 \\ 2 & 1 & 3 \\ 0 & 1 & 1 \end{bmatrix}$, then verify that $A^2 + A = A(A + I)$, where I is the identity matrix.

 [Watch Video Solution](#)

237. If $A = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$, then find $A^2 - 5A - 14I$.

 [Watch Video Solution](#)

238. If $P(x) = [\cos x \sin x - \sin x \cos x]$, then show that $P(x) P(y) = P(x + y) = P(y) P(x)$.

 [Watch Video Solution](#)

239. If $P = \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix}$ and $Q = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$, prove that $PQ = \begin{bmatrix} xa & 0 & 0 \\ 0 & yb & 0 \\ 0 & 0 & zc \end{bmatrix} = QP$

 [Watch Video Solution](#)

240. If $A = \begin{bmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{bmatrix}$, find $A^2 - 5A + 4I$ and hence find a matrix X such that $A^2 - 5A + 4I + X = O$.

 [Watch Video Solution](#)

241. If $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, prove that $A^n = \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}$ for all positive integers n .

 [Watch Video Solution](#)

242. If $A = \begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix}$, prove that $A^n = \begin{bmatrix} a^n & b\left(\frac{a^n-1}{a-1}\right) \\ 0 & 1 \end{bmatrix}$ for every positive integer n .

 [Watch Video Solution](#)

243. If $A = \begin{bmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{bmatrix}$, then prove by principle of mathematical induction that $A^n = \begin{bmatrix} \cos n \theta & i \sin n \theta \\ i \sin n \theta & \cos n \theta \end{bmatrix}$ for all $n \in N$.

 [Watch Video Solution](#)

244. If $A = \begin{bmatrix} \cos \alpha + \sin \alpha & \sqrt{2} \sin \alpha \\ -\sqrt{2} \sin \alpha & \cos \alpha - \sin \alpha \end{bmatrix}$, prove that $A^n = \begin{bmatrix} \cos n \alpha + \sin n \alpha & \sqrt{2} \sin n \alpha \\ -\sqrt{2} \sin n \alpha & \cos n \alpha - \sin n \alpha \end{bmatrix}$ for all $n \in N$.



Watch Video Solution

245. If $A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$, then use the principle of mathematical induction to show that $A^n = \begin{bmatrix} 1 & n & \frac{n(n+1)}{2} \\ 0 & 1 & n \\ 0 & 0 & 1 \end{bmatrix}$ for every positive integer n .



Watch Video Solution

246. If B, C are n rowed square matrices and if $A = B + C, BC = CB, C^2 = O$, then show that for every $n \in N, A^{n+1} = B^n(B + (n + 1)C)$.



Watch Video Solution

247. If $A = \text{diag}(a \ b \ c)$, show that $A^n = \text{diag}(a^n \ b^n \ c^n)$ for all positive integer n .



Watch Video Solution

248. If A is a square matrix, using mathematical induction prove that

$$(A^T)^n = (A^n)^T \text{ for all } n \in N.$$



Watch Video Solution

249. A matrix X has $a + b$ rows and $a + 2$ columns while the matrix Y has $b + 1$ rows and $a + 3$ columns. Both matrices XY and YX exist. Find a and b . Can you say XY and YX are of the same type? Are they equal.



Watch Video Solution

250. Give examples of matrices A and B such that $AB \neq BA$. (ii) A and B such that $AB = O$ but $A \neq O$, $B \neq O$



Watch Video Solution

251. Give examples of matrices A and B such that $AB = O$ but $BA \neq O$

.(ii) A , B and C such that $AB = AC$ but $B \neq C$, $A \neq O$.

 [Watch Video Solution](#)

252. Let A and B be square matrices of the same order. Does

$(A + B)^2 = A^2 + 2AB + B^2$ hold? If not, why?

 [Watch Video Solution](#)

253. If A and B are square matrices of the same order, explain, why in

general $(A + B)^2 \neq A^2 + 2AB + B^2$ (ii) $(A - B)^2 \neq A^2 - 2AB + B^2$

(iii) $(A + B)(A - B) \neq A^2 - B^2$.

 [Watch Video Solution](#)

254. Let A and B be square matrices of the order 3×3 . Is

$(AB)^2 = A^2B^2$? Give reasons.



[Watch Video Solution](#)

255. If A and B are square matrices of the same order such that $AB = BA$, then show that $(A + B)^2 = A^2 + 2AB + B^2$.



[Watch Video Solution](#)

256. Let $A = [111333]$, $B = [3152 - 24]$ and $C = [42 - 3550]$. Verify that $AB = AC$ though $B \neq C$, $A \neq O$.



[Watch Video Solution](#)

257. Three shopkeepers A , B and C go to a store to buy stationary. A purchases 12 dozen notebooks, 5 dozen pens and 6 dozen pencils. B purchases 10 dozen notebooks, 6 dozen pens and 7 dozen pencils. C purchases 11 dozen notebooks, 13 dozen pens and 8 dozen pencils. A notebook costs 40 paise, a pen costs Rs. 1.25 and a pencil costs 35 paise. Use matrix multiplication to calculate each individuals bill.



[Watch Video Solution](#)

258. The cooperative stores of a particular school has 10 dozen physics books, 8 dozen chemistry books and 5 dozen mathematics books. Their selling prices are Rs. 8.30, Rs. 3.45 and Rs. 4.50 each respectively. Find the total amount the store will receive from selling all the items.



[Watch Video Solution](#)

259. In a legislative assembly election, a political group hired a public relations firm to promote its candidate in three ways: telephone, house calls, and letters. The cost per contact (in paise) is given in matrix A as

Cost per contact



[Watch Video Solution](#)

260. A trust fund has Rs. 30000 that must be invested in two different types of bonds. The first bond pays 5% interest per year, and the second

bond pays 7% interest per year. Using matrix multiplication, determine how to divide Rs 30000 among the two types of bonds. If the trust fund must obtain an annual total interest of (i) Rs 1800 (ii) 2000.



[Watch Video Solution](#)

261. To promote making of toilets for women, an organisation tried to generate awareness attempt (i) house calls (ii) letters, and (iii) announcements. The cost for each mode per attempt is given below: (i) Rs 50 (ii) Rs 20 (iii) Rs 40 The number of attempts made in three villages X , Y and Z are given below: (i) (ii) (iii) X 400 300 100 Y 300 250 75 Z 500 400 150 Find the total cost incurred by the organisation for three villages separately, using matrices.



[Watch Video Solution](#)

262. There are two families A and B. In family A, there are 4 men, 6 women and 2 children : and in family B, there are 2 men, 2 women and 4 children. The recommended daily requirement of Calories is Men: 2400 ; Women:

1900 ; Children: 1800 Also daily requirement for protein is Men: 55 gm ; Women:45 gm and Children:33 gm Calculate the total requirement of calories and proteins for each of the two families..tsin matrix method.



[Watch Video Solution](#)

263. In a parliament election, a political party hired a public relations firm to promote its candidates in three ways telephone, house calls and letters. The cost per contact (in paisa) is given in matrix A as $A = [140200150]$ Telephone House calls Leers The number of contacts of each type made in two cities X and Y is given in the matrix B as

Telephone	House	call	Letters
$B = [1000$	500	50003000	1000
$10000]$			$10000]$

City X City Y

Find the total amount spent by the party in the two cities. What should one consider before casting his/her vote partys promotional activity or their social activities?



[Watch Video Solution](#)

264. The monthly incomes of Aryan and Babban are in the ratio 3 : 4 and their monthly expenditures are in the ratio 5 : 7. If each saves Rs. 15000 per month, find their monthly incomes using matrix method. This problem reflects which value?

 [Watch Video Solution](#)

265. A trust invested some money in two types of bonds. The first bond pays 10% interest and second bond pays 12% interest. The trust received Rs. 2,800 as interest. However, if trust had interchanged money in bonds, they would have got Rs. 100 less as interest. Using matrix method, find the amount invested by the trust.

 [Watch Video Solution](#)

266. If $A = \begin{bmatrix} - & 123 \end{bmatrix}$ and $B = \begin{bmatrix} -2 & -1 & -4 \end{bmatrix}$, verify that $(AB)^T = B^T A^T$

 [Watch Video Solution](#)

267. If $A = [\cos \theta \quad -\sin \theta \quad \sin \theta \quad \cos \theta]$, then find the values of θ satisfying the equation $A^T + A = I_2$.

 [Watch Video Solution](#)

268. If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ a & 2 & b \end{bmatrix}$ is a matrix satisfying $AA^T = 9I_3$, then find the values of a and b .

 [Watch Video Solution](#)

269. Find the values of x, y, z if the matrix $A = \begin{bmatrix} 0 & 2y & z \\ x & y & -z \\ x & -y & z \end{bmatrix}$ satisfy the equation $A^T A = I_3$

 [Watch Video Solution](#)

270. Let $A = [2 \ -3 \ -75]$ and $B = [102 \ -4]$, verify that $(2A)^T = 2A^T$

(ii) $(A + B)^T = A^T + B^T$ (iii) $(A - B)^T = A^T - B^T$ (iv)

$(AB)^T = B^T A^T$



[Watch Video Solution](#)

271. If $A = [352]$ and $B = [1 \ 0 \ 4]$, verify that $(AB)^T = B^T A^T$.



[Watch Video Solution](#)

272. Let $A = [1 \ -10213121]$ and $B = [123213011]$. Find A^T , B^T and

verify that: $(A + B)^T = A^T + B^T$ (ii) $(AB)^T = B^T A^T$ (iii)

$(2A)^T = 2A^T$



[Watch Video Solution](#)

273. If $A = [-245]$, $B = [1 \ 3 \ -6]$, verify that $(AB)^T = B^T A^T$



[Watch Video Solution](#)

274. If $A = \begin{bmatrix} 2 & 4 & -1 \\ -1 & 0 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 3 & 4 \\ -1 & 2 \\ 2 & 1 \end{bmatrix}$, find $(AB)^T$.

[Watch Video Solution](#)

275. For the matrices A and B , $A = [213410]$, $B = [1 - 10250]$ verify that $(AB)^T = B^T A^T$

[Watch Video Solution](#)

276. For two matrices A and B , verify that $(AB)^T = B^T A^T$, where $A = [1324]$, $B = [1425]$

[Watch Video Solution](#)

277. If $A^T = \begin{bmatrix} 3 & 4 \\ -1 & 2 \\ 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 2 & 1 \\ 1 & 2 & 3 \end{bmatrix}$, find $A^T - B^T$.

 [Watch Video Solution](#)

278. If $A = [\cos \alpha \sin \alpha \quad -\sin \alpha \cos \alpha]$, then verify that $A^T A = I_2$.

 [Watch Video Solution](#)

279. If $A = \begin{bmatrix} \sin \alpha & \cos \alpha \\ -\cos \alpha & \sin \alpha \end{bmatrix}$, verify that $A^T A = I_2$

 [Watch Video Solution](#)

280. Let $l_i, m_i, n_i; i = 1, 2, 3$ be the direction cosines of three mutually perpendicular vectors in space. Show that $\forall' = I_3$ where

$$A = \begin{bmatrix} l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \\ l_3 & m_3 & n_3 \end{bmatrix}$$

 [Watch Video Solution](#)

281. Show that the elements on the main diagonal of a skew-symmetric matrix are all zero.



[Watch Video Solution](#)

282. If the matrix $A = \begin{bmatrix} 0 & a & 3 & 2b - 1 \\ c & 1 & 0 & 10 \end{bmatrix}$ is skew-symmetric, find the values of a , b and c .



[Watch Video Solution](#)

283. Let A be a square matrix. Then prove that (i) $A + A^T$ is a symmetric matrix, (ii) $A - A^T$ is a skew-symmetric matrix and (iii) $A^T A$ and $A A^T$ are symmetric matrices.



[Watch Video Solution](#)

284. Prove that every square matrix can be uniquely expressed as the sum of a symmetric matrix and a skew-symmetric matrix.

 [Watch Video Solution](#)

285. If A and B are symmetric matrices, then show that AB is symmetric iff $AB = BA$ i.e. A and B commute.

 [Watch Video Solution](#)

286. Show that the matrix $B^T AB$ is symmetric or skew-symmetric according as A is symmetric or skew-symmetric.

 [Watch Video Solution](#)

287. Let A and B be symmetric matrices of same order. Then $A + B$ is a symmetric matrix, $AB - BA$ is a skew symmetric matrix and $AB + BA$

is a symmetric matrix

 [Watch Video Solution](#)

288. Express the matrix $A = \begin{bmatrix} 3 & 2 & 3 \\ 4 & 5 & 3 \\ 2 & 4 & 5 \end{bmatrix}$ as the sum of a symmetric and a skew-symmetric matrix.

 [Watch Video Solution](#)

289. Show that all positive integral powers of a symmetric matrix are symmetric.

 [Watch Video Solution](#)

290. Show that positive odd integral powers of a skew-symmetric matrix are skew-symmetric and positive even integral powers of a skew-symmetric matrix are symmetric.



[Watch Video Solution](#)

 [Watch Video Solution](#)

291. A matrix which is both symmetric as well as skew-symmetric is a null matrix.

 [Watch Video Solution](#)

292. If $A = [2345]$, prove that $A - A^T$ is a skew-symmetric matrix.

 [Watch Video Solution](#)

293. If $A = [3 \quad -4 \quad 1]$, show that $A - A^T$ is a skew symmetric matrix.

 [Watch Video Solution](#)

294. If the matrix $A = [52xyz - 34t - 7]$ is a symmetric matrix, find x , y , z and t .

 [Watch Video Solution](#)

 Watch Video Solution

295. Let $A = \begin{bmatrix} 3 & 2 & 7 & 1 & 4 & 3 \\ & & & & & -2 & 5 & 8 \end{bmatrix}$. Find matrices X and Y such that $X + Y = A$, where X is a symmetric and Y is a skew-symmetric matrix.

 Watch Video Solution

296. Express the matrix $A = \begin{bmatrix} 4 & 2 & -1 & 3 & 5 & 7 & 1 & -2 & 1 \end{bmatrix}$ as the sum of a symmetric and a skew-symmetric matrix.

 Watch Video Solution

297. Define a symmetric matrix. Prove that for $A = \begin{bmatrix} 2 & 4 & 5 & 6 \end{bmatrix}$, $A + A^T$ is a symmetric matrix where A^T is the transpose of A .

 Watch Video Solution

298. Express the matrix $A = [3 \ -4 \ 1 \ -1]$ as the sum of a symmetric and a skew-symmetric matrix.

 [Watch Video Solution](#)

299. Express the matrix $[3 \ -2 \ -4 \ 3 \ -2 \ -5 \ -1 \ 1 \ 2]$ as the sum of a symmetric and skew-symmetric matrix and verify your result.

 [Watch Video Solution](#)

300. If A is an $m \times n$ matrix and B is $n \times p$ matrix does AB exist? If yes, write its order.

 [Watch Video Solution](#)

301. If $A = [2 \ 1 \ 4 \ 4 \ 1 \ 5]$ and $B = [3 \ -1 \ 2 \ 2 \ 1 \ 3]$. Write the orders of AB and BA .

 [Watch Video Solution](#)

302. If $A = [4312]$ and $B = [-43]$, write AB .

 [Watch Video Solution](#)

303. If $A = [123]$, write AA^T .

 [Watch Video Solution](#)

304. Give an example of two non-zero 2×2 matrices A and B such that $AB = O$.

 [Watch Video Solution](#)

305. If $A = [2357]$, find $A + A^T$.

 [Watch Video Solution](#)

306. If $A = [i00i]$, write A^2 .

 [Watch Video Solution](#)

307. If $A = \begin{bmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{bmatrix}$, find x satisfying $0 < x < \frac{\pi}{2}$ when $A + A^T = I$

 [Watch Video Solution](#)

308. If $A = \begin{bmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{bmatrix}$, find AA^T

 [Watch Video Solution](#)

309. If $\begin{bmatrix} 1 & 0 \\ y & 5 \end{bmatrix} + 2 \begin{bmatrix} x & 0 \\ 1 & -2 \end{bmatrix} = I$, where I is 2×2 unit matrix. Find x and y .

 [Watch Video Solution](#)

310. If matrix $A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$ and $A^2 = kA$, then write the value of k

 [Watch Video Solution](#)

311. If $A = [1111]$ satisfies $A^4 = \lambda A$, then write the value of λ

 [Watch Video Solution](#)

312. If $A = [- 1000 - 1000 - 1]$, find A^2 .

 [Watch Video Solution](#)

313. If $A = [- 1000 - 1000 - 1]$, find A^3 .

 [Watch Video Solution](#)

314. If $A = \begin{bmatrix} -3 & 0 & -3 \end{bmatrix}$, find A^4 .



Watch Video Solution

315. If $\begin{bmatrix} x & 2 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} = 2$, find x



Watch Video Solution

316. If $A = [a_{ij}]$ is a 2×2 matrix such that $a_{ij} = i + 2j$, write A .



Watch Video Solution

317. Find a matrix A , if $A + \begin{bmatrix} 2 & 3 & -14 \end{bmatrix} = \begin{bmatrix} 3 & -6 & -38 \end{bmatrix}$



Watch Video Solution

318. If $A = [a_{ij}]$ is a square matrix such that $a_{ij} = i^2 - j^2$, then write whether A is symmetric or skew-symmetric.

 [Watch Video Solution](#)

319. For any square matrix write whether AA^T is symmetric or skew-symmetric.

 [Watch Video Solution](#)

320. If $A = [a_{ij}]$ is a skew-symmetric matrix, then write the value of

$$\sum_i a_{ii}.$$

 [Watch Video Solution](#)

321. If $A = [a_{ij}]$ is a skew-symmetric matrix, then write the value of

$$\sum_i \sum_j a_{ij}.$$



Watch Video Solution

322. If A and B are symmetric matrices, then write the condition for which AB is also symmetric.



Watch Video Solution

323. If B is a skew-symmetric matrix, write whether the matrix $AB A^T$ is symmetric or skew-symmetric.



Watch Video Solution

324. If B is a symmetric matrix, write whether the matrix $AB A^T$ is symmetric or skew-symmetric.



Watch Video Solution

325. If A is a skew-symmetric and $n \in \mathbb{N}$ such that $(A^n)^T = \lambda A^n$, write the value of λ

 [Watch Video Solution](#)

326. If A is a symmetric matrix and $n \in \mathbb{N}$, write whether A^n is symmetric or skew-symmetric or neither of these two.

 [Watch Video Solution](#)

327. If A is a skew-symmetric matrix and n is an even natural number, write whether A^n is symmetric or skew-symmetric matrix or neither of the two

 [Watch Video Solution](#)

328. If A and B are symmetric matrices of the same order, write whether $AB - BA$ is symmetric or skew-symmetric or neither of the two.

 [Watch Video Solution](#)

329. Write a square matrix which is both symmetric as well as skew-symmetric.

 [Watch Video Solution](#)

330. Find the values of x and y , if $2 \begin{bmatrix} 1 & 3 \\ 0 & x \end{bmatrix} + \begin{bmatrix} y & 0 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$.

 [Watch Video Solution](#)

331. If $\begin{bmatrix} x + 3 & 4 \\ y & -4x + y \end{bmatrix} = \begin{bmatrix} 5 & 4 \\ 3 & 9 \end{bmatrix}$, find x and y

 [Watch Video Solution](#)

332. Find the value of x and y from the following:

$$\begin{bmatrix} 2x & -y \\ 5 & 3y \end{bmatrix} = \begin{bmatrix} 6 & 5 \\ 3 & -2 \end{bmatrix}$$

 [Watch Video Solution](#)

333. Find the value of y , if $\begin{bmatrix} x - y & 2 \\ x & 5 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 3 & 5 \end{bmatrix}$.

 [Watch Video Solution](#)

334. Find the value of x , if $\begin{bmatrix} 3x + y & -y \\ 2y - x & 3 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -5 & 3 \end{bmatrix}$.

 [Watch Video Solution](#)

335. If matrix $A = [1, 2, 3]$, find AA^T .

 [Watch Video Solution](#)

336. If $\begin{bmatrix} 2x + y & 3y \\ 0 & 4 \end{bmatrix} = \begin{bmatrix} 6 & 0 \\ 6 & 4 \end{bmatrix}$, then find x .

 [Watch Video Solution](#)

337. If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, find $A + A^T$.

 [Watch Video Solution](#)

338. If $\begin{bmatrix} a + b & 2 \\ 5 & b \end{bmatrix} = \begin{bmatrix} 6 & 5 \\ 2 & 2 \end{bmatrix}$, then find a .

 [Watch Video Solution](#)

339. If A is a matrix of order 3×4 and B is a matrix of order 4×3 , find the order of the matrix of AB .

 [Watch Video Solution](#)

340. If $A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$ is identity matrix, then write the value of α .

 [Watch Video Solution](#)

341. If $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 2 & 5 \end{bmatrix} = \begin{bmatrix} 7 & 11 \\ 23 & k \end{bmatrix}$, then write the value of k .

 [Watch Video Solution](#)

342. If I is the identity matrix and A is a square matrix such that $A^2 = A$, then what is the value of $(I + A)^2 - 3A$?

 [Watch Video Solution](#)

343. If $A = \begin{bmatrix} 1 & 2 \\ 0 & 3 \end{bmatrix}$ is written as $B + C$, where B is a symmetric matrix and C is a skew-symmetric matrix, then find B .

 [Watch Video Solution](#)

344. If A is 2×3 matrix and B is a matrix such that $A^T B$ and BA^T both are defined, then what is the order of B ?

 [Watch Video Solution](#)

345. What is the total number of 2×2 matrices with each entry 0 or 1?

 [Watch Video Solution](#)

346. If $\begin{bmatrix} x & x - y \\ 2x + y & 7 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 8 & 7 \end{bmatrix}$, then find the value of y

 [Watch Video Solution](#)

347. If a matrix has 5 elements, write all possible orders it can have.

 [Watch Video Solution](#)

348. For a 2×2 matrix $A = [a_{ij}]$ whose elements are given by $a_{ij} = \frac{i}{j}$, write the value of a_{12} .

 [Watch Video Solution](#)

349. If $x \begin{bmatrix} 2 \\ 3 \end{bmatrix} + y \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 10 \\ 5 \end{bmatrix}$, find the value of x

 [Watch Video Solution](#)

350. If $\begin{bmatrix} 9 & -1 & 4 \\ -2 & 1 & 3 \end{bmatrix} = A + \begin{bmatrix} 1 & 2 & -1 \\ 0 & 4 & 9 \end{bmatrix}$, then find matrix A

 [Watch Video Solution](#)

351. If $\begin{bmatrix} a - b & 2a + c \\ 2a - b & 3c + d \end{bmatrix} = \begin{bmatrix} -1 & 5 \\ 0 & 13 \end{bmatrix}$, find the value of b

 [Watch Video Solution](#)

352. For what value of x is the matrix $A = \begin{bmatrix} 0 & 1 & -2 \\ 1 & 0 & 3 \\ x & 3 & 0 \end{bmatrix}$ a skew-symmetric matrix?

 [Watch Video Solution](#)

353. If matrix $A = \begin{bmatrix} 2 & -2 \\ -2 & 2 \end{bmatrix}$ and $A^2 = pA$, then write the value of p

 [Watch Video Solution](#)

354. If A is a square matrix such that $A^2 = A$, then write the value of $7A - (I + A)^3$, where I is the identity matrix.

 [Watch Video Solution](#)

355. If $2[3 \ 4 \ 5x] + [1y \ 0 \ 1] = [70 \ 10 \ 5]$, find $x - y$.

 [Watch Video Solution](#)

356. If $[x \ 1] \begin{bmatrix} 1 & 0 \\ -2 & 0 \end{bmatrix} = O$, find x

 [Watch Video Solution](#)

357. If $[a + 43b8 - 6] = [2a + 2b + 28a - 8b]$, write the value of $a - 2b$.

 [Watch Video Solution](#)

358. Write a 2×2 matrix which is both symmetric and skew-symmetric.

 [Watch Video Solution](#)

359. If $\begin{bmatrix} xy & 4 \\ z + 6 & x + y \end{bmatrix} = \begin{bmatrix} 8 & w \\ 0 & 6 \end{bmatrix}$, write the value of $(x + y + z)$.

 [Watch Video Solution](#)

360. Construct a 2×2 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by

$$a_{ij} = \begin{cases} \frac{|-3i + j|}{2}, & \text{if } i \neq j \\ j(i + j)^2, & \text{if } i = j \end{cases}$$

 [Watch Video Solution](#)

361. If $[x + yx - y] = [2143][1 - 2]$, then write the value of (x, y) .

 [Watch Video Solution](#)

362. Matrix $A = [02b - 23133a3 - 1]$ is given to be symmetric, find the values of a and b .

 [Watch Video Solution](#)

363. Write the number of all possible matrices of order 2×2 with each entry 1, 2 or 3.

 [Watch Video Solution](#)

364. If $[2 \ 1 \ 3][-10 \ -1 \ -110011][10 \ -1] = A$, then write the order of matrix A .

 [Watch Video Solution](#)

365. If $A = [3579]$ is written as $A = P + Q$, where P is symmetric and Q is skew-symmetric matrix, then write the matrix P .

 [Watch Video Solution](#)

366. If $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & -1 \end{bmatrix}$, then A^2 is equal to (a) a null matrix (b) a unit matrix (c) A (d) A

 [Watch Video Solution](#)

367. If $A = [i00i]$, $n \in N$, then A^{4n} equals $[0ii0]$ (b) $[0000]$ (c) $[1001]$ (d) $[0ii0]$



Watch Video Solution

368. If A and B are two matrices such that $AB = A$ and $BA = B$, then B^2 is equal to (a) B (b) A (c) 1 (d) 0



Watch Video Solution

369. If $AB = A$ and $BA = B$, where A and B are square matrices, then $B^2 = B$ and $A^2 = A$ (b) $B^2 \neq B$ and $A^2 = A$ (c) $A^2 \neq A$, $B^2 = B$ (d) $A^2 \neq A$, $B^2 \neq B$



Watch Video Solution

370. If A and B are two matrices such that $AB=B$ and $BA=A$, then

$$A^2 + B^2 =$$

 [Watch Video Solution](#)

371. If $\begin{bmatrix} \cos \frac{2\pi}{7} & -\sin \frac{2\pi}{7} \\ \sin \frac{2\pi}{7} & \cos \frac{2\pi}{7} \end{bmatrix}^k = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then the least positive integral value of k , is

 [Watch Video Solution](#)

372. If the matrix AB is zero, then It is not necessary that either $A = O$ or, $B = O$ (b) $A = O$ or $B = O$ (c) $A = O$ and $B = 0$ (d) all the above statements are wrong

 [Watch Video Solution](#)

373. Let $A = \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix}$, then A^n is equal to $\begin{bmatrix} a^n & 0 & 0 \\ 0 & a^n & 0 \\ 0 & 0 & a \end{bmatrix}$ (b)

$\begin{bmatrix} a^n & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix}$ (c) $\begin{bmatrix} a^n & 0 & 0 \\ 0 & a^n & 0 \\ 0 & 0 & a^n \end{bmatrix}$ (d) $\begin{bmatrix} na & 0 & 0 \\ 0 & na & 0 \\ 0 & 0 & na \end{bmatrix}$

 [Watch Video Solution](#)

374. If A, B are square matrices of order 3, A is non-singular and $AB = O$, then B is a (a) null matrix (b) singular matrix (c) unit matrix (d) non-singular matrix

 [Watch Video Solution](#)

375. If $A = \begin{bmatrix} n & 0 & 0 \\ 0 & n & 0 \\ 0 & 0 & n \end{bmatrix}$ and $B = \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix}$, then AB is equal to

(A) B (B) nB (C) B^n (D) $A + B$

 [Watch Video Solution](#)

376. If $A = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix}$, then A^n (where $n \in N$) equals $\begin{bmatrix} 1 & na \\ 0 & 1 \end{bmatrix}$ (b)

$\begin{bmatrix} 1 & n^2a \\ 0 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & na \\ 0 & 0 \end{bmatrix}$ (d) $\begin{bmatrix} n & na \\ 0 & n \end{bmatrix}$

 [Watch Video Solution](#)

377. If $A = [12x010001]$ and $B = [1 - 2y010001]$ and $AB = I_3$, then $x + y$ equals (a) 0 (b) -1 (c) 2 (d) none of these

 [Watch Video Solution](#)

378. If $A = [1 - 12 - 1]$, $B = [a1b - 1]$ and $(A + B)^2 = A^2 + B^2$, then values of a and b are $a = 4, b = 1$ (b) $a = 1, b = 4$ (c) $a = 0, b = 4$ (d) $a = 2, b = 4$

 [Watch Video Solution](#)

379. If $A = [\alpha\beta\gamma - \alpha]$ is such that $A^2 = I$, then $1 + \alpha^2 + \beta\gamma = 0$ (b) $1 - \alpha^2 + \beta\gamma = 0$ (c) $1 - \alpha^2 - \beta\gamma = 0$ (d) $1 + \alpha^2 - \beta\gamma = 0$



Watch Video Solution

380. If $S = [S_{ij}]$ is a scalar matrix such that $s_{ii} = k$ and A is a square matrix of the same order, then $AS = SA = ?$ (a) A^k (b) $k + A$ (c) kA (d) kS



Watch Video Solution

381. If a square matrix such that $A^2 = A$, then $(I + A)^3 - 7A$ is equal to (a) A (b) $I - A$ (c) I (d) $3A$



Watch Video Solution

382. If a matrix A is both symmetric and skew-symmetric, then A is a

A. Diagonal matrix

B. Zero matrix

C. Scalar matrix

D. Square matrix

Answer: B



[Watch Video Solution](#)

383. The matrix $\begin{bmatrix} 0 & 5 & -7 \\ -5 & 0 & 11 \\ 7 & -11 & 0 \end{bmatrix}$ is (a) a skew-symmetric matrix (b) a symmetric matrix (c) a diagonal matrix (d) an upper triangular matrix



[Watch Video Solution](#)

384. If A is a square matrix, then AA is a (a) skew-symmetric matrix (b) symmetric matrix (c) diagonal matrix (d) none of these



[Watch Video Solution](#)

385. If A and B are symmetric matrices, then ABA is (a) symmetric matrix (b) skew-symmetric matrix (c) diagonal matrix (d) scalar matrix



Watch Video Solution

386. If $A = \begin{bmatrix} 5 & x \\ y & 0 \end{bmatrix}$ and $A = A^T$, then

A. $x = 0, y = 5$

B. $x + y = 5$

C. $x = y$

D. none of these

Answer: C



Watch Video Solution

387. If A is 3×4 matrix and B is a matrix such that $A^T B$ and BA^T are both defined. Then, B is of the type 3×4 (b) 3×3 (c) 4×4 (d) 4×3

 [Watch Video Solution](#)

388. If $A = [a_{ij}]$ is a square matrix of even order such that $a_{ij} = i^2 - j^2$, then A is a skew-symmetric matrix and $|A| = 0$ (b) A is symmetric matrix and $|A|$ is a square (c) A is symmetric matrix and $|A| = 0$ (d) none of these

 [Watch Video Solution](#)

389. If $A = [\cos \theta \quad -\sin \theta \quad \sin \theta \quad \cos \theta]$, then $A^T + A = I_2$, if $\theta = n\pi$, $n \in Z$ (b) $\theta = (2n + 1)\frac{\pi}{2}$, $n \in Z$ (c) $\theta = 2n\pi + \frac{\pi}{3}$, $n \in Z$ (d) none of these

 [Watch Video Solution](#)

390. If $A = \begin{bmatrix} 2 & 0 & -3 \\ 4 & 3 & 1 \\ -5 & 7 & 2 \end{bmatrix}$ is expressed as the sum of a symmetric and

skew-symmetric matrix, then the symmetric matrix is (a) $\begin{bmatrix} 2 & 2 & -4 \\ 2 & 3 & 4 \\ -4 & 4 & 2 \end{bmatrix}$

(b) $\begin{bmatrix} 2 & 4 & -5 \\ 0 & 3 & 7 \\ -3 & 1 & 2 \end{bmatrix}$ (c) $\begin{bmatrix} 4 & 4 & -8 \\ 4 & 6 & 8 \\ -8 & 8 & 4 \end{bmatrix}$ (d) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

 [Watch Video Solution](#)

391. Out of the following matrices, choose that matrix which is a scalar

matrix: (a) $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ (c) $\begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ (d) $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$

 [Watch Video Solution](#)

392. The number of all possible matrices of order 3×3 with each entry 0 or 1 is (a) 27 (b) 18 (c) 81 (d) 512

 [Watch Video Solution](#)

393. Which of the given values of x and y make the following pairs of

matrices equal? $\begin{bmatrix} 3x + 7 & 5y + 1 \\ 2 & -3x \end{bmatrix}$ and, $\begin{bmatrix} 0 & y - 2 \\ 8 & 4 \end{bmatrix}$ a)
 $x = -\frac{1}{3}, y = 7$ (b) $y = 7, x = -\frac{2}{3}$ (c) $x = -\frac{1}{3}, y = -\frac{2}{5}$ (d)

Not possible to find



Watch Video Solution

394. If $A = \begin{bmatrix} 0 & 2 \\ 3 & -4 \end{bmatrix}$ and $kA = \begin{bmatrix} 0 & 3a \\ 2b & 24 \end{bmatrix}$, then the values of k, a, b ,

are respectively (a) -6, -12, -18 (b) -6, 4, 9 (c) -6, -4, -9 (d) -6, 12, 18



Watch Video Solution

395. If $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $J = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$, then B

equals a) $I \cos \theta + J \sin \theta$ (b) $I \sin \theta + J \cos \theta$ (c) $I \cos \theta - J \sin \theta$ (d)

$I \cos \theta + J \sin \theta$



Watch Video Solution

396. The trace of the matrix $A = \begin{bmatrix} 1 & -5 & 7 \\ 0 & 7 & 9 \\ 11 & 8 & 9 \end{bmatrix}$ is (a) 17 (b) 25 (c) 3 (d) 12

 [Watch Video Solution](#)

397. If $A = [a_{ij}]$ is a scalar matrix of order $n \times n$ such that $a_{ii} = k$ for all i , then trace of A is equal to nk (b) $n + k$ (c) $\frac{n}{k}$ (d) none of these

 [Watch Video Solution](#)

398. The matrix $A = [004040400]$ is a (a) square matrix (b) diagonal matrix (c) unit matrix (d) none of these

 [Watch Video Solution](#)

399. The number of possible matrices of order 3×3 with each entry 2 or 0 is (a) 9 (b) 27 (c) 81 (d) none of these

 [Watch Video Solution](#)

400. If $[2x + y \ 4x \ 5x - 74x] = [77y - 13yx + 6]$, then the value of $x + y$ is $x = 3, y = 1$ (b) $x = 2, y = 3$ (c) $x = 2, y = 4$ (d) $x = 3, y = 3$

 [Watch Video Solution](#)

401. If A is a square matrix such that $A^2 = I$, then $(A - I)^3 + (A + I)^3 - 7A$ is equal to A (b) $I - A$ (c) $I + A$ (d) $3A$

 [Watch Video Solution](#)

402. If A and B are two matrices of the order $3 \times m$ and $3 \times n$, respectively and $m = n$, then order of matrix $(5A - 2B)$ is (a) $m \times 3$ (b) 3×3 (c) $m \times n$ (d) $3 \times n$

 [Watch Video Solution](#)

403. If A is a matrix of order $m \times n$ and B is a matrix such that AB^T and $B^T A$ are both defined, then the order of matrix B is

A. $m \times n$

B. $n \times n$

C. $n \times m$

D. $m \times m$

Answer: A



[Watch Video Solution](#)

404. If A and B are matrices of the same order, then $AB^T - BA^T$ is a (a) skew-symmetric matrix (b) null matrix (c) unit matrix (d) symmetric matrix



[Watch Video Solution](#)

405. If matrix $A = ([a_{ij}])_{2 \times 2}$, where $a_{ij} = \begin{cases} 1, & \text{if } i \neq j \\ 0, & \text{if } i = j \end{cases}$, then A^2 is equal to (a) I (b) A (c) O (d) I

 Watch Video Solution

406. If $A = \frac{1}{\pi} \begin{bmatrix} \sin^{-1}(\pi x) & \tan^{-1}(\frac{x}{\pi}) \\ \sin^{-1}(\frac{x}{\pi}) & \cot^{-1}(\pi x) \end{bmatrix}$ and $B = \frac{1}{\pi} \begin{bmatrix} -\cos^{-1}(\pi x) & \tan^{-1}(\frac{x}{\pi}) \\ \sin^{-1}(\frac{x}{\pi}) & -\tan^{-1}(\pi x) \end{bmatrix}$, then $A - B$ is equal to (a) I (b) O (c) $2I$ (d) $\frac{1}{2}I$

 Watch Video Solution

407. If A and B are square matrices of the same order, then $(A + B)(A - B)$ is equal to (a) $A^2 - B^2$ (b) $A^2 - BA - AB - B^2$ (c) $A^2 - B^2 + BA - AB$ (d) $A^2 - BA + B^2 + AB$

 Watch Video Solution

408. If $A = \begin{bmatrix} 2 & -1 & 3 \\ -4 & 5 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 3 \\ 4 & -2 \\ 1 & 5 \end{bmatrix}$, then only AB is

defined (b) only BA is defined (c) AB and BA both are defined (d) AB and BA both are not defined

 [Watch Video Solution](#)

409. The matrix $A = \begin{bmatrix} 0 & -5 & 8 \\ 5 & 0 & 12 \\ -8 & -12 & 0 \end{bmatrix}$ is a (a) diagonal matrix (b) symmetric matrix (c) skew-symmetric matrix (d) scalar matrix

 [Watch Video Solution](#)

410. The matrix $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 4 \end{bmatrix}$ is (a) identity matrix (b) symmetric matrix (c) skew-symmetric matrix (d) diagonal matrix

 [Watch Video Solution](#)