

India's Number 1 Education App

PHYSICS

NCERT - NCERT PHYSICS(ENGLISH)

RAY OPTICS AND OPTICAL INSTRUMENTS

Solved Examples

1. Suppose that the lower half of a concave mirror's reflecting surface is covered with an

opaque non-reflecting material. What effect

will this have on the image of an object placed

in front of the mirror ?

Watch Video Solution

2. A mobile phone lies along the principal axis of a concave mirror as shown in Fig. Show by suitable diagram, the formation of its image. Explain why the magnification is not uniform, and distortion will occur depending on the location of the mobile with respect to the

mirror.



3. An object is placed (i)10cm (ii) 5cm in front of a convex mirror of radius of curvature 15cm
. Find the position, nature and magnification of the image in each case.

4. Suppose while sitting in a parked car, you notice a jogger approaching towards you in the rear view mirror of R = 2m. If the jogger is running at a speed of $5ms^{-1}$, how fast is the image of the jogger moving, when the jogger is

(a) 39 m

(b) 29 m

19 m and

(d) 9 m. away ?





5. The earth takes 24 hours to rotate once about its axis. How much time (in min) does the sun take to shift by 1° when viewed from the earth?

A. 2 min

B.4 min

C. 6 min

D. 8 min

Answer: B



6. Light from a point source in air falls on a spherical glass surface. If $\mu = 1.5$, and radius of curvature = 20cm, the distance of light source from the glass surface is 100cm, at what position will the image be formed ? (NCERT Solved Example)

Watch Video Solution

7. A magician during a show makes a glass lens $\mu = 1.5$ disappear in a through of liquid. What is the refractive index of the liquid ? Is the liquid water ?

View Text Solution

8. (i) If f=~+~0.5m, what is the power of the lens ?

(ii) The radii of curvature of the faces of a double convex lens are 9cm and 15cm. Its

```
focal length is 12cm. What is the refractive
index of glass ?
(iii) A convex lens has 20cm focal length in air.
What is the focal length in water ? (Refractive
index of air-water = 1.33, refractive index of
air-glass = 1.5).
```

9. Find the position of the image formed by the lens combination given in Fig.

Watch Video Solution



Exercise

1. A small candle 2.5*cm* in size is placed 27*cm* in front of a concave mirror of radius of curvature 36*cm*. At what distance from the mirror should a screen be placed in order to

receive a sharp image ? Describe the nature and size of the image. If the candle is moved closer to the mirror, how would the screen have to be moved ?



2. A 4.5*cm* needle is placed 12 cm away from a convex mirror of focal length 15 cm. Give the location of the image and the magnification. Describe what happens as the needle is moved farther from the mirror.



3. A tank is filled with water to a height of 12.5*cm*. The apparent depth of a needle lying at the bottom of the tank is measured by a microscope to be 9.4cm. What is the refractive index of water ? If water is replaced by a liquid of refractive index 1.63 upto the same height, by what distance would the microscope have to be moved to focus on the needle again?



4. Fig. (a) and (b) show refraction of an incident ray in air at 60° with the normal to a glass-air and water-air interface respectively. Predict the angle of refraction of an incident ray in water at 45° with the normal to a water glass interface. Take $.^{a} \mu_{g} = 1.32$.



Watch Video Solution

5. A small bulb (assumed to be a point source) is placed at the bottom of a tank containing water to a depth of 80cm. Find out the area of the surface of water through which light from the bulb can emerge. Take the value of refractive index of water to be 4/3.

Watch Video Solution

6. A prism is made of glass of unknown refractive index. A parallel beam of light is

incident on a face of the prism. By rotating the prism, the minimum angle of deviation is measured to be 40° . What is the refractive index of the prism ? If the prism is placed in water ($\mu = 1.33$), predict the new angle of minimum deviation of the parallel beam. The refracting angle of prism is 60° .



7. A double convex lens is made of glass of refractive index 1.55 with both faces of same

radius of curvature. Find the radius of curvature required, if focal length is 20cm.

A. 22 cm

B. 44 cm

C. 11 cm

D. 33 cm

Answer: A



8. A beam of light converges to a point P.A lens is placed in the path of the convergent beam 12cm from P. At what point does the beam converge if the lens is (a) a convex lens of focal length 20cm. (b) a

concave lens of focal length 16cm ?

Watch Video Solution

9. An object of size 3.0cm is placed 14cm in front of a concave lens of focal length 21cm.

Describe the image produced by the lens. What happens if the object is moved further from the lens ?

Watch Video Solution

10. What is the focal length of a convex lens of focal length 30cm in contact with a concave lens of focal length 20cm. Is the system a converging or a diverging lens ? Ignore thickness of the lenses.

B. 60 cm

C. -30 cm

D. -60 cm

Answer: D

Watch Video Solution

11. A compound microscope consists of an objects lens of focal length 2.0 cm and an eyepiece of focal length 20cm? If the system a

converging or a diverging lens? Ignore

thickness of the lenses.



12. A person with a normal near point (25cm) using a compound microscope with an objective of focal length 8.0mm and eye piece of focal length 2.5cm can bring an object placed 9.0cm from the objective in sharp focus. What is the separation between the two

lenses ? Calculate the magnifying power of the

microscope?



13. A small telescope has an objective lens of focal length 144*cm* and an eye-piece of focal length 6.0*cm*. What is the magnifying power of the telescope ? What is the separation between the objective and the eye-piece ?



14. (i) A giant refracting telescope at an observatory has an objective lens of focal length 15 m. If an eyepiece of focal length 1.0 cm is used, what is angular magnification of the telescope ? (ii) If this telescope is used to view the moon, what is the diameter of the image of the moon formed by the objective lens? the diameter of the moon is $3.48 \times 10^6 m$, and the radius of lunar orbit is $3.8 imes 10^8 m$.



15. Use the mirror equation to deduct that :
(a) an object between f and 2f of a concave mirror produces a real image beyond 2f.
(b) a convax mirror always produces a virtual image independent of the location of the object.
(c) the virtual image produced by a convex

(c) the virtual image produced by a convex mirror is always diminished in size and is located between the focus and the pole.
(d) an object placed between the pole and focus of a concave mirror produces a virtual and enlarged image.



16. A small pin fixed on a table top is viewed from above from a distance of 50cm. By what distance would the pin appear to be raised, if it be viewed from the same point through a 15cm. Thick glass slab held parallel to the table ? μ of glass 1.5 Does the answer depend on location of the slab ?



17. (a) Fig. shows a cross-section of a 'light pipe' made of a glass fibre of refractive index 1.68. The outer covering of the pipe is made of a material of refractive index 1.44. What is the axis of the pipe for which total reflection inside the pipe take place as shwon. (b) What is the answer if there is no outer

covering if the pipe?



18. Answer the following questions :

Does the apparent depth of a tank of water

change if viewed obliquely ? If so, does the

apparent depth increase or decrease ?



19. The image of a small electric bulb fixed on the wall of a room is to be obtained on the opposite wall 3m away by means of a large convex lens. What is the maximum possible focal length of the lens required for the purpose ?



20. A screen is placed 90cm from an object. The image of the object on the screen is formed by a convex lens at two different location separated by 20cm. Determine the focal length of the lens.



21. a) Determine the effective focal length of the combination of the two lenses in Exercise, if they are placed 8.0cm apart with their principal axes coincident. Does the answer depend on which side of the combination a beam of paralel light is incident? Is the notions of effective focal length of this system useful at all?

b) An object 1.5 cm in size is placed on the side of the convex lens in the arrangement a) above. The distance between the object and the convex lens is 40cm. Determine the magnification produced by the two-lens

system, and the size of the image.



22. At what angle should a ray of light be incident on the face of a prism of refracting angle 60° , so that it just suffers total internal reflection at the other face ? The refractive index of the prism is 1.524.



23. A cardsheet divided into squares each of size $1mm^2$ is being viewed at a distance of 9cm through a magnifying glass (a conerging lens of focal length 10cm) held close to the eye.

(a) What is the magnification produced by the lenas ? How much is the area of each square to the virtual image ?

(b) What is the angular magnification(magnifying power) of the lens ?(c) Is the magnification in (a) equal to the

magnifying power in (b) ? Explain

24. (i) At what distance should the lens be held
from the card sheet in order to view the
squares distinctly with the maximum possible
magnifying power ?
(ii) What is the magnification in this case ?
(iii) Is the magnification equal to magnifying
power in this case ? Explain.



25. What should be the distance between the object and magnifying glass if the virtual image of each square in the figure is to have an area of $6.25mm^2$. Would you be able to see the squares distinctly with your eyes very close to the magnifier ?

View Text Solution

26. a) The anlge subtended at the eye by an object is equal to the angle subtended at the eye by the virtual image produced by a

magnifying glass. In what sense then does amagnifying glass provide angularmagnifications?b) in viewing through a magnifying glass, one

lens. Does angular magnification change if the eye is moved back?

usually positions one's eyes very close to the

c) magnifying power of a simple microscopes is inversely proportional to the focal length of the lens. What then stops us from using a convex lens of smaller and smaller focal length and achieving greater and greater magnifying power? d) Why must both the objective and the eyepiece of a compound microscope have short focal lengths?

e) When viewing through a compound microscope, our eyes should be positioned not on the eyepiece but a short distance away from it fot best veiwing. Why? How much should be that short distance between the eye and eyepiece?

View Text Solution

27. An angular magnification (magnifying power) of 30X is desired using an objective of focal length 1.25cm and an eye piece of focal length 5cm. How will you set up the compound microscope ?

View Text Solution

28. A small telescope has an objective lens of focal length 140 cm and an eyepiece of focal length 5.0 cm. what is the magnifying power of

the telescope for viewing distant objects when(a) the telescope is in normal adjustment (i.e,when the final image is at infinity),(b) The final image is formed at the leastdistance of distinct vision (25 cm)

View Text Solution

29. (a) For the telescope described what is the separation between the objective lens and eye piece ?

(b) If this telescope is used to view a 100m tall

tower 3km away, what is the height of the image of the tower formed by the objective lens ?

(c) What is the height of the final image of the

tower if it is formed at 25 cm`?

View Text Solution

30. A Cassegrainian telescope uses two mirrors as shown in Fig. Such a telescope is built with the mirrors 20mm apart. If the radius of curvature of large mirror is 220mm

and the small mirror is 40mm, where will the

final image of an object at infinity be?



31. Light incident normally on a plane mirror attached to a galvanometer coil retraces backwards a shown in Fig. 6(a). 14. A current in

the coil produces a deflection of 3.5° in the mirror. What is the displacement of the reflected spot of light on a screen placed 1.5m away?



32. Fig. shows an equiconvex lens (of refractive index 1.5) in contact with a liquid layer on top of a plane mirror. A small needle with its tip on the principal axis is moved along the axis until its inverted image is found at the position of

the needle. The distance of the needle from the lens is measured to be 45.0*cm*. The liquid is removed and the experiment is repeated. The new distance is measured to be 30.0*cm*. What is the refractive index of the liquid ?





