



MATHS

BOOKS - CENGAGE MATHS (HINGLISH)

DIFFERENTIATION

Examples

1. If $y = \sqrt{\frac{1 - \cos 2x}{1 + \cos 2x}}$, $x \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right)$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

2. Find the derivative of $e^{\sqrt{x}}$ w.r.t. x using the first principle.

 [Watch Video Solution](#)

3. If $f(x) = x \tan^{-1} x$, find $f'(\sqrt{3})$ using the first principle.



Watch Video Solution

4. If $f(x) = [2x] \sin 3\pi x$ then prove that $f'(k^+) = 6k\pi(-1)^k$, (where $[.]$ denotes the greatest integer function and $k \in \mathbb{N}$).



Watch Video Solution

5. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfying $|f(x)| \leq x^2 \forall x \in \mathbb{R}$ be differentiable at $x = 0$. The find $f'(0)$.



Watch Video Solution

6. A function $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfies the equation $f(x+y) = f(x)f(y)$ for all $x, y \in \mathbb{R}$ and $f(x) \neq 0$ or all $x \in \mathbb{R}$. If $f(x)$ is differentiable at $x = 0$ and $f'(0) = 2$, then prove that $f'(x) = 2f(x)$.

 [Watch Video Solution](#)

7. If $y = \left(1 + x^{\frac{1}{4}}\right)\left(1 + x^{\frac{1}{2}}\right)\left(1 - x^{\frac{1}{4}}\right)$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

8. If $f(x) = x|x|$, then prove that $f'(x) = 2|x|$

 [Watch Video Solution](#)

9. If $y = 1 = \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!}$, show that $\frac{dy}{dx} - y + \frac{x^n}{n!} = 0$.

 [Watch Video Solution](#)

10. Find $\frac{dy}{dx}$ for $y = \sin^{-1}(\cos x)$, where $x \in (0, 2\pi)$.

 [Watch Video Solution](#)

11. Differentiate $\sin^{-1}\left(2x\sqrt{1-x^2}\right)$ with respect to x if $x \in \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$

 [Watch Video Solution](#)

12. $y = \tan^{-1}\left(\frac{a\cos x - b\sin x}{b\cos x + a\sin x}\right)$, where $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

 [Watch Video Solution](#)

13. $y = \sin^{-1}\left(\frac{x}{1+x^2}\right) + \cos^{-1}\left(\frac{x}{1+x^2}\right)$, where $x \in \mathbb{R}$

 [Watch Video Solution](#)

14. Find $\frac{dy}{dx}$ for $y = \sin^{-1}(x^2 + 1)$

 [Watch Video Solution](#)

15. If $y = \sqrt{\log \left\{ \sin \left(\frac{x^2}{3} - 1 \right) \right\}}$, then $f \in d \frac{dy}{dx}$.

 [Watch Video Solution](#)

16. Differentiate the function $f(x) = \sec(\tan(\sqrt{x}))$ with respect to x .

 [Watch Video Solution](#)

17. Find $\frac{dy}{dx} f$ or $y = \log(x + \sqrt{a^2 + x^2})$.

 [Watch Video Solution](#)

18. $y = \tan^{-1} \left(\frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} \right)$, where $x \in (-1, 1)$.

 [Watch Video Solution](#)

19. Find $\frac{dy}{dx}$ for $y = \tan^{-1} \sqrt{\frac{a-x}{a+x}}$, $a > x > -a$

 [Watch Video Solution](#)

20. If $y = \sin^{-1} \left[x\sqrt{1-x} - \sqrt{x}\sqrt{1-x^2} \right]$ and $x > 0$

 [Watch Video Solution](#)

21. If $y = \frac{\tan^{-1} 1}{1+x+x^2} + \frac{\tan^{-1} 1}{x^2+3x+3} + \frac{\tan^{-1} 1}{x^2+5x+7} + \dots$ upto n terms, then

find the value of $y'(0)$

 [Watch Video Solution](#)

22. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a one-one onto differentiable function, such that

$f(2) = 1$ and $f'(2) = 3$. The find the value of $\left(\left(\frac{d}{dx} (f^{-1}(x)) \right) \right)_{x=1}$

 [Watch Video Solution](#)

23. If $f(x) = \cos x \cos 2x \cos 4x \cos(8x) \dots \cos 16x$ then find $f' \left(\frac{\pi}{4} \right)$

 [Watch Video Solution](#)

24. If $\cos y = x \cos(a + y)$, with $\cos a \neq \pm 1$, prove that $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$.

 [Watch Video Solution](#)

25. Find $\frac{dy}{dx}$ or $y = x \sin x \log x$.

 [Watch Video Solution](#)

26. Evaluate $\lim_{h \rightarrow 0} \frac{(a + h)^2 \cdot \sin^{-1}(a + h) - a^2 \sin^{-1} a}{h}$.

 [Watch Video Solution](#)

27. Differentiate $y = \frac{e^x}{1 + \sin x}$

 [Watch Video Solution](#)

28. If $f(x) = \cos x \cos 2x \cos 4x \cos(8x) \dots \cos 16x$ then find $f\left(\frac{\pi}{4}\right)$

 [Watch Video Solution](#)

29. If $\cos y = x \cos(a + y)$, with $\cos a \neq \pm 1$, prove that $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$.

 [Watch Video Solution](#)

30. If $y = \sqrt{\frac{1-x}{1+x}}$, prove that $(1-x^2) \frac{dy}{dx} + y = 0$

 [Watch Video Solution](#)

31. Find the sum of the series $1 + 2x + 3x^2 + (n - 1)x^{n-2}$ using differentiation.

 [Watch Video Solution](#)

32. If $\sqrt{x} + \sqrt{y} = 4$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

33. If $xy + y^2 = \tan x + y$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

34. If $y = x + \frac{1}{x + \frac{1}{x + \frac{1}{x + \dots}}}$, prove that $\frac{dy}{dx} = \frac{y}{2y - x}$.

 [Watch Video Solution](#)

35. $\sec(x + y) = xy$

 [Watch Video Solution](#)

36. If $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots}}}$, prove that $\frac{dy}{dx} = \frac{\cos x}{2y - 1}$

 [Watch Video Solution](#)

37. If $\log(x^2 + y^2) = 2 \tan^{-1}\left(\frac{y}{x}\right)$, show that $\frac{dy}{dx} = \frac{x + y}{x - y}$

 [Watch Video Solution](#)

38. If $y = y(x)$ and it follows the relation $4xe^{xy} = y + 5\sin^2 x$, then $y'(0)$ is equal to _____

 [Watch Video Solution](#)

39. Find $\frac{dy}{dx}$ if $x = a(\theta - \sin\theta)$ and $y = a(1 - \cos\theta)$.

 [Watch Video Solution](#)

40. If $x = a \sec^3\theta$ and $y = a \tan^3\theta$, find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{3}$.

 [Watch Video Solution](#)

41. If $x = e^{\cos 2t}$ and $y = e^{\sin 2t}$, prove that $\frac{dy}{dx} = -\frac{y \log x}{x \log y}$.

 [Watch Video Solution](#)

42. Let $y = x^3 - 8x + 7$ and $x = f(t)$ if $\frac{f'(dy)}{dx} = 2$ and $x = 3a = 0$, then find the value of $\frac{dx}{dt} a = 0$.

 [Watch Video Solution](#)

43. Find the derivative of $\frac{\sqrt{x}(x+4)^{\frac{3}{2}}}{(4x-3)^{\frac{4}{3}}}$

 [Watch Video Solution](#)

44. If $x^m y^n = (x+y)^{m+n}$, prove that $\frac{dy}{dx} = \frac{y}{x}$.

 [Watch Video Solution](#)

45. Differentiate $(\log x)^{\cos x}$ with respect to x

 [Watch Video Solution](#)

46. If $f(x) = |x|^{|\sin x|}$, then find $f' \left(-\frac{\pi}{4} \right)$

 [Watch Video Solution](#)

47. If $y = x^x \wedge x \wedge (((((\infty))))))$, find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

48. If $f(x) = \lim_{h \rightarrow 0} \frac{(\sin(x+h))^{\log_e(x+h)} - (\sin x)^{\log_e x}}{h}$ then find $f(\pi/2)$.

 [Watch Video Solution](#)

49. If $x < 1$, provethat $\frac{1}{1+x} + \frac{2x}{1+x^2} + \frac{4x^3}{1+x^4} + \infty = \frac{1}{1-x}$

 [Watch Video Solution](#)

50. Differentiate $\log \sin x$ w.r.t. \sqrt{x} .

 [Watch Video Solution](#)

51. Differentiate $\tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$ with respect to $\tan^{-1}x$, when $x \neq 0$.

 [Watch Video Solution](#)

52. Find the derivative of $f(\tan x) \text{ wrt } g(\sec x) \text{ at } x = \frac{\pi}{4}$, where $f'(1) = 2$ and $g'(\sqrt{2}) = 4$.

 [Watch Video Solution](#)

53.

Let $f(x) = \begin{vmatrix} \cos x & \sin x & \cos x \\ \cos 2x & \sin 2x & 2\cos 2x \\ \cos 3x & \sin 3x & 3\cos 3x \end{vmatrix}$ Then find the value of $f(0)$ and $f(\pi/2)$.

 [Watch Video Solution](#)

54. $f(x) = \begin{vmatrix} \cos x & x & 1 \\ 2\sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix}$. Then find the value of $\lim_{x \rightarrow 0} \frac{f(x)}{x}$.

 [Watch Video Solution](#)

55. If $y = \cos^{-1}x$, find $\frac{d^2y}{dx^2}$ in terms of y alone.

 [Watch Video Solution](#)

56. If $y = (x^2 - 1)^m$, then the $(2m)$ th differential coefficient of y is

 [Watch Video Solution](#)

57. If $y = x \log \left\{ \frac{x}{(a + bx)} \right\}$, then show that $x^3 \frac{d^2y}{dx^2} = \left(x \frac{dy}{dx} - y \right)^2$.

 [Watch Video Solution](#)

58. If $(x - a)^2 + (y - b)^2 = c^2$, for some $c > 0$, prove that $\frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$ is a constant independent of a and b .

 [Watch Video Solution](#)

59. If $y = e^a \cos^{(-1)x}$, $-1 \leq x < 1$, show that $(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - a^2 y = 0$

 [Watch Video Solution](#)

60. If $x = a(\cos t + t \sin t)$ and $y = a(\sin t - t \cos t)$, $f \in d \frac{d^2y}{dx^2}$.

 [Watch Video Solution](#)

61. If g is inverse of f then prove that $f'(g(x)) = -g'(x)(f'(g(x)))^3$.

 [Watch Video Solution](#)

62. Let $f(x)$ and $g(x)$ be real valued functions such that $f(x)g(x)=1$,

$\forall x \in R$. If $f'(x)$ and $g'(x)$ exists $\forall x \in R$ and $f(x)$ and $g'(x)$

are never zero, then prove that $\frac{f'(x)}{f(x)} - \frac{g'(x)}{g(x)} = \frac{2f'(x)}{f(x)}$

 [Watch Video Solution](#)

63. Prove that $\phi(x) = \begin{vmatrix} f(x) & g(x) & h(x) \\ f'(x) & g'(x) & h'(x) \\ f''(x) & g''(x) & h''(x) \end{vmatrix}$ is a constant polynomial.

 [Watch Video Solution](#)

64. Let $f\left(\frac{x+y}{2}\right) = \frac{f(x)+f(y)}{2}f$ or *all real x and y* If $f'(0)$ exists and equals -1 and $f(0) = 1$, then $f \in df(2)$

 [Watch Video Solution](#)

65.

If $f(x) + f(y) = f\left(\frac{x+y}{1-xy}\right)$ for all $x, y \in R, (xy \neq 1)$, and $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 2$ then



[Watch Video Solution](#)

66. Let $f: R - \{0\} \rightarrow R$ be a function which is differentiable in its domain and satisfying the equation $f(x+y) = f(x) + f(y) + \frac{x+y}{xy} - \frac{1}{x+y}$, also $f'(1)=2$. Then find the function.



[Watch Video Solution](#)

67. Find function $f(x)$ which is differentiable and satisfy the relation $f(x+y) = f(x) + f(y) + (e^x - 1)(e^y - 1) \forall x, y \in R$, and $f(0) = 2$.



[Watch Video Solution](#)

68. If $f\left(\frac{x}{y}\right) = \frac{f(x)}{f(y)}$, $\forall y, f(y) \neq 0$ and $f(1) = 2$, find $f(x)$.

 [Watch Video Solution](#)

69. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function satisfying condition $f(x + y^3) = f(x) + [f(y)]^3$ or $\forall x, y \in \mathbb{R}$. If $f'(0) \geq 0$, find $f(10)$.

 [Watch Video Solution](#)

70. Let $f(x + y) = f(x) + f(y) + 2xy - 1$ for all real x and y and $f(x)$ be a differentiable function. If $f'(0) = \cos \alpha$, then prove that $f(x) > 0 \forall x \in \mathbb{R}$.

 [Watch Video Solution](#)

71. If $f(x) = (\log)_{x^2}(\log x)$, then $f'(x)$ at $x = e$ is (a) 0 (b) 1 (c) $\frac{1}{e}$ (d) $\frac{1}{2}e$

 [Watch Video Solution](#)

72. Given that $\cos\left(\frac{x}{2}\right) \cdot \cos\left(\frac{x}{4}\right) \cdot \cos\left(\frac{x}{8}\right) \dots = \frac{\sin x}{x}$ Prove that

$$\left(\frac{1}{2^2}\right)\sec^2\left(\frac{x}{2}\right) + \left(\frac{1}{2^4}\right)\sec^2\left(\frac{x}{4}\right) + \dots = \operatorname{cosec}^2 x - \frac{1}{x^2}$$

 [Watch Video Solution](#)

73. If $y = f(a^x)$ and $f'(\sin x) = (\log)_e x$, then $f \in d\frac{dy}{dx}$, if it exists, where $\pi/2$

 [Watch Video Solution](#)

74. If P_n is the sum of a GP upto n terms ($n \geq 3$), then prove that

$$(1-r)\frac{dP_n}{dr} = (1-n)P_n + nP_{n-1}, \text{ where } r \text{ is the common ratio of } GP$$

 [Watch Video Solution](#)

75. If $g(x) = \frac{f(x)}{(x-a)(x-b)(x-c)}$, where $f(x)$ is a polynomial of degree < 3 , then prove that

$$\frac{dg(x)}{dx} = \left| 1af(a)(x-a)^{-2} - 1bf(b)(x-b)^{-2} - 1cf(c)(x-c)^{-2} \right| + \left| a^2a1b^2b1c^2c1 \right|$$

 Watch Video Solution

76. If $x = \operatorname{cosec}\theta - \sin\theta$ and $y = \operatorname{cosec}^n\theta - \sin^n\theta$, then show that

$$(x^2 + 4) \left(\frac{dy}{dx} \right)^2 = n^2 (y^2 + 4)$$

 Watch Video Solution

77. If $y = \frac{ax^2}{(x-a)(x-b)(x-c)} + \frac{bx}{(x-b)(x-c)} + \frac{c}{x-c} + 1$, then prove that

$$\frac{y'}{y} = \frac{1}{x} \left[\frac{a}{a-x} + \frac{b}{b-x} + \frac{c}{c-x} \right]$$

 Watch Video Solution

78. Find the differential equation of the family of curves $y = Ae^{2x} + Be^{-2x}$, where A and B are arbitrary constants.

 [Watch Video Solution](#)

79. If $y = \left(\frac{1}{2}\right)^{n-1} \cos(n \cos^{-1} x)$, then prove that y satisfies the differential equation $(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + n^2y = 0$

 [Watch Video Solution](#)

80. Let $f(x)$ and $g(x)$ be two functions having finite nonzero third-order derivatives f''' and g''' for all $x \in \mathbb{R}$. If $f(x)g(x) = 1$ for all $x \in \mathbb{R}$, then prove that $f'f' - g'g' = 3(f'f - g'g)$

 [Watch Video Solution](#)

81. If a curve is represented parametrically by the equation

$$x = f(t) \text{ and } y = g(t) \text{ then prove that } \frac{d^2y}{dx^2} = - \left[\frac{g'(t)}{f'(t)} \right]^3 \left(\frac{d^2x}{dy^2} \right)$$

 [Watch Video Solution](#)

82. If $f\left(\frac{x+y}{3}\right) = \frac{2 + f(x) + f(y)}{3}$ for all real x and y and $f'(2) = 2$, then determine $y = f(x)$.

 [Watch Video Solution](#)

83. If $f(x) = \frac{f(x)}{y} + \frac{f(y)}{x}$ holds for all real x and y greater than 0 and $f(x)$ is a differentiable function for all $x > 0$ such that $f(e) = \frac{1}{e}$, then $f \in df(x)$.

 [Watch Video Solution](#)

84. If $\left| a_1 \sin x + a_2 \sin 2x + \dots + a_n \sin nx \right| \leq |\sin x|$ for $x \in \mathbb{R}$, then prove that $\left| a_1 + 2a_2 + \dots + na_n \right| \leq 1$

 [Watch Video Solution](#)

85. Suppose $p(x) = a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n$. If $|p(x)| \leq e^{x-1} - 1$ for all $x \geq 0$, prove that $\left| a_1 + 2a_2 + \dots + na_n \right| \leq 1$.

 [Watch Video Solution](#)

86. If $y = \sqrt{\frac{1 - \cos 2x}{1 + \cos 2x}}$, $x \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right)$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

87. Find the derivative of $e^{\sqrt{x}}$ w.r.t. x using the first principle.

 [Watch Video Solution](#)

88. If $f(x) = x \tan^{-1} x$, find $f'(\sqrt{3})$ using the first principle.

 [Watch Video Solution](#)

89. If $f(x) = [2x] \sin 3\pi x$ then prove that $f'(k^+) = 6k\pi(-1)^k$, (where $[.]$ denotes the greatest integer function and $k \in \mathbb{N}$).

 [Watch Video Solution](#)

90. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfying $|f(x)| \leq x^2 \forall x \in \mathbb{R}$ be differentiable at $x = 0$. The find $f'(0)$.

 [Watch Video Solution](#)

91. A function $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfies the equation $f(x+y) = f(x)f(y)$ for all $x, y \in \mathbb{R}$ and $f(x) \neq 0$ or $\forall x \in \mathbb{R}$. If $f(x)$ is differentiable at

$x = 0$ and $f'(0) = 2$, then prove that $f'(x) = 2f(x)$



Watch Video Solution

92. If $y = \left(1 + x^{\frac{1}{4}}\right)\left(1 + x^{\frac{1}{2}}\right)\left(1 - x^{\frac{1}{4}}\right)$, then find $\frac{dy}{dx}$.



Watch Video Solution

93. If $f(x) = x|x|$, then prove that $f'(x) = 2|x|$



Watch Video Solution

94. If $y = 1 = \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!}$, show that $\frac{dy}{dx} - y + \frac{x^n}{n!} = 0$.



Watch Video Solution

95. Find $\frac{dy}{dx}$ for $y = \sin^{-1}(\cos x)$, $x \in (0, \pi) \cup (\pi, 2\pi)$.



Watch Video Solution

96. Differentiate $\sin^{-1}\left(2x\sqrt{1-x^2}\right)$ with respect to x if $x \in \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$



Watch Video Solution

97. $y = \tan^{-1}\left(\frac{\cos x - b \sin x}{b \cos x + a \sin x}\right)$, where $\frac{\pi}{2} < x < \pi$



Watch Video Solution

98. $y = \sin^{-1}\left(\frac{x}{1+x^2}\right) + \cos^{-1}\left(\frac{x}{1+x^2}\right)$, where $x > 0$



Watch Video Solution

99. Find $\frac{dy}{dx}$ for $y = \sin(x^2 + 1)$.



Watch Video Solution

100. If $y = \sqrt{\log \left\{ \sin \left(\frac{x^2}{3} - 1 \right) \right\}}$, then $f \in d \frac{dy}{dx}$.

 [Watch Video Solution](#)

101. Differentiate the function $f(x) = \sec (\tan \sqrt{x})$ with respect to x ,

 [Watch Video Solution](#)

102. Find $\frac{dy}{dx}$ for $y = \log \left(x + \sqrt{a^2 + x^2} \right)$.

 [Watch Video Solution](#)

103. $y = \tan^{-1} \left(\frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} \right)$, where $x \in (-1, 1)$

 [Watch Video Solution](#)

104. Find $\frac{dy}{dx}$ for $y = \tan^{-1} \sqrt{\frac{a-x}{a+x}}$, $a > 0$

 [Watch Video Solution](#)

105. If $y = \sin^{-1} \left[x\sqrt{1-x} - \sqrt{x}\sqrt{1-x^2} \right]$ and $x = 0$

 [Watch Video Solution](#)

106. If $y = \frac{\tan^{-1} 1}{1+x+x^2} + \frac{\tan^{-1} 1}{x^2+3x+3} + \frac{\tan^{-1} 1}{x^2+5x+7} + \dots$ upto n terms, then find the value of $y'(0)$.

 [Watch Video Solution](#)

107. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a one-one onto differentiable function, such that

$f(2) = 1$ and $f'(2) = 3$. The find the value of $\left(\left(\frac{d}{dx} (f^{-1}(x)) \right) \right)_{x=1}$

 [Watch Video Solution](#)

108. If $f(x) = \cos x \cos 2x \cos 4x \cos(8x) \dots \cos 16x$ then find $f\left(\frac{\pi}{4}\right)$

 [Watch Video Solution](#)

109. If $\cos y = x \cos(a + y)$, with $\cos a \neq \pm 1$, prove that $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$

 [Watch Video Solution](#)

110. Find $\frac{dy}{dx}$ for $y = x \sin x \log x$

 [Watch Video Solution](#)

111. Evaluate $\lim_{h \rightarrow 0} \frac{(a + h)^2 \cdot \sin^{-1}(a + h) - a^2 \sin^{-1} a}{h}$

 [Watch Video Solution](#)

112. Differentiate $y = \frac{e^x}{1 + \sin x}$

 [Watch Video Solution](#)

113. If $f(x) = \cos x \cos 2x \cos 4x \cos(8x) \dots \cos 16x$ then find $f\left(\frac{\pi}{4}\right)$

 [Watch Video Solution](#)

114. If $\cos y = x \cos(a + y)$, with $\cos a \neq \pm 1$, prove that $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$

 [Watch Video Solution](#)

115. If $y = \sqrt{\frac{1-x}{1+x}}$, prove that $(1-x^2) \frac{dy}{dx} + y = 0$

 [Watch Video Solution](#)

116. Find the sum of the series $1 + 2x + 3x^2 + (n-1)x^{n-2}$ using differentiation.

 [Watch Video Solution](#)

117. If $\sqrt{x} + \sqrt{y} = 4$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

118. If $xy + y^2 = \tan x + y$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

119. If $y = x + \frac{1}{x + \frac{1}{x + \frac{1}{x + \dots}}}$, prove that $\frac{dy}{dx} = \frac{y}{2y - x}$.

 [Watch Video Solution](#)

120. If $\sec(x+y) = xy$, then find $\frac{dy}{dx}$

 [Watch Video Solution](#)

121. If $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots}}}$, prove that $\frac{dy}{dx} = \frac{\cos x}{2y - 1}$

 [Watch Video Solution](#)

122. If $\log(x^2 + y^2) = 2 \tan^{-1}\left(\frac{y}{x}\right)$, show that $\frac{dy}{dx} = \frac{x+y}{x-y}$

 [Watch Video Solution](#)

123. If $y = y(x)$ and it follows the relation $4xe^{xy} = y + 5\sin^2 x$, then $y'(0)$ is equal to _____

 [Watch Video Solution](#)

124. Find $\frac{dy}{dx}$ if $x = a(\theta - \sin\theta)$ and $y = a(1 - \cos\theta)$.

 [Watch Video Solution](#)

125. If $x = a \sec^3\theta$ and $y = a \tan^3\theta$, find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{3}$.

 [Watch Video Solution](#)

126. If $x = e^{\cos 2t}$ and $y = e^{\sin 2t}$, prove that $\frac{dy}{dx} = -\frac{y \log x}{x \log y}$

 [Watch Video Solution](#)

127. Let $y = x^3 - 8x + 7$ and $x = f(t)$ if $\frac{f(dy)}{dx} = 2$ and $x = 3a = 0$, then find the value of $\frac{dx}{dt} a = 0$.

 [Watch Video Solution](#)

128. Find the derivative of $\frac{\sqrt{x}(x+4)^{\frac{3}{2}}}{(4x-3)^{\frac{4}{3}}}$

 [Watch Video Solution](#)

129. If $x^m y^n = (x+y)^{m+n}$, provethat $\frac{dy}{dx} = \frac{y}{x}$.

 [Watch Video Solution](#)

130. Differentiate $(\log x)^{\cos x}$ with respect to x .

 [Watch Video Solution](#)

131. If $f(x) = |x|^{|\sin x|}$, then find $f' \left(-\frac{\pi}{4} \right)$

 [Watch Video Solution](#)

132. If $y = x^x \wedge x \wedge (((((\infty))))))$, find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

133. If $f(x) = \lim_{h \rightarrow 0} \frac{(\sin(x+h))^{\log_e(x+h)} - (\sin x)^{\log_e x}}{h}$ then find $f(\pi/2)$.

 [Watch Video Solution](#)

134. If $x < 1$, provethat $\frac{1}{1+x} + \frac{2x}{1+x^2} + \frac{4x^3}{1+x^4} + \infty = \frac{1}{1-x}$

 [Watch Video Solution](#)

135. Differentiate $\log \sin x$ w.r.t. \sqrt{x} .

 [Watch Video Solution](#)

136. Find the derivative of $f(\tan x) \text{ wrtg}(\sec x) \text{ at } x = \frac{\pi}{4}$, where $f'(1) = 2$ and $g'(\sqrt{2}) = 4$.

 [Watch Video Solution](#)

137.

Let $f(x) = \begin{vmatrix} \cos x & \sin x & \cos x \\ \cos 2x & \sin 2x & 2\cos 2x \\ \cos 3x & \sin 3x & 3\cos 3x \end{vmatrix}$ Then find the value of $f(0)$ and $f(\pi/2)$.

 [Watch Video Solution](#)

138. $f(x) = \begin{vmatrix} \cos x & x & 1 \\ 2\sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix}$ then find the value of

$$\lim_{x \rightarrow 0} \frac{f(x)}{x}$$

 [Watch Video Solution](#)

139. If $y = \cos^1 x$, find $\frac{d^2y}{dx^2}$ in terms of y alone.

 [Watch Video Solution](#)

140. if $y = (x^2 - 1)^m$, then the $(2m)$ th differential coefficient of y is

 [Watch Video Solution](#)

141. If $y = x \log \left\{ \frac{x}{(a + bx)} \right\}$, then show that $x^3 \frac{d^2y}{dx^2} = \left(x \frac{dy}{dx} - y \right)^2$.

 [Watch Video Solution](#)

142. If $(x - a)^2 + (y - b)^2 = c^2$, for some $c > 0$, prove that $\frac{\left[1 + \left(\frac{dy}{dx} \right)^2 \right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$ is a

constant independent of a and b .

 [Watch Video Solution](#)

143. If $y = e^a \cos^{(-1)x}$, $-1 \leq x < 1$, show that $(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - a^2y = 0$

 [Watch Video Solution](#)

144. If $x = a(\cos t + t \sin t)$ and $y = a(\sin t - t \cos t)$, find $\frac{d^2y}{dx^2}$

 [Watch Video Solution](#)

145. If g is inverse of f then prove that $f'(g(x)) = -g'(x)(f'(g(x)))^3$.

 [Watch Video Solution](#)

146. Let $f(x)$ and $g(x)$ be real valued functions such that $f(x)g(x) = 1$,

$\forall x \in R$. If $f'(x)$ and $g'(x)$ exists $\forall x \in R$ and $f'(x)$ and $g'(x)$

are never zero, then prove that $\frac{f'(x)}{f(x)} - \frac{g'(x)}{g(x)} = \frac{2f'(x)}{f(x)}$

 [Watch Video Solution](#)

147. If $f(x), g(x)$ and $h(x)$ are three polynomial of degree 2, then prove that $\varphi(x) = |f(x)g(x)h(x)f'(x)g'(x)h'(x)f''(x)g''(x)h''(x)|$ is a constant polynomial.

 Watch Video Solution

148. Let $f\left(\frac{x+y}{2}\right) = \frac{f(x)+f(y)}{2}$ or all real x and y . If $f'(0)$ exists and equals -1 and $f(0) = 1$, then $f \in C^1(2)$.

 Watch Video Solution

149.

If $f(x) + f(y) = f\left(\frac{x+y}{1-xy}\right)$ for all $x, y \in R, (xy \neq 1)$, and $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 2$ then

 Watch Video Solution

150. Find function $f(x)$ which is differentiable and satisfy the relation

$$f(x+y) = f(x) + f(y) + (e^x - 1)(e^y - 1) \forall x, y \in \mathbb{R}, \text{ and } f(0) = 2.$$

 [Watch Video Solution](#)

151. If $f\left(\frac{x}{y}\right) = \frac{f(x)}{f(y)}$, $\forall y, f(y) \neq 0$ and $f(1) = 2$, find $f(x)$.

 [Watch Video Solution](#)

152. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function satisfying condition

$$f(x+y^3) = f(x) + [f(y)]^3 \text{ for all } x, y \in \mathbb{R} \text{ If } f'(0) \geq 0, \text{ find } f(10)$$

 [Watch Video Solution](#)

153. Let $f(x+y) = f(x) + f(y) + 2xy - 1$ for all real x and y and $f(x)$ be a differentiable function. If $f'(0) = \cos \alpha$, then prove that $f(x) > 0 \forall x \in \mathbb{R}$



[Watch Video Solution](#)

[Watch Video Solution](#)

154. If $f(x) = \log_x(\log x)$, then find $f'(x)$ at $x = e$

[Watch Video Solution](#)

155. Given that $\frac{\cos x}{2} \cdot \frac{\cos x}{4} \cdot \frac{\cos x}{8} \dots = \frac{\sin x}{x}$ Then find the sum $\frac{1}{2^2} \frac{\sec^2 x}{2} + \frac{1}{2^4} \frac{\sec^2 x}{4} + \dots$

[Watch Video Solution](#)

156. If $y = f(a^x)$ and $f'(\sin x) = (\log)_e x$, then $f \in d \frac{dy}{dx}$, if it exists, where $\pi/2$

[Watch Video Solution](#)

157. If P_n is the sum of a GP upto n terms ($n \geq 3$), then prove that

$$(1 - r) \frac{dP_n}{dr} = (1 - n)P_n + nP_{n-1}, \text{ where } r \text{ is the common ratio of } GP$$



Watch Video Solution

158. If $x = \operatorname{cosec}\theta - \sin\theta$ and $y = \operatorname{cosec}^n\theta - \sin^n\theta$, then show that

$$(x^2 + 4) \left(\frac{dy}{dx} \right)^2 = n^2 (y^2 + 4)$$



Watch Video Solution

159. If $y = \frac{ax^2}{(x-a)(x-b)(x-c)} + \frac{bx}{(x-b)(x-c)} + \frac{c}{x-c} + 1$, then prove that

$$\frac{y'}{y} = \frac{1}{x} \left[\frac{a}{a-x} + \frac{b}{b-x} + \frac{c}{c-x} \right]$$



Watch Video Solution

160. Find the differential equation of the family of curves $y = Ae^{2x} + Be^{-2x}$, where A and B are arbitrary constants.



Watch Video Solution

161. If $y = \left(\frac{1}{2}\right)^{n-1} \cos(n \cos^{-1} x)$, then prove that y satisfies the differential equation $(1 - x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + n^2 y = 0$

 [Watch Video Solution](#)

162. If a curve is represented parametrically by the equation $x = f(t)$ and $y = g(t)$ then prove that $\frac{d^2 y}{dx^2} = - \left[\frac{g'(t)}{f'(t)} \right]^3 \left(\frac{d^2 x}{dy^2} \right)$

 [Watch Video Solution](#)

163. If $f\left(\frac{x+y}{3}\right) = \frac{2+f(x)+f(y)}{3}$ for all real x and y and $f'(2) = 2$, then determine $y = f(x)$.

 [Watch Video Solution](#)

164. If $f(x) = \frac{f(x)}{y} + \frac{f(y)}{x}$ holds for all real x and y greater than 0 and $f(x)$ is a differentiable function for all $x > 0$ such that $f(e) = \frac{1}{e}$, then $f \in df(x)$.

 [Watch Video Solution](#)

165. If $|a_1 \sin x + a_2 \sin 2x + \dots + a_n \sin nx| \leq |\sin x|$ for $x \in R$, then prove that $|a_1 + 2a_2 + 3a_3 + \dots + na_n| \leq 1$.

 [Watch Video Solution](#)

166. Suppose $p(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$. If $|p(x)| \leq e^{x-1} - 1$ for all $x \geq 0$, prove that $|a_1 + 2a_2 + \dots + na_n| \leq 1$.

 [Watch Video Solution](#)

1. Using the definition of derivative find the derivative of $\sqrt{\sin x}$

 [Watch Video Solution](#)

2. Find the derivative of $\sqrt{4-x}$ w.r.t. x using the first principle.

 [Watch Video Solution](#)

3. Let $f(x) = \frac{(2^x + 2^{-x}) \sin x \sqrt{\tan^{-1}(x^2 - x + 1)}}{(7x^2 + 3x + 1)^3}$. Then find the value of $f'(0)$.

 [Watch Video Solution](#)

4. Statement 1: Let $f: \vec{RR}$ be a real-valued function $\forall x, y \in R$ such that $|f(x) - f(y)| \leq |x - y|^3$. Then $f(x)$ is a constant function. Statement 2: If the derivative of the function w.r.t. x is zero, then function is constant.

 [Watch Video Solution](#)

5. Find the derivative of $\sqrt{4-x}$ w.r.t. x using the first principle.

 [Watch Video Solution](#)

6. Let $f(x) = \frac{(2^x + 2^{-x}) \sin x \sqrt{\tan^{-1}(x^2 - x + 1)}}{(7x^2 + 3x + 1)^3}$. Then find the value of $f'(0)$.

 [Watch Video Solution](#)

7. Statement 1: Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a real-valued function $\forall x, y \in \mathbb{R}$ such that $|f(x) - f(y)| \leq |x - y|^3$. Then $f(x)$ is a constant function. Statement 2: If the derivative of the function w.r.t. x is zero, then function is constant.

 [Watch Video Solution](#)

1. If $y = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$, then find $\frac{dy}{dx}$

 [Watch Video Solution](#)

2. $y = \tan^{-1}\frac{3x-x^3}{2x^2-1}$, $-\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$

 [Watch Video Solution](#)

3. If $y = \sec^{-1}\left(\frac{1}{2x^2-1}\right)$; 0

 [Watch Video Solution](#)

4. Find $\frac{dy}{dx}$ if $y = \frac{\tan^{-1}(4x)}{1+5x^2} + \frac{\tan^{-1}(2+3x)}{3-2x}$

 [Watch Video Solution](#)

5. Find $\frac{dy}{dx}$ if $y = \tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$, where $x \neq 0$

 [Watch Video Solution](#)

6. $y = \tan^{-1}\left(\frac{x}{1 + \sqrt{1-x^2}}\right)$

 [Watch Video Solution](#)

7. Find $\frac{dy}{dx}$ for the function: $y = \sin^{-1}\sqrt{1-x} + \cos^{-1}\sqrt{x}$

 [Watch Video Solution](#)

8. $y = \sqrt{\sin\sqrt{x}}$

 [Watch Video Solution](#)

9. $y = e^{\sin x^3}$

 [Watch Video Solution](#)

10. $y = \log \sqrt{\sin \sqrt{e^x}}$

 [Watch Video Solution](#)

11. Find $\frac{dy}{dx}$ for the function: $y = a^{\sin^{-1}x} \wedge 2$

 [Watch Video Solution](#)

12. Find $\frac{dy}{dx}$ if $y = \log \left\{ e^x \left(\frac{x-2}{x+2} \right)^{\frac{3}{4}} \right\}$

 [Watch Video Solution](#)

13. $y = \sin^{-1}[\sqrt{x - ax} - \sqrt{a - ax}]$



Watch Video Solution

14. Find $\frac{dy}{dx}$ for the functions: $y = x^3 e^x \sin x$



Watch Video Solution

15. Find $\frac{dy}{dx}$ for the function: $y = (\log)_e \sqrt{\frac{1 + \sin x}{1 - \sin x}}$, where $x = \frac{\pi}{3}$



Watch Video Solution

16. Find $\frac{dy}{dx}$ for the functions: $y = \frac{x + \sin x}{x + \cos x}$



Watch Video Solution

17. If $y = (1 + x)(1 + x^2)(1 + x^4) \dots (1 + x^{2^n})$ then $\frac{dy}{dx}$ at $x = 0$ is

 [Watch Video Solution](#)

18. $x\sqrt{1+y} + y\sqrt{1+x} = 0$ then $\frac{dy}{dx} =$

 [Watch Video Solution](#)

19. If g is the inverse function of and $f'(x) = \sin x$ then prove that $g'(x) = \operatorname{cosec}(g(x))$

 [Watch Video Solution](#)

20. $y = \sin^{-1} \frac{2x}{1+x^2}, -1 \leq x \leq 1$

 [Watch Video Solution](#)

$$21. y = \tan^{-1} \left[\frac{3x - x^3}{1 - 3x^2} \right], -\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$$

 [Watch Video Solution](#)

$$22. y = \sec^{-1} \frac{1}{2x^2 - 1}, 0 < x < \frac{1}{\sqrt{2}}$$

 [Watch Video Solution](#)

$$23. \text{ Find } \frac{dy}{dx} \text{ if } y = \frac{\tan^{-1}(4x)}{1 + 5x^2} + \frac{\tan^{-1}(2 + 3x)}{3 - 2x}$$

 [Watch Video Solution](#)

$$24. \text{ Find } \frac{dy}{dx} \text{ if } y = \tan^{-1} \left(\frac{\sqrt{1 + x^2} - 1}{x} \right), \text{ where } x \neq 0$$

 [Watch Video Solution](#)

$$25. y = \tan^{-1} \left(\frac{x}{1 + \sqrt{1 - x^2}} \right)$$

 [Watch Video Solution](#)

$$26. \text{ Find } \frac{dy}{dx} \text{ for the function: } y = \sin^{-1} \sqrt{1-x} + \cos^{-1} \sqrt{x}$$

 [Watch Video Solution](#)

$$27. y = \sqrt{\sin \sqrt{x}}$$

 [Watch Video Solution](#)

$$28. y = e^{\sin x^3}$$

 [Watch Video Solution](#)

$$29. y = \log \sqrt{\sin \sqrt{e^x}}$$



Watch Video Solution

$$30. \text{ Find } \frac{dy}{dx} \text{ for the function: } y = a^{\sin^{-1}x} \wedge 2$$



Watch Video Solution

$$31. \text{ Find } \frac{dy}{dx} \text{ if } y = \log \left\{ e^x \left(\frac{x-2}{x+2} \right)^{\frac{3}{4}} \right\}$$



Watch Video Solution

$$32. y = \sin^{-1} \left[\sqrt{x-ax} - \sqrt{a-ax} \right]$$



Watch Video Solution

33. Find $\frac{dy}{dx}$ for the functions: $y = x^3 e^x \sin x$

 [Watch Video Solution](#)

34. Find $\frac{dy}{dx}$ for the function: $y = (\log)_e \sqrt{\frac{1 + \sin x}{1 - \sin x}}$, where $x = \frac{\pi}{3}$

 [Watch Video Solution](#)

35. $y = \frac{x + \sin x}{x + \cos x}$ find $\frac{dy}{dx}$

 [Watch Video Solution](#)

36. If $y = (1 + x)(1 + x^2)(1 + x^4) \dots (1 + x^{2^n})$ then $\frac{dy}{dx}$ at $x = 0$ is

 [Watch Video Solution](#)

37. If $x\sqrt{1+y} + y\sqrt{1+x} = 0$, prove that $\frac{dy}{dx} = -\frac{1}{(x+1)^2}$.



Watch Video Solution

38. If g is the inverse function of $f(x) = \sin x$ then prove that $g'(x) = \operatorname{cosec}(g(x))$



Watch Video Solution

Exercise 3.3

1. If $x^3 + y^3 = 3axy$, find $\frac{dy}{dx}$



Watch Video Solution

2. If $y = b \tan^{-1} \left(\frac{x}{a} + \frac{\tan^{-1} y}{x} \right)$, $f \in d \frac{dy}{dx}$.



Watch Video Solution

3. If $\log_e(\log_e x - \log_e y) = e^{x^2 y}(1 - \log_e x)$, then find the value of $y'(e)$.

 **Watch Video Solution**

4. If $y = \sqrt{x + \sqrt{y + \sqrt{x + \sqrt{y + \dots}}}}$ then prove that $\frac{dy}{dx} = \frac{y^2 - x}{2y^3 - 2xy - 1}$

 **Watch Video Solution**

5. If $x^3 + y^3 = 3axy$, find $\frac{dy}{dx}$.

 **Watch Video Solution**

6. If $y = b \tan^{-1} \left(\frac{x}{a} + \frac{\tan^{-1} y}{x} \right)$, $f \in d \frac{dy}{dx}$.

 **Watch Video Solution**

7. If $\log_e(\log_e x - \log_e y) = e^{x^2 y}(1 - \log_e x)$, then find the value of $y'(e)$.

 [Watch Video Solution](#)

8. If $y = \sqrt{x + \sqrt{y + \sqrt{x + \sqrt{y + \dots}}}}$ then prove that $\frac{dy}{dx} = \frac{y^2 - x}{2y^3 - 2xy - 1}$

 [Watch Video Solution](#)

Exercise 3.4

1. If $x = \frac{2t}{1+t^2}$, $y = \frac{1-t^2}{1+t^2}$, then $f \in d \frac{dy}{dx} a = 2$.

 [Watch Video Solution](#)

2. If $x = \sqrt{a^{\sin^{-1} t}}$, $y = \sqrt{a^{\cos^{-1} t}}$ then show that, $\frac{dy}{dx} = -\frac{y}{x}$.

 [Watch Video Solution](#)

3. Find $\frac{dy}{dx}$ if $x = 3 \cos \theta - \cos 2\theta$ and $y = \sin \theta - \sin 2\theta$.

 [Watch Video Solution](#)

4. If $x = 3\cos\theta - 2\cos^3\theta$, $y = 3\sin\theta - 2\sin^3\theta$, then $\frac{dy}{dx}$ is

 [Watch Video Solution](#)

5. If $x = a \left(\cos t + \frac{1}{2} \log \tan^2 t \right)$ and $y = a \sin t$ then find $\frac{dy}{dx}$ at $t = \frac{\pi}{4}$

 [Watch Video Solution](#)

6. If $x = \frac{2t}{1+t^2}$, $y = \frac{1-t^2}{1+t^2}$, then find $\frac{dy}{dx}$ at $t = 2$.

 [Watch Video Solution](#)

7. If $x = \sqrt{a^{\sin^{-1}((-1)t)}}$, $y = \sqrt{a^{\cos^{-1}((-1)t)}}$, show that $\frac{dy}{dx} = -\frac{y}{x}$

 [Watch Video Solution](#)

8. Find $\frac{dy}{dx}$ if $x = \cos\theta - \cos 2\theta$

and $y = \sin\theta - \sin 2\theta$

 [Watch Video Solution](#)

9. Find $\frac{dy}{dx}$ if $x = 3\cos\theta - 2\cos^3\theta$, $y = 3\sin\theta - 2\sin^3\theta$.

 [Watch Video Solution](#)

10. If $x = a\left(\cos t + \frac{1}{2}\log \tan^2 t\right)$ and $y = a \sin t$, then find $\frac{dy}{dx}$ at $t = \frac{\pi}{4}$

 [Watch Video Solution](#)

Exercise 3.5

1. Differentiate $\sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}}$ with respect to x

 [Watch Video Solution](#)

2. If $x^y = e^{x-y}$, prove that $\frac{dy}{dx} = \frac{\log x}{(1 + \log x)^2}$

 [Watch Video Solution](#)

3. If $xy = e^{(x-y)}$, then find $\frac{dy}{dx}$

 [Watch Video Solution](#)

4. If $y^x = x^y$, then $f \in d \frac{dy}{dx}$

 [Watch Video Solution](#)

5. If $x = e^y + e^{(y \rightarrow \infty)}$, where $x > 0$, then $f \in d \frac{dy}{dx}$

 [Watch Video Solution](#)

6. Find $\frac{dy}{dx} f$ or $y = x^x$

 [Watch Video Solution](#)

7. Differentiate $(x \cos x)^x$ with respect to x

 [Watch Video Solution](#)

8. If $y = (\tan x)^{(\tan x)^{\tan x}}$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

9. Differentiate $\sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}}$ with respect to x

 [Watch Video Solution](#)

10. If $x^y = e^{x-y}$, prove that $\frac{dy}{dx} = \frac{\log x}{(1 + \log x)^2}$.

 [Watch Video Solution](#)

11. If $xy = e^{(x-y)}$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

12. If $y^x = x^y$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

13. If $x = e^y + e^{(y + \infty)}$, where $x > 0$, then $f \in d \frac{dy}{dx}$

 [Watch Video Solution](#)

14. Find $\frac{dy}{dx} f$ or $y = x^x$

 [Watch Video Solution](#)

15. Differentiate $(x \cos x)^x$ with respect to x

 [Watch Video Solution](#)

16. If $y = (\tan x)^{(\tan x)^{\tan x}}$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

1. Find the derivative of $\frac{\tan^{-1}(2x)}{1-x^2} \cdot \sin^{-1}(2x)$ wrt $\frac{1}{1+x^2}$

 [Watch Video Solution](#)

2. The differential coefficient of $\sec^{-1}\left(\frac{1}{2x^2-1}\right)$ w.r.t $\sqrt{1-x^2}$ is-

 [Watch Video Solution](#)

3. Differentiate $\frac{x}{\sin x}$ w.r.t $\sin x$.

 [Watch Video Solution](#)

4.

If $y = f(x^3)$, $z = g(x^5)$, $f(x) = \tan x$, and $g'(x) = \sec x$, then find the value of

 [Watch Video Solution](#)

5. Find the derivative of $\frac{\tan^{-1}(2x)}{1-x^2} \cdot \sin^{-1}(2x)$ wrt $\frac{1}{1+x^2}$

 [Watch Video Solution](#)

6. Find the derivative of $\sec^{-1}\left(\frac{1}{2x^2-1}\right)$ w.r.t. $\sqrt{1-x^2}$ at $x = \frac{1}{2}$.

 [Watch Video Solution](#)

7. Differentiate $\frac{x}{\sin x}$ w.r.t. $\sin x$.

 [Watch Video Solution](#)

8.

If $y = f(x^3)$, $z = g(x^5)$, $f'(x) = \tan x$, and $g'(x) = \sec x$, then find the value of

 [Watch Video Solution](#)

Exercise 3.7

1. If $f(x) = \begin{vmatrix} x + a^2 & ab & ac \\ ab & x + b^2 & bc \\ ac & bc & x + c^2 \end{vmatrix}$, then prove that

$$f(x) = 3x^2 + 2x(a^2 + b^2 + c^2).$$

 [Watch Video Solution](#)

2. Let $f(x) = \begin{vmatrix} \cos(x + x^2) & \sin(x + x^2) & -\cos(x + x^2) \\ \sin(x - x^2) & \cos(x - x^2) & \sin(x - x^2) \\ \sin 2x & 0 & \sin(2x^2) \end{vmatrix}$.

Find the value of $f'(0)$.

 [Watch Video Solution](#)

3. Let $g(x) = \begin{vmatrix} f(x+c) & f(x+2c) & f(x+3c) \\ f(c) & f(2c) & f(3c) \\ f'(c) & f'(2c) & f'(3c) \end{vmatrix}$,

where c is constant, then find $\lim_{x \rightarrow 0} \frac{g(x)}{x}$.

 **Watch Video Solution**

4. If $f(x) = \begin{vmatrix} x+a^2 & ab & ac \\ ab & x+b^2 & bc \\ ac & bc & x+c^2 \end{vmatrix}$, then prove that

$$f'(x) = 3x^2 + 2x(a^2 + b^2 + c^2).$$

 **Watch Video Solution**

5. Let $f(x) = \begin{vmatrix} \cos(x+x^2) & \sin(x+x^2) & -\cos(x+x^2) \\ \sin(x-x^2) & \cos(x-x^2) & \sin(x-x^2) \\ \sin 2x & 0 & \sin(2x^2) \end{vmatrix}$.

Find the value of $f'(0)$.



Watch Video Solution

$$6. \text{ Let } g(x) = \begin{vmatrix} f(x+c) & f(x+2c) & f(x+3c) \\ f(c) & f(2c) & f(3c) \\ f'(c) & f'(2c) & f'(3c) \end{vmatrix},$$

where c is constant, then find $\lim_{x \rightarrow 0} \frac{g(x)}{x}$.



Watch Video Solution

Exercise 3.8

$$1. \text{ If } f(x) = (1+x)^2, \text{ then the value of } f(x0) + f'(0) + \frac{f''(0)}{2!} + \frac{f'''(0)}{3!} + \frac{f^{(n)}(0)}{n!}.$$



Watch Video Solution

$$2. \text{ If } e^y(x+1) = 1, \text{ show that } \frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2.$$



Watch Video Solution

3. Prove that $\frac{d^n}{dx^n} (e^{2x} + e^{-2x}) = 2^n [e^{2x} + (-1)^n e^{-2x}]$

 [Watch Video Solution](#)

4. If $y = \sin(\sin x)$ and $\frac{d^2y}{dx^2} + \frac{dy}{dx} \tan x + f(x) = 0$, then find $f(x)$.

 [Watch Video Solution](#)

5. If $y = \log(1 + \sin x)$, prove that $y_4 + y_3 y_1 + y_2^2 = 0$.

 [Watch Video Solution](#)

6. If $f(x) = \left| x^n \left[2 \cos x \frac{\cos(n\pi)}{2} - 4 \sin x \frac{\sin(n\pi)}{2} \right] 8 \right|$ then find the value of $\frac{d^n}{dx^n} ([f(x)])_{x=0} \in \mathbb{Z}$

 [Watch Video Solution](#)

7. If $x = a\cos\theta, y = b\sin\theta$, then prove that $\frac{d^3y}{dx^3} = \frac{3b}{a^3} \operatorname{cosec}^4\theta \cot\theta$.

 [Watch Video Solution](#)

8. If $x = a\cos^3\theta, y = b\sin^3\theta, f \in d\frac{d^3y}{dx^3}$ at $\theta = 0$.

 [Watch Video Solution](#)

9. If $f(x) = (1+x)^2$, then the value of $f(x_0) + f'(0) + \frac{f''(0)}{2!} + \frac{f'''(0)}{3!} + \frac{f^{(n)}(0)}{n!}$.

 [Watch Video Solution](#)

10. If $e^y(x+1) = 1$, show that $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$.

 [Watch Video Solution](#)

11. Prove that $\frac{d^n}{dx^n} (e^{2x} + e^{-2x}) = 2^n [e^{2x} + (-1)^n e^{-2x}]$

 [Watch Video Solution](#)

12. If $y \sin (\sin x)$ and $\frac{d^2 y}{dx^2} + \frac{dy}{dx} \tan x + f(x) = 0$, then find $f(x)$.

 [Watch Video Solution](#)

13. If $y = \log (1 + \sin x)$, prove that $y_4 + y_3 y_1 + y_2^2 = 0$.

 [Watch Video Solution](#)

14. If $x = a \cos \theta, y = b \sin \theta$, then prove that $\frac{d^3 y}{dx^3} = \frac{3b}{a^3} \operatorname{cosec}^4 \theta \cot \theta$.

 [Watch Video Solution](#)

15. If $x = a \cos^3 \theta, y = b \sin^3 \theta, f \in d \frac{d^3 y}{dx^3}$ at $\theta = 0$.



Watch Video Solution

Exercise 3.9

1. Let $f(x + y) = f(x)f(y)$ for all x and y . Suppose $f(5) = 2$ and $f'(0) = 3$. Find $f'(5)$.



Watch Video Solution

2. Let $f(xy) = f(x)f(y) \forall x, y \in \mathbb{R}$ and f is differentiable at $x = 1$ such that $f'(1) = 1$. Also, $f(1) \neq 0, f(2) = 3$. Then find $f'(2)$.



Watch Video Solution

3. Let f be a function such that

$f(x + y) = f(x) + f(y)$ for all x and y and $f(x) = (2x^2 + 3x)g(x)$ for all x , where g

is continuous and $g(0) = 3$. Then find $f'(x)$.



Watch Video Solution

4. Let $g: \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function satisfying $g(x) = g(y)g(x-y) \forall x, y \in \mathbb{R}$ and $g'(0) = a$ and $g'(3) = b$. Then find the value of $g'(-3)$.



Watch Video Solution

5. Let $f(x^m y^n) = mf(x) + nf(y)$ for all $x, y \in \mathbb{R}^+$ and for all $m, n \in \mathbb{R}$. If $f'(x)$ exists and has the value $\frac{e}{x}$, then find $(\lim_{x \rightarrow 0} \frac{f(1+x)}{x})$.



Watch Video Solution

6. If $f\left(\frac{x+2y}{3}\right) = \frac{f(x)+2f(y)}{3} \forall x, y \in \mathbb{R}$ and $f'(0) = 1, f(0) = 2$, then find $f(x)$.



Watch Video Solution

7.

Prove that $\lim_{h \rightarrow 0} \frac{f(x+h) + f(x-h) - 2f(x)}{h^2} = f''(x)$ (without using L' Hospital's



Watch Video Solution

8. Let $f(x+y) = f(x)f(y)$ for all x and y . Suppose $f(5) = 2$ and $f'(0) = 3$. Find $f'(5)$.



Watch Video Solution

9. Let $f(xy) = f(x)f(y) \forall x, y \in \mathbb{R}$ and f is differentiable at $x = 1$ such that $f'(1) = 1$. Also, $f(1) \neq 0, f(2) = 3$. Then find $f'(2)$.



Watch Video Solution

10. Let f be a function such that $f(x+y) = f(x) + f(y)$ for all x and y and $f(x) = (2x^2 + 3x)g(x)$ for all x , where $g(x)$ is continuous and $g(0) = 3$, then

find $f'(x)$.



[Watch Video Solution](#)

11. Let $g: \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function satisfying $g(x) = g(y)g(x-y) \forall x, y \in \mathbb{R}$ and $g'(0) = a$ and $g'(3) = b$. Then find the value of $g'(-3)$.



[Watch Video Solution](#)

12. Let $f(x^m y^n) = mf(x) + nf(y)$ for all $x, y \in \mathbb{R}^+$ and for all $m, n \in \mathbb{R}$. If $f'(x)$ exists and has the value $\frac{e}{x}$, then find $(\lim_{x \rightarrow 0} \frac{f(1+x)}{x})$.



[Watch Video Solution](#)

13. If $f\left(\frac{x+2y}{3}\right) = \frac{f(x)+2f(y)}{3} \forall x, y \in \mathbb{R}$ and $f'(0) = 1, f(0) = 2$, then find $f(x)$.



Watch Video Solution

14. Prove that

$$\lim_{h \rightarrow 0} \frac{f(x+h) + f(x-h) - 2f(x)}{h^2} = f''(x) \text{ (without using L' Hospital's rule).}$$



Watch Video Solution

Exercise (Single)

1. If $y = a\sin x + b\cos x$, then $\left(\frac{dy}{dx}\right)^2 + y^2$ is

- A. function of x
- B. function of y
- C. function of x and y
- D. constant



Watch Video Solution

2. $\frac{d}{dx} \sqrt{\frac{1-\sin 2x}{1+\sin 2x}}$ is equal to,

A. $\sec^2 x$

B. $-\sec^2\left(\frac{\pi}{4} - x\right)$

C. $\sec^2\left(\frac{\pi}{4} + x\right)$

D. $\sec^2\left(\frac{\pi}{4} - x\right)$



Watch Video Solution

3. If $f(x) = |\cos x| + |\sin x|$, then $\frac{dy}{dx}$ at $x = \frac{2\pi}{3}$ is equal to

A. $\frac{1 - \sqrt{3}}{2}$

B. 0

C. $\frac{1}{2}(\sqrt{3} - 1)$

D. none of these



Watch Video Solution

4. If $f(x) = \left| \log_e |x| \right|$, then $f'(x)$ equals

A. $\frac{1}{|x|}$, where $x \neq 0$

B. $\frac{1}{x}$ for $|x| > 1$ and $-\frac{1}{x}$ for $|x| < 1$

C. $-\frac{1}{x}$ for $|x| > 1$ and $\frac{1}{x}$ for $|x| < 1$

D. $\frac{1}{x}$ for $x > 0$ and $-\frac{1}{x}$ for $x < 0$



Watch Video Solution

5. If $f(x) = \sqrt{1 - \sin 2x}$, then $f'(x)$ is equal to

A. $-(\cos x + \sin x)$, for $x \in (\pi/4, \pi/2)$

B. $\cos x + \sin x$ for $x \in (0, \pi/4)$

C. $-(\cos x + \sin x)$, for $x \in (0, \pi/4)$

D. $\cos x - \sin x$, for $x \in (\pi/4, \pi/2)$

Answer: C



Watch Video Solution

6. Instead of the usual definition of derivative $Df(x)$, if we define a new kind of derivative $D^*F(x)$ by the formula

$$D^*(x) = \lim_{h \rightarrow 0} \frac{f^2(x+h) - f^2(x)}{h}. \text{ where } f^2(x) \text{ means } [f(x)]^2 \text{ and if } f(x) = x$$

$\log x$, then $D^*f(x)|_{x=e}$ has the value

A. e

B. $2e$

C. $4e$

D. none of these



Watch Video Solution

7. If $y = \cot^{-1} \left[\frac{\sqrt{1 + \sin x} + \sqrt{1 - \sin x}}{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}} \right]$ ($0 < x < \pi/2$), then $\frac{dy}{dx} =$

A. $\frac{1}{2}$

B. $\frac{2}{3}$

C. 3

D. 1



Watch Video Solution

8. if $y = \frac{(a-x)\sqrt{a-x} - (b-x)\sqrt{x-b}}{\sqrt{a-x} + \sqrt{x-b}}$ then $\frac{dy}{dx}$ wherever it is defined is equal

to:

A. $\frac{x + (a + b)}{\sqrt{(a - x)(x - b)}}$

B. $\frac{2x - a - b}{2\sqrt{a - x}\sqrt{x - b}}$

C. $-\frac{(a + b)}{2\sqrt{(a - x)(x - b)}}$

D. $\frac{2x + (a + b)}{2\sqrt{(a - x)(x - b)}}$

Answer: B

 [Watch Video Solution](#)

9. the derivative of $y = (1 - x)(2 - x)\dots\dots\dots (n - x)$ at $x = 1$ is equal to

A. 0

B. $(-1)(n - 1)!$

C. $n! - 1$

D. $(-1)^{n-1}(n - 1)!$

 [Watch Video Solution](#)

10. If $y = \sqrt{\frac{1-x}{1+x}}$, then $(1-x^2)\frac{dy}{dx}$ is equal to

A. y^2

B. $1/y$

C. $-y$

D. $-y/x$



Watch Video Solution

11. If $y = \frac{\sqrt{a+x} - \sqrt{a-x}}{\sqrt{a+x} + \sqrt{a-x}}$, then $\frac{dy}{dx}$ is equal to (a) $\frac{ay}{x\sqrt{a^2-x^2}}$ (b) $\frac{ay}{\sqrt{a^2-x^2}}$

(c) $\frac{ay}{x\sqrt{a^2-x^2}}$ (d) none of these

A. $\frac{ay}{x\sqrt{a^2-x^2}}$

B. $\frac{ay}{\sqrt{a^2-x^2}}$

C. $\frac{ay}{x\sqrt{x^2-a^2}}$

D. none of these



Watch Video Solution

12. Let $u(x)$ and $v(x)$ be differentiable functions such that

$\frac{u(x)}{v(x)} = 7$. if $\frac{u'(x)}{v'(x)} = p$ and $\left(\frac{u(x)}{v(x)}\right)' = q$, then $\frac{p+q}{p-q}$ has the value equal

to

A. 1

B. 0

C. 7

D. -7



Watch Video Solution

13. If $\sin^{-1}\left(\frac{x^2 - y^2}{x^2 + y^2}\right) = \log a$, then $\frac{dy}{dx}$ is equal to $\frac{x}{y}$ (b) $\frac{y}{x^2} \frac{x^2 - y^2}{x^2 + y^2}$ (d) $\frac{y}{x}$

A. $\frac{x}{y}$

B. $\frac{y}{x^2}$

C. $\frac{x^2 - y^2}{x^2 + y^2}$

D. $\frac{y}{x}$



Watch Video Solution

14. Let $h(x)$ be differentiable for all x and let $f(x) = (kx + e^x)h(x)$, where k is some constant. If $h(0) = 5$, $h'(0) = -2$, and $f'(0) = 18$, then the value of k is 5 (b) 4 (c) 3 (d) 2.2.

A. 5

B. 4

C. 3

D. 2.2



Watch Video Solution

15. If $\lim_{t \rightarrow x} \frac{e^t f(x) - e^x f(t)}{(t-x)(f(x))^2} = 2$ and $f(0) = \frac{1}{2}$, then find the value of $f'(0)$.

A. 4

B. 2

C. 0

D. 1



Watch Video Solution

16. If $f(0) = 0, f'(0) = 2$ then the derivative of $y = f(f(f(x)))$ at $x = 0$ is

A. 2

B. 8

C. 16

D. 4

 [Watch Video Solution](#)

17. If $f(x) = \sqrt{1 + \cos^2(x^2)}$, then $f' \left(\frac{\sqrt{\pi}}{2} \right)$ is $\frac{\sqrt{\pi}}{6}$ (b) $-\sqrt{\pi/6}$ $1/\sqrt{6}$ (d) $\pi/\sqrt{6}$

A. $\sqrt{\pi/6}$

B. $-\sqrt{\pi/6}$

C. $1/\sqrt{6}$

D. $\pi/\sqrt{6}$

 [Watch Video Solution](#)

18. $\frac{d}{dx} \cos^{-1} \sqrt{\cos x}$ is equal to

A. $\frac{1}{2} \sqrt{1 + \sec x}$

B. $\sqrt{1 + \sec x}$

C. $-\frac{1}{2} \sqrt{1 + \sec x}$

D. $-\sqrt{1 + \sec x}$



Watch Video Solution

19. if $y = \tan^{-1} \left(\frac{2^x}{1 + 2^{2x+1}} \right)$ then $\frac{dy}{dx} \text{ at } x = 0$ is

A. 1

B. 2

C. $\ln 2$

D. $-\frac{1}{10} \ln 2$



Watch Video Solution

20. If $y = \left(x + \sqrt{x^2 + a^2}\right)^n$, then $\frac{dy}{dx}$ is $\frac{ny}{\sqrt{x^2 + a^2}}$ (b) $-\frac{ny}{\sqrt{x^2 + a^2}} - \frac{nx}{\sqrt{x^2 + a^2}}$ (d)

$$-\frac{nx}{\sqrt{x^2 + a^2}}$$

A. $\frac{ny}{\sqrt{x^2 + a^2}}$

B. $-\frac{ny}{\sqrt{x^2 + a^2}}$

C. $\frac{nx}{\sqrt{x^2 + a^2}}$

D. $-\frac{nx}{\sqrt{x^2 + a^2}}$



Watch Video Solution

21. if $y = \log_{\sin x} \tan x$ then $\left(\frac{dy}{dx}\right)_{\frac{\pi}{4}}$ is

A. $\frac{4}{\log 2}$

B. $-4\log 2$

C. $\frac{-4}{\log 2}$

D. none of these

 [Watch Video Solution](#)

22. $\frac{d}{dx} \left[\sin^2 \cot^{-1} \sqrt{\frac{1-x}{1+x}} \right]$ is

A. -1

B. $\frac{1}{2}$

C. $-\frac{1}{2}$

D. 1

 [Watch Video Solution](#)

23. The differential coefficient of $f(\log_e x)$ w. r. t. x , where $f(x) = \log_e x$, is (i)

$\frac{x}{\ln x}$ (ii) $\frac{\ln x}{x}$ (iii) $\frac{1}{x \ln x}$ (iv) $x \ln x$

A. $\frac{x}{\log_e x}$

B. $\frac{1}{x} \log_e x$

C. $\frac{1}{x \log_e x}$

D. none of these



Watch Video Solution

24. if $f(x) = \sqrt{2x^2 - 1}$ and $y = f(x^2)$ then $\frac{dy}{dx}$ at $x = 1$ is:

A. 2

B. 1

C. -2

D. none of these



Watch Video Solution

25. If $u = f(x^3)$, $v = g(x^2)$, $f'(x) = \cos x$, and $g'(x) = \sin x$, then $\frac{du}{dv}$ is

$\frac{3}{2}x \cos x^3 \operatorname{cosec} x^2$ $\frac{2}{3} \sin x^3 \sec x^2 \tan x$ (d) none of these

A. $\frac{3}{2}x \cos x^2 \operatorname{cosec} x^2$

B. $\frac{3}{2} \sin x^3 \sec x^2$

C. $\tan x$

D. none of these



Watch Video Solution

26. A function f , defined for all positive real numbers, satisfies the equation $f(x^2) = x^3$ for every $x > 0$. Then the value of $f'(4)$ is 12 (b) 3 (c)

3/2 (d) cannot be determined

A. 12

B. 3

C. 3/2

D. cannot be determined



Watch Video Solution

27. Let $f: (-5, 5) \rightarrow \mathbb{R}$ be a differentiable function of with $f(4) = 1, f'(4) = 1, f(0) = -1$ and $f'(0) = 1$. If $g(x) = \left(f(2f^2(x) + 2) \right)^2$, then $g'(0)$ equals

A. 4

B. -4

C. 8

D. -8



Watch Video Solution

28. The function $f(x) = e^x + x$, being differentiable and one-to-one, has a differentiable inverse $f^{-1}(x)$. The value of $\frac{d}{dx}(f^{-1})$ at the point $f(\log 2)$ is

$\frac{1}{\ln 2}$ (b) $\frac{1}{3}$ (c) $\frac{1}{4}$ (d) none of these

A. $\frac{1}{\ln 2}$

B. $\frac{1}{3}$

C. $\frac{1}{4}$

D. none of these



Watch Video Solution

29. If $f(x) = x + \tan x$ and f is the inverse of g , then $g'(x)$ is equal to

A. $\frac{1}{1 + [g(x) - x]^2}$

B. $\frac{1}{2 - [g(x) - x]^2}$

C. $\frac{1}{2 + [g(x) - x]^2}$

D. none of these

Answer: C



Watch Video Solution

30. If $f(x) = x^3 + 3x + 4$ and g is the inverse function of $f(x)$, then the value

of $\frac{d}{dx} \left(\frac{g(x)}{g(g(x))} \right)$ at $x = 4$ equals

A. $\frac{-1}{6}$

B. 6

C. $\frac{-1}{3}$

D. non-existent



Watch Video Solution

31. If $y = \frac{\sin^{-1}x}{\sqrt{1-x^2}}$, then $\frac{(1-x^2)dy}{dx}$ is equal to $x+y$ (b) $1+xy$ $1-xy$ (d) $xy-2$

A. $x+y$

B. $1+xy$

C. $1-xy$

D. $xy-2$



Watch Video Solution

32. If $y = \sqrt{\log x + \sqrt{\log x + \sqrt{\log x + \dots \infty}}}$, then $\frac{dy}{dx}$ is

A. $\frac{x}{2y-1}$

B. $\frac{x}{2y+1}$

C. $\frac{1}{x(2y-1)}$

D. $\frac{1}{x(1-2y)}$



Watch Video Solution

$$33. \frac{d}{dx} \left[\tan^{-1} \left(\frac{\sqrt{x}(3-x)}{1-3x} \right) \right] = \frac{1}{2(1+x)\sqrt{x}} \quad (b) \quad \frac{3}{(1+x)\sqrt{x}} \quad \frac{2}{(1+x)\sqrt{x}} \quad (d)$$

$$\frac{3}{2(1+x)\sqrt{x}}$$

A. $\frac{1}{2(1+x)\sqrt{x}}$

B. $\frac{3}{(1+x)\sqrt{x}}$

C. $\frac{2}{(1+x)\sqrt{x}}$

D. $\frac{3}{2(1+x)\sqrt{x}}$



Watch Video Solution

34. Suppose the function $f(x) - f(2x)$ has the derivative 5 at $x = 1$ and derivative 7 at $x = 2$. The derivative of the function $f(x) - f(4x)$ at $x = 1$ has the value equal to 19 (b) 9 (c) 17 (d) 14

A. 19

B. 9

C. 17

D. 14

 [Watch Video Solution](#)

35. If $y\sqrt{x^2 + 1} = \log(\sqrt{x^2 + 1} - x)$, show that $(x^2 + 1)\frac{dy}{dx} + xy + 1 = 0$

A. 0

B. 1

C. 2

D. none of these

 [Watch Video Solution](#)

36.

Let $e^y = \frac{\sqrt{1+\alpha} + \sqrt{1-\alpha}}{\sqrt{1+\alpha} - \sqrt{1-\alpha}}$ and $\tan \frac{x}{2} = \sqrt{\frac{1-\alpha}{1+\alpha}}$, $\alpha \in [-1, 0] \cup (0, 1]$. Then $\left(\frac{dy}{dx}\right)$

A. 1/2

B. 1

C. 2

D. 1/3



Watch Video Solution

37. The derivative of $\tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$ with respect to

$\tan^{-1}\left(\frac{2x\sqrt{1-x^2}}{1-2x^2}\right)$ at $x = 0$ is $\frac{1}{8}$ (b) $\frac{1}{4}$ (c) $\frac{1}{2}$ (d) 1

A. 1/8

B. $1/4$

C. $1/2$

D. 1



Watch Video Solution

38. If $\ln((e - 1)e^{xy} + x^2) = x^2 + y^2$ then $\left(\frac{dy}{dx}\right)_{1,0}$ is equal to

A. 0

B. 1

C. 2

D. 3



Watch Video Solution

39. If $y = (x^x)^x$ then $\frac{dy}{dx}$ is

A. $y \left[x^x (\log_e x) \log x + x^x \right]$

B. $y \left[x^x (\log_e x) \log x + x \right]$

C. $y \left[x^x (\log_e x) \log x + x^{-1} \right]$

D. $y \left[x^x (\log_e x) \log x + x^{-1} \right]$



Watch Video Solution

40. The first derivative of the function $\left[\cos^{-1} \left(\sin \sqrt{\frac{1+x}{2}} \right) + x^x \right]$ with

respect to x at $x = 1$ is

A. $3/4$

B. 0

C. $1/2$

D. $-1/2$



Watch Video Solution

41. $f(x) = x^x$, $x \in (0, \infty)$ and let $g(x)$ be inverse of $f(x)$, then $g(x)'$ must be

A. $x(1 + \log x)$

B. $x(1 + \log(x))$

C. $\frac{1}{x(1 + \log g(x))}$

D. non-existent



Watch Video Solution

42. If $y = ax^{n+1} + bx^{-n}$, then $x^2 \frac{d^2y}{dx^2}$ is equal to $n(n-1)y$ (b) $n(n+1)y$ (d) n^2y

A. $n(n-1)y$

B. $n(n+1)y$

C. ny

D. n^2y

 [Watch Video Solution](#)

43. If $y = ax^{n+1} + bx^{-n}$, then $x^2 \frac{d^2y}{dx^2}$ is equal to $n(n-1)y$ (b) $n(n+1)y$ (d) n^2y

A. $m^2(ae^{mx} - be^{-mx})$

B. 1

C. 0

D. none of these

 [Watch Video Solution](#)

44. Suppose $f(x) = e^{ax} + e^{bx}$, where $a \neq b$, and that

$f'(x) - 2f(x) - 15f(x) = 0$ for all x . Then the value of ab is equal to:

A. 25

B. 9

C. -15

D. -9



Watch Video Solution

45. $\frac{d^{20}y}{dx^{20}}(2\cos x \cos 3x) \text{ is equal to } 2^{20}(\cos 2x - 2^{20} \cos 3x)$

$2^{20}(\cos 2x + 2^{20} \cos 4x)$ $2^{20}(\sin 2x + 2^{20} \sin 4x)$ $2^{20}(\sin 2x - 2^{20} \sin 4x)$

A. $2^{20}(\cos 2x - 2^{20} \cos 3x)$

B. $2^{20}(\cos 2x + 2^{20} \cos 4x)$

$$C. 2^{20}(\sin 2x + 2^{20}\sin 4x)$$

$$D. 2^{20}(\sin 2x - 2^{20}\sin 4x)$$

 [Watch Video Solution](#)

$$46. \frac{d^n}{dx^n}(\log x) = \frac{(n-1)!}{x^n} \quad (b) \frac{n!}{x^n} \frac{(n-2)!}{x^n} \quad (d) (-1)^{n-1} \frac{(n-1)!}{x^n}$$

$$A. \frac{(n-1)!}{x^n}$$

$$B. \frac{n!}{x^n}$$

$$C. \frac{(n-2)!}{x^n}$$

$$D. (-1)^{n-1} \frac{(n-1)!}{x^n}$$

 [Watch Video Solution](#)

47. The n th derivative of the function $f(x) = \frac{1}{1-x^2}$ [where $x \in (-1, 1)$] at the point $x = 0$ where n is even is (b) $n!$ (c) $n^n C_2$ (d) $2^n C_2$

A. 0

B. $n!$

C. $b^n C_2$

D. $2^n C_2$



Watch Video Solution

48. If $y = x \log\left\{\frac{x}{(a+bx)}\right\}$, then show that $x^3 \frac{d^2y}{dx^2} = \left(x \frac{dy}{dx} - y\right)^2$.

A. $x \frac{dy}{dx} - y$

B. $\left(x \frac{dy}{dx} - y\right)^2$

C. $y \frac{dy}{dx} - x$

$$D. \left(y \frac{dy}{dx} - x \right)^2$$

 [Watch Video Solution](#)

49. If $ax^2 + 2hxy + by^2 = 1$, then $\frac{d^2y}{dx^2}$ is $\frac{h^2 - ab}{(hx + by)^2}$ (b) $\frac{ab - h^2}{(hx + by)^2} \frac{h^2 + ab}{(hx + by)^2}$

(d) none of these

A. $\frac{h^2 - ab}{(hx + by)^3}$

B. $\frac{ab - h^2}{(hx + by)^2}$

C. $\frac{h^2 + ab}{(hx + by)^2}$

D. none of these

 [Watch Video Solution](#)

50. If $y^{1/m} = (x + \sqrt{1+x^2})$, then $(1+x^2)y_2 + xy_1$ is (where y_r represents the r th derivative of y w.r.t. x)

A. m^2y

B. my^2

C. m^2y^2

D. none of these



Watch Video Solution

51. If $(\sin x)(\cos y) = \frac{1}{2}$, then $\frac{d^2y}{dx^2}$ at $\left(\frac{\pi}{4}, \frac{\pi}{4}\right)$ is -4 (b) -2 (c) -6 (d) 0

A. -4

B. -2

C. -6

D. 0



Watch Video Solution

52. A function f satisfies the condition $f(x) = f'(x) + f^x + f^x$, where $f(x)$ is a differentiable function indefinitely and dash denotes the order of derivative. If $f(x) = 1$, then $f(x)$ is $e^{\frac{x}{2}}$ (b) e^x (c) e^{2x} (d) e^{4x}

A. $e^{x/2}$

B. e^x

C. e^{2x}

D. e^{4x}



Watch Video Solution

53. Let $f(x)$ be a polynomial of degree 3 such that $f(3) = 1$, $f'(3) = -1$, $f^3 = 0$, and $f^3 = 12$. Then the value of $f'(1)$ is 12 (b) 23

(c) -13 (d) none of these

A. 12

B. 23

C. -13

D. none of these



Watch Video Solution

54. If $y^2 = ax^2 + bx + c$, then $y^3 \frac{d^2y}{dx^2}$ is (a) a constant (b) a function of x only (c) a function of y only (d) a function of x and y

A. a constant

B. a function of x only

C. a function of y only

D. a function of x and y



Watch Video Solution

55. If $y = \sin x + e^x$, then $\frac{d^2x}{dy^2} = \left(-\sin x + e^x\right)^{-1} \frac{\sin x - e^x}{\left(\cos x + e^x\right)^2}$ (c)

(d) $\frac{\sin x - e^x}{\left(\cos x + e^x\right)^3} \frac{\sin x + e^x}{\left(\cos x + e^x\right)^3}$

A. $\left(-\sin x + e^x\right)^{-1}$

B. $\frac{\sin x - e^x}{\left(\cos x + e^x\right)^2}$

C. $\frac{\sin x - e^x}{\left(\cos x + e^x\right)^3}$

D. $\frac{\sin x + e^x}{\left(\cos x + e^x\right)^3}$



Watch Video Solution

56. If $y = \sin px$ and y_n is the n^{th} derivative of y , then

$$\begin{vmatrix} y & y_1 & y_2 \\ y_3 & y_4 & y_5 \\ y_6 & y_7 & y_8 \end{vmatrix} \text{ is}$$

A. 1

B. 0

C. -1

D. none of these



Watch Video Solution

57. If $f^x = -f(x)$ and $g(x) = f'(x)$ and $F(x) = \left(f\left(\frac{x}{2}\right)\right)^2 + \left(g\left(\frac{x}{2}\right)\right)^2$ and given that $F(5) = 5$, then $F(10)$ is 5 (b) 10 (c) 0 (d) 15

A. 5

B. 10

C. 0

D. 15



Watch Video Solution

58. Let $y = 1n(1 + \cos x)^2$. Then the value of $\frac{d^2y}{dx^2} + \frac{2}{y}$ equal (b) $\frac{2}{1 + \cos x}$

$\frac{4}{1 + \cos x}$ (d) $\frac{-4}{(1 + \cos x)^2}$

A. 0

B. $\frac{2}{1 + \cos x}$

C. $\frac{4}{1 + \cos x}$

D. $\frac{-4}{(1 + \cos x)^2}$



Watch Video Solution

59. $x = t \cos t, y = t + \sin t$ Then $\frac{d^2x}{dy^2} a = \frac{\pi}{2}$ is $\frac{\pi+4}{2}$ (b) $-\frac{\pi+4}{2}$ -2 (d) none of these

A. $\frac{\pi+4}{2}$

B. $-\frac{\pi+4}{2}$

C. -2

D. none of these



Watch Video Solution

60. If $f(x) = (x-1)^4(x-2)^3(x-3)^2(x-4)$, then the value of $f''(1) + f'(2) + f(3) + f(4)$ equals

A. 0

B. 50

C. 324

D. 648

Answer: 648



Watch Video Solution

61. let $y = t^{10} + 1$, and $x = t^8 + 1$, then $\frac{d^2y}{dx^2}$ is

A. $\frac{5}{2}t$

B. $20t^8$

C. $\frac{5}{16t^6}$

D. none of these



Watch Video Solution

62. If $x = \log p$ and $y = \frac{1}{p}$, then

A. $\frac{d^2y}{dx^2} - 2p = 0$

B. $\frac{d^2y}{dx^2} + y = 0$

C. $\frac{d^2y}{dx^2} + \frac{dy}{dx} = 0$

D. $\frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$



Watch Video Solution

63. If $x = \phi(t)$, $y = \psi(t)$, then $\frac{d^2y}{dx^2}$ is $\frac{\phi' \psi'' - \psi' \phi''}{(\phi')^2}$ (b) $\frac{\phi' \psi'' - \psi' \phi''}{(\phi')^3}$ (c) $\phi' \psi''$ (d) $\psi' \phi''$

A. $\frac{\phi' \psi'' - \psi' \phi''}{(\phi')^2}$

B. $\frac{\phi' \psi'' - \psi' \phi''}{(\phi')^3}$

C. $\frac{\phi''}{\psi''}$

D. $\frac{\psi''}{\phi''}$



Watch Video Solution

64. If $f(x) = x^4 \tan(x^3) - x \ln(1 + x^2)$, then the value of $\frac{d^4(f(x))}{dx^4}$ at $x = 0$ is 0

(b) 6 (c) 12 (d) 24

A. 0

B. 6

C. 12

D. 24



Watch Video Solution

65. If graph of $y = f(x)$ is symmetrical about the y-axis and that of $y = g(x)$

is symmetrical about the origin and if $h(x) = f(x)g(x)$, then $\frac{d^3h(x)}{dx^3}$ at $x = 0$ is

cannot be determined (b) $f(0)g(0)$ 0 (d) none of these

A. cannot be determined

B. $f(0) \cdot g(0)$

C. 0

D. none of these

 [Watch Video Solution](#)

66. Let $g(x)$ be the inverse of an invertible function $f(x)$, which is differentiable for all real x . Then $g^{f(x)}$ equals. $-\frac{f^x}{(f'(x))^3}$ (b)

$\frac{f'(x)f^x - (f'(x))^3}{f'(x)}$ $\frac{f'(x)f^x - (f'(x))^2}{(f'(x))^2}$ (d) none of these

A. $-\frac{f'(x)}{(f'(x))^3}$

B. $\frac{f'(x)f'(x) - (f'(x))^3}{f'(x)}$

C. $\frac{f'(x)f'(x) - (f'(x))^2}{(f'(x))^2}$

D. none of these

 [Watch Video Solution](#)

67. $f(x) = e^x - e^{-x} - 2\sin x - \frac{2}{3}x^3$. Then the least value of n for which

$\frac{d^n}{dx^n}f(x) \big|_{x=0}$ is nonzero is

A. 5

B. 6

C. 7

D. 8



Watch Video Solution

68. Let $y = f(x)$ and $x = \frac{1}{z}$. If $\frac{d^2y}{dx^2} = \lambda(z^3)\frac{dy}{dz} + z^4\frac{d^2y}{dz^2}$, then the value of λ

is

A. 1

B. 2

C. $\frac{1}{2}$

D. $\frac{1}{4}$

 [Watch Video Solution](#)

69. Let $x=f(t)$ and $y=g(t)$, where x and y are twice differentiable function. If

$f'(0)=g'(0)=f''(0)=2$, $g''(0)=6$, then the value of $\left(\frac{d^2y}{dx^2}\right)_{t=0}$ is equal to

A. 0

B. 1

C. 2

D. 3

 [Watch Video Solution](#)

70. If $f(x)$ satisfies the relation $f\left(\frac{5x - 3y}{2}\right) = \frac{5f(x) - 3f(y)}{2} \forall x, y \in \mathbb{R}$, and $f(0) = 3$ and $f'(0) = 2$, then the period of $\sin(f(x))$ is 2π (b) π (c) 3π (d) 4π

A. 2π

B. π

C. 3π

D. 4π

 Watch Video Solution

71. A function $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfies

$$\cos x \left(f(2x + 2y) - f(2x - 2y) \right) = \cos x \sin y (f(2x + 2y) + f(2x - 2y)) \quad \text{if}$$

$$f'(0) = \frac{1}{2}, \text{ then } f'(x) = f(x) = 0 \quad 4f^x + f(x) = 0 \quad f^x + f(x) = 0 \quad 4f^x - f(x) = 0$$

A. $f'(x) = f(x) = 0$

B. $4f'(x) + f(x) = 0$

C. $f'(x) + f(x) = 0$

D. $4f'(x) - f(x) = 0$

 [Watch Video Solution](#)

72. If $y = a\sin x + b\cos x$, then $y^2 + \left(\frac{dy}{dx}\right)^2$ is a

A. function of x

B. function of y

C. function of x and y

D. constant

Answer: D

 [Watch Video Solution](#)

73. $\frac{d}{dx} \sqrt{\frac{1-\sin 2x}{1+\sin 2x}}$ is equal to

A. $\sec^2 x$

B. $-\sec^2\left(\frac{\pi}{4} - x\right)$

C. $\sec^2\left(\frac{\pi}{4} + x\right)$

D. $\sec^2\left(\frac{\pi}{4} - x\right)$



Watch Video Solution

74. If $f(x) = |\cos x| + |\sin x|$, then $\frac{dy}{dx}$ at $x = \frac{2\pi}{3}$ is equal to

A. $\frac{1 - \sqrt{3}}{2}$

B. 0

C. $\frac{1}{2}(\sqrt{3} - 1)$

D. none of these

Answer: C

 Watch Video Solution

75. If $f(x) = \left| \log_e |x| \right|$, then $f'(x)$ equals

A. $\frac{1}{|x|}$, where $x \neq 0$

B. $\frac{1}{x}$ for $|x| > 1$ and $-\frac{1}{x}$ for $|x| < 1$

C. $-\frac{1}{x}$ for $|x| > 1$ and $\frac{1}{x}$ for $|x| < 1$

D. $\frac{1}{x}$ for $x > 0$ and $-\frac{1}{x}$ for $x < 0$

 Watch Video Solution

76. If $f(x) = \sqrt{1 - \sin 2x}$, then $f'(x)$ is equal to

$-(\cos x + \sin x), f$ or $x \in \left(\frac{\pi}{4}, \frac{\pi}{2} \right)$ $\cos x + \sin x, f$ or $x \in \left(0, \frac{\pi}{4} \right)$

$-(\cos x + \sin x), f$ or $x \in \left(0, \frac{\pi}{4} \right)$ $\cos x - \sin x, f$ or $x \in \left(\frac{\pi}{4}, \frac{\pi}{2} \right)$

A. $-(\cos x + \sin x)$, for $x \in (\pi/4, \pi/2)$

B. $\cos x + \sin x$ for $x \in (0, \pi/4)$

C. $-(\cos x + \sin x)$, for $x \in (0, \pi/4)$

D. $\cos x - \sin x$, for $x \in (\pi/4, \pi/2)$



Watch Video Solution

77. Instead of the usual definition of derivative $Df(x)$, if we define a new kind of derivative $D^*F(x)$ by the formula

$$D^*(x) = \lim_{h \rightarrow 0} \frac{f^2(x+h) - f^2(x)}{h}. \text{ where } f^2(x) \text{ means } [f(x)]^2 \text{ and if } f(x) = x$$

$\log x$, then $D^*f(x)|_{x=e}$ has the value

A. e

B. $2e$

C. $4e$

D. none of these



Watch Video Solution

78. if $y = \frac{(a-x)\sqrt{a-x} - (b-x)\sqrt{x-b}}{\sqrt{a-x} + \sqrt{x-b}}$ then $\frac{dy}{dx}$ wherever it is defined is

equal to:

A. $\frac{x + (a + b)}{\sqrt{(a - x)(x - b)}}$

B. $\frac{2x - a - b}{2\sqrt{a - x}\sqrt{x - b}}$

C. $-\frac{(a + b)}{2\sqrt{(a - x)(x - b)}}$

D. $\frac{2x + (a + b)}{2\sqrt{(a - x)(x - b)}}$

Answer: B



Watch Video Solution

79. The derivative of $y = (1 - x)(2 - x)\dots\dots\dots (n - x)$ at $x = 1$ is equal to

A. 0

B. $(-1)(n-1)!$

C. $n! - 1$

D. $(-1)^{n-1}(n-1)!$

Answer: B



Watch Video Solution

80. If $y = \sqrt{\frac{1-x}{1+x}}$, then $(1-x^2)\frac{dy}{dx}$ is equal to

A. y^2

B. $1/y$

C. $-y$

D. $-y/x$

Answer: C



Watch Video Solution

81. If $y = \frac{\sqrt{a+x} - \sqrt{a-x}}{\sqrt{a+x} + \sqrt{a-x}}$, then $\frac{dy}{dx}$ is equal

A. $\frac{ay}{x\sqrt{a^2 - x^2}}$

B. $\frac{ay}{\sqrt{a^2 - x^2}}$

C. $\frac{ay}{x\sqrt{x^2 - a^2}}$

D. none of these

Answer: A

 Watch Video Solution

82. Let $\frac{u}{x}$ and $v(x)$ be differentiable functions such that

$$\frac{u(x)}{v(x)} = 7. \text{ If } \frac{f(u'(x))}{v'(x)} = p \text{ and } \left(\frac{u(x)}{v(x)}\right) = q, \text{ then } \frac{p+q}{p-q} \text{ has the value of to 1 (b)}$$

0 (c) 7 (d) -7

A. 1

B. 0

C. 7

D. -7

 Watch Video Solution

83. If $\sin^{-1}\left(\frac{x^2 - y^2}{x^2 + y^2}\right) = \log a$, then $\frac{dy}{dx}$ is equal to $\frac{x}{y}$ (b) $\frac{y}{x^2} \frac{x^2 - y^2}{x^2 + y^2}$ (d) $\frac{y}{x}$

A. $\frac{x}{y}$

B. $\frac{y}{x^2}$

C. $\frac{x^2 - y^2}{x^2 + y^2}$

D. $\frac{y}{x}$

 Watch Video Solution

84. Let $h(x)$ be differentiable for all x and let $f(x) = (kx + e^x)h(x)$, where k is some constant. If $h(0) = 5$, $h'(0) = -2$, and $f'(0) = 18$, then the value of k is 5 (b) 4 (c) 3 (d) 2.2.

A. 5

B. 4

C. 3

D. 2.2



Watch Video Solution

85. If $\lim_{t \rightarrow x} \frac{e^t f(x) - e^x f(t)}{(t-x)(f(x))^2} = 2$ and $f(0) = \frac{1}{2}$, then find the value of $f'(0)$.

A. 4

B. 2

C. 0

D. 1



Watch Video Solution

86. If $f(0) = 0$, $f'(0) = 2$, then the derivative of $y = f(f(f(x)))$ at $x = 0$ is 2 (b)

8 (c) 16 (d) 4

A. 2

B. 8

C. 16

D. 4



Watch Video Solution

87. If $f(x) = \sqrt{1 + \cos^2(x^2)}$, then $f'\left(\frac{\sqrt{\pi}}{2}\right)$ is $\frac{\sqrt{\pi}}{6}$ (b) $-\sqrt{\pi/6}$ 1/ $\sqrt{6}$ (d) $\pi/\sqrt{6}$

A. $\sqrt{\pi/6}$

B. $-\sqrt{\pi/6}$

C. $1/\sqrt{6}$

D. $\pi/\sqrt{6}$



Watch Video Solution

88. $\frac{d}{dx} \cos^{-1} \sqrt{\cos x}$, $0 < x < \frac{\pi}{2}$ is equal to

A. $\frac{1}{2} \sqrt{1 + \sec x}$

B. $\sqrt{1 + \sec x}$

C. $-\frac{1}{2} \sqrt{1 + \sec x}$

D. $-\sqrt{1 + \sec x}$



Watch Video Solution

89. If $t \tan y = \frac{2^x}{1 + 2^{2x+1}}$, then $\frac{dy}{dx} \tan x = 0$ is - $\frac{3}{10}$ (b) - $\frac{3}{10} \ln 2$ - $\frac{1}{10}$ (d) - $\frac{1}{10} \ln 2$

A. 1

B. 2

C. $\ln 2$

D. $-\frac{1}{10} \ln 2$



Watch Video Solution

90. If $y = \left(x + \sqrt{x^2 + a^2}\right)^n$, then $\frac{dy}{dx}$ is $\frac{ny}{\sqrt{x^2 + a^2}}$ (b) - $\frac{ny}{\sqrt{x^2 + a^2}} - \frac{nx}{\sqrt{x^2 + a^2}}$ (d) - $\frac{nx}{\sqrt{x^2 + a^2}}$

A. $\frac{ny}{\sqrt{x^2 + a^2}}$

B. $-\frac{ny}{\sqrt{x^2 + a^2}}$

C. $\frac{nx}{\sqrt{x^2 + a^2}}$

$$D. -\frac{nx}{\sqrt{x^2 + a^2}}$$

 [Watch Video Solution](#)

91. if $y = \log_{\sin x} \tan x$ then $\left(\frac{dy}{dx}\right)_{\frac{\pi}{4}}$ is

A. $\frac{4}{\log 2}$

B. $-4\log 2$

C. $\frac{-4}{\log 2}$

D. none of these

 [Watch Video Solution](#)

92. $\frac{d}{dx} \left[\sin^2 \cot^{-1} \sqrt{\frac{1-x}{1+x}} \right]$ is

A. -1

B. $\frac{1}{2}$

C. $-\frac{1}{2}$

D. 1

Answer: B



Watch Video Solution

93. The differential coefficient of $f(\log_e x)$ w. r. t. x , where $f(x) = \log_e x$, is (i)

$\frac{x}{\ln x}$ (ii) $\frac{\ln x}{x}$ (iii) $\frac{1}{x \ln x}$ (iv) $x \ln x$

A. $\frac{x}{\log_e x}$

B. $\frac{1}{x} \log_e x$

C. $\frac{1}{x \log_e x}$

D. none of these



Watch Video Solution

94. if $f(x) = \sqrt{2x^2 - 1}$ and $y = f(x^2)$ then $\frac{dy}{dx}$ at $x = 1$ is:

- A. 2
- B. 1
- C. -2
- D. none of these



Watch Video Solution

95. If $u = f(x^3)$, $v = g(x^2)$, $f'(x) = \cos x$, and $g'(x) = \sin x$, then $\frac{du}{dv}$ is

$\frac{3}{2}x \cos x^3 \operatorname{cosec} x^2$ $\frac{2}{3} \sin x^3 \sec x^2 \tan x$ (d) none of these

A. $\frac{3}{2}x \cos x^2 \operatorname{cosec} x^2$

B. $\frac{3}{2} \sin x^3 \sec x^2$

C. $\tan x$

D. none of these



Watch Video Solution

96. A function f , defined for all positive real numbers, satisfies the equation $f(x^2) = x^3 = x^3$ for every $x > 0$. Then the value of $f'(4)$ is

A. 12

B. 3

C. $3/2$

D. cannot be determined



Watch Video Solution

97. Let $f: (-5, 5) \rightarrow \mathbb{R}$ be a differentiable function of with $f(4) = 1$, $f'(4) = 1$, $f(0) = -1$ and $f'(0) = 1$. If $g(x) = \left(f(2f^2(x) + 2) \right)^2$, then

$g'(0)$ equals

A. 4

B. -4

C. 8

D. -8



Watch Video Solution

98. The function $f(x) = e^x + x$, being differentiable and one-to-one, has a differentiable inverse $f^{-1}(x)$. The value of $\frac{x}{dx} \left(f^{-1} \text{ at } f(\log 2) \right)$ is

A. $\frac{1}{\ln 2}$

B. $\frac{1}{3}$

C. $\frac{1}{4}$

D. none of these



Watch Video Solution

99. If $f(x) = x + \tan x$ and f is the inverse of g , then $g'(x)$ is equal to

A. $\frac{1}{1 + [g(x) - x]^2}$

B. $\frac{1}{2 - [g(x) - x]^2}$

C. $\frac{1}{2 + [g(x) - x]^2}$

D. none of these

Answer: C



Watch Video Solution

100. If $f(x) = x^3 + 3x + 4$ and g is the inverse function of $f(x)$, then the value

of $\frac{d}{dx} \left(\frac{g(x)}{g(g(x))} \right)$ at $x = 4$ equals

A. $\frac{-1}{6}$

B. 6

C. $\frac{-1}{3}$

D. non-existent

Answer: = $\frac{-1}{3}$



Watch Video Solution

101. If $y = \frac{\sin^{-1}x}{\sqrt{1-x^2}}$, then $\frac{(1-x^2)dy}{dx}$ is equal to $x+y$ (b) $1+xy$ $1-xy$ (d)

$xy - 2$

A. $x+y$

B. $1+xy$

C. $1-xy$

D. $xy-2$



Watch Video Solution

102. If $y = \sqrt{\log x + \sqrt{\log x + \sqrt{\log x + \dots \infty}}}$, then $\frac{dy}{dx}$ is

A. $\frac{x}{2y - 1}$

B. $\frac{x}{2y + 1}$

C. $\frac{1}{x(2y - 1)}$

D. $\frac{1}{x(1 - 2y)}$



Watch Video Solution

103. $\frac{d}{dx} \left[\tan^{-1} \left(\frac{\sqrt{x}(3-x)}{1-3x} \right) \right] =$

A. $\frac{1}{2(1+x)\sqrt{x}}$

B. $\frac{3}{(1+x)\sqrt{x}}$

C. $\frac{2}{(1+x)\sqrt{x}}$

D. $\frac{3}{2(1+x)\sqrt{x}}$

Answer: $= \frac{3}{2\sqrt{x}(1+x)}$

 [Watch Video Solution](#)

104. Suppose the function $f(x) - f(2x)$ has the derivative 5 at $x = 1$ and derivative 7 at $x = 2$. The derivative of the function $f(x) - f(4x)$ at $x = 1$ has the value equal to 19 (b) 9 (c) 17 (d) 14

A. 19

B. 9

C. 17

D. 14

 [Watch Video Solution](#)

105. If $y\sqrt{x^2 + 1} = \log(\sqrt{x^2 + 1} - x)$, show that $(x^2 + 1)\frac{dy}{dx} + xy + 1 = 0$

A. 0

B. 1

C. 2

D. none of these



Watch Video Solution

106.

Let $e^y = \frac{\sqrt{1+\alpha} + \sqrt{1-\alpha}}{\sqrt{1+\alpha} - \sqrt{1-\alpha}}$ and $\tan\frac{x}{2} = \sqrt{\frac{1-\alpha}{1+\alpha}}$, $\alpha \in [-1, 0] \cup (0, 1]$. Then $\left(\frac{dy}{dx}\right)$

A. 1/2

B. 1

C. 2

D. 1/3

Answer: 2



Watch Video Solution

107. The derivative of $\tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$ with respect to

$\tan^{-1}\left(\frac{2x\sqrt{1-x^2}}{1-2x^2}\right)$ at $x = 0$ is $\frac{1}{8}$ (b) $\frac{1}{4}$ (c) $\frac{1}{2}$ (d) 1

A. $1/8$

B. $1/4$

C. $1/2$

D. 1



Watch Video Solution

108. If $\ln\left((e - 1)e^{xy} + x^2\right) = x^2 + y^2$ then $\left(\frac{dy}{dx}\right)_{1,0}$ is equal to

A. 0

B. 1

C. 2

D. 3

Answer: 2



Watch Video Solution

109. If $y = (x^x)^x$ then $\frac{dy}{dx}$ is

A. $y \left[x^x (\log_e x) \log x + x^x \right]$

B. $y \left[x^x (\log_e x) \log x + x \right]$

C. $y \left[x^x (\log_e x) \log x + x^{-1} \right]$

D. $y \left[x^x (\log_e x) \log x + x^{-1} \right]$



Watch Video Solution

110. The first derivative of the function $\left[\cos^{-1} \left(\sin \sqrt{\frac{1+x}{2}} \right) + x^x \right]$ with respect to x at $x = 1$ is

A. $3/4$

B. 0

C. $1/2$

D. $-1/2$



Watch Video Solution

111. $f(x) = x^x, x \in (0, \infty)$ and let $g(x)$ be inverse of $f(x)$, then $g(x)'$ must be

A. $x(1 + \log x)$

B. $x(1 + \log(x))$

C. $\frac{1}{x(1 + \log(x))}$

D. non-existent

 [Watch Video Solution](#)

112. If $y = ax^{n+1} + bx^{-n}$, then $x^2 \frac{d^2y}{dx^2}$ is equal to $n(n-1)y$ (b) $n(n+1)y$ (d) n^2y

A. $n(n-1)y$

B. $n(n+1)y$

C. ny

D. n^2y

 [Watch Video Solution](#)

113. If $y = ax^{n+1} + bx^{-n}$, then $x^2 \frac{d^2y}{dx^2}$ is equal to $n(n-1)y$ (b) $n(n+1)y$ (d) n^2y

A. $m^2(ae^{mx} - be^{-mx})$

B. 1

C. 0

D. none of these



Watch Video Solution

114. Suppose $f(x) = e^{ax} + e^{bx}$, where $a \neq b$, and that $f''(x) - 2f'(x) - 15f(x) = 0$ for all x . Then the product ab is

A. 25

B. 9

C. -15

[Watch Video Solution](#)

115. $\frac{d^{20}y}{dx^{20}}(2\cos x \cos 3x) \text{ is equal to } 2^{20}(\cos 2x - 2^{20} \cos 3x)$

$$2^{20}(\cos 2x + 2^{20} \cos 4x) \quad 2^{20}(\sin 2x + 2^{20} \sin 4x) \quad 2^{20}(\sin 2x - 2^{20} \sin 4x)$$

A. $2^{20}(\cos 2x - 2^{20} \cos 3x)$

B. $2^{20}(\cos 2x + 2^{20} \cos 4x)$

C. $2^{20}(\sin 2x + 2^{20} \sin 4x)$

D. $2^{20}(\sin 2x - 2^{20} \sin 4x)$

[Watch Video Solution](#)

116. $\frac{d^n}{dx^n}(\log x) = \frac{(n-1)!}{x^n}$ (b) $\frac{n!}{x^n} \frac{(n-2)!}{x^n}$ (d) $(-1)^{n-1} \frac{(n-1)!}{x^n}$

A. $\frac{(n-1)!}{x^n}$

B. $\frac{n!}{x^n}$

C. $\frac{(n-2)!}{x^n}$

D. $(-1)^{n-1} \frac{(n-1)!}{x^n}$

 [Watch Video Solution](#)

117. If $y = x \log \left(\frac{x}{a+bx} \right)$, then $x^3 \frac{d^2y}{dx^2} =$

A. $x \frac{dy}{dx} - y$

B. $\left(x \frac{dy}{dx} - y \right)^2$

C. $y \frac{dy}{dx} - x$

D. $\left(y \frac{dy}{dx} - x \right)^2$

Answer: D

 [Watch Video Solution](#)

118. If $ax^2 + 2hxy + by^2 = 1$, then $\frac{d^2y}{dx^2}$ is $\frac{h^2 - ab}{(hx + by)^2}$ (b) $\frac{ab - h^2}{(hx + by)^2}$ $\frac{h^2 + ab}{(hx + by)^2}$

(d) none of these

A. $\frac{h^2 - ab}{(hx + by)^3}$

B. $\frac{ab - h^2}{(hx + by)^2}$

C. $\frac{h^2 + ab}{(hx + by)^2}$

D. none of these



Watch Video Solution

119. If $y^{1/m} = (x + \sqrt{1 + x^2})$, then $(1 + x^2)y_2 + xy_1$ is (where y_r represents the r th derivative of y w.r.t. x)

A. m^2y

B. my^2

C. m^2y^2

D. none of these

Answer: A



[Watch Video Solution](#)

120. If $(\sin x)(\cos y) = 1/2$, then d^2y/dx^2 at $(\pi/4, \pi/4)$ is

A. -4

B. -2

C. -6

D. 0



[Watch Video Solution](#)

121. A function f satisfies the condition $f(x) = f'(x) + f''(x) + f'''(x) + \dots$, where $f(x)$ is a differentiable function indefinitely and dash denotes the order the derivative. If $f(0) = 1$, then $f(x)$ is

A. $e^{x/2}$

B. e^x

C. e^{2x}

D. e^{4x}



Watch Video Solution

122. Let $f(x)$ be a polynomial of degree 3 such that $f(3) = 1$, $f'(3) = -1$, $f''(3) = 0$, and $f'''(3) = 12$. Then the value of $f(1)$ is

A. 12

B. 23

C. -13

D. none of these

Answer: 23



Watch Video Solution

123. If $y^2 = ax^2 + bx + c$, then $y^3 \frac{d^2y}{dx^2}$ is (a) a constant (b) a function of x only (c) a function of y only (d) a function of x and y

A. a constant

B. a function of x only

C. a function of y only

D. a function of x and y



Watch Video Solution

124. If $y = \sin x + e^x$, then $\frac{d^2x}{dy^2} =$

A. $(-\sin x + e^x)^{-1}$

B. $\frac{\sin x - e^x}{(\cos x + e^x)^2}$

C. $\frac{\sin x - e^x}{(\cos x + e^x)^3}$

D. $\frac{\sin x + e^x}{(\cos x + e^x)^3}$



Watch Video Solution

125. If $f''(x) = -f(x)$ and $g(x) = f'(x)$ and $F(x) = \left(f\left(\frac{x}{2}\right)\right)^2 + \left(g\left(\frac{x}{2}\right)\right)^2$

and given that $F(5) = 5$, then $F(10)$ is

A. 5

B. 10

C. 0

D. 15

 [Watch Video Solution](#)

126. Let $y = \sin(1 + \cos x)^2$. The value of $\frac{d^2y}{dx^2} + \frac{2}{e^{y/2}}$ equals

A. 0

B. $\frac{2}{1 + \cos x}$

C. $\frac{4}{1 + \cos x}$

D. $\frac{-4}{(1 + \cos x)^2}$

Answer: A

 [Watch Video Solution](#)

127. $x = t \cos t, y = t + \sin t$. Then $\frac{d^2x}{dy^2} a = \frac{\pi}{2}$ is $\frac{\pi+4}{2}$ (b) $-\frac{\pi+4}{2}$ -2 (d) none of these

A. $\frac{\pi+4}{2}$

B. $-\frac{\pi+4}{2}$

C. -2

D. none of these



Watch Video Solution

128. Let $y = t^{10} + 1$ and $x = t^8 + 1$. Then $\frac{d^2y}{dx^2}$ is

A. $\frac{5}{2}t$

B. $20t^8$

C. $\frac{5}{16t^6}$

D. none of these

Answer: C



Watch Video Solution

129. If $x = \log p$ and $y = \frac{1}{p}$, then

A. $\frac{d^2y}{dx^2} - 2p = 0$

B. $\frac{d^2y}{dx^2} + y = 0$

C. $\frac{d^2y}{dx^2} + \frac{dy}{dx} = 0$

D. $\frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$



Watch Video Solution

130. If $x = \phi(t)$, $y = \psi(t)$, then $\frac{d^2y}{dx^2}$ is $\frac{\phi' \psi'' - \psi' \phi''}{(\phi')^2}$ (b) $\frac{\phi' \psi'' - \psi' \phi''}{(\phi')^3}$ $\phi' \psi$ (d) $\psi' \phi$

A. $\frac{\phi' \psi'' - \psi' \phi''}{(\phi')^2}$

B. $\frac{\phi' \psi'' - \psi' \phi''}{(\phi')^3}$

C. $\frac{\phi''}{\psi''}$

D. $\frac{\psi''}{\phi''}$

 [Watch Video Solution](#)

131. If $f(x) = x^4 \tan(x^3) - x \ln(1 + x^2)$, then the value of $\frac{d^4(f(x))}{dx^4}$ at $x = 0$ is

0 (b) 6 (c) 12 (d) 24

A. 0

B. 6

C. 12

D. 24

 [Watch Video Solution](#)

132. Let $g(x)$ be the inverse of an invertible function $f(x)$, which is differentiable for all real x . Then $g''(f(x))$ equals

A. $-\frac{f'(x)}{(f'(x))^3}$

B. $\frac{f(x)f'(x) - (f(x))^3}{f'(x)}$

C. $\frac{f(x)f'(x) - (f'(x))^2}{(f'(x))^2}$

D. none of these



Watch Video Solution

133. $f(x) = e^x - e^{-x} - 2\sin x - \frac{2}{3}x^3$. Then the least value of n for which

$\frac{d^n}{dx^n}f(x) \big|_{x=0}$ is nonzero is

A. 5

B. 6

C. 7

D. 8

Answer: C



Watch Video Solution

134. Let $y = f(x)$ and $x = \frac{1}{z}$. If $\frac{d^2y}{dx^2} = \lambda z^3 \frac{dy}{dz} + z^4 \frac{d^2y}{dz^2}$, then the value of λ is

A. 1

B. 2

C. $\frac{1}{2}$

D. $\frac{1}{4}$

Answer: B



Watch Video Solution

135. Let $x=f(t)$ and $y=g(t)$, where x and y are twice differentiable function. If

$f'(0)=g'(0)=f''(0)=2$, $g''(0)=6$, then the value of $\left(\frac{d^2y}{dx^2}\right)_{t=0}$ is equal to

- A. 0
- B. 1
- C. 2
- D. 3



Watch Video Solution

136. If $f(x)$ satisfies the relation $f\left(\frac{5x-3y}{2}\right) = \frac{5f(x)-3f(y)}{2} \forall x, y \in R$, and

$f(0) = 3$ and $f'(0) = 2$, then the period of $\sin(f(x))$ is 2π (b) π (c) 3π (d) 4π

- A. 2π
- B. π

C. 3π

D. 4π

 [Watch Video Solution](#)

137. A function

$f: \mathbb{R} \rightarrow \mathbb{R}$ satisfies $\sin x \cos y (f(2x + 2y) - f(2x - 2y)) = \cos x \sin y (f(2x + 2y) + f(2x - 2y))$

" If $f'(0) = \frac{1}{2}$, then

A. $f'(x) = f(x) = 0$

B. $4f'(x) + f(x) = 0$

C. $f'(x) + f(x) = 0$

D. $4f'(x) - f(x) = 0$

 [Watch Video Solution](#)

Exercise (Multiple)

1. If $y = e^{\sqrt{x}} + e^{-\sqrt{x}}$, then $\frac{dy}{dx}$ is equal to (a) $\frac{e^{\sqrt{x}}}{2\sqrt{x}}$ (b) $\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2x}$ (c) $\frac{1}{2\sqrt{x}}\sqrt{y^2 - 4}$ (d)

$$\frac{1}{2\sqrt{x}}\sqrt{y^2 + 4}$$

A. $\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2\sqrt{x}}$

B. $\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2x}$

C. $\frac{1}{2\sqrt{x}}\sqrt{y^2 - 4}$

D. $\frac{1}{2\sqrt{x}}\sqrt{y^2 + 4}$



Watch Video Solution

2. Let $y = \sqrt{x + \sqrt{x + \sqrt{x + \infty}}}$, $\frac{dy}{dx}$ is equal to (a) $\frac{1}{2y - 1}$ (b) $\frac{x}{x + 2y}$ (c) $\frac{1}{\sqrt{1 + 4x}}$ (d)

$$\frac{y}{2x + y}$$

A. $\frac{1}{2y - 1}$

B. $\frac{x}{x + 2y}$

C. $\frac{1}{\sqrt{1 + 4x}}$

D. $\frac{y}{2x + y}$

 **Watch Video Solution**

3. If $f(\theta) = \tan\left(\sin^{-1}\sqrt{\frac{2}{3 + \cos 2\theta}}\right)$, then

A. $f\left(\frac{\pi}{4}\right) = 1$

B. $f\left(\frac{\pi}{4}\right) = \sqrt{2}$

C. $\frac{d(f(\theta))}{d(\cos\theta)}$ at $\theta = \frac{\pi}{4}$ is -2

D. $f\left(\frac{\pi}{4}\right) = \sqrt{2}$

 **Watch Video Solution**

4. $f(x) = |x^2 - 3x| + 2$. Then which of the following is/are true ?

A. $f(x) = 2x - 3$ for $x \in (0, 1) \cup (2, \infty)$

B. $f(x) = 2x + 3$ or $x \in (-\infty, -2) \cup (-1, 0)$

C. $f(x) = -2x - 3$ for $x \in (-2, -1)$

D. None of these



[Watch Video Solution](#)

5. Let $f(x) = \frac{\sqrt{x - 2\sqrt{x - 1}}}{\sqrt{x - 1} - 1}$. x then

A. $f(10) = 1$

B. $f(3/2) = -1$

C. domain of $f(x)$ is $x \geq 1$

D. range of $f(x)$ is $(-2, -1] \cup (2, \infty)$



Watch Video Solution

6. If $x^3 - 2x^2y^2 + 5x + y - 5 = 0$ and $y(1) = 1$, then $y'(1) = \frac{4}{3}$ (b) $y^1 = -\frac{4}{3}$
 $y^1 = -8\frac{22}{27}$ (d) $y'(1) = \frac{2}{3}$

A. $y'(1) = 4/3$

B. $y''(1) = -4/3$

C. $y'(1) = -8\frac{22}{27}$

D. $y'(1) = 2/3$



Watch Video Solution

7. If $y = x^{(\log x)^{\log(\log x)}}$ then $\frac{dy}{dx} =$

A. $\frac{y}{x} \left((\ln x^{x-1}) + 2 \ln x \ln(\ln x) \right)$

B. $\frac{y}{x} (\log x)^{\log(\log x)} (2 \log(\log x) + 1)$

$$C. \frac{y}{x \ln x} \left[(\ln x)^2 + 2 \ln(\ln x) \right]$$

$$D. \frac{y \log y}{x \log x} [2 \log(\log x) + 1]$$

Answer: B



Watch Video Solution

8. If $f(x - y)$, $f(x)f(y)$, and $f(x + y)$ are in A.P. for all x, y , and $f(0) \neq 0$, then

A. $f(4) = f(-4)$

B. $f(2) + f(-2) = 0$

C. $f(4) + f(-4) = 0$

D. $f(2) = f(-2)$



Watch Video Solution

9. If $y = \cos^{-1}\left(\frac{2x}{1+x^2}\right)$, then $\frac{dy}{dx}$ is $\frac{-2}{1+x^2}$ for all x (b) $\frac{-2}{1+x^2}$ for all $|x| < 1$

$\frac{2}{1+x^2}$ for $|x| > 1$ (d) none of these

A. $\frac{-2}{1+x^2}$ for all x

B. $\frac{-2}{1+x^2}$ for all $|x| < 1$

C. $\frac{2}{1+x^2}$ for $|x| > 1$

D. none of these



Watch Video Solution

10. $f_n(x) = e^{f_{n-1}(x)}$ for all $n \in N$ and $f_0(x) = x$, then $\frac{d}{dx}\{f_n(x)\}$ is

A. $f_n(x) \frac{d}{dx}\{f_{n-1}(x)\}$

B. $f_n(x)f_{n-1}(x)$

C. $f_n(x)f_{n-1}(x)\dots f_2(x) \cdot f_1(x)$

D. None of these



Watch Video Solution

11. Suppose f and g are functions having second derivatives f'' and g'' every where, if $f(x) \cdot g(x) = 1$ for all x and f', g'' are never zero then

$\frac{f'(x)}{f(x)} - \frac{g'(x)}{g(x)}$ equals

A. $\frac{-2f'(x)}{f(x)}$

B. $\frac{-2g'(x)}{g(x)}$

C. $\frac{-f'(x)}{f(x)}$

D. $\frac{2f'(x)}{f(x)}$



Watch Video Solution

12. If $y = e^{-x}\cos x$ and $y_n + k_n y = 0$ where $y_n = \frac{d^n y}{dx^n}$ and k_n are constant $n \in N$ then

A. $k_4 = 4$

B. $k_8 = -16$

C. $k_{12} = 20$

D. $k_{16} = -24$

Answer: B



Watch Video Solution

13. If a function is represented parametrically by the equations

$$x = \frac{1 + (\log)_e t}{t^2}; y = \frac{3 + 2(\log)_e t}{t}, \text{ then which of the following statements}$$

are true? $y^{x-2xy'} = y yy' = 2x(y')^2 + 1 xy' = 2y(y')^2 + 2 y^{y-4xy'} = (y')^2$

A. $y''(x - 2xy') = y$

$$B. yy' = 2x(y')^2 + 1$$

$$C. xy' = 2y(y')^2 + 2$$

$$D. y''(y - 4xy') = (y')^2$$



Watch Video Solution

14. If $y = \frac{x^2}{2} + \frac{1}{2}x\sqrt{x^2 + 1} + (\log)_e \sqrt{x + \sqrt{x^2 + 1}}$, prove that $2y = xy' + (\log)_e y'$, where y' denotes the derivative w.r.t x .

$$A. y' = x + \sqrt{x^2 + 1}$$

$$B. y' = \frac{1}{x + \sqrt{x^2 + 1}}$$

$$C. 2y = xy' + \log_e y'$$

$$D. 2y = xy' - \log_e y'$$



Watch Video Solution

15. A curve parametrically given by

$x = t + t^3$ and $y = t^2$, where $t \in R$. For what value(s) of t is $\frac{dy}{dx} = \frac{1}{2}$?

A. $1/3$

B. 2

C. 3

D. 1



Watch Video Solution

16. If $e^{\sin(x^2+y^3)} = \tan\frac{y^2}{4} + \sin^{-1}x$, then $y'(0)$ can be

A. $\frac{1}{3\sqrt{\pi}}$

B. $-\frac{1}{3\sqrt{\pi}}$

C. $-\frac{1}{5\sqrt{\pi}}$

D. $-\frac{1}{3\sqrt{5\pi}}$



Watch Video Solution

17. If $y = e^{\sqrt{x}} + e^{-\sqrt{x}}$, then $\frac{dy}{dx}$ is equal to

A. $\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2\sqrt{x}}$

B. $\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2x}$

C. $\frac{1}{2\sqrt{x}}\sqrt{y^2 - 4}$

D. $\frac{1}{2\sqrt{x}}\sqrt{y^2 + 4}$



Watch Video Solution

18. Let $y = \sqrt{x + \sqrt{x + \sqrt{x + \dots\infty}}}$, then $\frac{dy}{dx}$ is equal to

A. $\frac{1}{2y - 1}$

B. $\frac{x}{x + 2y}$

C. $\frac{1}{\sqrt{1+4x}}$

D. $\frac{y}{2x+y}$



Watch Video Solution

19. If $f(x - y)$, $f(x)f(y)$, and $f(x + y)$ are in A.P. for all x, y , and $f(0) \neq 0$, then

A. $f(4) = f(-4)$

B. $f(2) + f(-2) = 0$

C. $f(4) + f(-4) = 0$

D. $f(2) = f(-2)$



Watch Video Solution

20. If a function is represented parametrically by the equations

$$x = \frac{1 + (\log)_e t}{t^2}; y = \frac{3 + 2(\log)_e t}{t}, \text{ then which of the following statements}$$

are true? $y^{x-2xy'} = yyy' = 2x(y')^2 + 1$ $xy' = 2y(y')^2 + 2$ $y^{y-4xy'} = (y')^2$

A. $y'(x - 2xy') = y$

B. $yy' = 2x(y')^2 + 1$

C. $xy' = 2y(y')^2 + 2$

D. $y'(y - 4xy') = (y')^2$



Watch Video Solution

21. If $y = \frac{x^2}{2} + \frac{1}{2}x\sqrt{x^2 + 1} + \log_e \sqrt{x + \sqrt{x^2 + 1}}$, then

A. $y' = x + \sqrt{x^2 + 1}$

B. $y' = \frac{1}{x + \sqrt{x^2 + 1}}$

C. $2y = xy' + \log_e y'$

$$D. 2y = xy' - \log_e y'$$



Watch Video Solution

22. A curve parametrically given by

$x = t + t^3$ and $y = t^2$, where $t \in R$. For what value(s) of t is $\frac{dy}{dx} = \frac{1}{2}$?

A. $1/3$

B. 2

C. 3

D. 1



Watch Video Solution

Exercise (Comprehension)

1. $f(x)$ is a polynomial function, $f: R \rightarrow R$, such that $f(2x) = f'(x)f'(x)$. The value of $f(3)$ is
- A. 4
 - B. 12
 - C. 15
 - D. none of these

Answer: B



[Watch Video Solution](#)

2. $f(x)$ is a polynomial function, $f: R \rightarrow R$, such that $f(2x) = f'(x)f'(x)$. $f(x)$ is (A) one-one and onto (B) one-one and into (C) many-one and onto (D) many-one and into
- A. one-one and onto
 - B. one-one and into
 - C. many-one and onto

D. many-one and into



Watch Video Solution

3. $f(x)$ is a polynomial function, $f: R \rightarrow R$, such that $f(2x) = f'(x)f'(x)$.

Equation $f(x) = x$ has (A) three real and positive roots (B) three real and negative roots (C) one real root (D) three real roots such that sum of roots is zero

A. three real and positive roots

B. three real and negative roots

C. one real root

D. three real roots such that sum of roots is zero



Watch Video Solution

4. $f: \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = x^3 + x^2 f'(1) + x f''(2) + f'''(3)$ for all $x \in \mathbb{R}$.

The value of $f(1)$ is

A. 2

B. 3

C. -1

D. 4



Watch Video Solution

5. $f: \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = x^3 + x^2 f'(1) + x f''(2) + f'''(3)$ for all $x \in \mathbb{R}$.

$f(x)$ is

A. one-one and onto

B. one-one and into

C. many-one and onto

D. many-one and into



Watch Video Solution

6. $f: R \rightarrow R, f(x) = x^3 + x^2 f'(1) + x f''(2) + f'''(3)$ for all $x \in R$.

The value of $f(1) + f'(2) + f''(3)$ is

A. 0

B. -1

C. 2

D. 3



Watch Video Solution

7. Repeated roots : If equation $f(x) = 0$, where $f(x)$ is a polynomial function, has roots $\alpha, \alpha, \beta, \dots$ or α root is repeated root, then $f(x) = 0$ is

equivalent to $(x - \alpha)^2(x - \beta)\dots = 0$, from which we can conclude that $f(x) = 0$ or $2(x - \alpha)[(x - \beta)\dots] + (x - \alpha)^2[(x - \beta)\dots]' = 0$ or $(x - \alpha)[2\{(x - \beta)\dots\} + (x - \alpha)\dots] = 0$ has root α . Thus, if α root occurs twice in the, equation, then it is common in equations $f(x) = 0$ and $f'(x) = 0$. Similarly, if α root occurs thrice in equation, then it is common in the equations $f(x)=0$, $f'(x)=0$, and $f''(x)=0$. If $x-c$ is a factor of order m of the polynomial $f(x)$ of degree n ($1 < m < n$), then $x=c$ is a root of the polynomial [where $f^r(x)$ represent r th derivative of $f(x)$ w.r.t. x]

- A. $f^m(x)$
- B. $f^{m-1}(x)$
- C. $f^n(x)$
- D. none of these



[View Text Solution](#)

8. Repeated roots : If equation $f(x) = 0$, where $f(x)$ is a polynomial function, has roots $\alpha, \alpha, \beta, \dots$ or α root is repeated root, then $f(x) = 0$ is equivalent to $(x - \alpha)^2(x - \beta)\dots = 0$, from which we can conclude that $f(x) = 0$ or $2(x - \alpha)[(x - \beta)\dots] + (x - \alpha)^2[(x - \beta)\dots]' = 0$ or $(x - \alpha)[2\{(x - \beta)\dots\} + (x - \alpha)\dots] = 0$ has root α . Thus, if α root occurs twice in the equation, then it is common in equations $f(x) = 0$ and $f'(x) = 0$. Similarly, if α root occurs thrice in equation, then it is common in the equations $f(x)=0, f'(x)=0, \text{ and } f''(x)=0$.

If $a_1x^3 + b_1x^2 + c_1x + d_1 = 0$ and $a_2x^3 + b_2x^2 + c_2x + d_2 = 0$ have a pair of repeated roots common, then

$$\begin{vmatrix} 3a_1 & 2b_1 & c_1 \\ 3a_2 & 2b_2 & c_2 \\ a_2b_1 - a_1b_2 & c_1a_2 - c_2a_1 & d_1a_2 - d_2a_1 \end{vmatrix} =$$

- A. 0
- B. 1
- C. -1
- D. 2

 [View Text Solution](#)

9. Equation $x^n - 1 = 0, n > 1, n \in N$, has roots $1, a_1, a_2, \dots, a_{n-1}$.

The value of $(1 - a_1)(1 - a_2) \dots (1 - a_{n-1})$ is

A. $n^2/2$

B. n

C. $(-1)^n n$

D. none of these

 [Watch Video Solution](#)

10. Equation $x^n - 1 = 0, n > 1, n \in N$, has roots $1, a_1, a_2, \dots, a_{n-1}$.

The value of $\sum_{r=1}^{n-1} \frac{1}{2 - a_r}$ is

A. $\frac{2^{n-1}(n-2) + 1}{2^n - 1}$

B. $\frac{2^n(n-2) + 1}{2^n - 1}$

C. $\frac{2^{n-1}(n-1) - 1}{2^n - 1}$

D. none of these

 [Watch Video Solution](#)

11. Equation $x^n - 1 = 0, n > 1, n \in N$, has roots $1, a_1, a_2, \dots, a_{n-1}$.

The value of $\sum_{r=1}^{n-1} \frac{1}{1 - a_r}$ is

A. $\frac{n}{4}$

B. $\frac{n(n-1)}{2}$

C. $\frac{n-1}{2}$

D. none of these

 [Watch Video Solution](#)

12. $f(x) = x^2 + xg'(1) + g''(2)$ and $g(x) = f(1)x^2 + xf'(x) + f(x)$.

The value of $f(3)$ is

A. 1

B. 0

C. -1

D. -2



Watch Video Solution

13. $f(x) = x^2 + xg'(1) + g''(2)$ and $g(x) = f(1)x^2 + xf'(x) + f(x)$.

The value of $g(0)$ is

A. 0

B. -3

C. 2

D. none of these

Answer: C



Watch Video Solution

14. $f(x) = x^2 + xg'(1) + g''(2)$ and $g(x) = f(1)x^2 + xf'(x) + f(x)$.

The domain of the function $\sqrt{\frac{f(x)}{g(x)}}$ is

A. $(-\infty, 1] \cup (2, 3]$

B. $(-2, 0] \cup (1, \infty)$

C. $(-\infty, 0] \cup (2/3, 3]$

D. none of these



Watch Video Solution

15. $g(x + y) = g(x) + g(y) + 3xy(x + y) \forall x, y \in R$ and $g'(0) = -4$.

Number of real roots of the equation $g(x) = 0$ is

A. 2

B. 0

C. 1

D. 3



Watch Video Solution

16. $g(x + y) = g(x) + g(y) + 3xy(x + y) \forall x, y \in R$ and $g'(0) = -4$.

For which of the following values of x is $\sqrt{g(x)}$ not defined ?

A. $[-2, 0]$

B. $[2, \infty)$

C. $[-1, 1]$

D. none of these



Watch Video Solution

17. $g(x + y) = g(x) + g(y) + 3xy(x + y) \forall x, y \in R$ and $g'(0) = -4$.

The value of $g'(1)$ is

A. 0

B. 1

C. -1

D. none of these



Watch Video Solution

18. A curve is represented parametrically by the equations

$x = f(t) = a^{\ln(b^t)}$ and $y = g(t) = b^{-\ln(a^t)}$, $a, b > 0$ and $a \neq 1, b \neq 1$ Where $t \in \mathbb{R}$.

Which of the following is not a correct expression for $\frac{dy}{dx}$?

A. $\frac{1}{f(t)^2}$

B. $-(g(t))^2$

C. $\frac{-g(t)}{f(t)}$

D. $\frac{-f(t)}{g(t)}$



Watch Video Solution

19. A curve is represented parametrically by the equations

$$x = f(t) = a^{\ln(b^t)} \text{ and } y = g(t) = b^{-\ln(a^t)}, a, b > 0 \text{ and } a \neq 1, b \neq 1 \text{ Where } t \in \mathbb{R}$$

The value of $\frac{d^2y}{dx^2}$ at the point where $f(t)=g(t)$ is

A. 0

B. $\frac{1}{2}$

C. 1

D. 2



Watch Video Solution

20. A curve is represented parametrically by the equations

$$x = f(t) = a^{\ln(b^t)} \text{ and } y = g(t) = b^{-\ln(a^t)}, a, b > 0 \text{ and } a \neq 1, b \neq 1 \text{ Where } t \in \mathbb{R}$$

The value of $\frac{f(t)}{f(-t)} \cdot \frac{f'(-t)}{f'(-t)} + \frac{f(-t)}{f(-t)} \cdot \frac{f'(t)}{f'(t)} \forall t \in \mathbb{R}$ is

A. -2

B. 2

C. -4

D. 4



Watch Video Solution

21. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function satisfying

$$f(x+y) = f(x) + f(y) + x^2y + xy^2 \text{ for all real numbers } x \text{ and } y. \text{ If}$$

$\lim_{x \rightarrow 0} \frac{f(x)}{x} = 1$, then

The value of $f'(3)$ is

A. 8

B. 10

C. 12

D. 18



Watch Video Solution

22. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function satisfying

$f(x + y) = f(x) + f(y) + x^2y + xy^2$ for all real numbers x and y . If

$\lim_{x \rightarrow 0} \frac{f(x)}{x} = 1$, then

The value of $f(9)$ is

A. 240

B. 356

C. 252

D. 730

Answer: C



[Watch Video Solution](#)

23. If roots of an equation $x^n - 1 = 0$ are $1, a_1, a_2, \dots, a_{n-1}$ then the value of $(1 - a_1)(1 - a_2)(1 - a_3)\dots(1 - a_{n-1})$ will be (a) n (b) n^2 (c) n^n (d) 0

A. $n^2/2$

B. n

C. $(-1)^n n$

D. none of these



[Watch Video Solution](#)

Exercise (Matrix)

1. Match the following lists :



 [View Text Solution](#)

2. Match the following lists :



 [View Text Solution](#)

3. Match the following lists :



 [View Text Solution](#)

4. Match the values of x in List II where derivative of the function in List I is negative.



[View Text Solution](#)

5. Match the relation for derivatives given in List II with the relation given in List I and then choose the correct code.



- A. $a \ b \ c \ d$
 $q \ p \ s \ r$
- B. $a \ b \ c \ d$
 $s \ p \ q \ r$
- C. $a \ b \ c \ d$
 $r \ q \ s \ p$
- D. $a \ b \ c \ d$
 $q \ p \ r \ s$



[View Text Solution](#)

Exercise (Numerical)

1. $f'(x) = \varphi'(x) = f(x)$ for all x . Also, $f(3) = 5$ and $f'(3) = 4$. Then the value of $[f(10)]^2$ is _____

 [Watch Video Solution](#)

2. If $y=f(x)$ is an odd differentiable function defined on $(-\infty, \infty)$ such that $f(3) = -2$ then $f'(-3)$ equals -

 [Watch Video Solution](#)

3. If $x^3 + 3x^2 - 9x + c$ is of the form $(x - \alpha)^2(x - \beta)$ then c is equal to

 [Watch Video Solution](#)

4. If graph of $y = f(x)$ is symmetrical about the point $(5, 0)$ and $f'(7) = 3$, then the value of $f'(3)$ is _____

 [Watch Video Solution](#)

5. Let $g(x) = f(x)\sin x$, where $f(x)$ is a twice differentiable function on $(-\infty, \infty)$ such that $f(-\pi) = 1$. The value of $|g^{-\pi}|$ equals _____

 [Watch Video Solution](#)

6. Let $f(x) = (x-1)(x-2)(x-3)(x-n)$, $n \in N$, and $f(n) = 5040$. Then the value of n is _____

 [Watch Video Solution](#)

7. $y = f(x)$, where f satisfies the relation $f(x+y) = 2f(x) + xy(y) + y\sqrt{f(x)} \forall x, y \in R$ and $f'(0) = 0$. Then $f(6)$ is equal of

$f(-3)$ is _____

 [Watch Video Solution](#)

8. If function f satisfies the relation $f(x)xf'(-x) = f(-x)xf'(x)$ or $\forall x$, and $f(0) = 3$, and $f(3) = 3$, then the value of $f(-3)$ is _____

 [Watch Video Solution](#)

9. If $y = \frac{a + bx^{\frac{3}{2}}}{x^{\frac{5}{4}}}$ and $y' = 0$ at $x = 5$, then the value of $\frac{a^2}{b^2}$ is _____

 [Watch Video Solution](#)

10. Prove that $\frac{2^{\log_2 2^{1/4}x} - 3^{\log_{27} (x^2+1)^3} - 2x}{7^{4\log_{49}x} - x - 1} > 0, \forall x \in (0, \infty)$.

 [Watch Video Solution](#)

11. $\lim_{h \rightarrow 0} \frac{(e+h)^{\ln(e+h)} - e}{h}$ is-

 [Watch Video Solution](#)

12. If the function $f(x) = -4e^{\frac{1-x}{2}} + 1 + x + \frac{x^2}{2} + \frac{x^3}{3}$ and $g(x) = f^{-1}(x)$, then

the reciprocal of $g' \left(\frac{-7}{6} \right)$ is _____

 [Watch Video Solution](#)

13. Suppose that $f(0) = 0$ and $f'(0) = 2$, and $tg(x) = f(-x + f(f(x)))$. The value of $g'(0)$ is equal to -

 [Watch Video Solution](#)

14. $f(x)$ is polynomial of degree 4 with real coefficients such that $f(x)=0$ satisfied by $x=1, 2, 3$ only then $f'(1)f'(2)f'(3)$ is equal to -

 [Watch Video Solution](#)

15. A nonzero polynomial with real coefficients has the property that $f(x) = f'(x) \cdot f''(x)$. If a is the leading coefficient of $f(x)$, then the value of $1/a$ is -

 Watch Video Solution

16. A function is represented parametrically by the equations

$$x = \frac{1+t}{t^3}; y = \frac{3}{2t^2} + \frac{2}{t} \text{ Then the value of } \left| \frac{dy}{dx} - x \left(\frac{dy}{dx} \right)^3 \right| \text{ is } \text{-----}$$

 Watch Video Solution

17. Let $z = (\cos x)^{\sin x}$ and $y = \sin x$. Then the value of $2 \frac{d^2z}{dy^2} \text{ at } x = \frac{2\pi}{9}$ is _____.

 Watch Video Solution

18. Let $g(x) = \begin{cases} x^2 + x \tan x - x \tan 2x \\ ax + \tan x - \tan 3x \end{cases}$, $x \neq 0$, $x = 0$ if $g'(0)$ exists and is equal

to nonzero value b , then $52 \frac{b}{a}$ is equal to _____

 [Watch Video Solution](#)

19. Let $f(x) = x + \frac{1}{2x + \frac{1}{2x + \frac{1}{2x + \dots \infty}}}$.

Then the value of $f(50) \cdot f'(50)$ is -

 [Watch Video Solution](#)

20. Let $F(x) = f(x)g(x)h(x)$ for all real x , where $f(x)$, $g(x)$, and $h(x)$ are differentiable functions. At some point

x_0 , $F'(x_0) = 21F(x_0)$, $f'(x_0) = 4f(x_0)$, $g'(x_0) = -7g(x_0)$, then the value of $g'(1)$ is _____

 [Watch Video Solution](#)

21. If $y = \frac{\sqrt[3]{1+3x}\sqrt[4]{1+4x}\sqrt[5]{1+5x}}{\sqrt[7]{1+7x}\sqrt[8]{1+8x}}$, then $y'(0)$ is equal to -

 [Watch Video Solution](#)

22. $f'(x) = \phi'(x) = f(x)$ for all x . Also, $f(3) = 5$ and $f'(3) = 4$. Then the value of $[f(10)]^2$ is _____

 [Watch Video Solution](#)

23. If $y=f(x)$ is an odd differentiable function defined on $(-\infty, \infty)$ such that $f'(3) = -2$ then $f'(-3)$ equals -

 [Watch Video Solution](#)

24. If $x^3 + 3x^2 - 9x + c$ is of the form $(x - \alpha)^2(x - \beta)$ then c is equal to

 [Watch Video Solution](#)

25. If graph of $y = f(x)$ is symmetrical about the point $(5, 0)$ and $f'(7) = 3$, then the value of $f'(3)$ is _____

 [Watch Video Solution](#)

26. Let $f(x) = (x - 1)(x - 2)(x - 3)(x - n)$, $n \in N$, and $f(n) = 5040$. Then the value of n is _____

 [Watch Video Solution](#)

27. $y = f(x)$, where f satisfies the relation $f(x + y) = 2f(x) + xy(y) + y\sqrt{f(x)} \forall x, y \in R$ and $f'(0) = 0$. Then $f(6)$ is equal of $f(-3)$ is _____

 [Watch Video Solution](#)

28. If function f satisfies the relation $f(x)xf'(-x) = f(-x)xf'(x)$ or $\forall x$, and $f(0) = 3$, and $f(3) = 3$, then the value of $f(-3)$ is _____



Watch Video Solution

29. If $y = \frac{a + bx^{\frac{3}{2}}}{x^{\frac{5}{4}}}$ and $y' = 0$ at $x = 5$, then the value of $\frac{a^2}{b^2}$ is _____



Watch Video Solution

30. $\lim_{h \rightarrow 0} \frac{(e+h)^{\ln(e+h)} - e}{h}$ is-



Watch Video Solution

31. If the function $f(x) = -4e^{\frac{1-x}{2}} + 1 + x + \frac{x^2}{2} + \frac{x^3}{3}$ and $g(x) = f^{-1}(x)$, then the reciprocal of $g'\left(\frac{-7}{6}\right)$ is _____



Watch Video Solution

32. Suppose that $f(0) = 0$ and $f'(0) = 2$, and let $g(x) = f(-x + f(f(x)))$. The value of $g'(0)$ is equal to _____



Watch Video Solution

33. A nonzero polynomial with real coefficients has the property that $f(x) = f'(x) \cdot f'(x)$. If a is the leading coefficient of $f(x)$, then the value of $1/a$ is -



Watch Video Solution

34. A function is represented parametrically by the equations

$$x = \frac{1+t}{t^3}; y = \frac{3}{2t^2} + \frac{2}{t} \text{ Then the value of } \left| \frac{dy}{dx} - x \left(\frac{dy}{dx} \right)^3 \right| \text{ is } \underline{\hspace{2cm}}$$



Watch Video Solution

35. Let $z = (\cos x)^5$ and $y = \sin x$. Then the value of $2 \frac{d^2z}{dy^2}$ at $x = \frac{2\pi}{9}$ is _____.



Watch Video Solution

36. Let $g(x) = \begin{cases} \frac{x^2 + x \tan x - x \tan 2x}{ax + \tan x - \tan 3x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$ If $g'(0)$ exists and is equal to nonzero value b , then $52 \frac{b}{a}$ is equal to _____



Watch Video Solution

37. Let $f(x) = x + \frac{1}{2x + \frac{1}{2x + \frac{1}{2x + \dots}}}$ Compute the value of $f(50) \cdot f'(50)$



Watch Video Solution

38. Let $F(x) = f(x)g(x)h(x)$ for all real x , where $f(x)$, $g(x)$, and $h(x)$ are differentiable functions. At some point x_0 , $F'(x_0) = 21F(x_0)$, $f'(x_0) = 4f(x_0)$, $g'(x_0) = -7g(x_0)$, then the value of $g'(1)$ is _____

 [Watch Video Solution](#)

39. If $y = \frac{\sqrt[3]{1+3x}\sqrt[4]{1+4x}\sqrt[5]{1+5x}}{\sqrt[7]{1+7x}\sqrt[8]{1+8x}}$, then $y'(0)$ is equal to -

 [Watch Video Solution](#)

40. Let y be an implicit function of x defined by $x^{2x} - 2x^x \cot y - 1 = 0$. Then $y'(1)$ equals: 1 b. $\log 2$ c. $-\log 2$ d. -1

A. -1

B. 1

C. $\log 2$

D. $-\log 2$



Watch Video Solution

41. Let $f: (1, 1) \rightarrow \mathbb{R}$ be a differentiable function with $f(0) = 1$ and $f'(0) = 1$. Let

$g(x) = [f(2f(x) + 2)]^2$. Then $g'(0) =$ (1) 4 (2) 0 (3) 2 (4) 4

A. -2

B. 4

C. -4

D. 0



Watch Video Solution

JEE Previous Year

1. Let y be an implicit function of x defined by $x^{2x} - 2x^x \cot y - 1 = 0$. Then

$y'(1)$ equals: 1 b. $\log 2$ c. $-\log 2$ d. -1

A. -1

B. 1

C. $\log 2$

D. $-\log 2$

Answer: A



[Watch Video Solution](#)

2. Let $f: (1, 1) \rightarrow \mathbb{R}$ be a differentiable function with $f(0) = 1$ and $f'(0) = 1$. Let

$g(x) = [f(2f(x) + 2)]^2$. Then $g'(0) =$ (1) 4 (2) 0 (3) 2 (4) 4

A. -2

B. 4

C. -4

D. 0



Watch Video Solution

3. $\frac{d^2x}{dy^2}$ equals

A. $-\left(\frac{d^2y}{dx^2}\right)\left(\frac{dy}{dx}\right)^{-3}$

B. $\left(\frac{d^2y}{dx^2}\right)^{-1}$

C. $-\left(\frac{d^2y}{dx^2}\right)^{-1}\left(\frac{dy}{dx}\right)^{-3}$

D. $\left(\frac{d^2y}{dx^2}\right)\left(\frac{dy}{dx}\right)^{-2}$



Watch Video Solution

4. If $y = \sec(\tan^{-1}x)$, then $\frac{dy}{dx}$ at $x = 1$ is equal to: $\frac{1}{\sqrt{2}}$ (b) $\frac{1}{2}$ (c) 1 (d) $\sqrt{2}$

A. $1/2$

B. 1

C. $\sqrt{2}$

D. $1\sqrt{2}$



Watch Video Solution

5. If g is the inverse of a function f and $f'(x) = \frac{1}{1+x^5}$ then $g(x)$ is equal to

(1) $1+x^5$ (2) $5x^4$ (3) $\frac{1}{1+\{g(x)\}^5}$ (4) $1+\{g(x)\}^5$

A. $1+x^5$

B. $5x^4$

C. $\frac{1}{1+\{g(x)\}^5}$

D. $1+\{g(x)\}^5$



Watch Video Solution

6. If for $x \left(0, \frac{1}{4}\right)$, the derivative of $\tan^{-1}\left(\frac{6x\sqrt{x}}{1-9x^3}\right)$ is $\sqrt{x}g(x)$, then $g(x)$

equals: (1) $\frac{3x}{1-9x^3}$ (2) $\frac{3}{1+9x^3}$ (3) $\frac{9}{1+9x^3}$ (4) $\frac{3x\sqrt{x}}{1-9x^3}$

A. $\frac{3}{1+9x^3}$

B. $\frac{9}{1+9x^3}$

C. $\frac{3x\sqrt{x}}{1-9x^3}$

D. $\frac{3x}{1-9x^3}$



Watch Video Solution

7. $\frac{d^2x}{dy^2}$ equals

A. $-\left(\frac{d^2y}{dx^2}\right)\left(\frac{dy}{dx}\right)^{-3}$

B. $\left(\frac{d^2y}{dx^2}\right)^{-1}$

C. $-\left(\frac{d^2y}{dx^2}\right)^{-1}\left(\frac{dy}{dx}\right)^{-3}$

D. $\left(\frac{d^2y}{dx^2}\right)\left(\frac{dy}{dx}\right)^{-2}$



Watch Video Solution

8. If $y = \sec(\tan^{-1}x)$, then $\frac{dy}{dx}$ at $x = 1$ is equal to: $\frac{1}{\sqrt{2}}$ (b) $\frac{1}{2}$ (c) 1 (d) $\sqrt{2}$

A. $1/2$

B. 1

C. $\sqrt{2}$

D. $1\sqrt{2}$



Watch Video Solution

9. If g is the inverse of a function f and $f'(x) = \frac{1}{1+x^5}$ then $g(x)$ is equal to

(1) $1+x^5$ (2) $5x^4$ (3) $\frac{1}{1+\{g(x)\}^5}$ (4) $1+\{g(x)\}^5$

A. $1+x^5$

B. $5x^4$

C. $\frac{1}{1+\{g(x)\}^5}$

D. $1+\{g(x)\}^5$



Watch Video Solution

10. If for $x \left(0, \frac{1}{4}\right)$, the derivative of $\tan^{-1}\left(\frac{6x\sqrt{x}}{1-9x^3}\right)$ is $\sqrt{x}g(x)$, then $g(x)$

equals: $\frac{3x}{1-9x^3}$ (2) $\frac{3}{1+9x^3}$ (3) $\frac{9}{1+9x^3}$ (4) $\frac{3x\sqrt{x}}{1-9x^3}$

A. $\frac{3}{1 + 9x^3}$

B. $\frac{9}{1 + 9x^3}$

C. $\frac{3x\sqrt{x}}{1 - 9x^3}$

D. $\frac{3x}{1 - 9x^3}$



Watch Video Solution

11. Let $f(x) = x \sin \pi x$, $x > 0$ Then for all natural numbers n , $f(x)$ vanishes at

A. a unique point in the interval $\left(n, n + \frac{1}{2}\right)$

B. a unique point in the interval $\left(n + \frac{1}{2}, n + 1\right)$

C. a unique point in the interval $(n, n + 1)$

D. two points in the interval $(n, n + 1)$



Watch Video Solution

12. Let $f: (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function such that $f'(x) = 2 - \frac{f(x)}{x}$ for all $x \in (0, \infty)$ and $f(1) = 1$, then

A. $\lim_{x \rightarrow 0^+} f\left(\frac{1}{x}\right) = 1$

B. $\lim_{x \rightarrow 0^+} x f\left(\frac{1}{x}\right) = 2$

C. $\lim_{x \rightarrow 0^+} x^2 f'(x) = 0$

D. $|f(x)| \leq 2$ for all $x \in (0, 2)$

 [Watch Video Solution](#)

13. If $f(\theta) = \sin\left(\tan^{-1}\left(\frac{\sin\theta}{\sqrt{\cos 2\theta}}\right)\right)$, where $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$, then the value of $\frac{d}{d(\tan\theta)} f(\theta)$ is

 [Watch Video Solution](#)

14. The slope of the tangent to the curve $(y - x^5)^2 = x(1 + x^2)^2$ at the point (1, 3) is.

 [Watch Video Solution](#)

15. Let $f: R \rightarrow R$ be a differentiable function with $f(0) = 1$ and satisfying the equation $f(x + y) = f(x)f'(y) + f'(x)f(y)$ for all $x, y \in R$. Then, the value of $(\log)_e(f(4))$ is _____

 [Watch Video Solution](#)

Multiple Correct Answers Type

1. Let $f(x) = x \sin \pi x, x > 0$ Then for all natural numbers n , $f(x)$ vanishes at

A. a unique point in the interval $\left(n, n + \frac{1}{2}\right)$

B. a unique point in the interval $\left(n + \frac{1}{2}, n + 1\right)$

C. a unique point in the interval $(n, n + 1)$

D. two points in the interval $(n, n + 1)$

 [Watch Video Solution](#)

2. Let $f: (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function such that $f'(x) = 2 - \frac{f(x)}{x}$ for all $x \in (0, \infty)$ and $f(1) = 1$, then

A. $\lim_{x \rightarrow 0^+} f\left(\frac{1}{x}\right) = 1$

B. $\lim_{x \rightarrow 0^+} x f\left(\frac{1}{x}\right) = 2$

C. $\lim_{x \rightarrow 0^+} x^2 f'(x) = 0$

D. $|f(x)| \leq 2$ for all $x \in (0, 2)$

 [Watch Video Solution](#)

3. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $h: \mathbb{R} \rightarrow \mathbb{R}$ be differentiable functions such that $f(x) = x^3 + 3x + 2$, $g(f(x)) = x$ and $h(g(x)) = x$ for all $x \in \mathbb{R}$. Then, $h'(1)$ equals.

A. $g'(2) = \frac{1}{15}$

B. $h'(1) = 666$

C. $h(0) = 16$

D. $h(g(3)) = 36$

 [Watch Video Solution](#)

Matrix Match Type

1. Match the statements/expressions given in List I with the values given in List II.



- A. $a \ b \ c \ d$
 $s \ r \ q \ p$
- B. $a \ b \ c \ d$
 $q \ s \ r \ p$
- C. $a \ b \ c \ d$
 $s \ r \ p \ q$
- D. $a \ b \ c \ d$
 $q \ s \ p \ r$



[View Text Solution](#)

Numerical Value Type

1. If $f(\theta) = \sin\left(\tan^{-1}\left(\frac{\sin\theta}{\sqrt{\cos 2\theta}}\right)\right)$, where $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$, then the value of $\frac{d}{d(\tan\theta)} f(\theta)$ is



[Watch Video Solution](#)

2. The slope of the tangent to the curve $(y - x^5)^2 = x(1 + x^2)^2$ at the point $(1, 3)$ is.

 [Watch Video Solution](#)

3. Let $f: R \rightarrow R$ be a differentiable function with $f(0) = 1$ and satisfying the equation $f(x + y) = f(x)f'(y) + f'(x)f(y)$ for all $x, y \in R$. Then, the value of $(\log)_e(f(4))$ is _____

 [Watch Video Solution](#)