



MATHS

BOOKS - CENGAGE MATHS (HINGLISH)

INEQUALITIES INVOLVING MEANS

Single Correct Answer

1. If a, b, c be three positive numbers in $A.P.$ and

$$E = \frac{a + 8b}{2b - a} + \frac{8b + c}{2b - c}, \text{ then a value of } E \text{ can be}$$

A. 16

B. 15

C. 17

D. 21

Answer: D



Watch Video Solution

2. Let $ab = 1$, then the minimum value of $\frac{1}{a^4} + \frac{1}{4b^4}$ is

A. 1

B. 2

C. $1/4$

D. $1/2$

Answer: A



Watch Video Solution

3. If $x > 0, y > 0, z > 0$, the least value of

$x^{\log_e y - \log_e z} + y^{\log_e z - \log_e x} + z^{\log_e x - \log_e y}$ is

A. 3

B. 1

C. 5

D. 6

Answer: A



[Watch Video Solution](#)

4. Let $p, q, r \in R^+$ and $27pqr \geq (p + q + r)^3$ and $3p + 4q + 5r = 12$.

Then the value of $8p + 4q - 7r =$

A. 2

B. 3

C. 4

D. 5

Answer: D



[Watch Video Solution](#)

5. Minimum value of $f(x) = \cos^2 x + \frac{\sec x}{4}$, $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ is

A. $3/2$

B. $3/4$

C. $3/8$

D. none of these

Answer: B



[Watch Video Solution](#)

6. Let $x, y, z \in R^+$ and $2xy + 3yz + 4xz = 18$. If α, β and γ be the values of x, y and z respectively, for which xyz attains its maximum value, then the value of $2\alpha + \beta + \gamma =$

A. 4

B. 6

C. 8

D. 12

Answer: B



Watch Video Solution

7. The minimum value of $\frac{x^4 + y^4 + z^2}{xyz}$ for positive real numbers x, y, z is $\sqrt{2}$ $2\sqrt{2}$ $4\sqrt{2}$ $8\sqrt{2}$

A. $\sqrt{2}$

B. $2\sqrt{2}$

C. $4\sqrt{2}$

D. $8\sqrt{2}$

Answer: B



Watch Video Solution

8. If $n_1, n_2, n_3, \dots, n_{100}$ are positive real numbers such that

$$n_1 + n_2 + n_3 + \dots + n_{100} = 20 \quad \text{and}$$

$$k = n_1(n_2 + n_3 + n_4)(n_5 + n_6 + \dots + n_9)(n_{10} + \dots + n_{16}) \dots (\dots + n_{100})$$

, then k belongs to

A. $(0, 100]$

B. $(0, 128]$

C. $(0, 144]$

D. $(0, 1024]$

Answer: D



Watch Video Solution

9. If a, b, c are the sides of triangle, then the least value of

$$\frac{a}{c+a-b} + \frac{b}{a+b-c} + \frac{c}{b+c-a} \text{ is}$$

A. $1/3$

B. 1

C. 3

D. 6

Answer: B



Watch Video Solution

10. If $x > 0$, $\frac{x^n}{1 + x + x^2 + \dots + x^{2n}}$ is

A. $\leq \frac{1}{2n + 1}$

B. $< \frac{2}{2n + 1}$

C. $\geq \frac{1}{2n + 1}$

D. $> \frac{2}{2n + 1}$

Answer: A



Watch Video Solution

11. If positive quantities a, b, c are in $H. P.$, then which of the following is not true ?

A. $b > \frac{a + c}{2}$

B. $\frac{1}{a - b} - \frac{1}{b - c} > 0$

C. $ac > b^2$

D. none of these

Answer: D



[Watch Video Solution](#)

12. Given that x, y, z are positive real numbers such that $xyz = 32$, the minimum value of $\sqrt{(x + 2y)^2 + 2z^2} - 15$ is equal to

A. 6

B. 8

C. 9

D. 12

Answer: C



[Watch Video Solution](#)

13. If x, y, z are positive real numbers such that $x^2 + y^2 + z^2 = 7$ and $xy + yz + xz = 4$ then the minimum value of xy is

A. 1

B. $\frac{1}{2}$

C. $\frac{1}{4}$

D. $\frac{1}{8}$

Answer: C



[Watch Video Solution](#)

14. If x lies between -5 and 11 , then the greatest value of $(11 - x)^3(x + 5)^5$ is

A. $6^5 \cdot 10^3$

B. $6^3 \cdot 10^3$

C. $6^3 \cdot 10^4$

D. $6^3 \cdot 10^5$

Answer: D



Watch Video Solution

15. The least integral value of

$$f(x) = \frac{(x - 1)^7 + 3(x - 1)^6 + (x - 1)^5 + 1}{(x - 1)^5}, \forall x > 1 \text{ is}$$

A. 8

B. 6

C. 12

D. 18

Answer: B



Watch Video Solution

16. If a, b, c are positive real numbers and $2a + b + 3c = 1$, then the maximum value of $a^4 b^2 c^2$ is equal to

A. $\frac{1}{3 \cdot 4^8}$

B. $\frac{1}{9 \cdot 4^7}$

C. $\frac{1}{9 \cdot 4^8}$

D. $\frac{1}{27 \cdot 4^8}$

Answer: C



Watch Video Solution

17. If x, y, z be three positive numbers such that xyz^2 has the greatest value $\frac{1}{64}$, then the value of $\frac{1}{x} + \frac{1}{y} + \frac{1}{z}$ is

A. 6

B. 8

C. 10

D. 12

Answer: C



Watch Video Solution

18. If x_1, x_2 and x_3 are the positive roots of the equation $x^3 - 6x^2 + 3px - 2p = 0$, πnR , then the value of $\sin^{-1}\left(\frac{1}{x_1} + \frac{1}{x_2}\right) + \cos^{-1}\left(\frac{1}{x_2} + \frac{1}{x_3}\right) - \tan^{-1}\left(\frac{1}{x_3} + \frac{1}{x_1}\right)$ is equal to

A. $\frac{\pi}{4}$

B. $\frac{\pi}{2}$

C. $\frac{3\pi}{4}$

D. π

Answer: A



Watch Video Solution

Comprehension

1. If a, b, x, y are real number and $x, y > 0$, then $\frac{a^2}{x} + \frac{b^2}{y} \geq \frac{(a+b)^2}{x+y}$

so on solving it we have $(ay - bx)^2 \geq 0$.

Similarly, we can extend the inequality to three pairs of numbers, i.e,

$$\frac{a^2}{x} + \frac{b^2}{y} + \frac{c^2}{z} \geq \frac{(a+b+c)^2}{x+y+z}$$

Now use this result to solve the following questions.

The value of $\frac{a^2 + b^2}{a + b} + \frac{b^2 + c^2}{b + c} + \frac{a^2 + c^2}{a + c}$ is

A. $\geq (a + b + c)$

$$B. \geq \frac{1}{2}(a + b + c)$$

$$C. \frac{3}{2} \leq (a + b + c)$$

D. None of these

Answer: A



Watch Video Solution

2. If a, b, x, y are real number and $x, y > 0$, then $\frac{a^2}{x} + \frac{b^2}{y} \geq \frac{(a + b)^2}{x + y}$
so on solving it we have $(ay - bx)^2 \geq 0$.

Similarly, we can extend the inequality to three pairs of numbers, i.e,

$$\frac{a^2}{x} + \frac{b^2}{y} + \frac{c^2}{z} \geq \frac{(a + b + c)^2}{x + y + z}$$

Now use this result to solve the following questions.

If $abc = 1$, then the minimum value of

$$\frac{1}{a^3(b + c)} + \frac{1}{b^3(a + c)} + \frac{1}{c^3(a + b)}$$
 is

A. 3

B. $3/2$

C. 6

D. 9

Answer: B

 [Watch Video Solution](#)

Examples

1. Prove that $(ab + xy)(ax + by) > 4abxy$ ($a, b, x, y > 0$).

 [Watch Video Solution](#)

2. Prove that $b^2c^2 + c^2a^2 + a^2b^2 > abc(a + b + c)$, where $a, b, c > 0$.

 [Watch Video Solution](#)

3. Prove that $\frac{2}{b+c} + \frac{2}{c+a} + \frac{2}{a+b} > \left\langle \frac{1}{a} + \frac{1}{b} + \frac{1}{c}, \text{ where } a, b, c > 0 \right\rangle$.



Watch Video Solution

4. If $a, b, \text{ and } c$ are distinct positive real numbers such that $a + b + c = 1$, then prove that $\frac{(1+a)(1+b)(1+c)}{(1-a)(1-b)(1-c)} > 8$.



Watch Video Solution

5. Find the minimum value of $4^{\sin^{-1}(2x)} + 4^{\cos^{-1}(2x)}$.



Watch Video Solution

6. If $(\log)_2(a+b) + (\log)_2(c+d) \geq 4$. Then find the minimum value of the expression $a + b + c + \dots$



Watch Video Solution

7. Find all real solutions to $2^x + x^2 = 2 - \frac{1}{2^x}$.



[Watch Video Solution](#)

8. Find all positive real solutions to

$$4x + \frac{18}{y} = 14, 2y + \frac{9}{z} = 15, 9z + \frac{16}{x} = 17.$$



[Watch Video Solution](#)

9. Let A;G;H be the arithmetic; geometric and harmonic means between three given no. a;b;c then the equation having a;b;c as its root is

$$x^3 - 3Ax^2 + 3\frac{G^3}{H}x - G^3 = 0$$



[Watch Video Solution](#)

10. Let a_1, a_2, \dots be positive real numbers in geometric progression. For each n, let A_n, G_n, H_n , be respectively the arithmetic mean, geometric mean & harmonic mean of a_1, a_2, \dots, a_n . Find an

expression ,for the geometric mean of G_1, G_2, \dots, G_n in terms of $A_1, A_2, \dots, A_n, H_1, H_2, \dots, H_n$.

 [Watch Video Solution](#)

11. If $a + b + c = 1$, then prove that

$$\frac{8}{27abc} > \left\{ \frac{1}{a} - 1 \right\} \left\{ \frac{1}{b} - 1 \right\} \left\{ \frac{1}{c} - 1 \right\} > 8.$$

 [Watch Video Solution](#)

12. If $yz + zx + xy = 12$, where x, y, z are positive values, find the greatest value of xyz .

 [Watch Video Solution](#)

13. If a, b, c are positive, then prove that

$$a/(b + c) + b/(c + a) + c/(a + b) \geq 3/2.$$

 [Watch Video Solution](#)

14. Prove that $2^n > 1 + n\sqrt{2^{n-1}}$, $\forall n > 2$ where n is a positive integer.

 [Watch Video Solution](#)

15. If $S + a_1 + a_2 + a_3 + \dots + a_n$, $a_i \in R^+$ for $i = 1 \rightarrow n$, then prove

that
$$\frac{S}{S - a_1} + \frac{S}{S - a_2} + \dots + \frac{S}{S - a_n} \geq \frac{n^2}{n - 1}, \forall n \geq 2$$

 [Watch Video Solution](#)

16. If $a_1 + a_2 + a_3 + \dots + a_n = 1$, $\forall a_i > 0, i = 1, 2, 3, \dots, n$, then find the maximum value of $a_1 a_2 a_3 a_4 a_5 a_n$.

 [Watch Video Solution](#)

17. If a, b, c , are positive real numbers, then prove that (2004, 4M)

$$\{(1 + a)(1 + b)(1 + c)\}^7 > 7^7 a^4 b^4 c^4$$

 [Watch Video Solution](#)

18. Prove that $\frac{\sec^4 \alpha}{\tan^2 \beta} + \frac{\sec^4 \beta}{\tan^2 \alpha} \geq 8$. If each term in the expression is well defined.

 [Watch Video Solution](#)

19. Prove that

$$\left[\frac{x^2 + y^2 + z^2}{x + y + z} \right]^{x+y+z} > x^x y^y z^z > \left[\frac{x + y + z}{3} \right]^{x+y+z} \quad (x, y, z > 0)$$

 [View Text Solution](#)

20. Prove that $1^1 \times 2^2 \times 3^3 \times \dots \times n^n < \left(\frac{2n+1}{3} \right)^{\frac{n(+1)}{2}}$

 [Watch Video Solution](#)

21. Find the greatest value of x^2y^3 , where x and y lie in the first quadrant on the line $3x + 4y = 5$.

 [Watch Video Solution](#)

22. Find the maximum value of $(7 - x)^4(2 + x)^5$ when x lies between -2 and 7 .

 [Watch Video Solution](#)

23. Find the maximum value of xyz when $\frac{x}{1} + \frac{y^2}{4} + \frac{z^3}{27} = 1$, where $x, y, z > 0$.

 [Watch Video Solution](#)

24. If $a, c > 0$ such that $a^3 + b^3 = 2$, then show that $a + b \leq 2$.

 [Watch Video Solution](#)

25. If $m > 1, n \in \mathbb{N}$ show that

$$1m + 2m + 2^{2m} + 2^{3m} + \dots + 2^{nm-m} > n^{i-m}(2^n - 1)^m.$$

 [Watch Video Solution](#)

26. Prove that n in acute angled triangle

$$ABC, \sec A + \sec B + \sec C \geq 6.$$

 [Watch Video Solution](#)

27. Prove that $\frac{b^2 + c^2}{b + c} + \frac{c^2 + a^2}{c + a} + \frac{a^2 + b^2}{a + b} > a + b + c$, where $a, b, c > 0$.

 [Watch Video Solution](#)

28. Prove that $\frac{a^8 + b^8 + c^8}{a^3 b^3 c^3} > \frac{1}{a} + \frac{1}{b} + \frac{1}{c}$

 [Watch Video Solution](#)

29. If $a, b,$ and c are positive and $a + b + c = 6$, show that $(a + 1/b)^2 + (b + 1/c)^2 + (c + 1/a)^2 \geq 75/4$.

 [Watch Video Solution](#)

30. Let x_1, x_2, \dots, x_n be positive real numbers and we define

$$S = x_1 + x_2 + \dots + x_n.$$

Prove

that

$$(1 + x_1)(1 + x_2)\dots(1 + x_n) \leq 1 + S + \frac{S^2}{2!} + \frac{S^3}{3!} + \dots + \frac{S^n}{n!}$$

 [Watch Video Solution](#)

31. If $2x^3 + ax^2 + bx + 4 = 0$ (a and b are positive real numbers) has 3 real roots, then prove that $a + b \geq 6\left(2^{\frac{1}{3}} + 4^{\frac{1}{3}}\right)$

 [Watch Video Solution](#)

32. Calculate the greatest and least values of the function

$$f(x) = \frac{x^4}{x^8 + 2x^6 - 4x^4 + 8x^2 + 16}$$

 [View Text Solution](#)

33. If a, b, c are three distinct positive real numbers in G.P., then prove that

$$c^2 + 2ab > 3ac.$$

 [Watch Video Solution](#)

34. If x and y are real numbers such that $16^{x^2+y} + 16^{x+y^2} = 1$, then find the values of x and y .

 [Watch Video Solution](#)

35. In how many parts an integer $N \geq 5$ should be dissected so that the product of the parts is maximized.

 [Watch Video Solution](#)

36. If $x + y + z = 1$ and x, y, z are positive, then show that

$$\left(x + \frac{1}{x}\right)^2 + \left(y + \frac{1}{y}\right)^2 + \left(z + \frac{1}{z}\right)^2 > \frac{100}{3}$$

 [Watch Video Solution](#)

37. If a, b, c are three positive real numbers, then find minimum value of

$$\frac{a^2 + 1}{b + c} + \frac{b^2 + 1}{c + a} + \frac{c^2 + 1}{a + b}$$

 [Watch Video Solution](#)

Exercise 6 1

1. Prove that $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} \geq \frac{1}{\sqrt{(bc)}} + \frac{1}{\sqrt{(ca)}} + \frac{1}{\sqrt{(ab)}}$, where a, b, c

> 0

 [Watch Video Solution](#)

2. If a, b, c are real numbers such that $0 < a < 1, 0 < b < 1, 0 < c < 1, a + b + c = 2$, then prove that

$$\frac{a}{1-a} \frac{b}{1-b} \frac{c}{1-c} \geq 8$$

.

 [Watch Video Solution](#)

3. If x, y are positive real numbers and m, n are positive integers, then

prove that
$$\frac{x^n y^m}{(1+x^{2n})(1+y^{2m})} \leq \frac{1}{4}$$

 [Watch Video Solution](#)

4. If $a_i > 0$ ($i = 1, 2, 3, \dots, n$) prove that

$$\sum_{I \geq J \geq j \geq n} \sqrt{a_i a_j} \leq \frac{n-1}{2} (a_1 + a_2 + \dots + a_n)$$

 [Watch Video Solution](#)

5. Find the minimum value of $2^{\sin x} + 2^{\cos x}$

 [Watch Video Solution](#)

6. If $(\log)_{10}(x^3 + y^3) - (\log)_{10}(x^2 + y^2 - xy) \leq 2$, where x, y are positive real number, then find the maximum value of xy .

 [Watch Video Solution](#)

7. Prove that the greatest value of xy is $c^3 / \sqrt{2ab}$ if $a^2x^4 + b^4y^4 = c^6$.

 [Watch Video Solution](#)

8. If $x, y \in R^+$ such that $x + y = 8$, then find the minimum value of $\left(1 + \frac{1}{x}\right)\left(1 + \frac{1}{y}\right)$

 [Watch Video Solution](#)

Exercise 6 2

1. Prove that $\left(\frac{n+1}{2}\right)^n > n!$

 [Watch Video Solution](#)

2. If $a_1, a_2, \dots, a_n > 0$, then prove that

$$\frac{a_1}{a_2} + \frac{a_2}{a_3} + \frac{a_3}{a_4} + \dots + \frac{a_{n-1}}{a_n} + \frac{a_n}{a_1} > n$$

 [Watch Video Solution](#)

3. In ABC , prove that $\tan A + \tan B + \tan C \geq 3\sqrt{3}$, where A, B, C are acute angles.

 [Watch Video Solution](#)

4. If $n \geq 1$ is a positive integer, then prove that $3^n \geq 2^n + n \cdot 6^{\frac{n-1}{2}}$



[Watch Video Solution](#)

[View Text Solution](#)

5. If $abcd = 1$ where a, b, c, d are positive reals then the minimum value of $a^2 + b^2 + c^2 + d^2 + ab + ac + ad + bc + bd + cd$ is

[Watch Video Solution](#)

6. If $x, y, z > 0$ and $x + y + z = 1$, the prove that

$$\frac{2x}{1-x} + \frac{2y}{1-y} + \frac{2z}{1-z} \geq 3.$$

[Watch Video Solution](#)

Exercise 6 3

1. Prove that

$$\left[\frac{a^2 + b^2}{a + b} \right]^{a+b} > a^a b^b > \left\{ \frac{a + b}{2} \right\}^{a+b}, \text{ where } a, b > 0$$

[Watch Video Solution](#)

2. Prove that $a^p b^q > \left(\frac{ap + bq}{p + q} \right)^{p+q}$.

 [Watch Video Solution](#)

3. Prove that $px^{q-r} + qx^{r-p} + rx^{p-q} > p + q + r$, where p, q, r are distinct and $x \neq 1$.

 [Watch Video Solution](#)

4. Given are positive rational numbers a, b, c such that $a + b + c = 1$, then prove that $a^a b^b c^c + a^b b^a c^c + a^c b^a c^b \leq 1$.

 [Watch Video Solution](#)

5. If a and b are positive numbers such that $a^2 + b^2 = 4$, then find the maximum value of $a^2 b$.

 [Watch Video Solution](#)

6. Find the greatest value of $x^2y^3z^4$ if $x^2 + y^2 + z^2 = 1$, where x, y, z are positive.

 [Watch Video Solution](#)

Exercise 6 4

1. Prove that $a^4 + b^4 + c^4 > abc(a + b + c)$, where $a, b, c > 0$.

 [Watch Video Solution](#)

2. If $C_r = \frac{n!}{r!(n-r)!}$, then prove that

$$\sqrt{C_1} + \sqrt{C_2} + \dots + \sqrt{C_n} \sqrt{n(2^n - 1)}$$

 [Watch Video Solution](#)

3. If $a + b = 1$, $a > 0$, $b > 0$, prove that $\left(a + \frac{1}{a}\right)^2 + \left(b + \frac{1}{b}\right)^2 \geq \frac{25}{2}$



Watch Video Solution

4. In $\triangle ABC$, prove that $\sin A + \sin B + \sin C \geq \frac{3\sqrt{3}}{2}$



View Text Solution

Exercise Single

1. The minimum value of $\frac{x^4 + y^4 + z^4}{xyz}$ for positive real numbers x, y, z is

$$\sqrt{2} \quad 2\sqrt{2} \quad 4\sqrt{2} \quad 8\sqrt{2}$$

A. $\sqrt{2}$

B. $2\sqrt{2}$

C. $4\sqrt{2}$

D. $8\sqrt{2}$

Answer: B



Watch Video Solution

2. A rod of fixed length k slides along the coordinates axes, If it meets the axes at $A(a, 0)$ and $B(0, b)$, then the minimum value of

$$\left(a + \frac{1}{a}\right)^2 + \left(b + \frac{1}{b}\right)^2 \text{ is } k^2 = 4 + \frac{4}{k^2} k^2 + 4 + \frac{4}{k^2}$$

A. 0

B. 8

C. $k^2 - 4 + \frac{4}{k^2}$

D. $k^2 + 4 + \frac{4}{k^2}$

Answer: D



Watch Video Solution

3. The least value of $6 \tan^2 \varphi + 54 \cot^2 \varphi + 18$ is 54 when A.M. \geq GM. Is applicable for $6 \tan^2 \varphi, 54 \cot^2 \varphi, 18$ 54 when A.M. \geq GM. Is applicable for $6 \tan^2 \varphi, 54 \cot^2 \varphi, 18$ is added further 78 when $\tan^2 \varphi = \cot^2 \varphi$ (I) is correct, (II) is false (I) and (II) are correct (III) is correct None of the above are correct

A. (I) is correct, (II) is false

B. (I) and (II) are correct

C. (III) is correct

D. (III) is correct

Answer: B

 [Watch Video Solution](#)

4. If $ab^2c^3, a^2b^3c^4, a^3b^4c^5$ are in A.P. ($a, b, c > 0$), then the minimum value of $a + b + c$ is 1359

A. 1

B. 3

C. 5

D. 9

Answer: B

 [Watch Video Solution](#)

5. If $y = 3^{x-1} + 3^{-x-1}$, then the least value of y is $2\frac{2}{3}$ $\frac{3}{2}$

A. 2

B. 6

C. $\frac{2}{3}$

D. $\frac{3}{2}$

Answer: C

 [Watch Video Solution](#)

6. Minimum value of $(b + c)/a + (c + a)/b + (a + b)/c$ (for real positive numbers a, b, c) is

A. 1

B. 2

C. 3

D. 6

Answer: D



[Watch Video Solution](#)

7. If the product of n positive numbers is n^n , then their sum is a positive integer b. divisible by n equal to $n + 1/n$ never less than n^2

A. a positive integer

B. divisible by n

C. equal to $n + l/n$

D. never less than n^2

Answer: D



[Watch Video Solution](#)

8. The minimum value of $P = bcx + cay + abz$, when $xyz = abc$, is

A. $3abc$

B. $6abc$

C. abc

D. $4abc$

Answer: A



[Watch Video Solution](#)

9. If l, m, n are the three positive roots of the equation $x^3 - ax^2 + bx - 48 = 0$, then the minimum value of $(1/l) + (2/m) + (3/n)$ equals $1 \frac{3}{2}$ or $5/2$

A. 1

B. 2

C. $3/2$

D. $5/2$

Answer: C



[Watch Video Solution](#)

10. If positive numbers a, b, c are in H.P., then equation $x^2 - kx + 2b^{101} - a^{101} - c^{101} = 0 (k \in R)$ has both roots positive both roots negative one positive and one negative root both roots imaginary

A. both roots positive

B. both roots

C. one positive and one negative root

D. both roots imaginary

Answer: C



Watch Video Solution

11. For $x^2 - (a + 3)|x| + 4 = 0$ to have real solutions, the range of a is
 $(-\infty, -7] \cup [1, \infty)$ $(-3, \infty)$ $(-\infty, -7]$ $[1, \infty)$

A. $(-\infty, -7] \cup [1, \infty)$

B. $(-3, \infty)$

C. $(-\infty, -7]$

D. $[1, \infty)$

Answer: D



Watch Video Solution

12. If a, b, c are the sides of a triangle, then the minimum value of

$$\frac{a}{b+c-a} + \frac{b}{c+a-b} + \frac{c}{a+b-c}$$
 is equal to 3 6 9 12

- A. 3
- B. 6
- C. 9
- D. 12

Answer: A



[Watch Video Solution](#)

13. If $a, b, c, d \in R^+ - \{1\}$, then the minimum value of

$$\log_d a + \log_c b + \log_a c + \log_b d$$
 is

- A. 4
- B. 2

C. 1

D. none of these

Answer: A



Watch Video Solution

14. If $a, b, c \in R^+$, then $\frac{bc}{b+c} + \frac{ac}{a+c} + \frac{ab}{a+b}$ is always
 $\leq \frac{1}{2}(a+b+c) \geq \frac{1}{3}\sqrt{abc} \leq \frac{1}{3}(a+b+c) \geq \frac{1}{2}\sqrt{abc}$

A. $\leq \frac{1}{2}(a+b+c)$

B. $\geq \frac{1}{2}\sqrt{abc}$

C. $\leq \frac{1}{3}(a+b+c)$

D. $\geq \frac{1}{2}\sqrt{abc}$

Answer: A



Watch Video Solution

15. If $a, b, c \in R^+$ then $(a + b + c) \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right)$ is always $\geq 12 \geq 9$
 ≤ 12 none of these

A. ≥ 12

B. ≥ 9

C. ≤ 12

D. none of these

Answer: B



Watch Video Solution

16. If $a, b, c \in R^+$, then the minimum value of $a(b^2 + c^2) + b(c^2 + a^2) + c(a^2 + b^2)$ is equal to abc $2abc$ $3abc$ $6abc$

A. abc

B. $2abc$

C. $3abc$

D. $6abc$

Answer: D



[Watch Video Solution](#)

17. If $a, b, c, \in R^+$, and a, b, c, d are in H.P. then

A. $a + d > b + c$

B. $a + b > c + d$

C. $a + c > b + d$

D. none of these

Answer: A



[Watch Video Solution](#)

18. If $a, b, c, \in R^+$, such that $a + b + c = 18$, then the maximum value of a^2, b^3, c^4 is equal to

A. $2^{18} \times 3^2$

B. $2^{18} \times 3^3$

C. $2^{19} \times 3^2$

D. $2^{19} \times 3^3$

Answer: D



[Watch Video Solution](#)

19. $f(x) = \frac{(x-2)(x-1)}{x-3}, \forall x > 3$. The minimum value of $f(x)$ is equal to

A. $3 + 2\sqrt{2}$

B. $3 + 2\sqrt{3}$

C. $3\sqrt{2} + 2$

D. $3\sqrt{2} - 2$

Answer: A



[Watch Video Solution](#)

20. If $a > 0$, the least value of $(a^3 + a^2 + a + 1)^2$ is

A. $64a^2$

B. $16a^4$

C. $16a^3$

D. none of these

Answer: C



[Watch Video Solution](#)

Exercise Multiple

1. If A is the area and $2s$ the sum of the sides of a triangle, then

A. $A \leq \frac{s^2}{4}$

B. $A \leq \frac{s^2}{3\sqrt{3}}$

C. $A < \frac{s^2}{\sqrt{3}}$

D. none of these

Answer: A::B



Watch Video Solution

2. If x, y, z are positive numbers in A.P., then

A. $y^2 \geq xz$

B. $xy + yz \geq 2xz$

C. $\frac{x+y}{2y-x} + \frac{y+z}{2y-z} \geq 4$

D. none of these

Answer: A::C



Watch Video Solution

3. For positive real numbers a, b, c such that $a + b + c = p$, which one holds? $(p - a)(p - b)(p - c) \leq \frac{8}{27}p^3$ $(p - a)(p - b)(p - c) \geq 8abc$

$\frac{bc}{a} + \frac{ca}{b} + \frac{ab}{c} \leq p$ *none of these*

A. $(p - a)(p - b)(p - c) \leq \frac{8}{27}p^3$

B. $(p - a)(p - b)(p - c) > 8abc$

C. $\frac{bc}{a} + \frac{ca}{b} + \frac{ab}{c} \leq p$

D. none of these

Answer: A::B



Watch Video Solution

4. If first and $(2n - 1)^{th}$ terms of A.P., G.P. and H.P. are equal and their n th terms are a, b, c respectively, then

A. $a = b = c$

B. $a + c = b$

C. $a > b > c$

D. $ac - b^2 = 0$

Answer: C::D



Watch Video Solution

5. If $a > 0, b > 0, c > 0$ and $2a + b + 3c = 1$, then

A. $a^4 b^2 c^2$ is greatest then $a = \frac{1}{4}$

B. $a^4 b^2 c^2$ is greatest then $b = \frac{1}{4}$

C. $a^4 b^2 c^2$ is greatest then $c = \frac{1}{12}$

D. greatest value of $a^4 b^2 c^2$ is $\frac{1}{9.4^8}$

Answer: A::B::C::D



Watch Video Solution

Exercise Comprehension

1. If roots of the equation

$$f(x) = x^6 - 12x^5 + bx^4 + cx^3 + dx^2 + ex + 64 = 0$$
 are positive, then

Which has the greatest absolute value ?

A. b

B. c

C. d

D. e

Answer: C



Watch Video Solution

2. If roots of the equation

$$f(x) = x^6 - 12x^5 + bx^4 + cx^3 + dx^2 + ex + 64 = 0$$
 are positive, then

Which has the greatest absolute value ?

A. b

B. c

C. d

D. e

Answer: A



[Watch Video Solution](#)

3. If roots of the equation

$$f(x) = x^6 - 12x^5 + bx^4 + cx^3 + dx^2 + ex + 64 = 0$$
 are positive, then

remainder when $f(x)$ is divided by $x - 1$ is

A. 2

B. 1

C. 3

D. 10

Answer: B



[Watch Video Solution](#)

4. Equation $x^4 + ax^3 + bx^2 + cx + 1 = 0$ has real roots (a,b,c are non-negative).

Minimum non-negative real value of a is

A. 10

B. 9

C. 6

D. 4

Answer: D



[Watch Video Solution](#)

5. Equation $x^4 + ax^3 + bx^2 + cx + 1 = 0$ has real roots (a,b,c are non-negative).

Minimum non-negative real value of b is

A. 12

B. 15

C. 6

D. 10

Answer: C



[Watch Video Solution](#)

6. Equation $x^4 + ax^3 + bx^2 + cx + 1 = 0$ has real roots (a,b,c are non-negative). Minimum non-negative real value of c is

A. 10

B. 9

C. 6

D. 4

Answer: D

 [Watch Video Solution](#)

Exercise Numerical Jee Previous Year

1. For $x \geq 0$, the smallest value of the function $f(x) = \frac{4x^2 + 8x + 13}{6(1+x)}$,
is _____.

 [Watch Video Solution](#)

2. Let $x^2 - 3x + p = 0$ has two positive roots a and b , the minimum value of $\left(\frac{4}{a} + \frac{1}{b}\right)$ is _____.

 [Watch Video Solution](#)

3. If $x, y,$ and z are positive real numbers and $x = \frac{12 - yz}{y + z}$. The maximum value of (xyz) equals _____.

 [Watch Video Solution](#)

4. If $a, b,$ and c are positive and $9a + 3b + c = 90$, then the maximum value of $(\log a + \log b + \log c)$ is (base of the logarithm is 10) _____.

 [Watch Video Solution](#)

5. Given that x, y, z are positive reals such that $xyz = 32$. The minimum value of $x^2 + 4xy + 4y^2 + 2z^2$ is _____.

 [Watch Video Solution](#)

6. If $x, y, \in R^+$ satisfying $x + y = 3$, then the maximum value of x^2y is _____.

 [Watch Video Solution](#)

7. For any $x, y, \in R, xy > 0$. Then the minimum value of $\frac{2x}{y^3} + \frac{x^3y}{3} + \frac{4y^2}{9x^4}$ is _____.

 [Watch Video Solution](#)

8. Let a, b, c, d and e be positive real numbers such that $a + b + c + d + e = 15$ and $ab^2c^3d^4e^5 = (120)^3 \times 50$. Then the value of $a^2 + b^2 + c^2 + d^2 + e^2$ is _____.

 [Watch Video Solution](#)

9. Consider the system of equations $x_1 + x_2^2 + x_3^3 + x_4^4 + x_5^5 = 5$ and $x_1 + 2x_2 + 3x_3 + 4x_4 + 5x_5 = 15$ where x_1, x_2, x_3, x_4, x_5 are positive real numbers. Then numbers of $(x_1, x_2, x_3, x_4, x_5)$ is _____.

 [Watch Video Solution](#)

10. The least value of $a \in \mathbb{R}$ for which $4ax^2 + \frac{1}{x} \geq 1$, for all $x > 0$, is

A. $\frac{1}{64}$

B. $\frac{1}{32}$

C. $\frac{1}{27}$

D. $\frac{1}{25}$

Answer: C

 [Watch Video Solution](#)

11. The minimum value of the sum of real number a^{-5} , a^{-4} , $3a^{-3}$, 1 , a^8 , and a^{10} with $a > 0$ is



[Watch Video Solution](#)