



MATHS

BOOKS - CENGAGE MATHS (HINGLISH)

JEE 2019

Chapter 1 Coordinate System

1. If two vertices of a triangle are $(0, 2)$ and $(4, 3)$ and its orthocentre is $(0, 0)$ then the third vertex of the triangle lies in (a) I^{st} quadrant (b) 2^{nd} quadrant (c) 3^{rd} quadrant (d) 4^{th} quadrant

A. Fourth

B. Second

C. Third

D. First

Answer: B



Watch Video Solution

2. If the straight line $2x-3y+17=0$ is perpendicular to the line passing through the points $(7, 17)$ and $(15, \beta)$, then β equals

A. -5

B. $-\frac{35}{3}$

C. $\frac{35}{3}$

D. 5

Answer: D



Watch Video Solution

1. Consider the set of all lines $px+qy+r=0$ such that $3p+2q+4r=0$. Which one of the following statements is true ?

A. The lines are all parallel.

B. Each line passes through the origin.

C. The lines are not concurrent.

D. The lines are concurrent at the point $\left(\frac{3}{4}, \frac{1}{2}\right)$.

Answer: D



[Watch Video Solution](#)

2. The equations of two sides of a triangle are $3x - 2y + 6 = 0$ and $4x + 5y - 20 = 0$ and the orthocentre is (1,1). Find the equation of the third side.

A. $122t-26x-1675=0$

B. $26x+61y+1675=0$

C. $122y+26x+1675=0$

D. $26x-122y-1675=0$

Answer: D



Watch Video Solution

3. A line $4x + 3y = 24$ cut the x-axis at point A and cut the y-axis at point B then incentre of triangle OAB is (a) $(4, 4)$ (b) $(4, 3)$ (c) $(3, 4)$ (d) $(2, 2)$

A. $(3, 4)$

B. $(2,2)$

C. $(4,4)$

D. $(4,3)$

Answer: B



Watch Video Solution

4. A point P moves on line $2x - 3y + 4 = 0$ If $Q(1, 4)$ and $R(3, -2)$ are fixed points, then the locus of the centroid of $\triangle PQR$ is a line: (a) with slope $\frac{3}{2}$ (b) parallel to y-axis (c) with slope $\frac{2}{3}$ (d) parallel to x-axis

A. parallel to x-axis

B. with slope $\frac{2}{3}$

C. with slope $\frac{3}{2}$

D. parallel to y-axis

Answer: B



[Watch Video Solution](#)

5. Two sides of a parallelogram are along the lines $x+y=3$ and $x=y+3$. If its diagonals intersect at $(2, 4)$, then one of its vertices is

A. $(2, 6)$

B. $(2,1)$

C. (3, 5)

D. (6,3)

Answer: D



[Watch Video Solution](#)

6. If in parallelogram $ABDC$, the coordinate of A , B and C are respectively $(1, 2)$, $(3, 4)$ and $(2, 5)$, then the equation of the diagonal AD is

A. $5x+3y-11=0$

B. $3x-5y+7=0$

C. $3x+5y-13=0$

D. $5x-3y+1=0$

Answer: D



[Watch Video Solution](#)

7. If a straight line passing through the point $P(-3,4)$ is such that its intercepted portion between the coordinate axes is bisected at P , then its equation is

A. $x-y+7=0$

B. $3x-4y+25=0$

C. $4x+3y=0$

D. $4x-3y+24=0$

Answer: D

 [Watch Video Solution](#)

8. The tangent to the curve $y = x^2 - 5x + 5$, parallel to the line $2y = 4x + 1$, also passes through the point :

A. $\left(\frac{1}{4}, \frac{7}{2}\right)$

B. $\left(\frac{7}{2}, \frac{1}{4}\right)$

C. $\left(-\frac{1}{8}, 7\right)$

D. $\left(\frac{1}{8}, -7\right)$

Answer: D



Watch Video Solution

Chapter 4 Circle

1. 3 circles of radii a, b, c (a

A. $(1)(\sqrt{a}) = \frac{1}{\sqrt{b}} + \frac{1}{\sqrt{c}}$

B. a, b, c are in A.P.

C. $\sqrt{a}, \sqrt{b}, \sqrt{c}$ are in A. P.

D. $\frac{1}{\sqrt{b}} = \frac{1}{\sqrt{a}} + \frac{1}{\sqrt{c}}$

Answer: A



Watch Video Solution

2. If the circles $x^2 + y^2 - 16x - 20y + 164 = r^2$ and $(x - 4)^2 + (y - 7)^2 = 36$ intersect at two points then (a) $1 < r < 11$ (b) $r = 11$ (c) $r > 11$ (d) $0 < r < 1$

A. $0 < r < 1$

B. $1 < r < 11$

C. $r < 11$

D. $r = 11$

Answer: B



Watch Video Solution

3. If a circle C passing through $(4, 0)$ touches the circle $x^2 + y^2 + 4x - 6y - 12 = 0$ externally at a point $(1, -1)$, then the radius of the circle C is :-

A. $\sqrt{57}$

B. 4

C. $2\sqrt{5}$

D. 5

Answer: D



Watch Video Solution

4. A square is inscribed in a circle $x^2 + y^2 - 6x + 8y - 103 = 0$ such that its sides are parallel to co-ordinate axis then the distance of the nearest vertex to origin, is equal to (A) 13 (B) $\sqrt{127}$ (C) $\sqrt{41}$ (D) 1

A. 13

B. $\sqrt{137}$

C. 6

D. $\sqrt{41}$

Answer: D



Watch Video Solution

5. A line $2x + y = 1$ intersect co-ordinate axis at points A and B . A circle is drawn passing through origin and point A & B . If perpendicular from point A and B are drawn on tangent to the circle at origin then sum of perpendicular distance is (A) $\frac{5}{\sqrt{2}}$ (B) $\frac{\sqrt{5}}{2}$ (C) $\frac{\sqrt{5}}{4}$ (D) $\frac{5}{2}$

A. $\frac{\sqrt{5}}{4}$

B. $\frac{\sqrt{5}}{2}$

C. $2\sqrt{5}$

D. $24\sqrt{5}$

Answer: B



Watch Video Solution

6. Two circles with equal radii are intersecting at the points $(0, 1)$ and $(0, -1)$. The tangent at the point $(0, 1)$ to one of the circles passes through the centre of the other circle. Then the distance between the centres of these circles is.

A. 1

B. $\sqrt{2}$

C. $2\sqrt{2}$

D. 2

Answer: D



[Watch Video Solution](#)

7. A circle cuts the chord on x-axis of length $4a$. If this circle cuts the y-axis at a point whose distance from origin is $2b$. Locus of its centre is (A) Ellipse (B) Parabola (C) Hyperbola (D) Straight line

A. A hyperbola

B. A parabola

C. A straight line

D. An ellipse

Answer: B



Watch Video Solution

8. If a variable line $3x + 4y - \lambda = 0$ is such that the two circles $x^2 + y^2 - 2x - 2y + 1 = 0$ and $x^2 + y^2 - 18x - 2y + 78 = 0$ are on its opposite sides, then the set of all values of λ is the interval

A. [12,21]

B. (2, 17)

C. (23, 31)

D. [13, 23]

Answer: A



Watch Video Solution

9. Let $x^2 + y^2 - 2x - 2y - 2 = 0$ and $x^2 + y^2 - 6x - 6y + 14 = 0$ are two circles C_1, C_2 are the centre of circles and circles intersect at P, Q find the area of quadrilateral C_1PC_2Q (A) 12 (B) 6 (C) 8 (D) 4

A. 8

B. 6

C. 9

D. 4

Answer: D



Watch Video Solution

10. A circle of radius 'r' passes through the origin O and cuts the axes at A and B, Locus of the centroid of triangle OAB is

A. $(x^2 + y^2)^2 = 4Rx^2y^2$

B. $(x^2 + y^2)(x + y) = R^2xy$

C. $(x^2 + y^2)^3 = 4R^2x^2y^2$

D. $(x^2 + y^2)^2 = 4R^2x^2y^2$

Answer: C



[Watch Video Solution](#)

Chapter 5 Parabola

1. Equation of a common tangent to the circle $x^2 + y^2 - 6x = 0$ and the parabola $y^2 = 4x$ is

A. $2\sqrt{3}y = 12x + 1$

B. $2\sqrt{3}y = -x - 12$

C. $\sqrt{3}y = x + 3$

D. $\sqrt{3}y = 3x + 1$

Answer: C



Watch Video Solution

2. Axis of a parabola lies along x-axis. If its vertex and focus are at distances 2 and 4, respectively, from the origin on the positive x-axis, then which of the following points does not lie on it ?

A. (4, -4)

B. $(5, 2\sqrt{6})$

C. (8, 6)

D. $(6, 4\sqrt{2})$

Answer: C

[Watch Video Solution](#)

3. Let $A(4, -4)$ and $B(9,6)$ be points on the parabola $y^2 = 4x$. Let C be chosen on the on the arc AOB of the parabola where O is the origin such that the area of ΔACB is maximum. Then the area (in sq. units) of ΔACB is :

A. $31\frac{3}{4}$

B. 32

C. $30\frac{1}{2}$

D. $31\frac{1}{4}$

Answer: D

[Watch Video Solution](#)

4. If $y^2 = 4b(x - c)$ and $y^2 = 8ax$ having common normal then (a, b, c) is (a) $\left(\frac{1}{2}, 2, 0\right)$ (b) $(1, 1, 3)$ (c) $(1, 1, 1)$ (d) $(1, 3, 2)$

A. $(1, 1, 0)$

B. $\left(\frac{1}{2}, 2, 3\right)$

C. $\left(\frac{1}{2}, 2, 0\right)$

D. $(1, 1, 3)$

Answer: D



Watch Video Solution

5. The length of the common chord of the two circles $x^2 + y^2 - 4y = 0$ and $x^2 + y^2 - 8x - 4y + 11 = 0$ is

A. $2\sqrt{11}$

B. $3\sqrt{2}$

C. $6\sqrt{3}$

D. $8\sqrt{2}$

Answer: C

 [Watch Video Solution](#)

6. If the area of the triangle whose one vertex is at the vertex of the parabola, $y^2 + 4(x - a^2) = 0$ and the other two vertices are the points of intersection of the parabola and Y-axis, is 250 sq units, then a value of 'a' is

A. $5\sqrt{5}$

B. $(10)^{2/3}$

C. $5(2^{1/3})$

D. 5

Answer: D

 [Watch Video Solution](#)

7. Let $A(4, -4)$ and $B(9,6)$ be points on the parabola $y^2 = 4x$. Let C be chosen on the on the arc AOB of the parabola where O is the origin such

that the area of $\triangle ACB$ is maximum. Then the area (in sq. units) of $\triangle ACB$ is :

A. $\frac{125}{4}$

B. $\frac{125}{2}$

C. $\frac{625}{4}$

D. $\frac{75}{2}$

Answer: A



Watch Video Solution

8. A tangent is drawn to parabola $y^2 = 8x$ which makes angle θ with positive direction of x-axis. The equation of tangent is (A) $y = x \tan \theta + 2 \cot \theta$ (B) $y \cot \theta = x - 2 \tan \theta$ (C) $y \cot \theta = x + 2 \tan \theta$ (D) $y \cot \theta = x - \tan \theta$

A. $x = y \cot \theta + 2 \tan \theta$

B. $x = y \cot \theta - 2 \tan \theta$

$$C. y = x \tan \theta - 2 \cot \theta$$

$$D. y = x \tan \theta + 2 \cot \theta$$

Answer: A



Watch Video Solution

Chapter 6 Ellipse

1. If normals are drawn to the ellipse $x^2 + 2y^2 = 2$ from the point $(2, 3)$.

then the co-normal points lie on the curve

$$A. \frac{x^2}{2} + \frac{y^2}{4} = 1$$

$$B. \frac{x^2}{4} + \frac{y^2}{2} = 1$$

$$C. \frac{1}{2x^2} + \frac{1}{4y^2} = 1$$

$$D. \frac{1}{4x^2} + \frac{1}{2y^2} = 1$$

Answer: C



Watch Video Solution

2. Let the length of latus rectum of an ellipse with its major axis along x-axis and center at the origin, be 8. If the distance between the foci of this ellipse is equal to the length of the minor axis, then which of the following points lies on it: (a) $(4\sqrt{2}, 2\sqrt{2})$ (b) $(4\sqrt{3}, 2\sqrt{2})$ (c) $(4\sqrt{3}, 2\sqrt{3})$ (d) $(4\sqrt{2}, 2\sqrt{3})$

A. $(4\sqrt{3}, 2\sqrt{3})$

B. $(4\sqrt{3}, 2\sqrt{2})$

C. $(4\sqrt{2}, 2\sqrt{2})$

D. $(4\sqrt{2}, 2\sqrt{3})$

Answer: B



Watch Video Solution

3. Let S and S' be the foci of the ellipse and B be any one of the extremities of its minor axis. If $\Delta S'BS = 8sq.$ units, then the length of a latus rectum of the ellipse is

A. $2\sqrt{2}$

B. 2

C. 4

D. $4\sqrt{2}$

Answer: C



Watch Video Solution

Chapter 7 Hyperbola

1. If eccentricity of the hyperbola $\frac{x^2}{\cos^2 \theta} - \frac{y^2}{\sin^2 \theta} = 1$ is more than 2 when $\theta \in \left(0, \frac{\pi}{2}\right)$. Find the possible values of length of latus rectum (a) $(3, \infty)$ (b) $1, 3/2$ (c) $(2, 3)$ (d) $(-3, -2)$

A. (2,3)

B. (3, ∞)

C. (3/2, 2)

D. (1, 3/2)

Answer: B



Watch Video Solution

2. A hyperbola has its centre at the origin, passes through the point (4, 2) and has transverse axis of length 4 along the x-axis. Then the eccentricity of the hyperbola is

A. $\frac{2}{\sqrt{3}}$

B. $\frac{3}{2}$

C. $\sqrt{3}$

D. 2

Answer: A



Watch Video Solution

3. The equation of tangent to hyperbola $4x^2 - 5y^2 = 20$ which is parallel to $x - y = 2$ is (a) $x - y + 3 = 0$ (b) $x - y + 1 = 0$ (c) $x - y = 0$ (d) $x - y - 3 = 0$

A. $x - y + 9 = 0$

B. $x - y + 7 = 0$

C. $x - y + 1 = 0$

D. $x - y - 3 = 0$

Answer: C



Watch Video Solution

4. Let $S = \left\{ (x, y) \in \mathbb{R}^2 : \frac{y^2}{1+r} - \frac{x^2}{1-r} = 1 \right\}$, where $r \neq \pm 1$. Then S represents:

A. A hyperbola whose eccentricity is $\frac{2}{\sqrt{r+1}}$, where $0 < r < 1$.

B. An ellipse whose eccentricity is $\frac{1}{\sqrt{r+1}}$, where $r > 1$.

C. A hyperbola whose eccentricity is $\frac{2}{\sqrt{1-r}}$, where $0 < r < 1$.

D. An ellipse whose eccentricity is $\sqrt{\frac{2}{r+1}}$, where $r > 1$.

Answer: D



[Watch Video Solution](#)

5. Equation of a common tangent to the parabola $y^2 = 4x$ and the hyperbola $xy=2$ is

A. $x+2y+4=0$

B. $x-2y+4=0$

C. $x+y+1=0$

D. $4x+2y+1=0$

Answer: A



[Watch Video Solution](#)

6. If a hyperbola has length of its conjugate axis equal to 5 and the distance between its foci is 13, then the eccentricity of the hyperbola is

A. 2

B. $\frac{13}{6}$

C. $\frac{13}{8}$

D. $\frac{13}{12}$

Answer: D



[Watch Video Solution](#)

7. If the vertices of the parabola be at $(-2, 0)$ and $(2, 0)$ and one of the foci be at $(-3, 0)$ then which one of the following points does not lie on the hyperbola? (a) $(-6, 2\sqrt{10})$ (b) $(2\sqrt{6}, 5)$ (c) $(4, \sqrt{15})$ (d) $(6, 5\sqrt{2})$

A. $(4, \sqrt{15})$

B. $(-6, 2\sqrt{10})$

C. $(6, 5\sqrt{2})$

D. $(2, \sqrt{6}, 5)$

Answer: B



Watch Video Solution

Matching Column Type

Column I	Column II
(a) In a triangle ΔXYZ , let a, b and c be the lengths of the sides opposite to the angles X, Y and Z respectively. If $2(a^2 - b^2) = c^2$ and $\lambda = \frac{\sin(X - Y)}{\sin Z}$ then possible values of n for which $\cos(n\pi\lambda) = 0$ is (are)	(p) 1
(b) In a triangle ΔXYZ , let a, b and c be the lengths of the sides opposite to the angles X, Y and Z , respectively. If $1 + \cos 2X - 2 \cos 2Y = 2 \sin X \sin Y$, then possible value(s) of $\frac{a}{b}$ is (are)	(q) 2
(c) In R^2 , let $\sqrt{3}\hat{i} + \hat{j}, \hat{i} + \sqrt{3}\hat{j}$ and $\beta\hat{i} + (1 - \beta)\hat{j}$ be the position vectors of X, Y and Z with respect of the origin O , respectively. If the distance of Z from the bisector of the acute angle of \overline{OX} and \overline{OY} is $\frac{3}{\sqrt{2}}$, then possible value(s) of $ \beta $ is (are)	(r) 3
(d) Suppose that $F(\alpha)$ denotes the area of the region bounded by $x = 0, x = 2, y^2 = 4x$ and $y = \alpha x - 1 + \alpha x - 2 + \alpha x$, where $\alpha \in \{0, 1\}$. Then the value(s) of $F(\alpha) + \frac{8}{3}\sqrt{2}$, when $\alpha = 0$ and $\alpha = 1$, is (are)	(s) 5
	(t) 6

1.



[View Text Solution](#)

Integer Answer Type

1. Suppose that \vec{p}, \vec{q} and \vec{r} are three non-coplanar in R^3 , Let the components of a vector \vec{s} along \vec{p}, \vec{q} and \vec{r} be 4, 3, and 5, respectively, if the components this vector \vec{s} along $(-\vec{p} + \vec{q} + \vec{r}), (\vec{p} - \vec{q} + \vec{r})$ and $(-\vec{p} - \vec{q} + \vec{r})$ are x, y and z , respectively, then the value of $2x + y + z$ is

 Watch Video Solution

2. Let \vec{a} , \vec{b} and \vec{c} be three non coplanar unit vectors such that the angle between every pair of them is $\frac{\pi}{3}$. If $\vec{a} \times \vec{b} + \vec{b} \times \vec{c} = p\vec{a} + q\vec{b} + r\vec{c}$ where p , q , r are scalars then the value of $\frac{p^2 + 2q^2 + r^2}{q^2}$ is

 Watch Video Solution

Chapter 2 Multiple Correct Answers Type

1. Let \vec{x} , \vec{y} and \vec{z} be three vector each of magnitude $\sqrt{2}$ and the angle between each pair of them is $\frac{\pi}{3}$. if \vec{a} is a non - zero vector perpendicular to \vec{x} and $\vec{y} \times \vec{z}$ and \vec{b} is a non-zero vector perpendicular to \vec{y} and $\vec{z} \times \vec{x}$, then

A. $\vec{b} = \left(\vec{b} \cdot \vec{z} \right) \left(\vec{z} - \vec{x} \right)$

B. $\vec{a} = \left(\vec{a} \cdot \vec{y} \right) \left(\vec{y} - \vec{z} \right)$

$$C. \vec{a} \cdot \vec{b} = -(\vec{a} \cdot \vec{y})(\vec{b} \cdot \vec{z})$$

$$D. \vec{a} = (\vec{a} \cdot \vec{y})(\vec{z} - \vec{y})$$

Answer: A::B::C



Watch Video Solution

2. Let ΔPQR be a triangle Let

$\vec{a} = \overrightarrow{QR}$, $\vec{b} = \overrightarrow{RP}$ and $\vec{c} = \overrightarrow{PQ}$ if $|\vec{a}| = 12$, $|\vec{b}| = 4\sqrt{3}$ and $\vec{b} \cdot \vec{c}$

then which of the following is (are) true ?

$$A. \frac{|\vec{c}|^2}{2} - |\vec{a}| = 12$$

$$B. \frac{|\vec{c}|^2}{2} - |\vec{a}| = 30$$

$$C. \left| \vec{a} \times \vec{b} + \vec{c} \times \vec{a} \right| = 48\sqrt{3}$$

$$D. \vec{a} \cdot \vec{b} = -72$$

Answer: A::C::D



Watch Video Solution

Matching Column Type

1.

Column I	Column II
(p) Let $y(x) = \cos(3 \cos^{-1} x)$, $x \in [-1, 1]$, $x \neq \pm \frac{\sqrt{3}}{2}$.	(1) 1
Then $\frac{1}{y(x)} \left\{ (x^2 - 1) \frac{d^2 y(x)}{dx^2} + x \frac{dy(x)}{dx} \right\}$ equals	

(q) Let A_1, A_2, \dots, A_n ($n \geq 2$) be the vertices of a regular polygon of n sides with its centre at the origin. Let a_k be the position vector of the point A_k , $k = 1, 2, \dots, n$. If $\sum_{k=1}^{n-1} (a_k \times a_{k+1}) = \left \sum_{k=1}^{n-1} (a_k \cdot a_{k+1}) \right $, then the minimum value of n is	(2) 2
(r) If the normal from the point $P(h, 1)$ on the ellipse $\frac{x^2}{6} + \frac{y^2}{3} = 1$ is perpendicular to the line $x + y = 8$, then the value of h is	(3) 8
(s) Number of positive solutions satisfying the equation $\tan^{-1} \left(\frac{1}{2x+1} \right) + \tan^{-1} \left(\frac{1}{4x+1} \right) = \tan^{-1} \left(\frac{2}{x^2} \right)$ is	(4) 9

- A. (p) (q) (r) (s)
 (4) (3) (2) (1)
- B. (p) (q) (r) (s)
 (2) (4) (3) (1)
- C. (p) (q) (r) (s)
 (4) (3) (1) (2)
- D. (p) (q) (r) (s)
 (2) (4) (1) (3)

Answer: A



View Text Solution

Column I	Column II
(a) In R^2 , if the magnitude of the projection vector of the vector $\alpha \hat{i} + \beta \hat{j}$ on $\sqrt{3}\hat{i} + \hat{j}$ is $\sqrt{3}$ and if $\alpha = 2 + \sqrt{3}\beta$, then possible value(s) of $ \alpha $ is (are)	(p) 1
(b) Let a and b be real numbers such that the function $f(x) = \begin{cases} -3ax^2 - 2, & x < 1 \\ bx + a^2, & x \geq 1 \end{cases}$ is differentiable for all $x \in R$. Then possible value(s) of α is (are)	(q) 2
(c) Let $\omega \neq 1$ be a complex cube root of unity. If $(3 - 3\omega + 2\omega^2)^{4n+3} + (2 + 3\omega - 3\omega^2)^{4n+3} + (-3 + 2\omega + 3\omega^2)^{4n+3} = 0$, then possible value(s) of n is (are)	(r) 3
(d) Let the harmonic mean of two positive real numbers a and b be 4. If q is a positive real number such that $a, 5, q, b$ is an arithmetic progression, then the value(s) of $ q - a $ is (are)	(s) 4
	(t) 5

2.



View Text Solution

Chapter 3 Multiple Correct Answers Type

1. Let L be a straight line passing through the origin. Suppose that all the points on L are at a constant distance from the two planes $P_1: x + 2y - z + 1 = 0$ and $P_2: 2x - y + z - 1 = 0$. Let M be the locus

of the feet of the perpendiculars drawn from the points on L to the plane

P_1 . Which of the following points lie(s) on M?

A. $\left(0, -\frac{5}{6}, -\frac{2}{3}\right)$

B. $\left(-\frac{1}{6}, -\frac{1}{3}, \frac{1}{6}\right)$

C. $\left(-\frac{5}{6}, 0, \frac{1}{6}\right)$

D. $\left(-\frac{1}{3}, 0, \frac{2}{3}\right)$

Answer: A::B



Watch Video Solution

2. In R^3 , consider the planes $P_1: y = 0$ and $P_2, x + z = 1$. Let P_3 be a plane, different from P_1 and P_2 which passes through the intersection of P_1 and P_2 , If the distance of the point $(0,1,0)$ from P_3 is 1 and the distance of a point (α, β, γ) from P_3 is 2, then which of the following relation(s) is/are true?

A. $2\alpha + \beta + 2\gamma + 2 = 0$

$$B. 2\alpha + \beta + 2\gamma + 4 = 0$$

$$C. 2\alpha + \beta + 2\gamma - 10 = 0$$

$$D. 2\alpha + \beta + 2\gamma - 8 = 0$$

Answer: B::D



Watch Video Solution

Chapter 2

1. Let $\vec{u} = u_1\hat{i} + u_2\hat{j} + u_3\hat{k}$ be a unit vector in R^3 and $\vec{w} = \frac{1}{\sqrt{6}}(\hat{i} + \hat{j} + 2\hat{k})$, Given that there exists a vector \vec{v} in R^3 such that $|\vec{u} \times \vec{v}| = 1$ and $\vec{w} \cdot (\vec{u} \times \vec{v}) = 1$ which of the following statements is correct ?

A. there is exactly one choice for such \vec{v}

B. there are infinitely many choices for such \vec{v}

C. if \hat{u} lies in the xy -plane then $|u_1| = |u_2|$

D. if \hat{u} lies in the xz -plane then $2|u_1| = |u_3|$

Answer: B::C



Watch Video Solution

Single Correct Answer Type

1. the mirror image of point $(3, 1, 7)$ with respect to the plane $x - y + z = 3$ is P . then equation plane which is passes through the point P and contains the line $\frac{x}{1} = \frac{y}{2} = \frac{z}{1}$.

A. $x + y - 3z = 0$

B. $3x + z = 0$

C. $x - 4y + 7z = 0$

D. $2x - y = 0$

Answer: C



Watch Video Solution

Multiple Correct Answers Type

1. Consider a pyramid $OPQRS$ located in the first octant ($x \geq 0, y \geq 0, z \geq 0$) with O as origin, and OP and OR along the x -axis and the y -axis, respectively. The base $OPQR$ of the pyramid is a square with $OP = 3$. The point S is directly above the mid-point T of diagonal OQ such that $TS = 3$. Then

A. the acute angle between OQ and OS is $\pi/3$

B. the equation of the plane containing the triangle OQS is $x-y=0$

C. the length of the perpendicular from P to the plane containing the triangle OQS is $\frac{3}{\sqrt{2}}$

D. the perpendicular distance from O to the straight line containing RS is $\sqrt{\frac{15}{2}}$

Answer: b.,c.,d



Watch Video Solution

Chapter 2

1. Let O be the origin and let PQR be an arbitrary triangle. The point S is such that

$$\overline{OP} \cdot \overline{OQ} + \overline{OR} \cdot \overline{OS} = \overline{OR} \cdot \overline{OP} + \overline{OQ} \cdot \overline{OS} = \overline{OQ} \cdot \overline{OR} + \overline{OP} \cdot \overline{OS}$$

Then the triangle PQR has S as its

- A. centroid
- B. circumcentre
- C. incentre
- D. orthocenter

Answer: D



Watch Video Solution

1. Let O be the origin and $\vec{OX}, \vec{OY}, \vec{OZ}$ be three unit vector in the directions of the sides $\vec{QR}, \vec{RP}, \vec{PQ}$ respectively, of a triangle PQR .

$$|\vec{OX} \times \vec{OY}| =$$

A. $\sin (P + Q)$

B. $\sin 2R$

C. $\sin (P+R)$

D. $\sin (Q+R)$

Answer: A

 [Watch Video Solution](#)

2. Let O be the origin, and OX, OY, OZ be three unit vectors in the direction of the sides QR, RP, PQ , respectively of a triangle PQR . If the

triangle PQR varies, then the minimum value of

$\cos(P + Q) + \cos(Q + R) + \cos(R + P)$ is: $-\frac{3}{2}$ (b) $\frac{5}{3}$ (c) $\frac{3}{2}$ (d) $-\frac{5}{3}$

A. $-\frac{5}{3}$

B. $-\frac{3}{2}$

C. $\frac{3}{2}$

D. $\frac{5}{3}$

Answer: B



Watch Video Solution

Chapter 3

1. The equation of the plane passing through the point (1,1,1) and perpendicular to the planes $2x + y - 2z = 5$ and $3x - 6y - 2z = 7$, is

$$14x + 2y + 15z = 3 \quad 14x + 2y - 15z = 1 \quad 14x + 2y + 15z = 31$$

$$14x - 2y + 15z = 27$$

A. $14x + 2y + 15x = 31$

B. $14x + 2y - 15z = 1$

C. $14x + 2y + 15x = 3$

D. $14x - 2y + 15z = 27$

Answer: A



[Watch Video Solution](#)

Mcq

1. In a class 140 students numbered 1 to 140, all even numbered students opted Mathematics course, those whose number is divisible by 3 opted Physics course and those whose number is divisible by 5 opted Chemistry course. Then the number of students who did not opt for any of the three courses is (a) 38 (b) 1 (c) 42 (d) 102

A. 102

B. 42

C. 1

D. 38

Answer: D



[Watch Video Solution](#)

2. Let α and β be two roots of the equation $x^2 + 2x + 2 = 0$. Then $\alpha^{15} + \beta^{15}$ is equal to

A. 512

B. -512

C. -256

D. 256

Answer: C



[Watch Video Solution](#)

3. If both the roots of the quadratic equation $x^2 - mx + 4 = 0$ are real and distinct and they lie in the interval $[1, 5]$, then m lies in the interval

A. $(4, 5]$

B. $(3, 4)$

C. $(5, 6)$

D. $(-5, -4)$

Answer: A



[Watch Video Solution](#)

4. The number of all possible positive integral values of α for which the roots of the quadratic equation $6x^2 - 11x + \alpha = 0$ are rational numbers is : (a) 3 (b) 2 (c) 4 (d) 5

A. 2

B. 5

C. 3

D. 4

Answer: C



Watch Video Solution

5. Consider the quadratic equation $(c - 5)x^2 - 2cx + (c - 4) = 0$, $c \neq 5$. Let S be the set of all integral values of c for which one root of the equation lies in the interval (0, 2) and its other root lies in the interval (2, 3). Then the number of elements in S is

A. 11

B. 18

C. 10

D. 12

Answer: A



[Watch Video Solution](#)

6. The values of λ such that sum of the squares of the roots of the quadratic equation $x^2 + (3 - \lambda)x + 2 = \lambda$ has the least value is

A. 2

B. $\frac{4}{9}$

C. $\frac{15}{8}$

D. 1

Answer: A



[Watch Video Solution](#)

7. If one root is cube of the other of equation $81x^2 + kx + 256 = 0$ then value of k is equal to (A) 100 (B) -300 (C) -81 (D) 400

A. -81

B. 100

C. -300

D. 144

Answer: C

 [Watch Video Solution](#)

8. Let α and β be the roots of the quadratic equation $x^2 \sin \theta - x(\sin \theta \cos \theta + 1) + \cos \theta = 0$ ($0 < \theta < 45^\circ$), and $\alpha < \beta$.

Then $\sum_{n=0}^{\infty} \left(\alpha^n + \frac{(-1)^n}{\beta^n} \right)$ is equal to

A. $\frac{1}{1 - \cos \theta} + \frac{1}{1 + \sin \theta}$

B. $\frac{1}{1 + \cos \theta} + \frac{1}{1 - \sin \theta}$

C. $\frac{1}{1 - \cos \theta} - \frac{1}{1 + \sin \theta}$

D. $\frac{1}{1 + \cos \theta} - \frac{1}{1 - \sin \theta}$

Answer: A



Watch Video Solution

9. If ratio of the roots of the quadratic equation $3m^2x^2 + m(m - 4)x + 2 = 0$ is λ such that $\lambda + \frac{1}{\lambda} = 1$ then least value of m is (A) $-2 - 2\sqrt{3}$ (B) $-2 + 2\sqrt{3}$ (C) $4 + 3\sqrt{2}$ (D) $4 - 3\sqrt{2}$

A. $2 - \sqrt{3}$

B. $4 - 3\sqrt{2}$

C. $-2 + \sqrt{2}$

D. $4 - 2\sqrt{3}$

Answer: B



Watch Video Solution

10. The number of integral values of m for which the quadratic expression $(1 + 2m)x^2 - 2(1 + 3m)x + 4(1 + m)$, $x \in R$, is always positive is

- A. 8
- B. 7
- C. 6
- D. 3

Answer: B



[Watch Video Solution](#)

11. Let $A = \left\{ \theta \in \left(-\frac{\pi}{2}, \pi \right) : \frac{3 + 2i \sin \theta}{1 - 2 \sin \theta} \text{ is purely imaginary} \right\}$

Then the sum of the elements in A is

- A. $\frac{5\pi}{6}$
- B. $\frac{2\pi}{3}$
- C. $\frac{3\pi}{4}$

D. π

Answer: B



[Watch Video Solution](#)

12. Let Z_0 is the root of equation $x^2 + x + 1 = 0$ and $Z = 3 + 6i(Z_0)^{81} - 3i(Z_0)^{93}$ Then $\arg(Z)$ is equal to (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{3}$ (c) π (d) $\frac{\pi}{6}$

A. $\frac{\pi}{4}$

B. $\frac{\pi}{3}$

C. 0

D. $\frac{\pi}{6}$

Answer: A



[Watch Video Solution](#)

13. Let z_1 and z_2 be any two non-zero complex numbers such that

$$3|z_1| = 2|z_2|. \text{ If } z = \frac{3z_1}{2z_2} + \frac{2z_2}{3z_1}, \text{ then}$$

A. $|z| = \frac{1}{2} \sqrt{\frac{17}{2}}$

B. $\operatorname{Re}(z) = 0$

C. $|z| = \sqrt{\frac{5}{2}}$

D. $\operatorname{Im}(z) = 0$

Answer: D



Watch Video Solution

14. If $z = \left(\frac{\sqrt{3}}{2} + \frac{i}{2}\right)^5 + \left(\frac{\sqrt{3}}{2} - \frac{i}{2}\right)^5$, then prove that $\operatorname{Im}(z) = 0$

A. $\operatorname{R}(z) > 0$ and $\operatorname{I}(z) > 0$

B. $\operatorname{R}(z) < 0$ and $\operatorname{I}(z) > 0$

C. $\operatorname{R}(z) = -3$

D. $\operatorname{I}(z) = 0$

Answer: D



Watch Video Solution

15. Let $\left(-2 - \frac{1}{3}i\right)^3 = \frac{x + iy}{27}$ ($i = \sqrt{-1}$) where x and y are real numbers then $y-x$ equals

A. 85

B. 85

C. -91

D. 91

Answer: D



Watch Video Solution

16. Let $\frac{z - \alpha}{z + \alpha}$ is purely imaginary and $|z| = 2, \alpha \in R$ then α is equal to (A) 2 (B) 1 (C) $\sqrt{2}$ (D) $\sqrt{3}$

A. 1

B. 2

C. $\sqrt{2}$

D. $\frac{1}{2}$

Answer: B



Watch Video Solution

17. Let Z_1 and Z_2 be two complex numbers satisfying $|Z_1| = 9$ and $|Z_2 - 3 - 4i| = 4$. Then the minimum value of $|Z_1 - Z_2|$ is

A. 0

B. 1

C. $\sqrt{2}$

D. 2

Answer: A

 [Watch Video Solution](#)

18. Consider the statement : $P(n): n^2 - n + 41$ is prime." Then, which one of the following is true?

- A. $P(5)$ is false but $P(3)$ is true
- B. Both $P(3)$ and $P(5)$ are false
- C. $P(3)$ is false but $P(5)$ is true
- D. Both $P(3)$ and $P(5)$ are true

Answer: D

 [Watch Video Solution](#)

19. If a, b, c are three distinct real numbers in G.P. and $a + b + c = xb$, then prove that either $x < -1$ or $x > 3$.

A. 4

B. -3

C. -2

D. 2

Answer: D



[Watch Video Solution](#)

20. Let a_1, a_2, \dots, a_{30} be an A. P., $S = \sum_{i=1}^{30} a_i$ and $T = \sum_{i=1}^{15} a_{(2i-1)}$. If $a_5 = 27$ and $S - 2T = 75$, then a_{10} is equal to

A. 57

B. 47

C. 42

D. 52

Answer: D



[Watch Video Solution](#)

21. The sum of series

$$1 + 6 + \frac{9(1^2 + 2^2 + 3^2)}{7} + \frac{12(1^2 + 2^2 + 3^2 + 4^2)}{9} + \frac{15(1^2 + 2^2 + \dots + 5^2)}{11}$$

up to 15 terms is

A. 7820

B. 7830

C. 7520

D. 7510

Answer: A



[Watch Video Solution](#)

22. Let a , b and c be the 7th, 11th and 13th terms, respectively, of a non-constant A.P.. If these are also the three consecutive terms of a G.P., then

$\frac{a}{c}$ is equal to

A. $1/2$

B. 4

C. 2

D. $7/13$

Answer: B



Watch Video Solution

23. The sum of all two-digit positive numbers which when divided by 7 yield 2 or 5 as remainder is

A. 1365

B. 1256

C. 1465

D. 1356

Answer: D

 [Watch Video Solution](#)

24. If 5, $5r$ and $5r^2$ are the lengths of the sides of a triangle, then r cannot be equal to

A. $\frac{3}{2}$

B. $\frac{3}{4}$

C. $\frac{5}{4}$

D. $\frac{7}{4}$

Answer: D

 [Watch Video Solution](#)

25. The sum of an infinite geometric series with positive terms is 3 and the sums of the cubes of its terms is $\frac{27}{19}$. Then the common ratio of this series is

A. $\frac{4}{9}$

B. $\frac{2}{9}$

C. $\frac{2}{3}$

D. $\frac{1}{3}$

Answer: C



Watch Video Solution

26. Let $a_1, a_2, a_3, \dots, a_{10}$ are in G.P. if $\frac{a_3}{a_1} = 25$ then $\frac{a_9}{a_5}$ is equal to (A) 5^4 (B)

4.5^4 (C) 4.5^3 (D) 5^3

A. $2(5^2)$

B. $4(5^2)$

C. 5^4

D. 5^3

Answer: C

 [Watch Video Solution](#)

27. If 19^{th} term of a non-zero A.P. is zero, then $(49^{\text{th}} \text{ term}) : (29^{\text{th}} \text{ term})$ is

A. 3 : 1

B. 4 : 1

C. 2 : 1

D. 1 : 3

Answer: A

 [Watch Video Solution](#)

28. The product of three consecutive terms of a GP is 512. If 4 is added to each of the first and the second of these terms, the three terms now form an AP. Then the sum of the original three terms of the given GP is: (a) 36 (b) 32 (c) 24 (d) 28

A. 36

B. 24

C. 32

D. 28

Answer: D



Watch Video Solution

29. Let $S_k = \frac{1 + 2 + 3 + \dots + k}{k}$. If $S_1^2 + S_2^2 + \dots + S_{10}^2 = \frac{5}{12}A$, then A

is equal to

A. 303

B. 283

C. 156

D. 301

Answer: A

 [Watch Video Solution](#)

30. If the sum of the first 15 terms of the series $\left(\frac{3}{4}\right)^3 + \left(1\frac{1}{2}\right)^3 + \left(2\frac{1}{4}\right)^3 + 3^3 + \left(3\frac{3}{4}\right)^3 + \dots$ is equal to $225k$, then k is equal to

- A. 9
- B. 27
- C. 108
- D. 54

Answer: B

 [Watch Video Solution](#)

31. Let x, y be positive real numbers and m, n be positive integers, The maximum value of the expression

$$\frac{x^m y^n}{(1 + x^{2m})(1 + y^{2n})} \text{ is}$$

A. $\frac{1}{2}$

B. $\frac{1}{4}$

C. $\frac{m+n}{6mn}$

D. 1

Answer: B



Watch Video Solution

32. Consider a class of 5 girls and 7 boys. The number of different teams consisting of 2 girls and 3 boys that can be formed from this class, if there are two specific boys A and B, who refuse to be the members of the same team, is

A. 200

B. 300

C. 500

D. 350

Answer: B



[Watch Video Solution](#)

33. Let S be the set of all triangles in the xy -plane, each having one vertex at the origin and the other two vertices lie on coordinate axes with integral coordinates. If each triangle in S has area 50 sq. units, then the number of elements in the set S is

A. 9

B. 18

C. 32

D. 36

Answer: D



[Watch Video Solution](#)

34. The number of natural numbers less than 7,000 which can be formed by using the digits 0, 1, 3, 7, 9 (repetition of digits allowed) is equal to

- A. 250
- B. 374
- C. 372
- D. 375

Answer: B



[Watch Video Solution](#)

35. If set $A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15\}$, then find the number of onto functions from A to A such that $f(k)$ is a multiple of 3, whenever k is a multiple of 4.

4. (A) $6^5 \times 15!$ (B) $5^6 \times 15!$ (C) $6! \times 5!$ (D) $6! \times 15!$

- A. $(15)! \times 6!$
- B. $5^6 \times 15$

C. $5! \times 6!$

D. $6^5 \times (15)!$

Answer: A

 [Watch Video Solution](#)

36. Let $S = \{1, 2, 3, \dots, 100\}$. The number of non-empty subsets A to S such that the product of elements in A is even is

A. $2^{50}(2^{50} - 1)$

B. $2^{100} - 1$

C. $2^{50} - 1$

D. $2^{50} + 1$

Answer: A

 [Watch Video Solution](#)

37. Consider three boxes, each containing 10 balls labelled 1, 2, ..., 10. Suppose one ball is randomly drawn from each of the boxes denoted by n_i , the label of the ball drawn from the i^{th} box, ($i = 1, 2, 3$). Then, the number of ways in which the balls can be chosen such that $n_1 < n_2 < n_3$ is

- A. 82
- B. 240
- C. 164
- D. 120

Answer: D



[Watch Video Solution](#)

38. Let Z be the set of integers. If $A = \{x \in Z: 2^{(x+2)}(x^2 - 5x + 6) = 1\}$ and $B = \{x \in Z: -3 < 2x - 1 < 9\}$, then the number of subsets of the set $A \times B$ is

A. 2^{18}

B. 2^{10}

C. 2^{15}

D. 2^{12}

Answer: C



Watch Video Solution

39. There are m men and two women participating in a chess tournament. Each participant plays two games with every other participant. If the number of games played by the men between themselves exceeds the number of games played between the men and the women by 84, then the value of m is

A. 9

B. 11

C. 12

D. 7

Answer: C



Watch Video Solution

40. If the fractional part of the number $\frac{2^{403}}{15}$ is $\frac{k}{15}$ then k is equal to

A. 14

B. 6

C. 4

D. 8

Answer: D



Watch Video Solution

41. The coefficient of t^4 in $\left(\frac{1-t^6}{1-t}\right)^3$ (a) 18 (b) 12 (c) 9 (d) 15

A. 12

B. 15

C. 10

D. 14

Answer: B

 [Watch Video Solution](#)

42. If $\sum_{i=1}^{20} \left(\frac{{}^{20}C_{i-1}}{{}^{20}C_i + {}^{20}C_{i-1}} \right)^3 = \frac{k}{21}$, then k equals

A. 200

B. 50

C. 100

D. 400

Answer: C

 [Watch Video Solution](#)

43. If the third term in expansion of $(1 + x^{\log_2 x})^5$ is 2560 then x is equal to (a) $2\sqrt{2}$ (b) $\frac{1}{8}$ (c) $\frac{1}{4}$ (d) $4\sqrt{2}$

A. $2\sqrt{2}$

B. $\frac{1}{8}$

C. $4\sqrt{2}$

D. $\frac{1}{4}$

Answer: D



[Watch Video Solution](#)

44. The positive value of λ for which the coefficient of x^2 in the expression $x^2 \left(\sqrt{2} + \frac{\lambda}{x^2} \right)^{10}$ is 720 is

A. $\sqrt{5}$

B. 4

C. $2\sqrt{2}$

D. 3

Answer: B



Watch Video Solution

45. If $\sum_{r=0}^{25} ({}^{50}C_r {}^{50-r}C_{25-r}) = K({}^{50}C_{25})$, then K is equal to

A. $2^{25} - 1$

B. $(25)^2$

C. 2^{25}

D. 2^{24}

Answer: C



Watch Video Solution

46. If the middle term of the expansion of $\left(\frac{x^3}{3} + \frac{3}{x}\right)^8$ is 5670 then sum of all real values of x is equal to (A) 6 (B) 3 (C) 0 (D) 2

A. 6

B. 8

C. 0

D. 4

Answer: C



[Watch Video Solution](#)

47. The value of r for which

${}^{20}C_r, {}^{20}C_{r-1}, {}^{20}C_1 + {}^{20}C_2 + \dots + {}^{20}C_0, {}^{20}C_r$ is maximum, is

A. 20

B. 15

C. 11

D. 10

Answer: A



[Watch Video Solution](#)

48. Let $(x + 10)^{50} + (x - 10)^{50} = a_0 + a_1x + a_2x^2 + \dots + a_{50}x^{50}$ for all $x \in R$, then $\frac{a_2}{a_0}$ is equal to

A. 12.5

B. 12

C. 12.75

D. 12.25

Answer: D



[Watch Video Solution](#)

49. Let $S_n = 1 + q + q^2 + ? + q^n$ and
 $T_n = 1 + \left(\frac{q+1}{2}\right) + \left(\frac{q+1}{2}\right)^2 + ? + \left(\frac{q+1}{2}\right)^n$ If
 $\alpha T_{100} = {}^{101}C_1 + {}^{101}C_2 x S_1 + {}^{101}C_{101} x S_{100}$, then the value of α is equal
to (A) 2^{99} (B) 2^{101} (C) 2^{100} (D) -2^{100}

A. 2^{100}

B. 200

C. 2^{99}

D. 202

Answer: A



Watch Video Solution

50. Ratio of the 5^{th} term from the beginning to the 5^{th} term from the end
in the binomial expansion of $\left(2^{1/3} + \frac{1}{2(3)^{1/3}}\right)^{10}$ is

A. $1:4(16)^{\frac{1}{3}}$

B. $1:2(6)^{\frac{1}{3}}$

C. $2(36)^{\frac{1}{3}}:1$

D. $4(36)^{\frac{1}{3}}:1$

Answer: D



[Watch Video Solution](#)

51. If ${}^n C_4, {}^n C_5, {}^n C_6$ are in A.P., then find the value of n.

A. 14

B. 11

C. 9

D. 12

Answer: A



[Watch Video Solution](#)

52. Number of irrational terms in expansion of $\left(2^{\frac{1}{5}} + 3^{\frac{1}{10}}\right)^{60}$ is

A. 55

B. 49

C. 48

D. 54

Answer: D



[Watch Video Solution](#)

53. Two integers are selected at random from the set $\{1, 2, \dots, 11\}$. Given that the sum of selected numbers is even, the conditional probability that both the numbers are even is

A. $\frac{2}{5}$

B. $\frac{1}{2}$

C. $\frac{3}{5}$

D. $\frac{7}{10}$

Answer: A



Watch Video Solution

54. Let $S = \{1, 2, \dots, 20\}$ A subset B of S is said to be nice, if the sum of the elements of B is 203. Then the probability that a randomly chosen subset of S is nice is: (a) $\frac{7}{2^{20}}$ (b) $\frac{5}{2^{20}}$ (c) $\frac{4}{2^{20}}$ (d) $\frac{6}{2^{20}}$

A. $\frac{6}{2^{20}}$

B. $\frac{5}{2^{20}}$

C. $\frac{4}{2^{20}}$

D. $\frac{7}{2^{20}}$

Answer: B



Watch Video Solution

55. In a class of 60 students, 40 opted for NCC, 30 opted for NSS and 20 opted for both NCC and NSS. If one of these students is selected at random, then the probability that the student selected has opted neither for NCC nor for NSS is

A. $\frac{2}{3}$

B. $\frac{1}{6}$

C. $\frac{1}{3}$

D. $\frac{5}{6}$

Answer: B



[Watch Video Solution](#)

56. In a game, a man wins Rs 100 if he gets 5 or 6 on a throw of a fair die and loses Rs 50 for getting any other number on the die. If he decides to throw the die either till he gets a five or a six or to a maximum of three throws, then his expected gain/loss (in rupees) is: (a) $\frac{400}{3}$ gain (b) $\frac{400}{9}$ loss (c) 0 (d) $\frac{400}{3}$ loss

A. $\frac{400}{3}$ gain

B. $\frac{400}{3}$ loss

C. 0

D. $\frac{400}{9}$ loss

Answer: C



Watch Video Solution

57. If the Boolean expression $(p \oplus q) \wedge (\sim p \Theta q)$ is equivalent to $p \wedge q$, where $\oplus, \Theta \in \{ \vee, \wedge \}$, then the ordered pair (oplus, Theta) is

A. (\wedge, \vee)

B. (\vee, \vee)

C. (\wedge, \wedge)

D. (\vee, \wedge)

Answer: A



[View Text Solution](#)

58. The logical statement $[\sim(\sim p \vee q) \vee (p \wedge r)] \wedge (\sim q \wedge r)$ is equivalent to (a) $(\sim p \wedge \sim q) \wedge r$ (b) $\sim p \vee r$ (c) $(p \wedge r) \wedge \sim q$ (d) $(p \wedge \sim q) \vee r$

A. $(p \wedge r) \wedge \sim q$

B. $(\sim p \wedge \sim q) \wedge r$

C. $\sim p \vee r$

D. $(p \wedge \sim q) \vee r$

Answer: A



[Watch Video Solution](#)

59. Given three statements P: 5 is a prime number, Q:7 is a factor of 192, R:The LCM of 5 & 7 is 35 Then which of the following statements are true (a) $P \vee (\sim Q \wedge R)$ (b) $\sim P \wedge (\sim Q \wedge R)$ (c) $(P \vee Q) \wedge \sim R$ (d) $\sim P \wedge (\sim Q \wedge R)$

A. $(p \wedge q) \vee (\sim r)$

B. $(\sim p) \wedge (\sim q \wedge r)$

C. $(\sim p) \vee (q \wedge r)$

D. $p \vee (\sim q \wedge r)$

Answer: D



Watch Video Solution

60. If q is false and $(p \wedge q) \rightarrow r$

is also true then which of the following is true?

(A) $p \vee r$ (B) $p \wedge r$ (C) $(p \wedge r) \rightarrow (p \vee r)$ (D) $p \wedge \sim r$

A. $(p \vee r) \rightarrow (p \wedge r)$

B. $p \vee r$

C. $p \wedge r$

D. $(p \wedge r) \rightarrow (p \vee r)$

Answer: D



Watch Video Solution

61. The Boolean expression $\sim(p \vee q) \vee (\sim p \wedge q)$ is equivalent to (1) $\sim p$ (2) p
(3) q (4) $\sim q$

A. $p \wedge (\sim q)$

B. $p \vee (\sim q)$

C. $(\sim p) \wedge (\sim q)$

D. $p \wedge q$

Answer: C



Watch Video Solution

62. $(\sim p \vee \sim q)$ is logically equivalent to

A. $\sim p \wedge \sim q$

B. $p \wedge q$

C. $\sim p \wedge q$

D. $p \wedge \sim q$

Answer: A



Watch Video Solution

63. Average height & variance of 5 students in a class is 150 and 18 respectively. A new student whose height is 156cm is added to the group. Find new variance. (a) 20 (b) 22 (c) 16 (d) 14

A. 22

B. 20

C. 16

D. 18

Answer: B



Watch Video Solution

64. A data consists of n observations x_1, x_2, \dots, x_n . If $\sum_{i=1}^n (x_i + 1)^2 = 9n$ and $\sum_{i=1}^n (x_i - 1)^2 = 5n$, then the standard deviation of this data is

A. 5

B. $\sqrt{5}$

C. $\sqrt{7}$

D. 2

Answer: B



Watch Video Solution

65. The mean of five observations is 5 and their variance is 9.20. If three of the given five observations are 1, 3 and 8, then a ratio of other two observations is

A. 4:9

B. 6:7

C. 5:8

D. 10:3

Answer: A



[Watch Video Solution](#)

66. The mean and standard deviation of five observations x_1, x_2, x_3, x_4, x_5 are 10 and 3 respectively, then variance of the observation $x_1, x_2, x_3, x_4, x_5, -50$ is equal to (a) 437.5 (b) 507.5 (c) 537.5 (d) 487.5

A. 582.5

B. 507.5

C. 586.5

D. 509.5

Answer: B



Watch Video Solution

67. The outcome of each of 30 items was observed , 10 items gave an outcome $\frac{1}{2} - d$ each, 10 items gave outcome $\frac{1}{2}$ each and the remaining 10 items gave outcome $\frac{1}{2} + d$ each. If the variance of this outcome data is $\frac{4}{3}$, then $|d|$ equals

A. 2

B. $\frac{\sqrt{5}}{2}$

C. $\frac{2}{3}$

D. $\sqrt{2}$

Answer: D



[Watch Video Solution](#)

68. Contrapositive of the statement "If two numbers are not equal, then their squares are not equal." is

- A. If the squares of two numbers are equal, then the numbers are equal.
- B. If the squares of two numbers are equal, then the numbers are not equal.
- C. If the squares of two numbers are not equal, then the numbers are equal
- D. If the squares of two numbers are not equal, then the numbers are not equal.

Answer: A



[Watch Video Solution](#)

69. There are 30 white balls and 10 red balls in bag. 16 balls are drawn with replacement from the bag. If X be the number of white balls drawn

then the value of $\frac{\text{mean}(X)}{\text{standard deviation}(X)}$ is equal to (A) $4\sqrt{3}$ (B) $2\sqrt{3}$

(C) $3\sqrt{3}$ (D) $3\sqrt{2}$

A. 4

B. $\frac{4\sqrt{3}}{3}$

C. $4\sqrt{3}$

D. $3\sqrt{2}$

Answer: C

[Watch Video Solution](#)

70. If the sum of the deviations of 50 observations from 30 is 50, then the mean of these observations is

A. 50

B. 51

C. 30

D. 31

Answer: D



[Watch Video Solution](#)

71. Mean and variance of five observations are 4 and 5.2 respectively. If three of these observations are 3, 4, 4 then find absolute difference between the other two observations (A) 3 (B) 7 (C) 2 (D) 5

A. 1

B. 3

C. 7

D. 5

Answer: C



Watch Video Solution

72. The system of linear equations

$$x + y + z = 2$$

$$2x + 3y + 2z = 5$$

$$2x + 3y + (a^2 - 1)z = a + 1$$

A. has infinitely many solutions for $a = 4$

B. is inconsistent when $|a| = \sqrt{3}$

C. is inconsistent when $a = 4$

D. has a unique solution for $|a| = \sqrt{3}$

Answer: B



Watch Video Solution

73. If the system of linear equations $x - 4y + 7z = g$, $3y - 5z = h$, $-2x + 5y - 9z = k$ is consistent, then (a) $g + 2h + k = 0$ (b) $g + h + 2k = 0$ (c) $2g + h + k = 0$ (d) $g + h + k = 0$

A. $g + h + k = 0$

B. $2g + h + k = 0$

C. $g + h + 2k = 0$

D. $g + 2h + k = 0$

Answer: B



[Watch Video Solution](#)

74. If the system of equations

$$x + y + z = 5$$

$$x + 2y + 3z = 9$$

$$x + 3y + \alpha z = \beta$$

has infinitely many solutions, then $\beta - \alpha$ equals

A. 5

B. 18

C. 21

D. 8

Answer: D



Watch Video Solution

75. Let $a_1, a_2, a_3, \dots, a_{10}$ be in G.P. with $a_i > 0$ for $i=1, 2, \dots, 10$ and S be the set of pairs $(r, k), r, k \in \mathbb{N}$ (the set of natural numbers)

for which
$$\begin{vmatrix} \log_e a_1^r a_2^k & \log_e a_2^r a_3^k & \log_e a_3^r a_4^k \\ \log_e a_4^r a_5^k & \log_e a_5^r a_6^k & \log_e a_6^r a_7^k \\ \log_e a_7^r a_8^k & \log_e a_8^r a_9^k & \log_e a_9^r a_{10}^k \end{vmatrix} = 0.$$
 Then the number of

elements in S is

A. Infinitely many

B. 4

C. 10

D. 2

Answer: A



[View Text Solution](#)

76. If the system of linear equations

$$2x+2y+3z=a$$

$$3x-y+5z=b$$

$$x-3y+2z=c$$

where a, b and c are non-zero real numbers, has more than one solution, then

A. $b - c - a = 0$

B. $a + b + c = 0$

C. $b + c - a = 0$

D. $b - c + a = 0.$

Answer: A



Watch Video Solution

77. prove that
$$\begin{bmatrix} a - b - c & 2a & 2a \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{bmatrix} = (a + b + c)^3$$

A. $-(a + b + c)$

B. $2(a + b + c)$

C. abc

D. $-2(a + b + c)$

Answer: D



Watch Video Solution

78. An ordered pair (α, β) for which the system of linear equations $(1 + \alpha)x + \beta y + z = 2$, $\alpha x + (1 + \beta)y + z = 3$ and $\alpha x + \beta y + 2z = 2$ has unique solution is: (a) (2,4) (b) (-3,1) (c) (-4,2) (d) (1,-3)

A. (1, -3)

B. (-3, 1)

C. (2, 4)

D. (-4, 2)

Answer: C



Watch Video Solution

79. The set of all values of λ for which the system of linear equations

$$x - 2y - 2z = \lambda x$$

$$x + 2y + z = \lambda y$$

$$-x - y = \lambda z$$

has a non-trivial solution

A. contains more than two elements

B. is a singleton

C. is an empty set

D. contains exactly two elements

Answer: B

 [Watch Video Solution](#)

80. If $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$, then the matrix A^{-50} , when $\theta = \frac{\pi}{12}$, is equal to

- A. $\begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$
- B. $\begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}$
- C. $\begin{bmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}$
- D. $\begin{bmatrix} \frac{\sqrt{3}}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$

Answer: A

 [Watch Video Solution](#)

81. Matrix =
$$\begin{bmatrix} e^t & e^{-t}(\sin t - 2 \cos t) & e^{-t}(-2 \sin t - \cos t) \\ e^t & -e^{-t}(2 \sin t + \cos t) & e^{-t}(\sin t - 2 \cos t) \\ e^t & e^t \cos t & e^{-t} \sin t \end{bmatrix}$$
 is

invertible. (a) only if $t = \frac{\pi}{2}$ (b) only if $y = \pi$ (c) $t \in R$ (d) $t \notin R$

A. invertible only if $t = \frac{\pi}{2}$

B. not invertible for any $t \in R$

C. invertible for all $t \in R$

D. invertible only if $t = \pi$

Answer: C

 [Watch Video Solution](#)

82. Let $d \in R$ and $A = \begin{pmatrix} -2 & 4 + d & \sin \theta - 2 \\ 1 & \sin \theta + 2 & d \\ 5 & 2 \sin \theta - d & (-\sin \theta) + 2 + 2d \end{pmatrix}$

where $\theta \in [0, \pi]$. If the minimum value of $\det(A)$ is 8, then the value of d

is (a) -7 (b) -5 (c) $2(\sqrt{2} + 1)$ (d) $2(\sqrt{2} + 2)$

A. -7

B. $2(\sqrt{2} + 2)$

C. -5

D. $2(\sqrt{2} + 1)$

Answer: C

 [Watch Video Solution](#)

83. Let $A = \begin{bmatrix} 2 & b & 1 \\ b & b^2 + 1 & b \\ 1 & b & 2 \end{bmatrix}$ where $b > 0$. Then the minimum value of $\frac{\det.(A)}{b}$ is

A. $\sqrt{3}$

B. $-\sqrt{3}$

C. $-2\sqrt{3}$

D. $2\sqrt{3}$

Answer: D

 [Watch Video Solution](#)

[Watch Video Solution](#)

84. Let $A = \begin{bmatrix} 0 & 2q & r \\ p & q & -r \\ p & -q & r \end{bmatrix}$ If $AA^T = I_3$ then $|p| =$

A. $\frac{1}{\sqrt{2}}$

B. $\frac{1}{\sqrt{5}}$

C. $\frac{1}{\sqrt{6}}$

D. $\frac{1}{\sqrt{3}}$

Answer: A

[Watch Video Solution](#)

85. Let A and B be two invertible matrices of order 3×3 . If $\det. (ABA^T) = 8$ and $\det. (AB^{-1}) = 8$, then $\det. (BA^{-1}B^T)$ is equal to

A. 16

B. $\frac{1}{16}$

C. $\frac{1}{4}$

D. 1

Answer: B

 [Watch Video Solution](#)

86. Let $P = \begin{bmatrix} 1 & 0 & 0 \\ 4 & 1 & 0 \\ 16 & 4 & 1 \end{bmatrix}$ and I be the identity matrix of order 3. If

$Q = [q_{ij}]$ is a matrix, such that $P^{50} - Q = I$, then $\frac{q_{31} + q_{32}}{q_{21}}$ equals

A. 15

B. 9

C. 135

D. 10

Answer: D

 [Watch Video Solution](#)

87. If $A = \begin{bmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{bmatrix}$, then for all $\theta \in \left(\frac{3\pi}{4}, \frac{5\pi}{4}\right)$, $\det. (A)$ lies in the interval

A. $\left[\frac{5}{2}, 4\right)$

B. $\left(\frac{3}{2}, 3\right]$

C. $\left(0, \frac{3}{2}\right]$

D. $\left(1, \frac{5}{2}\right]$

Answer: B



Watch Video Solution

88. Two cards are drawn successively with replacement from a well-shuffled deck of 52 cards. Let X denote the random variable of number of aces obtained in the two drawn cards. Then $P(X = 1) + P(X = 2)$ equals

A. $52/169$

B. $\frac{25}{169}$

C. $\frac{49}{169}$

D. $\frac{24}{169}$

Answer: B



Watch Video Solution

89. An urn contains 5 red and 2 green balls. A ball is drawn at random from the urn. If the drawn ball is green, then a red ball is added to the urn and if the drawn ball is red, then a green ball is added to the urn, the original ball is not returned to the urn. Now, a second ball is drawn at random from it. The probability that the second ball is red is

A. $\frac{26}{49}$

B. $\frac{32}{49}$

C. $\frac{27}{49}$

D. $\frac{21}{49}$

Answer: B



[Watch Video Solution](#)

90. An unbiased coin is tossed. If the result is a head, a pair of unbiased dice is rolled and the number obtained by adding the numbers on two faces is noted. If the result is a tail, a card from a well-shuffled pack of 11 cards numbered 2, 3, 4, ..., 12 is picked and the number on the card is noted. What is the probability that the noted number is either 7 or 8?

A. $\frac{13}{36}$

B. $\frac{19}{36}$

C. $\frac{19}{72}$

D. $\frac{15}{72}$

Answer: C



[Watch Video Solution](#)

91. If the probability of hitting a target by a shooter, in any shot is $\frac{1}{3}$, then the minimum number of independent shots at the target required by him so that the probability of hitting the target at least once is greater than $\frac{5}{6}$ is

A. 6

B. 5

C. 4

D. 3

Answer: B



[Watch Video Solution](#)

92. In a random experiment, a fair die is rolled until two fours are obtained in succession. The probability that the experiment will end in the fifth throw of the die is equal to

A. $\frac{150}{6^5}$

- B. $\frac{175}{6^5}$
- C. $\frac{200}{6^5}$
- D. $\frac{225}{6^5}$

Answer: B



Watch Video Solution

Chapter 1

1. For $x \in \mathbb{R} - \{0, 1\}$, let $f_1(x) = \frac{1}{x}$, $f_2(x) = 1 - x$ and $f_3(x) = \frac{1}{1 - x}$ be three given functions. If a function, $J(x)$ satisfies $(f_2 \circ J \circ f_1)(x) = f_3(x)$ then $J(x)$ is equal to :

- A. $f_3(x)$
- B. $f_1(x)$
- C. $f_2(x)$

D. $\frac{1}{x} f_3(x)$

Answer: A



Watch Video Solution

2. Let $A = \{x \in \mathbb{R} : x \text{ is not a positive integer}\}$ define a function $f: A \rightarrow \mathbb{R}$ such that $f(x) = \frac{2x}{x-1}$. Then f is

- A. injective but not surjective
- B. not injective
- C. surjective but not injective
- D. neither injective nor surjective

Answer: A



Watch Video Solution

3. Let N be the set of numbers and two functions f and g be defined as

$f, g: N \rightarrow N$ such that

$$f(n) = \begin{cases} \frac{n+1}{2} & \text{if } n \text{ is odd} \\ \frac{n}{2} & \text{if } n \text{ is even} \end{cases}$$

and $g(n) = n - (-1)^n$. Then, $f \circ g$ is

- A. both one-one and onto
- B. one-one but not onto
- C. neither one-one nor onto
- D. onto but not one-one

Answer: D



[Watch Video Solution](#)

4. Let $f: R \rightarrow R$ be defined by $f(x) = \frac{x}{1+x^2}$, $x \in R$. Then the range of f is

- A. $(-1,1) - \{0\}$

B. $\left[-\frac{1}{2}, \frac{1}{2}\right]$

C. $R - \left[-\frac{1}{2}, \frac{1}{2}\right]$

D. $R - [-1, 1]$

Answer: B



Watch Video Solution

5. Let a function $f: (0, \infty) \rightarrow [0, \infty)$ be defined by $f(x) = \left|1 - \frac{1}{x}\right|$.

Then f is

A. injective only

B. not injective but it is surjective

C. both injective nor surjective

D. injective only

Answer: B



Watch Video Solution

Chapter 2

1. $\lim_{y \rightarrow 0} \frac{\sqrt{1 + \sqrt{1 + y^4}} - \sqrt{2}}{y^4}$

A. exists and equals $\frac{1}{4\sqrt{2}}$

B. does not exist

C. exists and equals $\frac{1}{2\sqrt{2}}$

D. exists and equals $\frac{1}{2\sqrt{2}(\sqrt{2} + 1)}$

Answer: A



Watch Video Solution

2. For each $x \in \mathbb{R}$, let $[x]$ be the greatest integer less than or equal to x .

Then $\lim_{x \rightarrow 1^+} \frac{x([x] + |x|)\sin[x]}{|x|}$ is equal to

A. $-\sin 1$

B. 0

C. 1

D. $\sin 1$

Answer: A



Watch Video Solution

3. For each $t \in \mathbb{R}$, let $[t]$ be the greatest integer less than or equal to t .

Then

$$\lim_{x \rightarrow 1^+} \frac{(1 - |x| + \sin|1 - x|)\sin\left(\frac{\pi}{2}[1 - x]\right)}{|1 - x|[1 - x]}$$

A. equals -1

B. equals 1

C. does not exist

D. equals 0

Answer: D



Watch Video Solution

4. let $[x]$ denote the greatest integer less than or equal to x .

Then $\lim_{x \rightarrow 0} \frac{\tan(\pi \sin^2 x) + (|x - \sin(x[x])|)^2}{x^2}$

- A. equals π
- B. equals 0
- C. equals $\pi + 1$
- D. does not exist

Answer: D



Watch Video Solution

5. $\lim_{x \rightarrow 0} \frac{x \cot(4x)}{\sin^2 x \cot^2(2x)}$ is equal to

A. 2

B. 0

C. 4

D. 1

Answer: D



Watch Video Solution

6. $\lim_{x \rightarrow \frac{\pi}{4}} \frac{\cot^3 x - \tan x}{\cos\left(x + \frac{\pi}{4}\right)}$ is

A. 4

B. $8\sqrt{2}$

C. 8

D. $4\sqrt{2}$

Answer: C



Watch Video Solution

7. $\lim_{x \rightarrow 1^-} \frac{\sqrt{\pi} - \sqrt{2 \sin^{-1} x}}{\sqrt{1-x}}$ is equal to

A. $\frac{1}{\sqrt{2\pi}}$

B. $\frac{\sqrt{\pi}}{2}$

C. $\sqrt{\frac{2}{\pi}}$

D. $\sqrt{\pi}$

Answer: C



Watch Video Solution

Chapter 3

1. If $x = 3 \tan t$ and $y = 3 \sec t$, then the value of $\frac{d^2y}{dx^2}$ at $t = \frac{\pi}{4}$ is

A. $\frac{3}{2\sqrt{2}}$

B. $\frac{1}{3\sqrt{2}}$

C. $\frac{1}{6}$

D. $\frac{1}{6\sqrt{2}}$

Answer: D



Watch Video Solution

2. Let $f: R \rightarrow R$ be a function such that $f(x) = x^3 + x^2 f'(1) + x f''(2) + f'''(3)$, $x \in R$. Then $f(2)$ equals

A. 8

B. -2

C. -4

D. 30

Answer: B



Watch Video Solution

3. If $x \log_e(\log_e x) - x^2 + y^2 = 4 (y > 0)$, then dy/dx at $x = e$ is equal to

A. $\frac{e}{4 + e^2}$

B. $\frac{(1 + 2e)}{2\sqrt{4 + e^2}}$

C. $\frac{(2e - 1)}{2\sqrt{4 + e^2}}$

D. $\frac{(1 + 2e)}{\sqrt{4 + e^2}}$

Answer: C



Watch Video Solution

4. for $x > 1$ if $(2x)^{2y} = 4e^{2x-2y}$ then $(1 + \log_e 2x)^2 \frac{dy}{dx}$

A. $\log_e 2x$

B. $\frac{x \log_e 2x + \log_e 2}{x}$

C. $x \log_e 2x$

D. $\frac{x \log_e 2x - \log_e 2}{x}$

Answer: D



Watch Video Solution

5. Let f be a differentiable function such that $f(1) = 2$ and $f'(x) = f(x)$ for all $x \in \mathbb{R}$. If $h(x) = f(f(x))$, then $h'(1)$ is equal to

A. $4e$

B. $4e^2$

C. $2e$

D. $2e^2$

Answer: A



Watch Video Solution

1. $f(x) = \begin{cases} 5 - x & |x| \leq 1 \\ a + bx & |x| > 1 \end{cases}$

- A. continuous if $a = 5$ and $b = 5$
- B. continuous if $a = -5$ and $b = 10$
- C. continuous if $a = 0$ and $b = 5$
- D. not continuous for any values of a and b

Answer: D



[Watch Video Solution](#)

2. Let $f(x) = \begin{cases} \max \{ |x|, x^2 \}, & |x| \leq 2 \\ 8 - 2|x|, & 2 < |x| \leq 4 \end{cases}$. Let S be the set of points

in the interval $(-4, 4)$ at which f is not differentiable. Then S

- A. is an empty set
- B. equals $\{-2, -1, 1, 2\}$
- C. equals $\{-2, -1, 0, 1, 2\}$

D. equals $\{-2,2\}$

Answer: C

 [Watch Video Solution](#)

3. Let $f: (-1, 1) \rightarrow \mathbb{R}$ be a function defined by $f(x) = \max. \left\{ -|x|, -\sqrt{1-x^2} \right\}$. If K is the set of all points at which f is not differentiable, then K has set of all points at which f is not differentiable, then K has exactly

- A. three elements
- B. one element
- C. five elements
- D. two elements

Answer: A

 [Watch Video Solution](#)

4. Let $f(x) = \begin{cases} -1 & -2 \leq x < 0 \\ x^2 & 0 \leq x < 2 \end{cases}$ if $g(x) = |f(x)| + f(|x|)$ then $g(x)$ in $(-2, 2)$ (A) not continuous (B) not differential at one point (C) differential at all points (D) not differential at two points

- A. Differentiable at all points
- B. not differentiable at two points
- C. Not continuous
- D. not differentiable at one point

Answer: D



[Watch Video Solution](#)

5. Let K be the set of all values of x , where the function $f(x) = \sin|x| - |x| + 2(x - \pi)\cos|x|$ is not differentiable.

Then, the set K is equal to

- A. $\{\pi\}$

B. $\{0\}$

C. ϕ (an empty set)

D. $\{0, \pi\}$

Answer: C



Watch Video Solution

6. Let S be the set of all points in $(-\pi, \pi)$ at which the Then, S is a subset of which of the following?

A. $\left\{ -\frac{3\pi}{4}, -\frac{\pi}{4}, \frac{3\pi}{4}, \frac{\pi}{4} \right\}$

B. $\left\{ -\frac{3\pi}{4}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{4} \right\}$

C. $\left\{ -\frac{\pi}{2}, -\frac{\pi}{4}, \frac{\pi}{4}, \frac{\pi}{2} \right\}$

D. $\left\{ -\frac{\pi}{4}, 0, \frac{\pi}{4} \right\}$

Answer: A



Watch Video Solution

Chapter 5

1. if θ denotes the acute angle between the curves, $y = 10 - x^2$ and $y = 2 + x^2$ at a point of their intersection, then $|\tan \theta|$ is equal to

A. $4/9$

B. $7/17$

C. $8/17$

D. $8/15$

Answer: D

 [Watch Video Solution](#)

2. The tangent to the curve $y = xe^{x^2}$ passing through the point $(1, e)$ also passes through the point

A. $\left(\frac{4}{3}, 2e\right)$

B. $(2, 3e)$

C. $\left(\frac{5}{3}, 2e\right)$

D. $(3, 6e)$

Answer: A



Watch Video Solution

3. A helicopter flying along the path $y = 7 + x^{\frac{3}{2}}$, A soldier standint at point $\left(\frac{1}{2}, 7\right)$ wants to hit the helicopter when it is closest from him, then minimum distance is equal to (a) $\frac{1}{6} \frac{\sqrt{2}}{3}$ (b) $\frac{1}{2}$ (c) $\frac{1}{3} \sqrt{\frac{2}{3}}$ (d) $\sqrt{\frac{5}{2}}$

A. $\frac{1}{2}$

B. $\frac{1}{3} \sqrt{\frac{7}{3}}$

C. $\frac{1}{6} \sqrt{\frac{7}{3}}$

D. $\frac{\sqrt{5}}{6}$

Answer: C



Watch Video Solution

Chapter 6

1. The maximum volume (in cu.m) of the right circular cone having slant height 3 m is

A. $3\sqrt{3}\pi$

B. 6π

C. $2\sqrt{3}\pi$

D. $\frac{4}{3}\pi$

Answer: C



Watch Video Solution

2. The shortest distance between the point $\left(\frac{3}{2}, 0\right)$ and the curve $y = \sqrt{x}$, ($x > 0$), is

A. $\frac{\sqrt{5}}{2}$

B. $\frac{5}{4}$

C. $\frac{3}{2}$

D. $\frac{\sqrt{3}}{2}$

Answer: A



[Watch Video Solution](#)

3. If x satisfies the condition $f(x) = \{x : x^2 + 30 \leq 11x\}$ then maximum value of function $f(x) = 3x^3 - 18x^2 - 27x - 40$ is equal to (A) -122 (B) 122 (C) 222 (D) -222

A. 122

B. -222

C. -122

D. 222

Answer: A



Watch Video Solution

4. Let $f(x) = -\frac{x}{\sqrt{a^2 + x^2}} - \frac{d - x}{\sqrt{b^2 + (d - x)^2}}$, $x \in R$, where a , b and

d are non-zero real constants. Then,

A. f is a decreasing function of x

B. f is neither increasing nor decreasing function of x

C. f' is not a continuous function of x

D. f is an increasing function of x

Answer: D



Watch Video Solution

5. Let a parabola be $y = 12 - x^2$. Find the maximum area of rectangle whose base lie on x-axis and two points lie on parabola. (A) 8 (B) 4 (C) 32 (D) 34

A. $20\sqrt{2}$

B. $18\sqrt{2}$

C. 32

D. 36

Answer: C



Watch Video Solution

6. Let $f(x) = x^3 - 3(a - 2)x^2 + 3ax + 7$ and $f(x)$ is increasing in $(0, 1]$ and decreasing is $[1, 5)$, then roots of the equation $\frac{f(x) - 14}{(x - 1)^2} = 0$ is (A) 1 (B) 3 (C) 7 (D) -2

A. 6

B. 5

C. 7

D. -7

Answer: C



Watch Video Solution

Chapter 7

1. If $x^2 \neq n\pi - 1, n \in N$. Then, the value of

$\int x \sqrt{\frac{2 \sin(x^2 + 1) - \sin 2(x^2 + 1)}{2 \sin(x^2 + 1) + \sin 2(x^2 + 1)}} dx$ is equal to:

A. $\log_e \left| \left(\sec. \frac{x^2 - 1}{2} \right) \right| + c$

B. $\log_e \left| \frac{1}{2} \sec^2. (x^2 - 1) \right| + c$

C. $\frac{1}{2} \log_e \left| \sec^2. \left(\frac{x^2 - 1}{2} \right) \right| + c$

D. $\frac{1}{2} \log_e \left| \sec. (x^2 - 1) \right| + c$

Answer: A



Watch Video Solution

2. If $f(x) = \int \frac{5x^8 + 7x^6}{(x^2 + 1 + 2x^7)^2} dx$, ($x \geq 0$), and $f(0) = 0$, then the value of $f(1)$ is

A. $-\frac{1}{2}$

B. $\frac{1}{2}$

C. $-\frac{1}{4}$

D. $\frac{1}{4}$

Answer: D



Watch Video Solution

3. Let $n \geq 2$ be a natural number and $0 < \theta < \frac{\pi}{2}$, Then,

$\int \frac{(\sin^n \theta - \sin \theta)^{\frac{1}{n}} \cos \theta}{\sin^{n+1} \theta} d\theta$ is equal to (where C is a constant of

integration)

$$\text{A. } \frac{n}{n^2 - 1} \left(1 - \frac{1}{\sin^{n+1} \theta} \right)^{\frac{n+1}{n}} + C$$

$$\text{B. } \frac{n}{n^2 + 1} \left(1 - \frac{1}{\sin^{n-1} \theta} \right)^{\frac{n+1}{n}} + C$$

$$\text{C. } \frac{n}{n^2 - 1} \left(1 - \frac{1}{\sin^{n-1} \theta} \right)^{\frac{n+1}{n}} + C$$

$$\text{D. } \frac{n}{n^2 - 1} \left(1 + \frac{1}{\sin^{n-1} \theta} \right)^{\frac{n+1}{n}} + C$$

Answer: C



Watch Video Solution

4. If $\int x^5 e^{4x^2} dx = \frac{1}{48} e^{4x^2} (f(x)) + c$, where c is constant of integration

then $f(x)$ equals to (a) $-4x^3 - 1$ (b) $-1 - 2x^3$ (c) $4x^3 + 1$ (d) $1 - 2x^3$

A. $-4x^3 - 1$

B. $4x^3 + 1$

C. $-2x^3 - 1$

D. $-2x^3 + 1$

Answer: A



Watch Video Solution

5. If $\int \frac{\sqrt{1-x^2}}{x^4} dx = A(x) \left(\sqrt{1-x^2}\right)^m + C$, for a suitable chosen integer m and a function $A(x)$, where C is a constant of integration, then $(A(x))^m$ equals

A. $\frac{-1}{3x^3}$

B. $\frac{-1}{27x^9}$

C. $\frac{1}{9x^4}$

D. $\frac{1}{27x^6}$

Answer: B



Watch Video Solution

6. If $\int \frac{x+1}{\sqrt{2x-1}} dx = f(x)\sqrt{2x-1} + C$, where C is a constant of integration, then $f(x)$ is equal to

A. $\frac{1}{3}(x+4)$

B. $\frac{1}{3}(x+1)$

C. $\frac{2}{3}(x+2)$

D. $\frac{2}{3}(x-4)$

Answer: A



Watch Video Solution

7. The integral $\int \cos(\log_e x) dx$ is equal to: (where C is a constant of integration)

A. $\frac{x}{2} [\sin(\log_e x) - \cos(\log_e x)] + C$

B. $\frac{x}{2} [\cos(\log_e x) + \sin(\log_e x)] + C$

C. $x [\cos(\log_e x) + \sin(\log_e x)] + C$

$$D. x[\cos(\log_e x - \sin(\log_e x))] + C$$

Answer: B



Watch Video Solution

$$8. \int \frac{3x^{13} + 2x^{11}}{(2x^4 + 3x^2 + 1)^4} dx$$

$$A. \frac{x^4}{(2x^4 + 3x^2 + 1)^3} + C$$

$$B. \frac{x^{12}}{6(2x^4 + 3x^2 + 1)^3} + C$$

$$C. \frac{x^4}{6(2x^4 + 3x^2 + 1)^3} + C$$

$$D. \frac{x^{12}}{(2x^4 + 3x^2 + 1)^3} + C$$

Answer: B



Watch Video Solution

1. The value of $\int_0^{\pi} |\cos x|^3 dx$ is

A. $2/3$

B. 0

C. $-4/3$

D. $4/3$

Answer: D



[Watch Video Solution](#)

2. Let f be a differentiable function from R to R such that

$|f(x) - f(y)| \leq 2||x - y|^{3/2}$, for all $x, y \in R$. If $f(0) = 1$, then

$\int_0^1 f^2(x) dx$ is equal to

A. 0

B. $\frac{1}{2}$

C. 2

D. 1

Answer: D

 [Watch Video Solution](#)

3. $\int_0^{\frac{\pi}{3}} \frac{\tan \theta}{\sqrt{2k \sec \theta}} d\theta = 1 - \frac{1}{\sqrt{2}}$, ($k > 0$), then the value of k is

A. 2

B. $\frac{1}{2}$

C. 4

D. 1

Answer: A

 [Watch Video Solution](#)

4. Let $I = \int_a^b (x^4 - 2x^2) dx$. If it is minimum, then the ordered pair (a, b) is

A. $(-\sqrt{2}, 0)$

B. $(-\sqrt{2}, \sqrt{2})$

C. $(0, \sqrt{2})$

D. $(\sqrt{2}, -\sqrt{2})$

Answer: B

 [Watch Video Solution](#)

5. The value of $\int_{-\pi/2}^{\pi/2} \frac{dx}{[x] + [\sin x] + 4}$ where $[t]$ denotes the greatest integer less or equal to t , is

A. $\frac{1}{12}(7\pi + 5)$

B. $\frac{3}{10}(4\pi - 3)$

C. $\frac{1}{12}(7\pi - 5)$

D. $\frac{3}{20}(4\pi - 3)$

Answer: D



Watch Video Solution

6. If $\int_0^1 f(t) dt = x^2 + \int_0^1 t^2 f(t) dt$, then $f'(1/2)$ is

A. $\frac{6}{25}$

B. $\frac{24}{25}$

C. $\frac{18}{25}$

D. $\frac{4}{5}$

Answer: B



Watch Video Solution

7. The value of the integral $\int_{-2}^2 \frac{\sin^2 x}{-2\left[\frac{x}{\pi}\right] + \frac{1}{2}} dx$ (where $[x]$ denotes the greatest integer less than or equal to x) is

A. 4

B. $4 - \sin 4$

C. $\sin 4$

D. 0

Answer: D



Watch Video Solution

8. The integral $\int_{\pi/6}^{\pi/4} \frac{dx}{\sin 2x (\tan^5 x + \cot^5 x)}$ equals

A. $\frac{1}{10} \left(\frac{\pi}{4} - \tan^{-1} \left(\frac{1}{9\sqrt{3}} \right) \right)$

B. $\frac{1}{5} \left(\frac{\pi}{4} - \tan^{-1} \left(\frac{1}{3\sqrt{3}} \right) \right)$

C. $\frac{\pi}{10}$

D. $\frac{1}{20} - \tan^{-1} \left(\frac{1}{9\sqrt{3}} \right)$

Answer: A



Watch Video Solution

9. Let f and g be continuous functions on $[0, a]$ such that $f(x) = f(a - x)$ and $g(x) + g(a - x) = 4$ then $\int_0^a f(x)g(x)dx$ is equal to

A. $4 \int_0^a f(x)dx$

B. $2 \int_0^a f(x)dx$

C. $-3 \int_0^a f(x)dx$

D. $\int_0^a f(x)dx$

Answer: B



Watch Video Solution

10. The integral $\int_1^e \left\{ \left(\frac{x}{e}\right)^{2x} - \left(\frac{e}{x}\right)^x \right\} \log_e x dx$ is equal to

A. $\frac{1}{2} - e - \frac{1}{e^2}$

B. $\frac{3}{2} - \frac{1}{e} - \frac{1}{2e^2}$

C. $-\frac{1}{2} + \frac{1}{e} - \frac{1}{2e^2}$

D. $\frac{3}{2} - e - \frac{1}{2e^2}$

Answer: D



Watch Video Solution

11. $\lim_{n \rightarrow \infty} \left(\frac{n}{n^2 + 1^2} + \frac{n}{n^2 + 2^2} + \frac{n}{n^2 + 3^2} + \dots + \frac{1}{2n} \right)$ is equal to

A. $\frac{\pi}{4}$

B. $\tan^{-1}(2)$

C. $\tan^{-1}(3)$

D. $\frac{\pi}{2}$

Answer: B



Watch Video Solution

1. The area (in sq. units) bounded by the parabola $y = x^2 - 1$, the tangent at the point (2,3) to it and the y-axis is

A. $\frac{14}{3}$

B. $\frac{56}{3}$

C. $\frac{8}{3}$

D. $\frac{32}{3}$

Answer: C



[Watch Video Solution](#)

2. The area (in sq. units) of the region

$$A = \{(x, y) : 0 \leq y \leq x|x| + 1 \text{ and } -1 \leq x \leq x\} \text{ is}$$

A. $\frac{1}{3}$

B. $\frac{1}{3}$

C. 2

D. $\frac{4}{3}$

Answer: C

 [View Text Solution](#)

3. If the area enclosed between the curves $y = kx^2$ and $x = ky^2$, where $k > 0$, is 1 square unit. Then k is: (a) $\frac{1}{\sqrt{3}}$ (b) $\frac{\sqrt{3}}{2}$ (c) $\frac{2}{\sqrt{3}}$ (d) $\sqrt{3}$

A. $\frac{1}{\sqrt{3}}$

B. $\frac{2}{\sqrt{3}}$

C. $\frac{\sqrt{3}}{2}$

D. $\sqrt{3}$

Answer: A

 [Watch Video Solution](#)

4. The area of the region bounded by the curve $x^2 = 4y$ and the straight line $x = 4y - 2$ is

A. $\frac{5}{4}$

B. $\frac{9}{8}$

C. $\frac{3}{4}$

D. $\frac{7}{8}$

Answer: B



[Watch Video Solution](#)

5. The area (in sq. units) in the first quadrant bounded by the parabola $y = x^2 + 1$, the tangent to it at the point $(2, 5)$ and the coordinate axes is

A. $\frac{14}{3}$

B. $\frac{187}{24}$

C. $\frac{37}{24}$

D. $\frac{8}{3}$

Answer: C



Watch Video Solution

6. The area (in sq. units) of the region bounded by the parabola $y = x^2 + 2$ and the lines $y = x + 1$, $x = 0$ and $x = 3$, is

A. $\frac{15}{4}$

B. $\frac{15}{2}$

C. $\frac{21}{2}$

D. $\frac{17}{4}$

Answer: B



Watch Video Solution

Chapter 10

1. If $y = y(x)$ is the solution of the differential equation, $x \frac{dy}{dx} + 2y = x^2$ satisfying $y(1) = 1$, then $y\left(\frac{1}{2}\right)$ is equal to

A. $\frac{4}{64}$

B. $\frac{13}{16}$

C. $\frac{49}{16}$

D. $\frac{1}{4}$

Answer: C



Watch Video Solution

2. Let $f: [0, 1] \rightarrow R$ be such that $f(xy) = f(x) \cdot f(y)$, for all $x, y \in [0, 1]$ and $f(0) \neq 0$. If $y = y(x)$ satisfies the

differential equation, $\frac{dy}{dx} = f(x)$ with $y(0) = 1$, then

$y\left(\frac{1}{4}\right) + y\left(\frac{3}{4}\right)$ is equal to

A. 4

B. 3

C. 5

D. 2

Answer: B



[Watch Video Solution](#)

3.

If

$$\frac{dy}{dx} + \frac{3}{\cos^2 x} y = \frac{1}{\cos^2 x}, x \in \left(-\frac{\pi}{3}, \frac{\pi}{3}\right) \text{ and } y\left(\frac{\pi}{4}\right) = \frac{4}{3}, \text{ then } y\left(-\frac{\pi}{4}\right)$$

equals

A. $\frac{1}{3} + e^6$

B. $\frac{1}{3}$

C. $-\frac{1}{4}$

D. $\frac{1}{3} + e^3$

Answer: A



Watch Video Solution

4. Let f be differentiable function such that

$$f'(x) = 7 - \frac{3}{4} \frac{f(x)}{x}, (x > 0) \text{ and } f(1) \neq 4 \text{ Then } \lim_{x \rightarrow 0^+} x f\left(\frac{1}{x}\right)$$

A. exists and equals 4

B. does not exist

C. exists and equals 0

D. exists and equals $4/7$

Answer: A



Watch Video Solution

5. The curve amongst the family of curves, represented by the differential equation $(x^2 - y^2)dx + 2xydy = 0$ which passes through (1,1) is

- A. a circle with centre on the y-axis
- B. a circle with centre on the x-axis
- C. an ellipse with major axis along the y-axis
- D. a hyperbola with transverse axis along the

Answer: B



[Watch Video Solution](#)

6. If $y(x)$ is the the solution of the differential equation

$$\frac{dy}{dx} + \left(\frac{2x+1}{x}\right)y = e^{-2x}, x > 0, \text{ where } y(1) = \frac{1}{2}e^{-2}, \text{ then}$$

- A. $y(x)$ is decreasing in $(0,1)$
- B. $y(x)$ is decreasing in $\left(\frac{1}{2}, 1\right)$
- C. $y(\log_e 2) = \frac{\log_e 2}{4}$

$$D. y(\log_e 2) = \log_2 4$$

Answer: B



Watch Video Solution

7. The solution of the differential equation, $\frac{dy}{dx} = (x - y)^2$,

when $y(1) = 1$, is

A. $\log_e \left| \frac{2 - y}{2 - x} \right| = 2(y - 1)$

B. $\log_e \left| \frac{2 - x}{2 - y} \right| = x - y$

C. $-\log_e \left| \frac{1 + x - y}{1 - x + y} \right| = x + y - 2$

D. $-\log_e \left| \frac{1 - x + y}{1 + x - y} \right| = 2(x - 1)$

Answer: D



Watch Video Solution

8. Let $y = y(x)$ be the solution of the differential equation $x \frac{dy}{dx} + y = x \log_e x$, ($x > 1$). If $2y(2) = \log_e 4 - 1$, then $y(e)$ is equal to

A. $\frac{e^2}{4}$

B. $\frac{e}{4}$

C. $-\frac{e}{2}$

D. $-\frac{e^2}{2}$

Answer: B



Watch Video Solution

9. If a curve passes through the point $(1, -2)$ and has slope of the tangent at any point (x, y) on it as $\frac{x^2 - 2y}{x}$, then the curve also passes through the point

A. $(-\sqrt{2}, 1)$

B. $(\sqrt{3}, 0)$

C. $(-1, 2)$

D. $(3, 0)$

Answer: B



Watch Video Solution