



MATHS

BOOKS - CENGAGE MATHS (HINGLISH)

PROGRESSION AND SERIES

Single correct Answer

1. If $3x^2 - 2ax + (a^2 + 2b^2 + 2c^2) = 2(ab + bc)$, then a, b, c can be in

A. *A. P.*

B. *G. P.*

C. *H. P.*

D. None of these

Answer: A



Watch Video Solution

2. If $x = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$, $y = \frac{1}{1^2} + \frac{3}{2^2} + \frac{1}{3^2} + \frac{3}{4^2} + \dots$ and $z = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$ then

A. x, y, z are in *A. P.*

B. $\frac{y}{6}, \frac{x}{3}, \frac{z}{2}$ are in *A. P.*

C. $\frac{y}{6}, \frac{x}{3}, \frac{z}{2}$ are in *A. P.*

D. $6y, 3x, 2z$ are in *H. P.*

Answer: B

 Watch Video Solution

3. For $a, b, c \in \mathbb{R} - \{0\}$, let $\frac{a+b}{1-ab}, b, \frac{b+c}{1-bc}$ are in *A. P.* If α, β are the roots of the quadratic equation

$2acx^2 + 2abcx + (a+c) = 0$, then the value of $(1+\alpha)(1+\beta)$ is

A. 0

B. 1

C. -1

D. 2

Answer: B



[Watch Video Solution](#)

4. If $a_1, a_2, a_3, \dots, a_{87}, a_{88}, a_{89}$ are the arithmetic means between 1 and

89, then $\sum_{r=1}^{89} \log(\tan(a_r)^\circ)$ is equal to

A. 0

B. 1

C. $\log_2 3$

D. $\log 5$

Answer: A



[Watch Video Solution](#)

5. Let a_1, a_2, \dots and b_1, b_2, \dots be arithmetic progression such that $a_1 = 25, b_1 = 75$ and $a_{100} + b_{100} = 100$, then the sum of first hundred term of the progression $a_1 + b_1, a_2 + b_2, \dots$ is equal to

A. 1000

B. 100000

C. 10000

D. 24000

Answer: C



[Watch Video Solution](#)

6. The sum of 25 terms of an *A. P.*, whose all the terms are natural numbers, lies between 1900 and 2000 and its 9^{th} term is 55. Then the first term of the *A. P.* is

A. 5

B. 6

C. 7

D. 8

Answer: C



Watch Video Solution

7. If the first, fifth and last terms of an $A. P.$ is l, m, p , respectively, and sum of the $A. P.$ is $\frac{(l + p)(4p + m - 5l)}{k(m - l)}$ then k is

A. 2

B. 3

C. 4

D. 5

Answer: A



Watch Video Solution

8. If $a_1, a_2, a_3, \dots, a_{15}$ are in A.P. and $a_1 + a_8 + a_{15} = 15$, then $a_2 + a_3 + a_8 + a_{13} + a_{14}$ is equal to

A. 25

B. 35

C. 10

D. 15

Answer: A



Watch Video Solution

9. If a_1, a_2, a_3, \dots are in A.P. and $a_i > 0$ for each i , then

$\sum_{i=1}^n \frac{n}{a_{i+1}^{2/3} + a_{i+1}^{1/3} a_i^{1/3} + a_i^{2/3}}$ is equal to

A. $\frac{n}{a_n^{2/3} + a_n^{1/3} + a_1^{2/3}}$

B.
$$\frac{n + 1}{a_n^{2/3} + a_n^{1/3} + a_1^{2/3}}$$

C.
$$\frac{n - 1}{a_n^{2/3} + a_n^{1/3} \cdot a_1^{1/3} + a_1^{2/3}}$$

D. None of these

Answer: C



Watch Video Solution

10. Between the numbers 2 and 20, 8 means are inserted. Then their sum is

A. 88

B. 44

C. 176

D. None of these

Answer: A



Watch Video Solution

11. Let $a_1, a_2, a_3, \dots, a_{4001}$ is an *A. P.* such that

$$\frac{1}{a_1 a_2} + \frac{1}{a_2 a_3} + \dots + \frac{1}{a_{4000} a_{4001}} = 10$$

$$a_2 + a_{400} = 50.$$

Then $|a_1 - a_{4001}|$ is equal to

A. 20

B. 30

C. 40

D. None of these

Answer: B



[Watch Video Solution](#)

12. An *A. P.* consist of even number of terms $2n$ having middle terms equal to 1 and 7 respectively. If n is the maximum value which satisfy

$t_1 t_{2n} + 713 \geq 0$, then the value of the first term of the series is

A. 17

B. -15

C. 21

D. -23

Answer: D



Watch Video Solution

13. If the sum of the first 100 terms of an AP is -1 and the sum of even terms lying in first 100 terms is 1, then which of the following is not true ?

A. Common difference of the sequence is $\frac{3}{50}$

B. First term of the sequence is $\frac{-149}{50}$

C. 100^{th} term = $\frac{74}{25}$

D. None of these

Answer: D

 [Watch Video Solution](#)

14. Given the sequence of numbers $x_1, x_2, x_3, x_4, \dots, x_{2005}$,
$$\frac{x_1}{x_1 + 1} = \frac{x_2}{x_2 + 3} = \frac{x_3}{x_3 + 5} = \dots = \frac{x_{2005}}{x_{2005} + 4009}$$
, the nature of the
sequence is

A. *A. P.*

B. *G. P.*

C. *H. P.*

D. None of these

Answer: A

 [Watch Video Solution](#)

15. If $b - c, bx - cy, bx^2 - cy^2$ ($b, c \neq 0$) are in *G. P.*, then the value of
$$\left(\frac{bx + cy}{b + c} \right) \left(\frac{bx - cy}{b - c} \right)$$
 is

A. x^2

B. $-x^2$

C. $2y^2$

D. $3y^2$

Answer: A



Watch Video Solution

16. If a_1, a_2, a_3, \dots are in *G. P.*, where $a_i \in C$ (where C stands for set of complex numbers) having r as common ratio such that

$$\sum_{k=1}^n a_{2k-1} \sum_{k=1}^n a_{2k+3} \neq 0, \text{ then the number of possible values of } r \text{ is}$$

A. 2

B. 3

C. 4

D. 5

Answer: C



[Watch Video Solution](#)

17. If a, b, c are real numbers forming an $A. P.$ and $3 + a, 2 + b, 3 + c$ are in $G. P.$, then minimum value of ac is

A. -4

B. -6

C. 3

D. None of these

Answer: B



[Watch Video Solution](#)

18. a, b, c, d are in increasing $G. P.$ If the AM between a and b is 6 and the AM between c and d is 54 , then the AM of a and b is

A. 15

B. 48

C. 44

D. 42

Answer: D



[Watch Video Solution](#)

19. The numbers a, b, c are in $A. P.$ and $a + b + c = 60$. The numbers $(a - 2), b, (c + 3)$ are in $G. P.$ Then which of the following is not the possible value of $a^2 + b^2 + c^2$?

A. 1208

B. 1218

C. 1298

D. None of these

Answer: B



[Watch Video Solution](#)

20. a, b, c are positive integers forming an increasing $G. P.$ and $b - a$ is a perfect cube and $\log_6 a + \log_6 b + \log_6 c = 6$, then $a + b + c =$

A. 100

B. 111

C. 122

D. 189

Answer: D



[Watch Video Solution](#)

21. The first three terms of a geometric sequence are x, y, z and these have the sum equal to 42. If the middle term y is multiplied by $5/4$, the

numbers $x, \frac{5y}{4}, z$ now form an arithmetic sequence. The largest possible value of x is

- A. 6
- B. 12
- C. 24
- D. 20

Answer: C



[Watch Video Solution](#)

22. An infinite $G. P.$ has 2^{nd} term x and its sum is 4. Then x belongs to

- A. $(0, 2]$
- B. $(1, 8)$
- C. $(-8, 1]$
- D. none of these

Answer: C



[Watch Video Solution](#)

23. In a GP , the ratio of the sum of the first eleven terms of the sum of the last even terms is $1/8$ and the ratio of the sum of all the terms without the first nine to the sum of all terms without the last nine is 2. Then the number of terms in the GP is

- A. 40
- B. 38
- C. 36
- D. 34

Answer: B



[Watch Video Solution](#)

24. The number of ordered pairs (x, y) , where $x, y \in \mathbb{N}$ for which $4, x, y$ are in *H. P.*, is equal to

A. 1

B. 2

C. 3

D. 4

Answer: C



[Watch Video Solution](#)

25. If $a + c, a + b, b + c$ are in *G. P.* and a, c, b are in *H. P.* where $a, b, c > 0$, then the value of $\frac{a + b}{c}$ is

A. 3

B. 2

C. $\frac{3}{2}$

D. 4

Answer: B



Watch Video Solution

26. If a, b, c are in $H. P.$, b, c, d are in $G. P.$ and c, d, e are in $A. P.$, then the value of e is

A. $\frac{ab^2}{(2a - b)^2}$

B. $\frac{a^2b}{(2a - b)^2}$

C. $\frac{a^2b^2}{(2a - b)^2}$

D. None of these

Answer: A



Watch Video Solution

27. If $x > 1, y > 1, z > 1$ are in *G. P.*, then $\log_{ex} e, \log_{ey} e, \log_{ez} e$ are in

A. *A. P.*

B. *H. P.*

C. *G. P.*

D. none of these

Answer: B



[Watch Video Solution](#)

28. If x, y, z are in *G. P.* ($x, y, z > 1$), then $\frac{1}{2x + \log_e x}, \frac{1}{4x + \log_e y},$
 $\frac{1}{6x + \log_{ez} z}$ are in

A. *A. P.*

B. *G. P.*

C. *H. P.*

D. none of these

Answer: C



[Watch Video Solution](#)

29. The arithmetic mean of two positive numbers is 6 and their geometric mean G and harmonic mean H satisfy the relation $G^2 + 3H = 48$. Then the product of the two numbers is

A. 24

B. 32

C. 48

D. 54

Answer: B



[Watch Video Solution](#)

30. If x, y, z be three numbers in $G. P.$ such that 4 is the $A. M.$ between x and y and 9 is the $H. M.$ between y and z , then y is

- A. 4
- B. 6
- C. 8
- D. 12

Answer: B



[Watch Video Solution](#)

31. If harmonic mean of $\frac{1}{2}, \frac{1}{2^2}, \frac{1}{2^3}, \dots, \frac{1}{2^{10}}$ is $\frac{\lambda}{2^{10} - 1}$, then $\lambda =$

- A. $10 \cdot 2^{10}$
- B. 5
- C. $5 \cdot 2^{10}$
- D. 10

Answer: B



[Watch Video Solution](#)

32. An aeroplane flies around squares whose all sides are of length 100 miles. If the aeroplane covers at a speed of $100mph$ the first side, $200mph$ the second side $300mph$ the third side and $400mph$ the fourth side. The average speed of aeroplane around the square is

A. $190mph$

B. $195mph$

C. $192mph$

D. $200mph$

Answer: C



[Watch Video Solution](#)

33. The sum of the series $1 + \frac{9}{4} + \frac{36}{9} + \frac{100}{16} + \dots$ infinite terms is

A. 446

B. 746

C. 546

D. 846

Answer: A



[Watch Video Solution](#)

34. The sum $2 \times 5 + 5 \times 9 + 8 \times 13 + \dots$ 10 terms is

A. 4500

B. 4555

C. 5454

D. None of these

Answer: B



Watch Video Solution

35. The sum of n terms of series
 $ab + (a + 1)(b + 1) + (a + 2)(b + 2) + \dots + (a + (n - 1))(b + (n - 1))$
if $ab = \frac{1}{6}$ and $(1 + b) = \frac{1}{3}$ is

A. $\frac{n}{6}(1 - 2n)^2$

B. $\frac{n}{6}(1 + n - 2n^2)$

C. $\frac{n}{6}(1 - 2n + 2n^2)$

D. None of these

Answer: C



Watch Video Solution

36. $\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{a^{i+j+k}}$ is equal to (where $|a| > 1$)

A. $(a - 1)^{-3}$

B. $\frac{3}{a - 1}$

C. $\frac{3}{a^3 - 1}$

D. None of these

Answer: A



Watch Video Solution

37. The coefficient of x^{1274} in the expansion of $(x + 1)(x - 2)^2(x + 3)^3(x - 4)^4 \dots (x + 49)^{49}(x - 50)^{50}$ is

A. 1275

B. -1275

C. $-\sum_{i=1}^{50} i^2$

D. $-\sum_{i=1}^{50} i^2$

Answer: B



Watch Video Solution

38. If the positive integers are written in a triangular array as shown below,



then the row in which the number 2010 will be, is

A. 65

B. 61

C. 63

D. 65

Answer: C



View Text Solution

39. The value of $\sum_{i=1}^n \sum_{j=1}^i \sum_{k=1}^j = 220$, then the value of n equals

A. 11

B. 12

C. 10

D. 9

Answer: C

 [Watch Video Solution](#)

40. The sum $\sum_{k=1}^{10} \sum_{\substack{j=1 \\ i \neq j \neq k}}^{10} \sum_{i=1}^{10} 1$ is equal to

A. 240

B. 720

C. 540

D. 1080

Answer: B

41. The sum $\sum_{k=1}^{10} \sum_{\substack{j=1 \\ i < j < k}}^{10} \sum_{i=1}^{10} 1$ is equal to

- A. 120
- B. 240
- C. 360
- D. 720

Answer: A

42. If the sum to infinity of the series $1 + 4x + 7x^2 + 10x^3 + \dots$, is $\frac{35}{16}$, where $|x| < 1$, then 'x' equals to

- A. $19/7$
- B. $1/5$

C. $1/4$

D. None of these

Answer: B



[Watch Video Solution](#)

43. The value of $\sum_{n=1}^{\infty} (-1)^{n+1} \left(\frac{n}{5^n} \right)$ equals

A. $\frac{5}{12}$

B. $\frac{5}{24}$

C. $\frac{5}{36}$

D. $\frac{5}{16}$

Answer: C



[Watch Video Solution](#)

44. Find the sum of the infinite series $\frac{1}{9} + \frac{1}{18} + \frac{1}{30} + \frac{1}{45} + \frac{1}{63} + \dots$

A. $\frac{1}{3}$

B. $\frac{1}{4}$

C. $\frac{1}{5}$

D. $\frac{2}{3}$

Answer: A



[Watch Video Solution](#)

45. If $\sum_{r=1}^{r=n} \frac{r^4 + r^2 + 1}{r^4 + r} = \frac{675}{26}$, then n equal to

A. 10

B. 15

C. 25

D. 30

Answer: C



Watch Video Solution

46. The sequence $\{x_k\}$ is defined by $x_{k+1} = x_k^2 + x_k$ and $x_1 = \frac{1}{2}$. Then

$\left[\frac{1}{x_1 + 1} + \frac{1}{x_2 + 1} + \dots + \frac{1}{x_{100} + 1} \right]$ (where $[.]$ denotes the greatest

integer function) is equal to

A. 0

B. 2

C. 4

D. 1

Answer: D



Watch Video Solution

47. The absolute value of the sum of first 20 terms of series, if

$$S_n = \frac{n+1}{2} \text{ and } \frac{T_{n-1}}{T_n} = \frac{1}{n^2} - 1, \text{ where } n \text{ is odd, given } S_n \text{ and } T_n$$

denotes sum of first n terms and n^{th} terms of the series

A. 340

B. 430

C. 230

D. 320

Answer: B



[Watch Video Solution](#)

48.

if

$$S_n = (1^2 - 1 + 1)(1!) + (2^2 - 2 + 1)(2!) + \dots + (n^2 - n + 1)(n!),$$

then $S_{50} =$

A. 52!

B. $1 + 49 \times 5!$

C. $52! - 1$

D. $50 \times 51! - 1$

Answer: B



Watch Video Solution

49. If $S_n = \frac{1.2}{3!} + \frac{2.2^2}{4!} + \frac{3.2^2}{5!} + \dots +$ up to n terms, then sum of infinite terms is

A. $\frac{4}{\pi}$

B. $\frac{3}{e}$

C. $\frac{\pi}{r}$

D. 1

Answer: D



Watch Video Solution

50. There is a certain sequence of positive real numbers. Beginning from the third term, each term of the sequence is the sum of all the previous terms. The seventh term is equal to 1000 and the first term is equal to 1. The second term of this sequence is equal to

A. 246

B. $\frac{123}{2}$

C. $\frac{123}{4}$

D. 124

Answer: B



[Watch Video Solution](#)

51. The sequence $\{x_1, x_2, \dots, x_{50}\}$ has the property that for each k , x_k is k less than the sum of other 49 numbers. The value of $96x_{20}$ is

A. 300

B. 315

C. 1024

D. 0

Answer: B

 [Watch Video Solution](#)

52. Let $a_0 = 0$ and $a_n = 3a_{n-1} + 1$ for $n \geq 1$. Then the remainder obtained dividing a_{2010} by 11 is

A. 0

B. 7

C. 3

D. 4

Answer: A

 [Watch Video Solution](#)

53. Suppose $a_1, a_2, a_3, \dots, a_{2012}$ are integers arranged on a circle. Each number is equal to the average of its two adjacent numbers. If the sum of all even indexed numbers is 3018, what is the sum of all numbers ?

- A. 0
- B. 9054
- C. 12072
- D. 6036

Answer: D

 [Watch Video Solution](#)

54. The sum of the series $\frac{9}{5^2 \cdot 2 \cdot 1} + \frac{13}{5^3 \cdot 3 \cdot 2} + \frac{17}{5^4 \cdot 4 \cdot 3} + \dots$ upto infinity

A. 1

B. $\frac{9}{5}$

C. $\frac{1}{5}$

D. $\frac{2}{5}$

Answer: C



Watch Video Solution

Comprehension

1. The 1^{st} , 2^{nd} and 3^{rd} terms of an arithmetic series are a , b and a^2 where ' a ' is negative. The 1^{st} , 2^{nd} and 3^{rd} terms of a geometric series are a , a^2 and b respectively.

The sum of infinite geometric series is

A. $\frac{-1}{2}$

B. $\frac{-3}{2}$

C. $\frac{-1}{3}$

D. None of these

Answer: C



Watch Video Solution

2. The 1^{st} , 2^{nd} and 3^{rd} terms of an arithmetic series are a , b and a^2 where ' a ' is negative. The 1^{st} , 2^{nd} and 3^{rd} terms of a geometric series are a , a^2 and b respectively.

The sum of the 40 terms of the arithmetic series is

A. $\frac{545}{2}$

B. 220


C. 250

D. $\frac{575}{2}$

Answer: A



Watch Video Solution



3.

Let $ABCD$ is a unit square and each side of the square is divided in the ratio $\alpha:(1-\alpha)$ ($0 < \alpha < 1$) . These points are connected to obtain another square. The sides of new square are divided in the ratio $\alpha:(1-\alpha)$ and points are joined to obtain another square. The process is continued indefinitely. Let a_n denote the length of side and A_n the area of the n^{th} square

If $\alpha = \frac{1}{3}$, then the least value of n for which $A_n > \frac{1}{10}$ is

A. 4


B. 5

C. 6

D. 7

Answer: B

 [View Text Solution](#)



4.

Let $ABCD$ is a unit square and each side of the square is divided in the ratio $\alpha : (1 - \alpha)$ ($0 < \alpha < 1$). These points are connected to obtain another square. The sides of new square are divided in the ratio

$\alpha: (1 - \alpha)$ and points are joined to obtain another square. The process is continued indefinitely. Let a_n denote the length of side and A_n the area of the n^{th} square

The value of α for which $\sum_{n=1}^{\infty} A_n = \frac{8}{3}$ is/are

A. $\frac{1}{3}, \frac{2}{3}$

B. $\frac{1}{4}, \frac{3}{4}$


C. $\frac{1}{5}, \frac{4}{5}$

D. $\frac{1}{2}$

Answer: B



[View Text Solution](#)



5.

Let $ABCD$ is a unit square and each side of the square is divided in the ratio $\alpha : (1 - \alpha)$ ($0 < \alpha < 1$). These points are connected to obtain another square. The sides of new square are divided in the ratio $\alpha : (1 - \alpha)$ and points are joined to obtain another square. The process is continued indefinitely. Let a_n denote the length of side and A_n the area of the n^{th} square

The value of α for which side of n^{th} square equal to the diagonal of $(n + 1)^{\text{th}}$ square is

A. $\frac{1}{3}$

B. $\frac{1}{4}$

C. $\frac{1}{2}$

D. $\frac{1}{\sqrt{2}}$

Answer: C



[View Text Solution](#)

6. Let $f(n)$ denote the n^{th} terms of the sequence of 3, 6, 11, 18, 27, and $g(n)$ denote the n^{th} terms of the sequence of 3, 7, 13, 21, Let $F(n)$ and $G(n)$ denote the sum of n terms of the above sequences, respectively. Now answer the following:

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} =$$

A. 0

B. 1

C. 2

D. ∞

Answer: B



Watch Video Solution

7. Let $f(n)$ denote the n^{th} terms of the sequence of 3, 6, 11, 18, 27, ... and $g(n)$ denote the n^{th} terms of the sequence of 3, 7, 13, 21, Let $F(n)$ and $G(n)$ denote the sum of n terms of the above sequences, respectively. Now answer the following:

$$\lim_{n \rightarrow \infty} \frac{F(n)}{G(n)} =$$

A. 2

B. 1

C. 0

D. ∞

Answer: B



Watch Video Solution

Multiple Correct Answer

1. Let a, x, b be in $A. P$, a, y, b be in $G. P$ and a, z, b be in $H. P$. If $x = y + 2$ and $a = 5z$, then

A. $y^2 = xz$

B. $x > y > z$

C. $a = 9, b = 1$

D. $a = 1/4, b = 9/4$

Answer: A::B::C



[Watch Video Solution](#)

2. If $A_1, A_2, A_3, G_1, G_2, G_3$, and H_1, H_2, H_3 are the three arithmetic, geometric and harmonic means between two positive numbers a and b ($a > b$), then which of the following is/are true ?

A. $2G_1G_3 = H_2(A_1 + A_3)$

B. $A_2H_2 = G_2^2$

C. $A_2G_2 = H_2^2$

D. $2G_1A_1 = H_1(A_1 + A_3)$

Answer: A::B



Watch Video Solution

3. Given that α, γ are roots of the equation $Ax^2 - 4x + 1 = 0$ and β, δ are roots of the equation $Bx^2 - 6x + 1 = 0$. If α, β, γ and δ are in *H. P.*, then

A. $A = 5$

B. $A = 3$

C. $B = 8$

D. $B = -8$

Answer: B

 Watch Video Solution

4. If $\frac{1}{a} + \frac{1}{c} = \frac{1}{2b - a} + \frac{1}{2b - c}$, then

A. a, b, c are in *A. P.*

B. $a, \frac{b}{2}, c$ are in *A. P.*

C. $a, \frac{b}{2}, c$ are in *H. P.*

D. $a, 2b, c$ are in *H. P.*

Answer: A::D

 Watch Video Solution

Examples

1. Write down the sequence whose n th term is $2^n/n$ and (ii)

$$[3 + (-1)^n] / 3^n$$

 Watch Video Solution

2. Find the sequence of the numbers defined by

$$a_n = \begin{cases} \frac{1}{n} & \text{when } n \text{ is odd} \\ -\frac{1}{n} & \text{when } n \text{ is even} \end{cases}$$

 [Watch Video Solution](#)

3. Write the first three terms of the sequence defined by

$$a_1 = 2, a_{n+1} = \frac{2a_n + 3}{a_n + 2}.$$

 [Watch Video Solution](#)

4. The Fibonacci sequence is defined by

$$1 = a_1 = a_2 \text{ and } a_n = a_{n-1} + a_{n-2}, n > 2. \text{ Find } \frac{a_{n+1}}{a_n}, \text{ for } n = 5.$$

 [Watch Video Solution](#)

5. A sequence of integers $a_1 + a_2 + \dots + a_n$ satisfies $a_{n+2} = a_{n+1} - a_n$ for $n \geq 1$. Suppose the sum of first 999 terms is 1003 and the sum of the first 1003 terms is -99. Find the sum of the first 2002 terms.

 [Watch Video Solution](#)

6. Show that the sequence 9,12,15,18,... is an A.P. Find its 16th term and the general term.

 [Watch Video Solution](#)

7. Show that the sequence $\log a, \log(ab), \log(ab^2), \log(ab^3), \dots$ is an A.P. Find its n th term.

 [Watch Video Solution](#)

8. In a certain A.P., 5 times the 5th term is equal to 8 times the 8th term.

Then prove that its 13th term is 0.

 [Watch Video Solution](#)

9. Find the term of the series $25, 22, \frac{3}{4}, 20\frac{1}{2}, 18\frac{1}{4}$ which is numerically the smallest.

 [Watch Video Solution](#)

10. If p th, q th and r th terms of an A.P. are a, b, c , respectively, then show that

(i) $a(q-r)+b(r-p)+c(p-q)=0$

(ii) $(a-b)r+(b-c)p+(c-a)q=0$

 [Watch Video Solution](#)

11. Consider two A.P. s: $S_1 : 2, 7, 12, 17, 500\text{terms}$ and $S_2 : 1, 8, 15, 22, 300\text{terms}$ Find the number of common term. Also find the last common term.



[Watch Video Solution](#)

12. If $a_1, a_2, a_3, \dots, a_n$ are in A.P., where $a_i > 0$ for all i , show that

$$\frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_1} + \sqrt{a_3}} + \dots + \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} = \frac{n-1}{\sqrt{a_1} + \sqrt{a_n}}.$$



[Watch Video Solution](#)

13. If p, q and r ($p \neq q$) are terms (not necessarily consecutive) of an A.P., then prove that there exists a rational number k such that $\frac{r-q}{q-p} = k$.
hence, prove that the numbers $\sqrt{2}, \sqrt{3}$ and $\sqrt{5}$ cannot be the terms of a single A.P. with non-zero common difference.



[Watch Video Solution](#)

$$\frac{1}{\sqrt{b} + \sqrt{c}}, \frac{1}{\sqrt{c} + \sqrt{a}}, \frac{1}{\sqrt{a} + \sqrt{b}}$$

(iii) $a\left(\frac{1}{b} + \frac{1}{c}\right), b\left(\frac{1}{c} + \frac{1}{a}\right), c\left(\frac{1}{a} + \frac{1}{b}\right)$



[View Text Solution](#)

18. If the sum of three numbers in A.P., is 24 and their product is 440, find the numbers.



[Watch Video Solution](#)

19. Divide 32 into four parts which are in A.P. such that the ratio of the product of extremes to the product of means is 7:15.



[Watch Video Solution](#)

20. The digits of a positive integer, having three digits, are in A.P. and their sum is 15. The number obtained by reversing the digits is 594 less than the original number. Find the number.



[Watch Video Solution](#)

21. If eleven A.M. s are inserted between 28 and 10, then find the number of integral A.M. s.



[Watch Video Solution](#)

22. Between 1 and 31 are inserted m arithmetic mean so that the ratio of the 7th and $(m - 1)th$ means is 5:9. Find the value of m .



[Watch Video Solution](#)

23. find the sum of all three digit natural numbers which are divisible by 7



[Watch Video Solution](#)

24. Find the number of terms in the series $20, 19\frac{1}{3}, 18\frac{2}{3}, \dots$ the sum of which is 300. Explain the answer.

 [Watch Video Solution](#)

25. Find the degree of the expression $(1 + x)(1 + x^6)(1 + x^{11})(1 + x^{101})$.

 [Watch Video Solution](#)

26. Find the sum of first 24 terms of the A.P. a_1, a_2, a_3, \dots , if it is known that $a_1 + a_5 + a_{10} + a_{15} + a_{20} + a_{24} = 225$.

 [Watch Video Solution](#)

27. If S_1 is the sum of an AP of 'n' odd number of terms and S_2 be the sum of the terms of series in odd places of the same AP then $\frac{S_1}{S_2} =$

 [Watch Video Solution](#)

28. If the sequence $a_1, a_2, a_3, \dots, a_n$ is an A.P., then prove that

$$a_1^2 - a_2^2 + a_3^2 - a_4^2 + \dots + a_{2n-1}^2 - a_{2n}^2 = \frac{n}{2n-1} (a_1^2 - a_{2n}^2)$$

 [Watch Video Solution](#)

29. If the arithmetic progression whose common difference is nonzero the sum of first $3n$ terms is equal to the sum of next n terms. Then, find the ratio of the sum of the $2n$ terms to the sum of next $2n$ terms.

 [Watch Video Solution](#)

30. The sum of n terms of two arithmetic progressions are in the ratio $5n + 4 : 9n + 6$. Find the ratio of their 18th terms.

 [Watch Video Solution](#)

31. If n arithmetic means are inserted between 2 and 38, then the sum of the resulting series is obtained as 200. Then find the value of n .

 [Watch Video Solution](#)

32. The third term of a geometric progression is 4. Then find the product of the first five terms.

 [Watch Video Solution](#)

33. किसी गुणोत्तर श्रेणी का प्रथम पद 1 है | तीसरे एवं पाँचवें पदों का योग 90 हो तो गुणोत्तर श्रेणी का सार्व अनुपात ज्ञात किजिए |

 [Watch Video Solution](#)

34. If $\frac{a + bx}{a - bx} = \frac{b + cx}{b - cx} = \frac{c + dx}{c - dx} (x \neq 0)$, then show that a, b, c and d are in G.P.

 [Watch Video Solution](#)

35. The fourth, seventh, and the last term of a G.P. are 10, 80, and 2560, respectively. Find the first term and the number of terms in G.P.

 [Watch Video Solution](#)

36. If a, b, c, d and p are distinct real numbers such that $(a^2 + b^2 + c^2)p^2 - 2(ab + bc + cd)p + (b^2 + c^2 + d^2) \leq 0$, then prove that a, b, c, d are in G.P.

 [Watch Video Solution](#)

37. Does there exist a geometric progression containing 27, 8 and 12 as three of its terms? If it exists, then how many such progressions are possible?

 [Watch Video Solution](#)

38. In a sequence of $(4n + 1)$ terms, the first $(2n + 1)$ terms are in A.P. whose common difference is 2, and the last $(2n + 1)$ terms are in G.P. whose common ratio is 0.5 if the middle terms of the A.P. and G.P. are equal, then the middle terms of the sequence is $\frac{n \cdot 2n + 1}{2^{2n} - 1}$ b. $\frac{n \cdot 2n + 1}{2^n - 1}$
 c. $n \cdot 2^n$ d. none of these

 [Watch Video Solution](#)

39. For what value of n , $\frac{a^{n+1} + b^{n+1}}{a^n + b^n}$ is the arithmetic mean of a and b ?

 [Watch Video Solution](#)

40. If $(p + q)$ th term of a G.P. is a and its $(p - q)$ th term is b where $a, b \in \mathbb{R}^+$, then its p th term is $\sqrt{\frac{a^3}{b}}$ b. $\sqrt{\frac{b^3}{a}}$ c. \sqrt{ab} d. none of these

 [Watch Video Solution](#)

41. Find four numbers in G.P. whose sum is 85 and product is 4096.



Watch Video Solution

42. Three non-zero numbers $a, b, \text{ and } c$ are in A.P. Increasing a by 1 or increasing c by 2, the numbers are in G.P. Then find b .



Watch Video Solution

43. If a, b, c are in A.P., b, c, d are in G.P. and $\frac{1}{c}, \frac{1}{d}, \frac{1}{e}$ are in A.P. prove that a, c, e are in G.P.



Watch Video Solution

44. If G is the geometric mean of x and y then prove that

$$\frac{1}{G^2 - x^2} + \frac{1}{G^2 - y^2} = \frac{1}{G^2}$$



Watch Video Solution

45. Insert four G.M.s between 2 and 486.

 [Watch Video Solution](#)

46. If A.M. and G.M. between two numbers is in the ratio $m:n$ then prove

that the numbers are in the ratio

$$\left(m + \sqrt{m^2 - n^2}\right) : \sqrt{\left(m - m^2 - n^2\right)}.$$

 [Watch Video Solution](#)

47. If a be one A.M and G_1 and G_2 be then geometric means between b

and c then $G_1^3 + G_2^3 =$

 [Watch Video Solution](#)

48. Determine the number of terms in G.P. $\langle a_n \rangle$, if $a_1=3, a_n=96a_{n-1}, S_n=189$.



[Watch Video Solution](#)



Watch Video Solution

49. if S is the sum , P the product and R the sum of reciprocals of n terms in G . P. prove that $P^2 R^n = S^n$



Watch Video Solution

50. Find the sum to n terms of the sequence $(x + 1/x)^2, (x^2 + 1/x)^2, (x^3 + 1/x)^2, ,$



Watch Video Solution

51. Prove that the sum to n terms of the series $11 + 103 + 1005 + \dots + \frac{10}{9}(10^n - 1) + n^2$.



Watch Video Solution

52. Find the sum of the following series up to n terms: (i)

$$5 + 55 + 555 + \dots \quad \text{(ii)}$$

$$.6 + .66 + .666 + \dots$$

 [Watch Video Solution](#)

53. Find the sum $1 + (1 + 2) + (1 + 2 + 2^2) + (1 + 2 + 2^2 + 2^3) + \dots$

To n terms.

 [Watch Video Solution](#)

54. If the sum of the n terms of a G.P. is $(3^n - 1)$, then find the sum of the series whose terms are reciprocal of the given G.P..

 [Watch Video Solution](#)

55. Prove that in a sequence of numbers 49,4489,444889,44448889 in which every number is made by inserting 48-48 in the middle of previous as indicated, each number is the square of an integer.



[Watch Video Solution](#)

56. If f is a function satisfying $f(x + y) = f(x) \times f(y)$ for all $x, y \in N$ such that $f(1) = 3$ and $\sum_{x=1}^n f(x) = 120$, find the value of n .



[Watch Video Solution](#)

57. Using the sum of G.P., prove that $a^n + b^n$ ($a, b \in N$) is divisible by $a+b$ for odd natural numbers n . Hence prove that $1^{99} + 2^{99} + \dots + 100^{99}$ is divisible by 10100



[Watch Video Solution](#)

58. Sum the following geometric series to infinity:

$$(\sqrt{2} + 1) + 1 + (\sqrt{2} - 1) + \infty \quad \frac{1}{2} + \frac{1}{3^3} + \frac{1}{2^3} + \frac{1}{3^4} + \frac{1}{2^5} + \frac{1}{3^6} + \infty$$

 [Watch Video Solution](#)

59. The sum of infinite number of terms in G.P. is 20 and the sum of their squares is 100. Then find the common ratio of G.P.

 [Watch Video Solution](#)

60. If each term of an infinite G.P. is twice the sum of the terms following it, then find the common ratio of the G.P.

 [Watch Video Solution](#)

61. If

$$x = a + \frac{a}{r} + \frac{a}{r^2} + \infty, y = b - \frac{b}{r} + \frac{b}{r^2} + \infty, \text{ and } z = c + \frac{c}{r^2} + \frac{c}{r^4} + \infty$$

prove that $\frac{xy}{z} = \frac{ab}{.}$

 [Watch Video Solution](#)

62. After striking a floor a certain ball rebounds $\left(\frac{4}{5}\right)^{th}$ of the height from which it has fallen. Find the total distance that it travels before coming to rest, if it is gently dropped from a height of 120metres.

 [Watch Video Solution](#)

63. If an infinite G.P. has 2nd term x and its sum is 4, then prove that $\xi_n(-8, 1] - \{0\}$

 [View Text Solution](#)

64. If the 20th term of a H.P. is 1 and the 30th term is $-1/17$, then find its largest term.

 [Watch Video Solution](#)

65. If $\frac{a-x}{px} = \frac{a-y}{qy} = \frac{a-z}{r}$ and $p, q, \text{ and } r$ are in A.P., then prove that x, y, z are in H.P.



Watch Video Solution

66. If $a, b, \text{ and } d$ are in H.P., then prove that $(b+c+d)/a, (c+d+a)/b, (d+a+b)/c$ and $(a+b+c)/d$, are in A.P.



Watch Video Solution

67. The m th term of a H.P is n and the n th term is m . Prove that its r th term is mn/r .



Watch Video Solution

68. If $a > 1$, $b > 1$ and $c > 1$ are in G.P., then show that $\frac{1}{1 + \log_e a}$, $\frac{1}{1 + \log_e b}$ and $\frac{1}{1 + \log_e c}$ are in H.P.

 [Watch Video Solution](#)

69. If a, b, c are in G.P and $a + x, b + x, c + x$ are in H.P, then the value of x is (a, b, c are distinct numbers)

 [Watch Video Solution](#)

70. If first three terms of the sequence $1/16, a, b, c, 1/16$ are in geometric series and last three terms are in harmonic series, then find the values of a and b .

 [Watch Video Solution](#)

71. if $(m + 1)th$, $(n + 1)th$ and $(r + 1)th$ term of an AP are in GP. and m , n and r in HP. . find the ratio of first term of A.P to its common difference

 [Watch Video Solution](#)

72. Insert four H.M.s between $2/3$ and $2/13$.

 [Watch Video Solution](#)

73. If nine arithmetic means and nine harmonic means are inserted between 2 and 3 alternatively, then prove that $A + 6/H = 5$ (where A is any of the A.M.'s and H the corresponding H.M.).

 [Watch Video Solution](#)

74. Let a, b be positive real numbers. If a, A_1, A_2, b be are in arithmetic progression a, G_1, G_2, b are in geometric progression, and a, H_1, H_2, b

are in harmonic progression, show that

$$\frac{G_1 G_2}{H_1 H_2} = \frac{A_1 + A_2}{H_1 + H_2} = \frac{(2a + b)(a + 2b)}{9ab}$$

 [Watch Video Solution](#)

75. The A.M. and H.M. between two numbers are 27 and 122, respectively, then find their G.M.

 [Watch Video Solution](#)

76. If the A.M. between two numbers exceeds their G.M. by 2 and the GM. Exceeds their H.M. by $\frac{8}{5}$, find the numbers.

 [Watch Video Solution](#)

77. Find the sum

$$2017 + \frac{1}{4} \left(2016 + \frac{1}{4} \left(2015 + \dots + \frac{1}{4} \left(2 + \frac{1}{4}(1) \right) \dots \right) \right)$$

 [Watch Video Solution](#)

78. The sum of 50 terms of the series $1 + 2\left(1 + \frac{1}{50}\right) + 3\left(1 + \frac{1}{50}\right)^2 + \dots$ is given by 2500 b. 2550 c. 2450 d. none of these

 [Watch Video Solution](#)

79. Find the sum to infinity of the series $1 - 3x + 5x^2 + 7x^3 + \dots \infty$ when $|x| < 1$.

 [Watch Video Solution](#)

80. The sum of the infinite series $1 + \left(1 + \frac{1}{5}\right)\left(\frac{1}{2}\right) + \left(1 + \frac{1}{5} + \frac{1}{5^2}\right)\left(\frac{1}{2^2}\right) + \dots$

 [Watch Video Solution](#)

81. If the sum to infinity of the series $3 + (3 + d)\frac{1}{4} + (3 + 2d)\frac{1}{4^2} + \dots$ is $\frac{44}{9}$, then find d .

 [Watch Video Solution](#)

82. Find the sum to infinity of the series $1^2 + 2^2 + 3^2 + 4^2 + \dots$.

 [Watch Video Solution](#)

83. Find the sum $2 \times 5 + 5 \times 9 + 8 \times 13 + 11 \times 17 + \dots$ n terms.

 [Watch Video Solution](#)

84. Find the sum of the series

$$1 \times n + 2(n - 1) + 3 \times (n - 2) + \dots + (n - 1) \times 2 + n \times 1.$$

 [Watch Video Solution](#)

85. For and odd integer $n \geq 1$, $n^3 - (n - 1)^3 + \dots$

$$+ (-1)^{n-1} 1^3$$

 [Watch Video Solution](#)

86. Find the sum of the series $\frac{1^3}{1} + \frac{1^3 + 2^3}{1 + 3} + \frac{1^3 + 2^3 + 3^3}{1 + 3 + 5} + \dots$ up to n terms.

 [Watch Video Solution](#)

87. Find the sum of first n terms of the series $1^3 + 3 \times 2^2 + 3^3 + 3 \times 4^2 + 5^3 + 3 \times 6^2 + \dots$ when n is even n is odd

 [Watch Video Solution](#)

88. If $\sum_{r=1}^n T_r = n(2n^2 + 9n + 13)$, then find the sum $\sum_{r=1}^n \sqrt{T_r}$.

 [Watch Video Solution](#)

89. Find the sum to n terms of the series $3 + 15 + 35 + 63 +$

 [Watch Video Solution](#)

90. Find the sum of the following series to n terms
 $5 + 7 + 13 + 31 + 85 +$

 [Watch Video Solution](#)

91. Find the $\sum_{k=1}^{\infty} \sum_{n=1}^{\infty} \frac{k}{2^{n+k}}$.

 [Watch Video Solution](#)

92. The sum of the products of the ten numbers $\pm 1, \pm 2, \pm 3, \pm 4, \pm 5$ taking two at a time is:

 [Watch Video Solution](#)

93. Find the $\sum_{0 \leq i < j \leq n} 1$.

 [View Text Solution](#)

94. Let the terms $a_1, a_2, a_3, \dots, a_n$ be in G.P. with common ratio r . Let S_k denote the sum of first k terms of this G.P. Prove that

$$S_{m-1} \times S_m = \frac{r+1}{r} \sum_{i=1}^m \sum_{j=i}^m a_i a_j$$

 [View Text Solution](#)

95. Find the sum $1 + \frac{1}{1+2} + \frac{1}{1+2+3} + \dots + \frac{1}{1+2+3+\dots+n}$.

 [Watch Video Solution](#)

96. Find the sum of the series:

$$\frac{1}{(1 \times 3)} + \frac{1}{(3 \times 5)} + \frac{1}{(5 \times 7)} + \dots + \frac{1}{(2n-1)(2n+1)}$$

 [Watch Video Solution](#)

97. Find the sum to n terms of the series
 $3/(1^2 \times 2^2) + 5/(2^2 \times 3^2) + 7/(3^2 \times 4^2) + \dots$

 [Watch Video Solution](#)

98. Find the sum to n terms of the series:

$$\frac{1}{1 + 1^2 + 1^4} + \frac{1}{1 + 2^2 + 2^4} + \frac{1}{1 + 3^2 + 3^4} + \dots$$

 [Watch Video Solution](#)

99. Find the sum $\sum_{r=1}^n \frac{r}{(r+1)!}$. Also, find the sum of infinite terms.

 [Watch Video Solution](#)

100. Find the sum $\sum_{r=1}^n \frac{1}{r(r+1)(r+2)(r+3)}$

Also, find $\sum_{r=1}^{\infty} \frac{1}{r(r+1)(r+2)(r+3)}$

 [Watch Video Solution](#)

101. Find the sum $\sum_{r=1}^n r(r+1)(r+2)(r+3)$.

 [Watch Video Solution](#)

102. Find the sum of the series $\sum_{r=11}^{99} \left(\frac{1}{r\sqrt{r+1} + (r+1)\sqrt{r}} \right)$

 [Watch Video Solution](#)

103. Find the sum of the series

$$\frac{1}{3^2 + 1} + \frac{1}{4^2 + 2} + \frac{1}{5^2 + 3} + \frac{1}{6^2 + 4} + \infty$$

 [Watch Video Solution](#)

104. Find the sum of first 100 terms of the series whose general term is given by $T_r = (r^2 + 1)r!$.

 [Watch Video Solution](#)

105. Find the sum of the series $\frac{2}{1 \times 3} + \frac{5}{2 \times 3} \times 2 + \frac{10}{3 \times 4} \times 2^2 + \frac{17}{4 \times 5} \times 2^3 + \dots \rightarrow n$ terms.

 [Watch Video Solution](#)

106. Along a road lie an odd number of stones placed at intervals of 10 metres. These stones have to be assembled around the middle stone. A person can carry only one stone at a time. A man carried the job with one of the end stones by carrying them in succession. In carrying all the stones he covered a distance of 3 km. Find the number of stones.

 [Watch Video Solution](#)

107. Prove that $x = \underbrace{1111, \dots}_{91 \text{ times}}$ is composite number.

 [Watch Video Solution](#)

108. If a, b, c are distinct positive real numbers in G.P and $\log_c a, \log_b c, \log_a b$ are in A.P, then find the common difference of this A.P

 [Watch Video Solution](#)

109. The values of xyz is $\frac{15}{2}$ or $\frac{18}{5}$ according as the series a, x, y, z, b is an *AP* or *HP*. Find the values of a & b assuming them to be positive integer.

 [Watch Video Solution](#)

110. Let $p (> 0)$ be the first of the n arithmetic means between two numbers and $q (> 0)$ the first of n harmonic means between the

same numbers. Then prove that

$$q \notin \left(p, \left(\frac{n+1}{n-1} \right)^2 p \right) \text{ and } p \notin \left(\left(\frac{n-1}{n+1} \right)^2 q, q \right)$$

 [View Text Solution](#)

111. If $S_n = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}$ ($n \in N$), then prove that

$$S_1 + S_2 + \dots + S_{(n-1)} = (nS((n)) - n) \text{ or } (nS((n-1)) - n + 1)$$

 [Watch Video Solution](#)

112. The value of the expression

$$1. (2 - \omega). (2 - \omega^2) + 2. (3 - \omega)(3 - \omega^2) + \dots + (n - 1)(n - \omega)(n - \omega^2),$$

where ω is an imaginary cube root of unity, is.....

 [Watch Video Solution](#)

113. Find the value of $\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \frac{1}{3^i 3^j 3^k}$.
($\in e j \neq k$)



Watch Video Solution

114. Find the sum $\sum_{j=1}^{10} \sum_{i=1}^{10} i \times 2^j$



Watch Video Solution

115. Coefficient of $x^{18} \in (1 + x + 2x^2 + 3x^3 + \dots + 18x^{18})^2$ equal to 995
b. 1005 c. 1235 d. none of these



Watch Video Solution

116. Let a_1, a_2, \dots, a_n be real numbers such that

$$\sqrt{a_1} + \sqrt{a_2 - 1} + \sqrt{a_3 - 2} + \dots + \sqrt{a_n - (n - 1)} = \frac{1}{2}(a_1 + a_2 + \dots + a_n)$$

then find the value of $\sum_{i=1}^{100} a_i$



Watch Video Solution

117. A sequence of numbers $A_n = 1, 2, 3$ is defined as follows : $A_1 = \frac{1}{2}$ and for each $n \geq 2$, $A_n = \left(\frac{2n-3}{2n}\right)A_{n-1}$, then prove that

$$\sum_{k=1}^n A_k < 1, n \geq 1$$

 [Watch Video Solution](#)

118. If $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous such that $f(x) - f\left(\frac{x}{2}\right) = \frac{4x^2}{3}$ for all $x \in \mathbb{R}$ and $f(0)=0$, find the value of $f\left(\frac{3}{2}\right)$.

 [Watch Video Solution](#)

119. Find the value of $\frac{\sum_{r=1}^n \frac{1}{r}}{\sum_{r=1}^n \frac{k}{(2n-2k+1)(2n-k+1)}}$.

 [View Text Solution](#)

120. Find the sum $\sum_{n=1}^{\infty} \frac{6^n}{(3^n - 2^n)(3^{n+1} - 2^{n+1})}$

 [Watch Video Solution](#)

Exercise 5.1

1. Write the first five terms of the following sequence and obtain the corresponding series.

$$a_1 = a_2 = 2, a_n = a_{n-1} - 1, n > 2$$

 [Watch Video Solution](#)

2. If $a_{n+1} = \frac{1}{1 - a_n}$ for $n \geq 1$ and $a_3 = a_1$. then find the value of $(a_{2001})^{2001}$.

 [Watch Video Solution](#)

3. Let $\{a_n\}(n \geq 1)$ be a sequence such that $a_1 = 1$, and $3a_{n+1} - 3a_n = 1$ or $a_n \ln \geq 1$. Then find the value of

a_{2002} .



Watch Video Solution

Exercise 5.2

1. If the p th term of an A.P. is q and the q th term is p , then find its r th term.



Watch Video Solution

2. If x is a positive real number different from 1, then prove that the numbers $\frac{1}{1 + \sqrt{x}}$, $\frac{1}{1 - x}$, $\frac{1}{1 - \sqrt{x}}$, ... are in A.P. Also find their common difference.



Watch Video Solution

3. एक समांतर श्रेणी के प्रथम चार पदों का योगफल 56 है | अंतिम चार पदों का योगफल 112 है | यदि इसका प्रथम पद 11 है, तो पदों की संख्या ज्ञात किजिए |



[Watch Video Solution](#)

4. The fourth power of the common difference of an arithmetic progression with integer entries is added to the product of any four consecutive of it. Prove that the resulting sum is the squares of an integer.



[Watch Video Solution](#)

5. Divide 28 into four parts in an A.P. so that the ratio of the product of first and third with the product of second and fourth is 8:15.



[Watch Video Solution](#)

6. If $(b - c)^2, (c - a)^2, (a - b)^2$ are in A.P., then prove that $\frac{1}{b - c}, \frac{1}{c - a}, \frac{1}{a - b}$ are also in A.P.



[Watch Video Solution](#)

7. Find the number of common terms to the two sequences 17,21,25,...,417 and 16,21,26,...,466.



Watch Video Solution

8. If a, b, c, d are distinct integers in A. P. Such that $d = a^2 + b^2 + c^2$, then $a + b + c + d$ is



Watch Video Solution

9. यदि $\frac{a^n + b^n}{a^{n-1} + b^{n-1}}$, a तथा b के मध्य समांतर माध्य हो तो n का मान ज्ञात कीजिए।



Watch Video Solution

10. n arithmetic means are inserted between x and $2y$ and then between $2x$ and y . If the r th means in each case be equal, then find the ratio x/y .



Watch Video Solution

Exercise 5.3

1. If $S_n = nP + \frac{n(n-1)}{2}Q$, where S_n denotes the sum of the first n terms of an A.P., then find the common difference.

 Watch Video Solution

2. Solve the equation
 $(x + 1) + (x + 4) + (x + 7) + \dots + (x + 28) = 155$.

 Watch Video Solution

3. If the sum of the first ten terms of an A.P. is four times the sum of its first five terms, the ratio of the first term to the common difference is:

 Watch Video Solution

4. If the sum of n , $2n$, $3n$ terms of an AP are S_1 , S_2 , S_3 respectively. Prove that $S_3 = 3(S_2 - S_1)$

 [Watch Video Solution](#)

5. Let S_n denote the sum of first n terms of an A.P. If $S_{2n} = 3S_n$, then find the ratio S_{3n}/S_n .

 [Watch Video Solution](#)

6. The ratio of the sum of m and n terms of an A.P. is $m^2 : n^2$. Show that the ratio of the m th and n th terms is $(2m - 1) : (2n - 1)$.

 [Watch Video Solution](#)

7. Find the sum to n terms of the series $1^2 + 2^2 + 3^2 - 4^2 + 5^2 - 6^2 + \dots$





[Watch Video Solution](#)

8. The interior angles of a polygon are in arithmetic progression. The smallest angle is 120° and the common difference is 5° . Find the number of sides of the polygon.



[Watch Video Solution](#)

9. 150 workers were engaged to finish a piece of work in a certain number of days. Four workers dropped the second day, four more workers dropped the third day and so on. It takes 8 more days to finish the work now. Find the number of days in which the work was completed.



[Watch Video Solution](#)

Exercise 5.4

1. The first and second term of a G.P. are x^{-4} and x^n respectively. If x^{52} is the 8^{th} term, then find the value of n .

 [Watch Video Solution](#)

2. If $a, b, \text{ and } c$ are respectively, the p th, q th, and r th terms of a G.P., show that $(q - r)\log a + (r - p)\log b + (p - q)\log c = 0$.

 [Watch Video Solution](#)

3. If $p, q, \text{ and } r$ are in A.P., show that the p th, q th, and r th terms of any G.P. are in G.P.

 [Watch Video Solution](#)

4. यदि a, b, c, d गुणोत्तर श्रेणी में है, तो सिद्ध किजिए कि $(a^n + b^n), (b^n + c^n), (c^n + d^n)$ गुणोत्तर श्रेणी में है।

[Watch Video Solution](#)

5. Let T_r denote the r th term of a G.P. for $r = 1, 2, 3$, If for some positive integers m and n , we have $T_m = 1/n^2$ and $T_n = 1/m^2$, then find the value of $T_{m+n/2}$.

[Watch Video Solution](#)

6. If a, b, c , and d are in G.P., show that $(ab + bc + cd)^2 = (a^2 + b^2 + c^2)(b^2 + c^2 + d^2)$.

[Watch Video Solution](#)

7. The sum of three numbers in GP. Is 56. If we subtract 1, 7, 21 from these numbers in that order, we obtain an arithmetic progression. Find the numbers.

[Watch Video Solution](#)

8. If x , y , and z are p th, q th, and r th terms, respectively, of an A.P. and also of a G.P., then $x^{y-z}y^{z-x}z^{x-y}$ is equal to xyz b. 0 c. 1 d. none of these

 [Watch Video Solution](#)

9. The product of the three numbers in G.P. is 125 and sum of their product taken in pairs is $\frac{175}{2}$. Find them.

 [Watch Video Solution](#)

10. Find the product of three geometric means between 4 and $\frac{1}{4}$.

 [Watch Video Solution](#)

11. Find two numbers whose arithmetic mean is 34 and the geometric mean is 16.

 [Watch Video Solution](#)

12. If the arithmetic means of two positive number a and b ($a > b$) is twice their geometric mean, then find the ratio $a: b$

 [Watch Video Solution](#)

13. Let $a_1, a_2, a_3 \dots$ and $b_1, b_2, b_3 \dots$ be two geometric progressions with $a_1 = 2\sqrt{3}$ and $b_1 = \frac{52}{9}\sqrt{3}$ If $3a_{99}b_{99} = 104$ then find the value of $a_1b_1 + a_2b_2 + \dots + a_nb_n$

 [Watch Video Solution](#)

Exercise 5.5

1. किसी गुणोत्तर श्रेणी के पदों की संख्या सम है | यदि उसके सभी पदों का योगफल, विषम स्थान पर रखे पदों के योगफल का 5 गुना है, तो सार्व अनुपात ज्ञात किजिए |

 [Watch Video Solution](#)

2. If the sum of n terms of a G.P. is $3\frac{3^{n+1}}{4^{2n}}$, then find the common ratio.

 [Watch Video Solution](#)

3. $(\underbrace{666\dots6}_{n\text{-digits}})^2 + (\underbrace{888\dots8}_{n\text{-digits}})$ is equal to

 [Watch Video Solution](#)

4. Find the sum of n terms of series
 $(x + y) + (x^2 + xy + y^2) + (x^3 + x^2y + xy^2 + y^3) + \dots$

 [Watch Video Solution](#)

5. Find the sum of n terms of the series $4/3 + 10/9 + 28/27 + \dots$

 [Watch Video Solution](#)

6. If $p(x) = (1 + x^2 + x^4 + \dots + x^{2n-2}) / (1 + x + x^2 + \dots + x^{n-1})$ is a polynomial in x , then find possible value of n .

 [Watch Video Solution](#)

7.

Let

$$A_n = \left(\frac{3}{4}\right) - \left(\frac{3}{4}\right)^2 + \left(\frac{3}{4}\right)^3 + \dots + (-1)^{n-1} \left(\frac{3}{4}\right)^n \text{ and } B_n = 1 -$$

n_0 , so that $B_n > A_n$ for $n > 0$.

 [Watch Video Solution](#)

8. If the sum of the series $\sum_{n=0}^{\infty} r^n$, $|r| \leq 1$ is s , then find the sum of the series $\sum_{n=0}^{\infty} r^{2n}$, $|r| \leq 1$

 [Watch Video Solution](#)

9. Prove that $6^{1/2} \times 6^{1/4} \times 6^{1/8} \dots = 6$.



Watch Video Solution

10. The sum to n terms of series

$$1 + \left(\frac{1}{2} + \frac{1}{2^2}\right) + 1 + \left(\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4}\right) + \dots$$



Watch Video Solution

Exercise 5.6

1. The 8th and 14th term of a H.P. are $\frac{1}{2}$ and $\frac{1}{3}$, respectively. Find its 20th term. Also, find its general term.



Watch Video Solution

2. If the first two terms of a H.P. are $\frac{2}{5}$ and $\frac{12}{23}$ respectively. Then, largest term is



Watch Video Solution

3. If a, b, c are in G.P. and $a - b, c - a, \text{ and } db - c$ are in H.P., then prove that $a + 4b + c$ is equal to 0.

 [Watch Video Solution](#)

4. If x, y and z are in A.P ax, by and cz in G.P and a, b, c in H.P then prove that

$$\frac{x}{z} + \frac{z}{x} = \frac{a}{c} + \frac{c}{a}$$

 [Watch Video Solution](#)

5. If a, b, c and the d are in H.P then find the vlaue of $\frac{a^{-2} - d^{-2}}{b^{-2} - c^{-2}}$

 [Watch Video Solution](#)

6. If $x = \sum_{n=0}^{\infty} a^n, y = \sum_{n=0}^{\infty} b^n, z = \sum_{n=0}^{\infty} c^n$ where $a, b, \text{ and } c$ are in A.P and $|a| < 1, |b| < 1$ and $|c| < 1$ then prove that x, y and z are in H.P

 [Watch Video Solution](#)

 Watch Video Solution

7. If $x, 1, \text{ and } z$ are in A.P. and $x, 2, \text{ and } z$ are in G.P., then prove that $x, 4, z$ are in H.P.

 Watch Video Solution

8. If $a, a_1, a_2, a_3, a_{2n}, b$ are in A.P. and $a, g_1, g_2, g_3, \dots, g_{2n}, b$ are in G.P. and h is the H.M. of a and b , then prove that

$$\frac{a_1 + a_{2n}}{g_1 g_{2n}} + \frac{a_2 + a_{2n-1}}{g_1 g_{2n-1}} + \dots + \frac{a_n + a_{n+1}}{g_n g_{n+1}} = \frac{2n}{h}$$

 Watch Video Solution

9. If the sum of the roots of the quadratic equation $ax^2 + bx + c = 0$ is equal to the sum of the squares of their reciprocals, then prove that $\frac{a}{c}, \frac{b}{a}, \text{ and } \frac{c}{b}$ are in H.P.

 Watch Video Solution

10. The A.M. of two given positive numbers is 2. If the larger number is increased by 1, the G.M. of the numbers becomes equal to the A.M. of the given numbers. Then find the H.M.

 [Watch Video Solution](#)

11. The harmonic mean between two numbers is $21/5$, their A.M. ' A ' and G.M. ' G ' satisfy the relation $3A + G^2 = 36$. Then find the sum of square of numbers.

 [Watch Video Solution](#)

Exercise 5.7

1. If $\alpha (\neq 1)$ is a n th root of unity then $S = 1 + 3\alpha + 5\alpha^2 + \dots$ upto n terms is equal to

 [Watch Video Solution](#)

2. Find the sum of n terms of the series $1 + \frac{4}{5} + \frac{7}{5^2} + 10 + 5^3 + \dots$

 [Watch Video Solution](#)

3. Find the sum $\frac{3}{2} - \frac{5}{6} + \frac{7}{18} - \frac{9}{54} + \dots$

 [Watch Video Solution](#)

4. Find the sum $\frac{1^2}{2} + \frac{3^2}{2^2} + \frac{5^2}{2^3} + \frac{7^2}{2^4} + \dots \infty$

 [Watch Video Solution](#)

Exercise 5.8

1. Find the sum to n terms of the series :

$$1 \times 2 \times 3 + 2 \times 3 \times 4 + 3 \times 4 \times 5 + \dots$$


[Watch Video Solution](#)

[Watch Video Solution](#)

2. Find the sum of the series $1^2 + 3^2 + 5^2 + \dots \rightarrow n$ terms.

A. $\frac{n(2n - 1)(2n + 1)}{3}$

B. $\frac{n(2n + 1)(2n + 1)}{3}$

C. $\frac{n(2n - 1)(2n - 1)}{3}$

D. $\frac{n(2n + 1)(2n - 1)}{3}$

Answer: A

[Watch Video Solution](#)

3. Find the sum of the series $31^3 + 32^3 + \dots + 50^3$.

[Watch Video Solution](#)

4. Find the sum $1^2 + (1^2 + 2^2) + (1^2 + 2^2 + 3^2) + \dots$ up to 22nd term.



[Watch Video Solution](#)

5. The sum of the first n terms of the series $1^2 + 2.2^2 + 3^2 + 2.4^2 + \dots$ is $\frac{n(n+1)^2}{2}$ when n is even. Then the sum if n is odd, is



[Watch Video Solution](#)

6. Find the sum $11^2 - 1^2 + 12^2 - 2^2 + 13^2 - 3^2 + \dots + 20^2 - 10^2$



[Watch Video Solution](#)

7. Find the sum $3 + 7 + 14 + 24 + 37 + \dots$.20 terms



[Watch Video Solution](#)

8. Find the sum $\sum_{j=1}^n \sum_{i=1}^n I \times 3^j$



[Watch Video Solution](#)

9. If for sequence $\langle a_n \rangle$ sum of n terms $S_n = 2n^2 + 3n$ then find the

$$\text{sum } \sum_{1 \leq i < j \leq 10} a_i a_j$$

 [View Text Solution](#)

10. Find the value of $\sum_{1 \leq i \leq j} i \times \left(\frac{1}{2}\right)^j$

 [View Text Solution](#)

Exercise 5.9

1. Find the sum of infinite series

$$\frac{1}{1 \times 3 \times 5} + \frac{1}{3 \times 5 \times 7} + \frac{1}{5 \times 7 \times 9} + \dots$$

 [Watch Video Solution](#)

2. If $\sum_{r=1}^n T_r = \frac{n}{8}(n+1)(n+2)(n+3)$ then find $\sum_{r=1}^n \frac{1}{T_r}$

 [Watch Video Solution](#)

3. Find the sum $\sum_{r=1}^{\infty} \frac{3n^2 + 1}{(n^2 - 1)^3}$

 [Watch Video Solution](#)

4. Find the sum $\sum_{r=1}^{\infty} \frac{r}{r^4 + \frac{1}{4}}$

 [Watch Video Solution](#)

5. Find the sum

$$\frac{3}{1! + 2! + 3!} + \frac{4}{2! + 3! + 4!} + \dots + \frac{1000}{998! + 999! + 1000!}$$

 [Watch Video Solution](#)

6.

Let

$$S = \frac{\sqrt{1}}{1 + \sqrt{1} + \sqrt{2}} + \frac{\sqrt{2}}{1 + \sqrt{2} + \sqrt{3}} + \frac{\sqrt{3}}{1 + \sqrt{3} + \sqrt{4}} + \dots + \frac{\sqrt{n}}{1 + \sqrt{n} + \sqrt{n+1}}$$

Then find the value of n.



7. Find the sum $\frac{1 \times 2}{3!} + \frac{(2 \times 2)^2}{4!} + \frac{(3 \times 2)^3}{5!} + \dots + \frac{(20 \times 2)^{30}}{22!}$



8. Find the sum $\sum_{r=1}^{\infty} \frac{r-2}{(r+2)(r+3)(r+4)}$



9. Find the sum of the series

$$1 + 2(1-x) + 3(1-x)(1-2x) + \dots + n(1-x)(1-2x)(1-3x)[1 - (n-x)]$$



Exercise (Single)

1. If a, b, c are in A.P., then $a^3 + c^3 - 8b^3$ is equal to

A. $2abc$

B. $3abc$

C. $4abc$

D. $-6abc$

Answer: D



Watch Video Solution

2. If three positive real numbers a, b, c are in AP such that $abc=4$, then the minimum value of b is

A. $2^{1/3}$

B. $2^{2/3}$

C. $2^{1/2}$

D. $2^{3/2}$

Answer: B

 [Watch Video Solution](#)

3. If $\log_2(5 \cdot 2^x + 1)$, $\log_4(2^{1-x} + 1)$ and 1 are in A.P, then x equals

A. $\log_2 5$

B. $1 - \log_5 2$

C. $\log_5 2$

D. $1 - \log_2 5$

Answer: D

 [Watch Video Solution](#)

4. The largest term common to the sequences $1, 11, 21, 31, \dots \rightarrow 100$ terms and $31, 36, 41, 46, \dots \rightarrow 100$ terms is 381 b. 471 c. 281 d. none of these

A. 381

B. 471

C. 281

D. 521

Answer: D



[Watch Video Solution](#)

5. In any A.P. if sum of first six terms is 5 times the sum of next six terms then which term is zero?

A. 10 th

B. 11 th

C. 12 th

D. 13 th

Answer: B



Watch Video Solution

6. If the sides of a right angled triangle are in A.P then the sines of the acute angles are

A. $\frac{3}{5}, \frac{4}{5}$

B. $\frac{1}{\sqrt{3}}, \sqrt{\frac{2}{3}}$

C. $\frac{1}{2}, \frac{\sqrt{3}}{2}$

D. none of these

Answer: A



Watch Video Solution

7. If $a, \frac{1}{b},$ and $\frac{1}{p}, q, \frac{1}{r}$ from two arithmetic progressions of the common difference, then a, q, c are in A.P. if p, b, r are in A.P. b. $\frac{1}{p}, \frac{1}{b}, \frac{1}{r}$ are in A.P. c. p, b, r are in G.P. d. none of these

A. p, b, r are in A.P

B. $\frac{1}{p}, \frac{1}{b}, \frac{1}{r}$ are $\in A. P$

C. p, b, r are in G.P

D. none of these

Answer: B



Watch Video Solution

8. Suppose that $F(n + 1) = \frac{2f(n) + 1}{2}$ for $n = 1, 2, 3, \dots$ and $f(1) = 2$ Then

$F(101)$ equals = ?

A. 50

B. 52

C. 54

D. none of these

Answer: B



[Watch Video Solution](#)

9. Consider an *A. P.* a_1, a_2, a_3, \dots such that $a_3 + a_5 + a_8 = 11$ and $a_4 + a_2 = -2$ then the value of $a_1 + a_6 + a_7$ is.....

A. -8

B. 5

C. 7

D. 9

Answer: C



[Watch Video Solution](#)

10. If a_1, a_2, a_3, \dots are in A.P., then a_p, a_q, a_r are in A.P. if p, q, r are in

A. A.P

B. G.P

C. H.P

D. none of these

Answer: A



[Watch Video Solution](#)

11. Let $\alpha, \beta \in R$. If α, β^2 are the roots of quadratic equation $x^2 - px + 1 = 0$ and α^2, β are the roots of quadratic equation $x^2 - qx + 8 = 0$, then the value of r if $\frac{r}{8}$ is the arithmetic mean of p and q , is $\frac{83}{2}$ b. 83 c. $\frac{83}{8}$ d. $\frac{83}{4}$

A. $\frac{83}{2}$

B. 83

C. $\frac{83}{8}$

D. $\frac{83}{4}$

Answer: B



[Watch Video Solution](#)

12. If the sum of m terms of an A.P. is same as the sum of its n terms, then the sum of its $(m+n)$ terms is

A. mn

B. $-mn$

C. $1/mn$

D. 0

Answer: D



[Watch Video Solution](#)

13. If S_n , denotes the sum of n terms of an $A.P.$, then

$$S_{n+3} - 3S_{n+2} + 3S_{n+1} - S_n =$$

A. $2s_n$

B. S_{n+1}

C. $3S_n$

D. 0

Answer: D



Watch Video Solution

14. The first term of an A.P. is a and the sum of first p terms is zero, show

that the sum of its next q terms is $\frac{a(p+q)q}{p-1}$.

A. $\frac{-a(p+q)p}{q+1}$

B. $\frac{a(q+q)p}{P+1}$

C. $\frac{-a(p+q)q}{p-1}$

D. none of these

Answer: C



[Watch Video Solution](#)

15. If S_n denotes the sum of first n terms of an A.P. and

$\frac{S_{3n} - S_{n-1}}{S_{2n} - S_{2n-1}} = 31$, then the value of n is 21 b. 15 c.16 d. 19

A. 21

B. 15

C. 16

D. 19

Answer: B



[Watch Video Solution](#)

16. The number of terms of an A.P. is even, the sum of odd terms is 24, of the even terms is 3, and the last term exceeds the first by $10\frac{1}{2}$ find the number of terms and the series.

A. 8

B. 4

C. 6

D. 10

Answer: A



Watch Video Solution

17. The number of terms of an A.P. is even, the sum of odd terms is 24, of the even terms is 3, and the last term exceeds the first by $10\frac{1}{2}$ find the number of terms and the series.

A. 8

B. 4

C. 6

D. 10

Answer: D



Watch Video Solution

18. Concentric circles of radii $1, 2, 3, \dots, 100\text{cm}$ are drawn. The interior of the smallest circle is colored red and the angular regions are colored alternately green and red, so that no two adjacent regions are of the same color. Then, the total area of the green regions in sq. cm is equal to 1000π b. 5050π c. 4950π d. 5151π

A. 1000π

B. 5050π

C. 4950π

D. 5151π

Answer: B



Watch Video Solution

19. If $a_1, a_2, a_3, \dots, a_{2n+1}$ are in A.P then

$\frac{a_{2n+1} - a_1}{a_{2n+1} + a_1} + \frac{a_{2n} - a_2}{a_{2n} + a_2} + \dots + \frac{a_{n+2} - a_n}{a_{n+2} + a_n}$ is equal to

A. $\frac{n(n+1)}{2} \times \frac{a_2 - a_1}{a_{n+1}}$

B. $\frac{n(n+1)}{2}$

C. $(n+1)(a_2 - a_1)$

D. none of these

Answer: A



Watch Video Solution

20. If a_1, a_2, \dots, a_n are in A.P. with common difference $d \neq 0$, then the sum of the series $\sin d[\sec a_1 \sec a_2 + \dots + \sec a_{n-1} \sec a_n]$ is

A. $\cos eca_n - \cos eca$

B. $\cot a_n - \cot a$

C. $\sec a_n - \sec a_1$

D. $\tan a_n - \tan a_1$

Answer: D



Watch Video Solution

21. ABC is a right-angled triangle in which $\angle B = 90^\circ$ and $BC = a$. If n points L_1, L_2, \dots, L_n on AB is divided in $n+1$ equal parts and $L_1M_1, L_2M_2, \dots, L_nM_n$ are line segments parallel to BC and M_1, M_2, \dots, M_n are on AC, then the sum of the lengths of $L_1M_1, L_2M_2, \dots, L_nM_n$ is

A. $\frac{a(n+1)}{2}$

B. $\frac{a(n-1)}{2}$

C. $\frac{an}{2}$

D. none of these

Answer: C



View Text Solution

22. If a, b, c, d are in G.P, then $(b - c)^2 + (c - a)^2 + (d - b)^2$ is equal to `

A. $(a - d)^2$

B. $(ad)^2$

C. $(a + d)^2$

D. $(a/d)^2$

Answer: A



Watch Video Solution

23. Let $\{t_n\}$ be a sequence of integers in G.P. in which $t_4:t_6 = 1:4$ and $t_2 + t_5 = 216$. Then t_1 is

A. 12

B. 14

C. 16

D. none of these

Answer: A



[Watch Video Solution](#)

24. if x , $2y$ and $3z$ are in AP where the distinct numbers x , y and z are in gp.

Then the common ratio of the GP is

A. 3

B. $\frac{1}{3}$

C. 2

D. $\frac{1}{2}$

Answer: B



Watch Video Solution

25. If $a, b,$ and c are in A.P and $b-a, c-b$ and a are in G.P then $a:b:c$ is

A. $1:2:3$

B. $1:3:5$

C. $2:3:4$

D. $1:2:4$

Answer: A



Watch Video Solution

26. If the sides of a triangle are in G.P., and its largest angle is twice the smallest, then the common ratio r satisfies the inequality

A. $0 < r < \sqrt{2}$

B. $1 < r < \sqrt{2}$

C. $1 < r < 2$

D. none of these

Answer: B



[Watch Video Solution](#)

27. If x, y, z are in G.P and $a^x = b^y = c^z$, then

A. $\log_b a = \log_a c$

B. $\log_c b = \log_a c$

C. $\log_b a = \log_b c$

D. none of these

Answer: C



Watch Video Solution

28. The number of terms common between the series $1 + 2 + 4 + 8 \dots$ to 100 terms and $1 + 4 + 7 + 10 + \dots$ to 100 terms is

A. 6

B. 4

C. 5

D. none of these

Answer: C



Watch Video Solution

29. If $a^2 + b^2$, $ab + bc$, and $db^2 + c^2$ are in G.P., then a, b, c are in a. A.P. b. G.P. c. H.P. d. none of these

A. A.P.

B. G.P

C. H.P

D. none of these

Answer: B



Watch Video Solution

30. In a G.P. the first, third, and fifth terms may be considered as the first, fourth, and sixteenth terms of an A.P. Then the fourth term of the A.P., knowing that its first term is 5, is 10 b. 12 c. 16 d. 20

A. 10

B. 12

C. 16

D. 20

Answer: D



Watch Video Solution

31. If the p th, q th and r th terms of an AP are in G.P then the common ratio of the GP is

A. $p \frac{r}{q^2}$

B. $\frac{r}{p}$

C. $\frac{q+r}{p+q}$

D. $\frac{q-r}{p-q}$

Answer: D



Watch Video Solution

32. If p th, q th, r th and s th terms of an AP are in GP then show that $(p-q)$, $(q-r)$, $(r-s)$ are also in GP

A. A.P

B. G.P

C. H.P

D. none of these

Answer: B



Watch Video Solution

33. If $a, b, \text{ and } c$ are in G.P. and x, y , respectively, are the arithmetic means between a, b , and b, c , then the value of $\frac{a}{x} + \frac{c}{y}$ is 1 b. 2 c. $1/2$ d. none of these

A. 1

B. 2

C. $1/2$

D. none of these

Answer: B



Watch Video Solution

34. If a, b and c are in A.P., and p and p' are respectively, A.M. and G.M. between a and b while q and q' are, respectively, the A.M. and G.M. between b and c , then $p^2 + q^2 = p'^2 + q'^2$ b. $pq = p'q'$ c. $p^2 - q^2 = p'^2 - q'^2$ d. none of these

A. $p^2 + q^2 = P'^2 + q'^2$

B. $pq = p'q'$

C. $p^2 - q^2 = p'^2 - q'^2$

D. none of these

Answer: C



Watch Video Solution

35. If $(1 + x)(1 + x^2)(1 + x^4) \dots (1 + x^{128}) = \sum_{r=0}^n x^r$, then n is equal is

A. 256

B. 255

C. 254

D. none of these

Answer: B



[Watch Video Solution](#)

36. If $(1 - p)(1 + 3x + 9x^2 + 27x^3 + 81x^4 + 243x^5) = 1 - p^6 p \neq 1$, then the value of $\frac{p}{\xi}$ is $\frac{1}{3}$ b. 3 c. $\frac{1}{2}$ d. 2

A. $\frac{1}{3}$

B. 3

C. $\frac{1}{2}$

D. 2

Answer: B



[Watch Video Solution](#)

37. Consider the ten numbers $ar, ar^2, ar^3, \dots, ar^{10}$. If their sum is 18 and the sum of their reciprocals is 6, then the product of these ten numbers is 81 b. 243 c. 343 d. 324

A. 81

B. 243

C. 343

D. 324

Answer: B



[Watch Video Solution](#)

38. If $x, y, \text{ and } z$ are distinct prime numbers, then $x, y, \text{ and } z$ may be in A.P. but not in G.P. $x, y, \text{ and } z$ may be in G.P. but not in A.P. $x, y, \text{ and } z$ can neither be in A.P. nor in G.P. none of these

A. x, y and z may be in A.P but not in G.P

B. x, y and z may be in G.P but not in A.P

C. x, y and z can neither be in

D. none of these

Answer: A



Watch Video Solution

39.

Let

$$a = 1111(\text{55 digits}), b = 1 + 10 + 10^2 + \dots + 10^4, c = 1 + 10^5 + 10^{10} + 10^{25} + \dots$$

then $a = b + c$ $a = bc$ $b = ac$ $c = ab$

A. $a+b+c$

B. $a=bc$

C. $b=ac$

D. $c=ab$

Answer: B



Watch Video Solution

40. Let a_n be the n th term of a G.P of positive numbers .Let

$\sum_{n=1}^{100} a_{2n} = \alpha$ and $\sum_{n=1}^{100} a_{an-1} = \beta$ then the common ratio is

A. α / β

B. β / α

C. $\sqrt{\alpha / \beta}$

D. $\sqrt{\beta / \alpha}$

Answer: A



Watch Video Solution

41. The sum of 20 terms of a series of which every term is 2 times the term before it ,and every odd term is 3 times the term before it the first term being unity is

A. $\left(\frac{2}{7}\right)(6^{10} - 1)$

B. $\left(\frac{3}{7}\right)(6^{10} - 1)$

C. $\left(\frac{3}{5}\right)(6^{10} - 1)$

D. none of these

Answer: C



Watch Video Solution

42. Let $a \in (0, 1)$ satisfies the equation

$$a^{2008} - 2a + 1 = 0 \text{ values}(s) \rightarrow S \text{ is } 2010 \text{ b. } 2009 \text{ c. } 2008 \text{ d. } 2$$

A. 2010

B. 2009

C. 2008

D. 2

Answer: A



[Watch Video Solution](#)

43. In a geometric series , the first term is a and common ratio is r . If S_n denotes the sum of the n terms and $U_n = \sum_{n=1}^n S_n$, then $rS_n + (1 - r)U_n$ equals

A. 0

B. n

C. na

D. nar

Answer: C



[Watch Video Solution](#)

44. Let $S \subset (0, \pi)$ denote the set of values of x satisfying the equation

$$8^{1 + |\cos x| + \cos^2 x + |\cos^{3x}| \rightarrow \infty} = 4^3 . \text{ Then, } S = \{ \pi/3 \} \text{ b. } \{ \pi/3, 2\pi/3 \} \text{ c.}$$

$$\{ -\pi/3, 2\pi/3 \} \text{ d. } \{ \pi/3, 2\pi/3 \}$$

A. $\{ \pi/3 \}$

B. $\{ \pi/6, 5\pi/6 \}$

C. $\{ \pi/3, 5\pi/6 \}$

D. $\{ \pi/3, 2\pi/3 \}$

Answer: D



Watch Video Solution

45. If $|a| < 1$ and $|b| < 1$ then the sum of the series

$$1 + (1 + a)b + (1 + a + a^2)b^2 + (1 + a + a^2 + a^3)b^3 + \dots \text{ is}$$

A. $\frac{1}{(1 - a)(1 - b)}$

B. $\frac{1}{(1-a)(1-ab)}$

C. $\frac{1}{(1-b)(1-ab)}$

D. $\frac{1}{(1-a)(1-b)(1-ab)}$

Answer: C



Watch Video Solution

46. The value of $0.2^{\log \sqrt{5} \frac{1}{4} + \frac{1}{8} + \frac{1}{16}}$ is 4 b. $\log 4$ c. $\log 2$ d. none of these

A. 4

B. $\log 4$

C. $\log 2$

D. none of these

Answer: A



Watch Video Solution

47.

If

$x = 9^{1/3} \times 9^{1/9} \times 9^{1/27} \times \dots$, $y = 4^{1/3} \times -4^{1/9} \times 4^{1/27} \times \dots$, and

$z = \sum_{r=1}^{\infty} (1+i)^r$ then $\arg(x+yz)$ is equal to

A. 0

B. $\pi - \tan^{-1}\left(\frac{\sqrt{2}}{3}\right)$ C. $-\tan^{-1}\left(\frac{\sqrt{2}}{3}\right)$ D. $-\tan^{-1}\left(\frac{2}{\sqrt{3}}\right)$

Answer: C


[Watch Video Solution](#)

48. The value of x that satisfies the relation

$$x = 1 - x + x^2 - x^3 + x^4 - x^5 + \dots \infty$$

A. $2\cos 36^\circ$ B. $2\cos 144^\circ$

C. $2\sin 18^\circ$

D. $2\cos 18^\circ$

Answer: C



Watch Video Solution

49. If S denotes the sum to infinity and S_n the sum of n terms of the series $1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$, such that $S - S_n < \frac{1}{1000}$ then the least value of n is

A. 8

B. 9

C. 10

D. 11

Answer: D



Watch Video Solution

50. The first term of an infinite geometric series is 21. The second term and the sum of the series are both positive integers. The possible value(s) of the second term can be

A. 12

B. 14

C. 18

D. none of these

Answer: D



[Watch Video Solution](#)

51. The sum of an infinite G.P. is 57 and the sum of their cubes is 9457, find the G.P.

A. $1/3$

B. $2/3$

C. $1/6$

D. none of these

Answer: B



Watch Video Solution

52. If S_p denotes the sum of the series $1 + r^p + r^{2p} + \dots \rightarrow \infty$ and s_p the sum of the series $1 - r^{2p} + r^{3p} + \dots \rightarrow \infty, |r| < 1$, then $S_p + s_p$ in term of S_{2p} is $2S_{2p}$ b. 0 c. $\frac{1}{2}S_{2p}$ d. $-\frac{1}{2}S_{2p}$

A. $2S_{2p}$

B. 0

C. $\frac{1}{2}S_{2p}$

D. $-\frac{1}{2}S_{2p}$

Answer: A



Watch Video Solution

53. If the sum to infinity of the series $1 + 2r + 3r^2 + 4r^3 + \dots$ is $9/4$, then value of r is $1/2$ b. $1/3$ c. $1/4$ d. none of these

A. $1/2$

B. $1/3$

C. $1/4$

D. none of these

Answer: B



Watch Video Solution

54. Find the sum of the series $1 + \frac{4}{5} + \frac{7}{5^2} + \frac{10}{5^3} + \dots$

A. $7/16$

B. $5/16$

C. $105/64$

D. $35/16$

Answer: D



Watch Video Solution

55. The sum of $0.2 + 0.004 + 0.00006 + 0.0000008 + \dots$ to ∞ is

A. $\frac{200}{891}$

B. $\frac{2000}{9801}$

C. $\frac{1000}{9801}$

D. $\frac{2180}{9801}$

Answer: D



Watch Video Solution

56. The positive integer n for which $2 \times 2^2 + 3 \times 2^3 + 4 \times 2^4 + \dots + n \times 2^n = 2^{n+10}$ is 510 b. 511 c. 512 d. 513

A. 510

B. 511

C. 512

D. 513

Answer: D



[Watch Video Solution](#)

57. If ω is a complex n th root of unity, then $\sum_{r=1}^n ar^r + b\omega^{r-1}$ is equal to

A. $(n(n+1))a \frac{1}{a}$

B. $\frac{nb}{1-n}$

C. $\frac{na}{\omega-1}$

D. none of these

Answer: C



Watch Video Solution

58. ABCD is a square of length a , $a \in \mathbb{N}$, $a > 1$. Let L_1, L_2, L_3, \dots be points on BC such that $BL_1L_2 = L_2L_3 = \dots = 1$ and $M_1, M_2M_3 \dots$ be points on CD such that $CM_1 = M_1M_2 = M_2M_3 = \dots = 1$. Then $\sum_{n=1}^{a-1} (AL_n^2 + L_nM_n^2)$ is equal to

A. $\frac{1}{2}a(a-1)^2$

B. $\frac{1}{2}(a-1)(2a-1)(4a-1)$

C. $\frac{1}{2}a(a-1)^2$

D. none of these

Answer: C



Watch Video Solution

59. The 15th term of the series $2\frac{1}{2} + 1\frac{7}{13} + 1\frac{1}{9} + \frac{20}{23} + \dots$ is

A. $\frac{10}{39}$

B. $\frac{10}{21}$

C. $\frac{10}{23}$

D. none of these

Answer: A



[Watch Video Solution](#)

60. If a_1, a_2, \dots, a_n are in H.P then

$$\frac{a_1}{a_2 + a_3 + \dots + a_n}, \frac{a_2}{a_1 + a_3 + \dots + a_n}, \dots, \frac{a_n}{a_1 + a_2 + \dots + a_{n-1}}$$

are in

A. A.P

B. G.P

C. H.P

D. none of these

Answer: C



Watch Video Solution

61. If $a_1, a_2, a_3 \dots a_n$ are in H.P and $f(k) = (\sum_{r=1}^n a_r) - a_k$ then

$\frac{a_1}{f(1)}, \frac{a_2}{f(2)}, \dots, \frac{a_n}{f(n)}$ are in

A. A.P

B. G.P

C. H.P

D. none of these

Answer: C



Watch Video Solution

62. If $a, b, \text{ and } c$ are in A.P. $p, q, \text{ and } r$ are in H.P., and $ap, bq, \text{ and } cr$ are in G.P., then $\frac{p}{r} + \frac{r}{p}$ is equal to $\frac{a}{c} - \frac{c}{a}$ b. $\frac{a}{c} + \frac{c}{a}$ c. $\frac{b}{q} + \frac{q}{b}$ d. $\frac{b}{q} - \frac{q}{b}$

A. A.P

B. G.P

C. G.P

D. none of these

Answer: D



Watch Video Solution

63. If $a, b, \text{ and } c$ are in A.P. $p, q, \text{ and } r$ are in H.P., and $ap, bq, \text{ and } cr$ are in G.P., then $\frac{p}{r} + \frac{r}{p}$ is equal to $\frac{a}{c} - \frac{c}{a}$ b. $\frac{a}{c} + \frac{c}{a}$ c. $\frac{b}{q} + \frac{q}{b}$ d. $\frac{b}{q} - \frac{q}{b}$

A. $\frac{a}{c} - \frac{c}{a}$

B. $\frac{a}{c} + \frac{c}{a}$

C. $\frac{b}{q} + \frac{q}{b}$

D. $\frac{b}{q} - \frac{q}{b}$

Answer: B



Watch Video Solution

64. $a, b, c, d \in R^+$ such that a, b and c are in H.P and ap, bq , and cr are in G.P then $\frac{p}{r} + \frac{t}{p}$ is equal to

A. $ab=cd$

B. $ac=bd$

C. $bc=ad$

D. none of these

Answer: C



View Text Solution

65. If in a progression $a_1, a_2, a_3, \dots, (a_r - a_{r+1})$ bears a constant ratio with $a_r \times a_{r+1}$, then the terms of the progression are in a. A.P b. G.P. c. H.P. d. none of these

A. A.P

B. G.P

C. H.P

D. none of these

Answer: C



[Watch Video Solution](#)

66. If $a, b,$ and c are in G.P then $a+b, 2b$ and $b+c$ are in

A. A.P

B. G.P

C. H.P

D. none of these

Answer: C



[Watch Video Solution](#)

67. If a, x, b are in A.P., a, y, b are in G.P. and a, z, b are in H.P. such that $x=9z$ and $x > 0, b > 0$, then

A. $|y| = 3z$

B. $x = 3|y|$

C. $2y = x + z$

D. none of these

Answer: B



[Watch Video Solution](#)

68. Let $n \in \mathbb{N}$, $n > 25$. Let A, G, H denote the arithmetic mean, geometric mean, and harmonic mean of 25 and n . The least value of n for which $A, G, H \in \{25, 26, n\}$ is a. 49 b. 81 c. 169 d. 225

A. 49

B. 81

C. 169

D. 225

Answer: D



[Watch Video Solution](#)

69. If A.M., G.M., and H.M. of the first and last terms of the series of 100, 101, 102, ..., $n - 1$, n are the terms of the series itself, then the value of n is (100

A. 200

B. 300

C. 400

D. 500

Answer: C



[Watch Video Solution](#)

70. If H_1, H_2, \dots, H_{20} are 20 harmonic means between 2 and 3, then

$$\frac{H_1 + 2}{H_1 - 2} + \frac{H_{20} + 3}{H_{20} - 3} =$$

A. 20

B. 21

C. 40

D. 38

Answer: C



[Watch Video Solution](#)

71. If the sum of n terms of an A.P is $cn(n-1)$ where $c \neq 0$ then the sum of the squares of these terms is

A. $c^2n(n+1)^2$

B. $\frac{2}{3}c^2n(n-1)(2n-1)$

C. $\frac{2c^2}{3}n(n+1)(2n+1)$

D. none of these

Answer: B



Watch Video Solution

72.

If

$$b_i = 1 - a_i, na = \sum_{i=1}^n a_i, nb = \sum_{i=1}^n b_i \text{ then } \sum_{i=1}^n a_i b_i + \sum_{i=1}^n (a_i - a)^2 =$$

A. ab

B. $-nab$

C. $(n + 1)ab$

D. nab

Answer: D



[View Text Solution](#)

73. The sum $1 + 3 + 7 + 15 + 31 + \dots \rightarrow 100$ terms is $2^{100} - 102b$ b.

$2^{99} - 101$ c. $2^{101} - 102$ d. none of these

A. $2^{100} - 102$

B. $2^{99} - 101$

C. $2^{101} - 102$

D. none of these

Answer: C



[Watch Video Solution](#)

74. Consider the sequence 1,2,2,4,4,4,8,8,8,8,8,8,... Then 1025th terms will be 2^9 b. 2^{11} c. 2^{10} d. 2^{12}

A. 2^9

B. 2^{11}

C. 2^{10}

D. 2^{12}

Answer: C



Watch Video Solution

75. The value of $\sum_{i=1}^n \sum_{j=1}^i \sum_{k=1}^j = 220$, then the value of n equals

A. 11

B. 12

C. 10

D. 9

Answer: C



Watch Video Solution

76. If $1^2 + 2^2 + 3^2 + \dots + 2003^2 = (2003)(4007)(334)$ and $(1)(2003) + (2)(2002) + (3)(2001) + \dots + (2003)(1) = (2003)(334)(x)$, the nx equals

A. 2005

B. 2004

C. 2003

D. 2001

Answer: A



Watch Video Solution

77. If t_n denotes the n th term of the series $2+3+6+11+18+\dots$. Then t_{50} is

A. $49^2 - 1$

B. 49^2

C. $50^2 + 1$

D. $49^2 + 2$

Answer: D

 [Watch Video Solution](#)

78. The sum of series $\sum_{r=0}^r (-1)^r (n + 2r)^2$ (where n is even) is

A. $-n^2 + 2n$

B. $-4n^2 + 2n$

C. $-n^2 + 3n$

D. $-n^2 + 4n$

Answer: B

 [Watch Video Solution](#)

79. If $(1^2 - t_1) + (2^2 - t_2) + \dots + (n^2 - t_n) = \frac{n(n^2 - 1)}{3}$ then t_n is

equal to

A. n^2

B. $2n$

C. $n^2 - 2n$

D. none of these

Answer: D



Watch Video Solution

80. If $(1 + 3 + 5 + \dots + p) + (1 + 3 + 5 + \dots + q) = (1 + 3 + 5 + \dots + r)$

where each set of parentheses contains the sum of consecutive odd integers as shown, the smallest possible value of $p + q + r$ (where $p > 6$)

is 12 b. 21 c. 45 d. 54

A. 12

B. 21

C. 45

D. 54

Answer: B



Watch Video Solution

81. If $H_n = 1 + 2 + \dots + \frac{1}{n}$, then the value of $S_n = 1 + \frac{3}{2} + \frac{5}{3} + \dots + \frac{99}{50}$ is $H_{50} + 50$ b. $100 - H_{50}$ c. $49 + H_{50}$ d. $H_{50} + 100$

A. $H_{50} + 50$

B. $100 - H_{50}$

C. $49 + H_{50}$

D. $H_{50} + 100$

Answer: B



Watch Video Solution

82. The sum to 50 terms of the series

$$\frac{3}{1^2} + \frac{5}{1^2 + 2^2} + \frac{7}{1^2 + 2^2 + 3^2} + \dots + \dots is$$

A. $\frac{100}{17}$

B. $\frac{150}{17}$

C. $\frac{200}{51}$

D. $\frac{50}{17}$

Answer: A



Watch Video Solution

83. Let $S = \frac{4}{19} + \frac{44}{(19)^2} + \frac{444}{(19)^3} + \dots \infty$ then find the value of S

A. $40/9$

B. $38/81$

C. $36/171$

D. none of these

Answer: B



Watch Video Solution

84. If $1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} + \dots = \frac{\pi}{4}$, then value of $\frac{1}{1 \times 3} + \frac{1}{5 \times 7} + \frac{1}{9 \times 11} + \dots$ is $\pi/8$ b. $\pi/6$ c. $\pi/4$ d. $\pi/36$

A. $\pi/8$

B. $\pi/6$

C. $\pi/4$

D. $\pi/36$

Answer: A



Watch Video Solution

85. If $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots \rightarrow \infty = \frac{\pi^2}{6}$, then $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$ equals
a. $\pi^2/8$ b. $\pi^2/12$ c. $\pi^2/3$ d. $\pi^2/2$

A. $\pi^2/8$

B. $\pi^2/8$

C. $\pi/3$

D. $\pi^2/2$

Answer: A



Watch Video Solution

86. $\lim_{n \rightarrow \infty} \frac{\sum_{r=1}^n r}{1 \times 3 \times 5 \times 7 \times 9 \times \dots \times (2r+1)}$ is equal to

A. $\frac{1}{3}$

B. $\frac{3}{2}$

C. $\frac{1}{2}$

D. none of these

Answer: C



[Watch Video Solution](#)

87. The greatest interger by which $1 + \sum_{r=1}^{30} r \times r!$ is divisible is

A. composite number

B. odd number

C. divisible by 3

D. none of these

Answer: D



[Watch Video Solution](#)

88. If $\sum_{r=1}^n r^4 = I(n)$, then $\sum_{r=1}^n (r+1)^n (2r-1)^4$ is equal to

A. $I(2n) - I(n)$

B. $I(2n) - 16I(n)$

C. $I(2n) - 8I(n)$

D. $I(2n) - 4I(n)$

Answer: B



Watch Video Solution

89. Value of $\left(1 + \frac{1}{3}\right)\left(1 + \frac{1}{3^2}\right)\left(1 + \frac{1}{3^4}\right)\left(1 + \frac{1}{3^8}\right)\dots$ is equal to 3 b.

$\frac{6}{5}$ c. $\frac{3}{2}$ d. none of these

A. 3

B. $\frac{6}{5}$

C. $\frac{3}{2}$

D. none of these

Answer: C



[Watch Video Solution](#)

90. If x_1, x_2, \dots, x_{20} are in H.P and $x_1, 2, x_{20}$ are in G.P then $\sum_{r=1}^{19} x_r r_{x+1}$

A. 76

B. 80

C. 84

D. none of these

Answer: A



[Watch Video Solution](#)

91. The value of $\sum_{r=1}^n (a + r + ar)(-a)^r$ is equal to

A. $(-1)^n [n + 1)a^{n+1} - a]$

B. $(-1)^n (n + 1)a^{n+1}$

C. $(-1)^n \frac{(n + 2)a^{n+1}}{2}$

D. $(-1)^n \frac{na^n}{2}$

Answer: B



Watch Video Solution

92. The sum of series $\frac{x}{1-x^2} + \frac{x^2}{1-x^4} + \frac{x^4}{1-x^8} + \dots$ to infinite terms, if $|x| < 1$, is $\frac{x}{1-x}$ b. $\frac{1}{1-x}$ c. $\frac{1+x}{1-x}$ d. 1

A. $\frac{x}{1-x}$

B. $\frac{1}{1-x}$

C. $\frac{1+x}{1-x}$

D. 1

Answer: A



Watch Video Solution

93. The sum of 20 terms of the series whose r th term is given by

$$T(n) = (-1)^n \frac{n^2 + n + 1}{n!} \text{ is } \frac{20}{19!} \text{ b. } \frac{21}{20!} - 1 \text{ c. } \frac{21}{20!} \text{ d. none of these}$$

A. $\frac{20}{19!}$

B. $\frac{21}{20!} - 1$

C. $\frac{21}{20!}$

D. none of these

Answer: B



Watch Video Solution

Exercise (Multiple & Comprehension)

1. For an increasing A.P. a_1, a_2, a_n if $a_1 = a_2 + a_3 + a_5 = -12$ and $a_1 a_3 a_5 = 80$, then which of the following is/are true? $a_1 = -10$ b.

$$a_2 = -1 \quad c. a_3 = -4 \quad d. a_5 = +2$$

A. $a_1 = -10$

B. $a_2 = -1$

C. $a_3 = -4$

D. $a_5 = +2$

Answer: A::C::D



Watch Video Solution

2. If the sum of n terms of an A.P. is given by $S_n = a + bn + cn^2$, where a, b, c are independent of n , then $a = 0$
common difference of A.P. must be $2b$
common difference of A.P. must be $2c$
first term of A.P. is $b + c$

A. $a=0$

B. common difference of A.P must be $2b$

C. common difference of A.P must $2c$

D. first term of A.P is $b+c$

Answer: A::C::D



Watch Video Solution

3. If a, b, c and d are four unequal positive numbers which are in A.P then

A. $\frac{1}{a} + \frac{1}{d} > \frac{1}{b} + \frac{1}{c}$

B. $\frac{1}{a} + \frac{1}{d} < \frac{1}{b} + \frac{1}{c}$

C. $\frac{1}{b} + \frac{1}{c} > \frac{4}{a+d}$

D. $\frac{1}{a} + \frac{1}{d} = \frac{1}{b} + \frac{1}{c}$

Answer: A::C



Watch Video Solution

4. Which of the following can be terms (not necessarily consecutive) of any A.P.? a. 1,6,19 b. $\sqrt{2}$, $\sqrt{50}$, $\sqrt{98}$ c. $\log 2$, $\log 16$, $\log 128$ d. $\sqrt{2}$, $\sqrt{3}$, $\sqrt{7}$

A. 1,6,19

B. $\sqrt{2}$, $\sqrt{50}$, $\sqrt{98}$

C. $\log 2$, $\log 16$, $\log 128$

D. $\sqrt{2}$, $\sqrt{3}$, $\sqrt{7}$

Answer: A::B::C



[Watch Video Solution](#)

5. In a arithmetic progression whose first term is α and common difference is β , $\alpha, \beta \neq 0$ the ratio r of the sum of the first n terms to the sum of n terms succeeding them, does not depend on n . Then which of the following is /are correct ?

A. $\alpha : \beta = 2 : 1$

B. If α and β are roots of the equation $ax^2 + bx + c = 0$ then

$$2b^2 = 9ac$$

C. The sum of infinite G.P. $1 + r + r^2 + \dots$ is $3/2$

D. If $\alpha = 1$, then sum of 10 terms of A.P is 100

Answer: B::C::D

 [Watch Video Solution](#)

6. If $a^2 + 2bc, b^2 + 2ca, c^2 + 2ab$ are in A.P. then :-

A. $(a - b)(c - a), (a - b)(b - c), (b - c)(c - a)$ are in A.P

B. $b-c, c-a, a-b$ are in H.P

C. $a+b, b+c, c+a$ are in H.P

D. a^2, b^2, c^2 are in H.P

Answer: A::B

 [Watch Video Solution](#)

7. If sum of an infinite G.P. $p, 1, 1/p, 1/p^2, \dots = 9/2$. Is then value of p is

A. 2

B. $3/2$

C. 3

D. $9/2$

Answer: B::C



Watch Video Solution

8. The terms of an infinitely decreasing decreasing G.P having common ratio r in which all the terms are positive, the first term is 4, and the difference between the third and fifth terms is $32/81$ then

A. $r = 1/3$

B. $r = 2\sqrt{2}/3$

C. Sum of infinite terms is 6

D. none of these

Answer: A::B::C



Watch Video Solution

9. Let $a_1, a_2, a_3, \dots, a_n$ be in G.P such that $3a_1 + 7a_2 + 3a_3 - 4a_5 = 0$

Then common ratio of G.P can be

A. 2

B. $\frac{3}{2}$

C. $\frac{5}{2}$

D. $-\frac{1}{2}$

Answer: B::D



Watch Video Solution

10. If $p(x) = \frac{1 + x^2 + x^4 + \dots + x^{2n-2}}{1 + x + x^2 + \dots + x^{n-1}}$ is a polynomial in x , the number of factors can be 5 b. 10 c. 20 d. 17

A. 5

B. 10

C. 20

D. 17

Answer: A::D



Watch Video Solution

11. If $n > 1$, the value of the positive integer m for which $n^m + 1$ divides $a = 1 + n + n^2 + \dots + n^{63}$ is/are 8 b. 16 c. 32 d. 64

A. 8

B. 16

C. 32

D. 64

Answer: A::B::C



Watch Video Solution

12. The next term of the G.P. $x, x^2 + 2$, and $x^3 + 10$ is $\frac{729}{16}$ b. 6 c. 0 d. 54

A. $\frac{729}{16}$

B. 6

C. 0

D. 54

Answer: A::D



Watch Video Solution

13. If $1 + 2x + 3x^2 + 4x^3 + \dots \infty \geq 4$ then

- A. least value of x is $1/2$
- B. greatest value of x is $4/3$
- C. least value of x is $2/3$
- D. greatest value of x does not exist

Answer: A:D

 [Watch Video Solution](#)

14. Let S_1, S_2, \dots be squares such that for each $n \geq 1$, the length of a side of S_n equals the length of a diagonal of S_{n+1} . If the length of a side of S_1 is 10cm , then for which of the following value of n is the area of S_n less than 1 sq. cm ? a. 5 b. 7 c. 9 d. 10

- A. 7
- B. 8
- C. 9
- D. 10

Answer: B::C::D



Watch Video Solution

15. If a, b and c are in G.P and x and y , respectively, be arithmetic means between a, b and b, c then

A. $\frac{a}{x} + \frac{c}{y} = 2$

B. $\frac{a}{x} + \frac{c}{y} = \frac{c}{a}$

C. $\frac{1}{x} + \frac{1}{y} = \frac{2}{b}$

D. $\frac{1}{x} + \frac{1}{y} = \frac{2}{a}$

Answer: A::C



Watch Video Solution

16. Consider a sequence $\{a_n\}$ with $a_1=2$ & $a_n = \frac{a_{n-1}^2}{a_{n-2}}$ for all $n \geq 3$ terms of the sequence being distinct .Given that a_2 and a_5 are positive

integers and $a_5 \leq 162$, then the possible values (s) of a_5 can be

A. 162

B. 64

C. 32

D. 2

Answer: A::C



[Watch Video Solution](#)

17. The numbers 1, 4, 16 can be three terms (not necessarily consecutive) of no A.P. only on G.P. infinite number o A.P.s infinite number of G.P.s

A. no. A.P

B. only one G.P

C. infinite number of A.P's

D. infinite nuber of G.P' s

Answer: C::D



Watch Video Solution

18. The sum of an infinite geometric series is 162 and the sum of its first n terms is 160. If the inverse of its common ratio is an integer, then which of the following is not a possible first term? 108 b. 144 c. 160 d. none of these

A. 108

B. 120

C. 144

D. 160

Answer: A::C::D



Watch Video Solution

19. If $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are in A.P and $a, b, -2c$, are in G.P where a, b, c are non-zero then

A. $a^3 + b^3 + c^3 = 3abc$

B. $-2a, b, -2c$ are in A.P

C. $a^2, b^2, 4c^2$ are in G.P

D.

Answer: A::B::C::D



Watch Video Solution

20. Sum of an infinite G.P is 2 and sum of its two terms is 1.If its second terms is negative then which of the following is /are true ?

A. one of the possible values of the first terms is $(2 - \sqrt{2})$

B. one of the possible values of the first terms is $(2 + \sqrt{2})$

C. one of the possible values of the common ratio is $(\sqrt{2} - 1)$

D. one of the possible values of the common ratio is $\frac{1}{\sqrt{2}}$

Answer: A::B::D



Watch Video Solution

21. For $0 < \phi < \pi/2$, if $x = \sum_{n=0}^{\infty} \cos^{2n} \phi$, $y = \sum_{n=0}^{\infty} \sin^{2n} \phi$ and $z = \sum_{n=0}^{\infty} \cos^{2n} \phi$ then

A. $xyz = xz + y$

B. $xyz = xy + z$

C. $xyz = z + y + z$

D. $xyz = yz + x$

Answer: B::C



Watch Video Solution

22. For the series,

$$S = 1 + \frac{1}{1+3}(1+2)^2 + \frac{1}{(1+3+5)}(1+2+3)^2 + \frac{1}{(1+3+5+7)}(1$$

A. 7th term is 16

B. 7th term is 18

C. Sum of first 10 terms is $\frac{505}{4}$

D. Sum of first 10 terms is $\frac{405}{4}$

Answer: A:C



Watch Video Solution

23. If $\sum_{r=1}^n r(r+1)(2r+3) = an^4 + bn^3 + cn^2 + dn + e$ then

A. $a-b=d-c$

B. $e=0$

C. $a, b - 2/3, c - 1$ are in $\in A. P$

D. $(b + d) / a$ is an integer

Answer: A::B::C::D



Watch Video Solution

24. If $S_n = 1^2 - 2^2 + 3^2 - 4^2 + 5^2 - 6^2 + \dots$, then $S_{40} = -820$ b.

$S_{2n} > S_{2n+2}$ c. $S_{51} = 1326$ d. $S_{2n+1} > S_{2n-1}$

A. $S_{40} = -820$

B. $S_{2n} > S_{2n+2}$

C. $S_{51} = 1326$

D. $S_{2n+1} > S_{2n-1}$

Answer: A::B::C::D



Watch Video Solution

25. $\frac{1}{\sqrt{2} + \sqrt{5}} + \frac{1}{\sqrt{5} + \sqrt{8}} + \frac{1}{\sqrt{8} + \sqrt{11}} + n$ terms is equal to $\frac{\sqrt{3n+2} - \sqrt{2}}{3}$ b. $\frac{n}{\sqrt{2+3n} + \sqrt{2}}$ c. less than n d. less than $\sqrt{\frac{n}{3}}$

A. $\frac{(\sqrt{3n+2}) - \sqrt{2}}{3}$

B. $\frac{n}{\sqrt{2+3n} + \sqrt{2}}$

C. less than n

D. less than $\sqrt{\frac{n}{3}}$

Answer: A::B::C



Watch Video Solution

26. Given that $x + y + z = 15$ when a, x, y, z, b are in A.P and $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = \frac{5}{3}$ when a, x, y, z, b are in H.P. Then

A. G.M of a and b is 3

B. one possible value of $a + 2b$ is 11

C. A.M of a and b is 6

D. greatest value of a-b is 8

Answer: A::B::D

 [Watch Video Solution](#)

27. If a, b and c are in H.P., then the value of $\frac{(ac + ab - bc)(ab + bc - ac)}{(abc)^2}$ is

A. $\frac{(a + c)(3a - c)}{4a^2c^2}$

B. $\frac{2}{bc} - \frac{1}{b^2}$

C. $\frac{2}{bc} - \frac{1}{b^2}$

D. $\frac{(a - c)(3a + c)}{4a^2c^2}$

Answer: A::B

 [Watch Video Solution](#)

28. If p, q and r are in A.P then which of the following is / are true ?

- A. p th, q th and r th terms of A.P are in A.P
- B. p th, q th, and r th terms of G.P are in G.P
- C. p th , q th , and r th terms of H.P are in H.P
- D. none of these

Answer: A::B::C



[View Text Solution](#)

29. If $x^2 + 9y^2 + 25z^2 = xyz \left(\frac{15}{2} + \frac{5}{y} + \frac{3}{z} \right)$, then $x, y,$ and z are in

H.P. b. $\frac{1}{x}, \frac{1}{y}, \frac{1}{z}$ are in A.P. c. x, y, z are in G.P. d. $\frac{1}{a} + \frac{1}{d} = \frac{1}{b} = \frac{1}{c}$

A. x, y and z are in H.P

B. $\frac{1}{x}, \frac{1}{y}, \frac{1}{z}$ are in G.P

C. x, y, z are in G.P

D. $\frac{1}{x}, \frac{1}{y}, \frac{1}{z}$ are in G.P

Answer: A::C



Watch Video Solution

30. If $A_1, A_2, G_1, G_2, ;$ and H_1, H_2 are two arithmetic, geometric and harmonic means respectively, between two quantities a and b , then ab is equal to A_1H_2 b. A_2H_1 c. G_1G_2 d. none of these

A. $A_H - 2$

B. A_2H_1

C. G_1G_2

D. none of these

Answer: A::B::C



Watch Video Solution

31. If $\frac{1}{b-a} + \frac{1}{b-c} = \frac{1}{a} + \frac{1}{c}$, then $a, b, \text{ and } c$ are in H.P. $a, b, \text{ and } c$ are in A.P. $b = a + c$ $3a = b + c$

A. $a, b, \text{ and } c$ are in H.P

B. $a, b, \text{ and } c$ are in A.P

C. $b = a + c$

D. $3a = b + c$

Answer: A::B

 [Watch Video Solution](#)

32. If a, b, c are three distinct numbers in G.P., b, c, a are in A.P and a, bc, abc , in H.P then the possible value of b is

A. $3 + 4\sqrt{2}$

B. $3 - 4\sqrt{2}$

C. $4 + 3\sqrt{2}$

$$D. 4 - 3\sqrt{2}$$

Answer: C::D



Watch Video Solution

33. If a, b, c are in A.P and a^2, b^2, c^2 are in H.P then which is of the following is /are possible ?

A. $ax^2 + bx + c = 0$

B. $ax^2bx + c = 0$

C. $a, b - \frac{c}{2}$ form a G.P

D. $a - b, \frac{c}{2}$ from a G.P

Answer: A::C



Watch Video Solution

34. If first and $(2n - 1)^{th}$ terms of A.P., G.P. and H.P. are equal and their n th terms are a, b, c respectively, then

A. $a=b=c$

B. $a \geq be \geq c$

C. $a + b = c$

D. $ac - b^2 = 0$

Answer: B::D



Watch Video Solution

35. Let $E = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots$ Then, $E < 3$ b. $E > 3/2$ c. $E > 2$ d.

$E < 2$

A. $E < 3$

B. $E > 3/2$

C. $E > 2$

D. $E < 2$

Answer: A::B::D



Watch Video Solution

36. Sum of certain consecutive odd positive intergers is $57^2 - 13^2$

The least value of the an interger is

A. 40

B. 37

C. 44

D. 51

Answer: C



Watch Video Solution

37. Sum of certain consecutive odd positive intergers is $57^2 - 13^2$

The least value of the an interger is

A. 22

B. 27

C. 31

D. 43

Answer: B



[Watch Video Solution](#)

38. Sum of certain consecutive odd positive intergers is $57^2 - 13^2$

The least value of the an interger is

A. divisible by 7

B. divisible by 11

C. divisible by 9

D. none of these

Answer: D

 [Watch Video Solution](#)

39. Consider three distinct real numbers a, b, c in a G.P with $a^2 + b^2 + c^2 = t^2$ and $a+b+c = \alpha t$. The sum of the common ratio and its reciprocal is denoted by S .

Complete set of α^2 is

A. $\left(\frac{1}{3}, 3\right)$

B. $\left[\frac{1}{3}, 3\right]$

C. $\left(\frac{1}{3}, 3\right) - \{1\}$

D. $\left(-\infty, \frac{1}{3}\right) \cap (3, \infty)$

Answer: C

 [View Text Solution](#)

40. Consider three distinct real numbers a, b, c in a G.P with $a^2 + b^2 + c^2 = t^2$ and $a+b+c = \alpha t$. The sum of the common ratio and its reciprocal is denoted by S .

Complete set of α^2 is

A. $(-2, 2)$

B. $(-\infty, -2) \cup (2, \infty)$

C. $(-1, 1)$

D. $(-\infty, -1) \cup (1, \infty)$

Answer: B



[View Text Solution](#)

41. If a, b and c also represent the sides of a triangle and a, b, c are in g.p

then the complete set of $\alpha^2 = \frac{r^2 + r + 1}{r^2 - r + 1}$ is

A. $\left(\frac{1}{3}, 3\right)$

B. $(2, 3)$

C. $\left[\frac{1}{3}, 2\right]$

D. $\left(\frac{\sqrt{5+3}}{2}, 3\right)$

Answer: D



Watch Video Solution

42. In a n increasing G.P. , the sum of the first and the last term is 66, the product of the second and the last but one is 128 and the sum of the terms is 126. How many terms are there in the progression?

A. 9

B. 8

C. 12

D. 6

Answer: D



[Watch Video Solution](#)

43. In a n increasing G.P. , the sum of the first and the last term is 66, the product of the second and the last but one is 128 and the sum of the terms is 126. How many terms are there in the progression?

A. 64

B. 128

C. 256

D. 729

Answer: B



[Watch Video Solution](#)

44. In a G.P the sum of the first and last terms is 66, the product of the second and the last but one is 126, and the sum of the terms is 128 In any case, the difference of the least and greatest terms is

A. 78

B. 126

C. 126

D. none of these

Answer: D



[View Text Solution](#)

45. Four different integers form an increasing A.P .One of these numbers is equal to the sum of the squares of the other three numbers. Then

The product of all numbers is

A. -2

B. 1

C. 0

D. 2

Answer: C



Watch Video Solution

46. The sum of all the four numbers is

A. 3

B. 0

C. 4

D. 2

Answer: D



View Text Solution

47. The common difference of the divisible by

A. 1

B. 3

C. 2

D. -2

Answer: A



[View Text Solution](#)

48. Consider the sequence in the form of group $(1),(2,2)(3,3,3),(4,4,4,4),$
 $(5,5,5,5,5.....)$

The 2000^{th} term of the sequence is not divisible by

A. 3

B. 9

C. 7

D. none of these

Answer: D



[View Text Solution](#)

49. Consider the sequence in the form of group $(1),(2,2)(3,3,3),(4,4,4,4),$
 $(5,5,5,5,5.....)$

The sum of first 2000 terms is

- A. 84336
- B. 96324
- C. 78466
- D. none of these

Answer: A



[View Text Solution](#)

50. Consider the sequence in the form of group $(1),(2,2)(3,3,3),(4,4,4,4),$
 $(5,5,5,5,5.....)$

A. 1088

B. 1008

C. 1040

D. none of these

Answer: B



[View Text Solution](#)

51. There are two sets A and B each of which consists of three numbers in A.P. whose sum is 15 and where D and d are the common differences such that $D - d = 1$. If $\frac{p}{q} = \frac{7}{8}$, where p and q are the product of the numbers, respectively, and $d > 0$ in the two sets .

The sum of the products of the numbers in set A taken two at a time is

A. 51

B. 71

C. 74

D. 86

Answer: B



[View Text Solution](#)

52. There are two sets A and B each of which consists of three numbers in A.P. whose sum is 15 and where D and d are the common differences such that $D - d = 1$. If $\frac{p}{q} = \frac{7}{8}$, where p and q are the product of the numbers, respectively, and $d > 0$ in the two sets .

The sum of the product of the numbers in set B taken two at a time is

A. 74

B. 64

C. 73

D. 81

Answer: A



[Watch Video Solution](#)

53. There are two sets M_1 and M_2 each of which consists of three numbers in arithmetic sequence whose sum is 15. Let d_1 and d_2 be the common differences such that $d_1 = 1 + d_2$ and $8p_1 = 7p_2$ where p_1 and p_2 are the product of the numbers respectively in M_1 and M_2 . If $d_2 > 0$ then find the value of $\frac{p_2 - p_1}{d_1 + d_2}$

A. 20

B. 30

C. 15

D. 25

Answer: C



Watch Video Solution

54. Let $A_1, A_2, A_3, \dots, A_m$ be the arithmetic means between -2 and 1027 and $G_1, G_2, G_3, \dots, G_n$ be the geometric means between 1 and

1024 .The product of gerometric means is 2^{45} and sum of arithmetic means is 1024×171

The value of $\sum_{r=1}^n G_r$ is



[View Text Solution](#)

55. Let $A_1, A_2, A_3, \dots, A_m$ be the arithmetic means between -2 and 1027 and $G_1, G_2, G_3, \dots, G_n$ be the gemetric means between 1 and 1024 .The product of gerometric means is 2^{45} and sum of arithmetic means is 1024×171

The n umber of arithmetic means is

A. 442

B. 342

C. 378

D. none of these

Answer: B



[View Text Solution](#)

56. Let $A_1, A_2, A_3, \dots, A_m$ be the arithmetic means between -2 and 1027 and $G_1, G_2, G_3, \dots, G_n$ be the geometric means between 1 and 1024. The product of geometric means is 2^{45} and sum of arithmetic means is 1024×171

The number $2A_{171}, G_5^2 + 1, 2A_{121}$

A. A.P

B. G.P

C. H.P

D. none of these

Answer: A



[View Text Solution](#)

57. Two consecutive numbers from 1, 2, 3, ..., n are removed, then arithmetic mean of the remaining numbers is $\frac{105}{4}$ then $\frac{n}{10}$ must be

equal to

A. [45,55]

B. [52,60]

C. [41,49]

D. none of these

Answer: A



Watch Video Solution

58. Two consecutive numbers from $1, 2, 3, \dots, n$ are removed. The arithmetic mean of the remaining numbers is $105/4$.

The removed numbers

A. lie between 10 and 20

B. are less than 1500

C. are less than 1500

D. none of these

Answer: C



[View Text Solution](#)

59. Two consecutive numbers from $1, 2, 3, \dots, n$ are removed. The arithmetic mean of the remaining numbers is $105\frac{1}{4}$

The sum of all numbers

A. exceeds 1600

B. is less than 1500

C. lies between 1300 and 1500

D. none of these

Answer: B



[Watch Video Solution](#)

60. Two arithmetic progressions have the same numbers. The ratio of the last term of the first progression to the first term of the second progression is equal to the ratio of the last term of the second progression to the first term of first progression is equal to 4. The ratio of the sum of the n terms of the first progression to the sum of the n terms of the first progression to the sum of the n terms of the second progression is equal to 2.

The ratio of their common difference is

A. 12

B. 24

C. 26

D. 9

Answer: C



[View Text Solution](#)

61. Two arithmetic progressions have the same numbers. The ratio of the last term of the first progression to the first term of the second progression is equal to the ratio of the last term of the second progression to the first term of first progression is equal to 4. The ratio of the sum of the n terms of the first progression to the sum of the n terms of the second progression is equal to 2.

A. $6/5$

B. $7/2$

C. $9/5$

D. none of these

Answer: B



View Text Solution

62. Two arithmetic progressions have the same numbers. The ratio of the last term of the first progression to the first term of the second progression is equal to the ratio of the last term of the second progression to the first term of first progression is equal to 4. The ratio of the sum of the n terms of the first progression to the sum of the n terms of the first progression to the sum of the n terms of the second progression is equal to 2.

The ratio of their first term is

A. $2/7$

B. $3/5$

C. $4/7$

D. $2/5$

Answer: A



View Text Solution

63. Find three numbers a, b, c between 2 & 18 such that; O their sum is 25
@ the numbers 2, a, b are consecutive terms of an AP & Q.3 the numbers
 $b, c, 18$ are consecutive terms of a GP

A. 500

B. 450

C. 720

D. 480

Answer: D



Watch Video Solution

64. Find three numbers a, b, c between 2 & 18 such that; O their sum is 25
@ the numbers 2, a, b are consecutive terms of an AP & Q.3 the numbers
 $b, c, 18$ are consecutive terms of a GP

A. real and positive

B. real and negative

C. imaginary

D. real and of opposite sign

Answer: C



[Watch Video Solution](#)

65. If a, b and c are roots of the equation $x^3 + qx^2 + rx + s = 0$

then the value of r is

A. 184

B. 196

C. 224

D. none of these

Answer: B



[View Text Solution](#)

EXERCISE (MULTIPLE CORRECT ANSWER TYPE)

1. In the 20 th row of the triangle



A. last term = 210

B. first term = 191

C. sum = 4010

D. sum =4200

Answer: A::B::C



[View Text Solution](#)

EXERCISE (MATRIX MATCH TYPE)

1. If α and β are roots of the equation $x^2 - 8x + 4 = 0$, then match the following lists :



- A. $a \ b \ c \ d$
(1) $r \ p \ q \ s$
- B. $a \ b \ c \ d$
(2) $q \ s \ p \ r$
- C. $a \ b \ c \ d$
(3) $r \ q \ p \ q$
- D. $a \ b \ c \ d$
(4) $s \ p \ q \ r$

Answer: A::B::C::D



View Text Solution

2. Match the following lists :



- A. $a \ b \ c \ d$
(1) $r \ p \ q \ s$
- B. $a \ b \ c \ d$
(2) $q \ s \ p \ r$

- C. $a \ b \ c \ d$
(3) $r \ q \ p \ q$
- D. $a \ b \ c \ d$
(4) $s \ p \ q \ r$

Answer: A::B::C::D



View Text Solution

3. Match the following lists



- A. $a \ b \ c \ d$
(1) $r \ p \ q \ s$
- B. $a \ b \ c \ d$
(2) $q \ s \ p \ r$
- C. $a \ b \ c \ d$
(3) $r \ q \ p \ q$
- D. $a \ b \ c \ d$
(4) $s \ p \ q \ r$

Answer: C



View Text Solution

Exercise (Numerical)

1. Let a, b, c, d be four distinct real numbers in A.P. Then half of the smallest positive value of k satisfying $a(a - b) + k(b - c)^2 = (c - a)^3 = 2(a - x) + (b - d)^2 + (c - d)^3$ is _____.



[Watch Video Solution](#)

2. Let fourth term of an arithmetic progression be 6 and m^{th} term be 18. If A.P has integral terms only then the numbers of such A.P s is _____.



[Watch Video Solution](#)

3. The 5th and 8th terms of a geometric sequence of real numbers are $7!$ and $8!$ respectively. If the sum to first n terms of the G.P. is 2205, then n equals _____.



Watch Video Solution

4. Let $a_1, a_2, a_3, \dots, a_{101}$ are in G.P with $a_{101} = 25$ and $\sum_{i=1}^{201} a_i = 625$
 Then the value of $\sum_{i=1}^{201} \frac{1}{a_i}$ equals _____.



Watch Video Solution

5. Let $a, b > 0$, let $5a - b, 2a + b, a + 2b$ be in A.P. and
 $(b + 1)^2, ab + 1, (a - 1)^2$ are in G.P., then the value of $(a^{-1} + b^{-1})$ is
 _____.



Watch Video Solution

6. Let $a + ar_1 + ar_1^2 + \dots + \infty$ and $a + ar_2 + ar_2^2 + \dots + \infty$ be two
 infinite series of positive numbers with the same first term. The sum of
 the first series is r_1 and the sum of the second series r_2 . Then the value
 of $(r_1 + r_2)$ is _____.



Watch Video Solution

 Watch Video Solution

7. If the equation $x^3 + ax^2 + bx + 216 = 0$ has three real roots in G.P., then b/a has the value equal to _____.

 Watch Video Solution

8. Let $a_n = 16, 4, 1, \dots$ be a geometric sequence. Define P_n as the product of the first n terms. The value of $\sum_{n=1}^{\infty} n\sqrt{P_n}$ is _____.

 Watch Video Solution

9. The terms a_1, a_2, a_3 from an arithmetic sequence whose sum is 18. The terms $a_1 + 1, a_2, a_3 + 2$, in that order, form a geometric sequence. Then the absolute value of the sum of all possible common differences of the A.P. is _____.

 Watch Video Solution

10. Let the sum of first three terms of G.P. with real terms be $\frac{13}{12}$ and their product is -1 . If the absolute value of the sum of their infinite terms is S , then the value of $7S$ is _____.



[Watch Video Solution](#)

11. The first term of an arithmetic progression is 1 and the sum of the first nine terms equal to 369. The first and the ninth term of a geometric progression coincide with the first and the ninth term of the arithmetic progression. Find the seventh term of the geometric progression.



[Watch Video Solution](#)

12. A person drops a ball from an 80 m tall building and each time the ball bounces, it rebounds to $p\%$ of its previous height. If the ball travels a total distance of 320 m, then the value of p is



[Watch Video Solution](#)

13. The digits in units's place of number $\frac{10^{2013} - 1}{10^{33} - 1}$ is. _____.

 [Watch Video Solution](#)

14. The number of positive integral ordered pairs of (a, b) such that $6, a, b$ are in harmonic progression is _____.

 [Watch Video Solution](#)

15. If the roots of $10x^3 - nx^2 - 54x - 27 = 0$ are in harmonic oprogresion, then n eqauls _____.

 [Watch Video Solution](#)

16. Given a, b, c are in A.P., b, c, d are in G.P and c, d, e are in H.P .If $a=2$ and $e=18$, then the sum of all possible value of c is _____.

 [Watch Video Solution](#)

17. Let S_k be sum of an indinite G.P whose first term is 'K' and common ratio is $\frac{1}{k+1}$. Then $\sum_{k=1}^{10} S_k$ is equal to _____.

 [Watch Video Solution](#)

18. The value of the sum $\sum_{i=1}^{20} i \left(\frac{1}{i} + \frac{1}{i+1} + \frac{1}{i+2} + \dots + \frac{1}{2} \right)$ is _____.

 [Watch Video Solution](#)

19. The difference between the sum of the first k terms of the series $1^3 + 2^3 + 3^3 + \dots + n^3$ and the sum of the first k terms of $1 + 2 + 3 + \dots + n$ is 1980 . The value of k is :

 [Watch Video Solution](#)

20. The value of the $\sum_{n=0}^{\infty} \frac{2n+3}{3^n}$ is equal to _____.



[Watch Video Solution](#)

21. The sum of the infinite Arithmetic-Geometric progression $3, 4, 4, \dots$ is _____.



[Watch Video Solution](#)

22. $\sum_{r=1}^{50} \frac{r^2}{r^2 + (11-r)^2}$ is equal to _____.



[Watch Video Solution](#)

23. If $\sum_{r=1}^{50} \frac{2}{r^2 + (11-r)^2}$, then the value of n is _____



[Watch Video Solution](#)

24. Let $\langle a_n \rangle$ be an arithmetic sequence of 99 terms such that sum of its odd numbered terms is 1000 then the value of

$$\sum_{r=1}^{50} (-1)^{\frac{r(r+1)}{2}} \cdot a_{2r-1} \text{ is } \underline{\hspace{2cm}}.$$



Watch Video Solution

25. Find the sum of series upto n terms

$$\left(\frac{2n+1}{2n-1}\right) + 3\left(\frac{2n+1}{2n-1}\right)^2 + 5\left(\frac{2n+1}{2n-1}\right)^3 + \dots$$



Watch Video Solution

26. Let $S = \sum_{n=1}^{999} \frac{1}{(\sqrt{n} + \sqrt{n+1})(4\sqrt{n} + 4\sqrt{n+1})}$, then S equals _____.



Watch Video Solution

27. Let S denote sum of the series $\frac{3}{2^3} + \frac{4}{2^4 \cdot 3} + \frac{5}{2^6 \cdot 3} + \frac{6}{2^7 \cdot 5} + \dots$ Then the value of S^{-1} is _____.

 [Watch Video Solution](#)

28. The sum $\frac{7}{2^2 \times 5^2} + \frac{13}{5^2 \times 8^2} + \frac{19}{8^2 \times 11^2} + \dots$ 10 terms is S , then the value of $1024(S)$ is _____.

 [Watch Video Solution](#)

JEE Main Previous Year

1. The sum to infinity of the series $1 + \frac{2}{3} + \frac{6}{3^2} + \frac{10}{3^3} + \frac{14}{3^4} + \dots$ is (1)

2 (2) 3 (3) 4 (4) 6

A. 2

B. 3

C. 4

D. 6

Answer: B



[Watch Video Solution](#)

2. A person is to count 4500 currency notes. Let a_n , denote the number of notes he counts in the n th minute if $a_1 = a_2 = a_3 = \dots = a_{10} = 150$ and a_{10}, a_{11}, \dots are in an AP with common difference -2 , then the time taken by him to count all notes is :- (1) 24 minutes 10 11 (2) 34 minutes (3) 125 minutes (4) 135 minutes

A. 135 min

B. 24 min

C. 34 min

D. 125 min

Answer: C



[Watch Video Solution](#)

3. A man saves Rs. 200 in each of the first three months of his service. In each of the subsequent months In each of their mupienent montha his saving increases by Rs, 40 more than the saving of immediately previous month. His total saving s from the start of service will be Rs. 11040 after

A. 21 months

B. 18 months

C. 19 months

D. 20 months

Answer: A



[Watch Video Solution](#)

4. Statement 1 :

The sum of the series $1+(1+2+4)+(4+6+9)+(9+12+16)+\dots+(361+380+400)$ is 8000

Statement 1:

$$\sum_{k=1}^n (k^3 - (k-1)^3) = n^3, \text{ for any natural number } n.$$

- A. Statement 1 is false ,statement 2 is true
- B. Statement 1 is true ,statement 2 is true , statement 2 is a correct explanation for statement 1.
- C. Statement 1 is true, statements 2 is true statement 2 is not a correct explanation for statement 1
- D. Statement 1 is true, statement 2 is false

Answer: B

 [Watch Video Solution](#)

5. If 100 times the 100^{th} term of an AP with non zero common difference equals the 50 times its 50^{th} term, then the 150^{th} term of this AP is (1) 150 (2) 150 times its 50^{th} term (3) 150 (4) zero

A. – 150

B. 150 times its 50 th term

C. 150

D. Zero

Answer: D



Watch Video Solution

6. The sum of first 20 terms of the sequence $0.7, 0.77, 0.777, ..$, is (1)

$$\frac{7}{9}(99 - 10^{-20}) \quad (2) \quad \frac{7}{81}(179 + 10^{-20}) \quad (3) \quad \frac{7}{9}(99 + 10^{-20}) \quad (3)$$

$$\frac{7}{81}(179 - 10^{-20})$$

A. $\frac{7}{81}(179 - 10)^{20}$

B. $\frac{7}{9}(99 - 10^{20})$

C. $\frac{7}{81}(179 + 10^{-20})$

D. $\frac{7}{9}(99 + 10^{-20})$

Answer: C



Watch Video Solution

7. If $(10)^9 + 2(11)^1(10)^8 + 3(11)^2(10)^7 + \dots + 10(11)^9 = k(10)^9$,

then k is equal to :

A. $\frac{121}{10}$

B. $\frac{441}{100}$

C. 100

D. 110

Answer: C



Watch Video Solution

8. If m is the A.M. of two distinct real numbers l and n ($l, n > 1$) and G_1, G_2 and G_3 are three geometric means between l and n , then $G_1^4 + 2G_2^4 + G_3^4$ equals, (1) $4l^2 mn$ (2) $4l^m \wedge 2 mn$ (3) $4lmn^2$ (4) $4l^2 m^2 n^2$

A. $4l^2 mn$

B. $4lm^2 n$

C. $4lmn^2$

D. $4l^2 m^n \wedge 2$

Answer: B



Watch Video Solution

9. The sum of the first 9 terms of the series

$$\frac{1^3}{1} + \frac{1^3 + 2^3}{1 + 3} + \frac{1^3 + 2^3 + 3^3}{1 + 3 + 5} \dots \text{ is :}$$

A. 71

B. 96

C. 142

D. 192

Answer: B



Watch Video Solution

10. If the 2nd , 5th and 9th terms of a non-constant A.P. are in G.P., then

the common ratio of this G.P. is : (1) $\frac{8}{5}$ (2) $\frac{4}{3}$ (3) 1 (4) $\frac{7}{4}$

A. $\frac{4}{3}$

B. 1

C. $\frac{7}{4}$

D. $\frac{8}{5}$

Answer: A

 [Watch Video Solution](#)

11. If the sum of the first ten terms of the series,
 $\left(1\frac{3}{5}\right)^2 + \left(2\frac{2}{5}\right)^2 + \left(3\frac{1}{5}\right)^2 + 4^2 + \left(4\frac{4}{5}\right)^2 + \dots$, is $\frac{16}{5}m$, then m
is equal to

A. 101

B. 100

C. 99

D. 102

Answer: A

 [Watch Video Solution](#)

12. If, for a positive integer n , the quadratic equation,
 $x(x+1) + (x-1)(x+2) + \dots + (x+n-1)(x+n) = 10n$ has two
consecutive integral solutions, then n is equal to : 10 (2) 11 (3) 12 (4) 9

A. 11

B. 12

C. 9

D. 10

Answer: A

 [Watch Video Solution](#)

JEE Advanced Previous Year

1. For any three positive real numbers a , b and c , $9(25a^2 + b^2) + 25(c^2 - 3ac) = 15b(3a + c)$ Then: (1) b , c and a are in G.P. (2) b , c and a are in A.P. (3) a , b and c are in A.P. (4) a , b and c are in G.P.

A. a, b and c are in G.P.

B. b, c and a are in G.P.

C. b, c and a are in A.P.

D. a,b and c are in A.P

Answer: C



Watch Video Solution

2. Let $a, b, c \in R$. If $f(x) = ax^2 + bx + c$ is such that $a + b + c = 3$ and $f(x + y) = f(x) + f(y) + xy, \forall x, y \in R$, then $\sum_{n=1}^{10} f(n)$ is equal to

A. 255

B. 330

C. 165

D. 190

Answer: B



Watch Video Solution

3. Let A be the sum of the first 20 terms and B be the sum of the first 40 terms of the series $1^2 + 2.2^2 + 3^2 + 2.4^2 + 5^2 + 2.6^2 + \dots$. If $B - 2A = 100\lambda$ then λ is equal to (1) 232 (2) 248 (3) 464 (4) 496

A. 496

B. 232

C. 248

D. 464

Answer: C



[Watch Video Solution](#)

4. Let $a_1, a_2, a_3, \dots, a_{49}$ be in A.P. Such that $\sum_{k=0}^{12} a_{4k+1} = 416$ and $a_9 + a_{43} = 66$. If $a_1^2 + a_2^2 + \dots + a_{17} = 140m$ then m is equal to

A. 33

B. 66

C. 68

D. 34

Answer: D



[Watch Video Solution](#)

5. Let a_1, a_2, a_3, \dots be a harmonic progression with $a_1 = 5$ and $a_{20} = 25$. The least positive integer n for which $a_n < 0$, is

A. 22

B. 23

C. 24

D. 25

Answer: D



[Watch Video Solution](#)

6. The value of $\sum_{k=1}^{13} \frac{1}{\sin\left(\frac{\pi}{4} + \frac{(k-1)\pi}{6}\right)\sin\left(\frac{\pi}{4} + \frac{k\pi}{6}\right)}$ is equal to

A. $3 - \sqrt{3}$

B. $2(3 - \sqrt{3})$

C. $2(3 + \sqrt{3})$

D. $2(\sqrt{3} - 1)$

Answer: C



Watch Video Solution

7. Let $b_i > 1$ for $i = 1, 2, \dots, 101$. Suppose $\log_e b_1, \log_e b_2, \dots, \log_e b_{101}$ are in Arithmetic Progression (A.P.) with the common difference $\log_e 2$. Suppose a_1, a_2, \dots, a_{101} are in A.P. such that $a_1 = b_1$ and $a_{51} = b_{51}$. If $t = b_1 + b_2 + \dots + b_{51}$ and $s = a_1 + a_2 + \dots + a_{51}$ then

A. $s > t$ and $a_{101} > b_{101}$

B. $s > t$ and $a_{101} < b_{101}$

C. $s < t$ and $a_{101} > b_{101} > b_{101}$

D. $s < t$ and $a_{101} < b_{101}$

Answer: B



Watch Video Solution

8. Let $S_n = \sum_{k=1}^{4n} (-1)^{\frac{k(k+1)}{2}} k^2$. Then S_n can take value (s)

A. 1056

B. 1088

C. 1120

D. 1332

Answer: A::D



Watch Video Solution

9. Let $S_k, k = 1, 2, \dots, 100$ denote the sum of the infinite geometric series whose first term is $\frac{k-1}{k!}$ and the common ratio is $\frac{1}{k}$ then the value of $\frac{(100)^2}{100!} + \sum_{k=1}^{100} |(k^2-3k+1)S_k|$ is _____.

 [Watch Video Solution](#)

10. Let $a_1, a_2, a_3, \dots, a_{11}$ be real numbers satisfying $a_1 = 15, 27 - 2a_2 > 0$ and $a_k = 2a_{k-1} - a_{k-2}$ for $k = 3, 4, \dots, 11$. If $\frac{a_1^2 + a_2^2 + \dots + a_{11}^2}{11} = 90$ then find the value of $\frac{a_1 + a_2 + \dots + a_{11}}{11}$.

 [Watch Video Solution](#)

11. Let $a_1, a_2, a_3, \dots, a_{100}$ be an arithmetic progression with $a_1 = 3$ and $S_p = \sum_{i=1}^p a_i, 1 \leq p \leq 100$. For any integer n with $1 \leq n \leq 20, \leq tm = 5n$. If $\frac{S_m}{S_n}$ does not depend on n then a_2 is _____.

 [Watch Video Solution](#)

12. A pack contains n cards numbered from 1 to n . Two consecutive numbered cards are removed from the pack and the sum of the numbers on the remaining cards is 1224. If the smaller of the numbers on the removed cards is k , then $k - 20 =$ _____.



[Watch Video Solution](#)

13. Let a, b, c be positive integers such that $\frac{b}{a}$ is an integer. If a, b, c are in GP and the arithmetic mean of a, b, c , is $b+2$ then the value of $\frac{a^2 + a - 14}{a + 1}$ is



[Watch Video Solution](#)

14. Suppose that all the terms of an arithmetic progression (A.P.) are natural numbers. If the ratio of the sum of the first seven terms to the sum of the first eleven terms is 6: 11 and the seventh term lies in between 130 and 140, then the common difference of this A.P. is



[Watch Video Solution](#)

15. The sides of a right angled triangle are in arithmetic progression. If the triangle has area 24, then what is the length of its smallest side?



[Watch Video Solution](#)

16. Let X be the set consisting of the first 2018 terms of the arithmetic progression 1, 6, 11, ; and Y be the set consisting of the first 2018 terms of the arithmetic progression 9, 16, 23, .. . Then, the number of elements in the set $X \cup Y$ is _____.



[Watch Video Solution](#)

ARCHIVES (MATRIX MATCH TYPE)

1. Match the statements /expression given in List I with the values given in List II.



[View Text Solution](#)