



## MATHS

### BOOKS - CENGAGE MATHS (HINGLISH)

## TRIGONOMETRIC RATIOS AND TRANSFORMATION FORMULAS

### Examples

1. Prove that 
$$\frac{\sin(B - C)}{\cos B \cos C} + \frac{\sin(C - A)}{\cos C \cos A} + \frac{\sin(A - B)}{\cos A \cos B} = 0$$



Watch Video Solution

2. Eliminate  $x$  from equations  $\sin(a + x) = 2b$  and  $\sin(a - x) = 2c$ .



Watch Video Solution

3. Let  $A, B, C$  be the three angles such that  $A + B + C = \pi$ ,  $\tan A \cdot \tan B = 2$ , then find the value of  $\frac{\cos A \cos B}{\cos C}$

 [Watch Video Solution](#)

4. If  $\sin \alpha \sin \beta - \cos \alpha \cos \beta + 1 = 0$ , then prove that  $1 + \cot \alpha \tan \beta = 0$ .

 [Watch Video Solution](#)

5. If  $\sin \alpha \cos \beta = -\frac{1}{2}$  then find the range of values of  $\cos \alpha \sin \beta$

 [Watch Video Solution](#)

6. Show that  $\cos^2 \theta + \cos^2 \theta(\alpha + \theta) - 2 \cos \alpha \cos \theta \cos(\alpha + \theta)$  is independent of  $\theta$ .

 [Watch Video Solution](#)

7. If  $3 \tan \theta \tan \phi = 1$ , then prove that  $2 \cos(\theta + \phi) = \cos(\theta - \phi)$ .

 [Watch Video Solution](#)

8. If  $\triangle ABC$ , if  $\cot A + \cot B + \cot C = 0$  then find the value of  $\cos A \cos B \cos C$ .

 [Watch Video Solution](#)

9. If  $\alpha, \beta, \gamma \in \left(0, \frac{\pi}{2}\right)$ , then prove that  $\frac{\sin(\alpha + \beta + \gamma)}{\sin \alpha + \sin \beta + \sin \gamma} < 1$ .

 [Watch Video Solution](#)

10. Let  $\alpha, \beta$  and  $\gamma$  satisfy `0

 [Watch Video Solution](#)

11. If in triangle  $ABC$ ,  $\angle C = 45^\circ$  then find the range of the values of  $\sin^2 A + \sin^2 B$ .

 [Watch Video Solution](#)

12. Prove that: 
$$\sum_{k=1}^{100} \sin(kx) \cos(101 - k)x = 50 \sin(101x)$$

 [Watch Video Solution](#)

13. Find the maximum value of  $4 \sin^2 x + 3 \cos^2 + \frac{\sin x}{2} + \frac{\cos x}{2}$ .

 [Watch Video Solution](#)

14. Prove that  $5 \cos \theta + 3 \cos\left(\theta + \frac{\pi}{3}\right) + 3$  lies between  $-4$  and  $10$ .

 [Watch Video Solution](#)

15. Find the maximum vertical distance between the graphs  $y = 2 + 3 \sin x$  and  $y = 4 \cos x - 3$ .

 [Watch Video Solution](#)

16. Find the range of the expression  $27^{\cos 2x} 81^{\sin 2x}$

 [Watch Video Solution](#)

17. Find the range of  $f(x) = \frac{1}{(\cos x - 3)^2 + (\sin x + 4)^2}$

 [Watch Video Solution](#)

18. find the range of function  $f(x) = \sin\left(x + \frac{\pi}{6}\right) + \cos\left(x - \frac{\pi}{6}\right)$

 [Watch Video Solution](#)

19. If  $\sin^2(\theta - \alpha)\cos\alpha = \cos^2(\theta - \alpha)\sin\alpha = m \sin\alpha \cos\alpha$ , then prove that  $|m| \geq \frac{1}{\sqrt{2}}$

 [Watch Video Solution](#)

20. In  $\triangle ABC$ , if  $\sqrt{3}\sin C = 2\sec A - \tan A$ , then prove that triangle is right angled.

 [Watch Video Solution](#)

21. If  $\tan\alpha = \frac{m}{m+1}$  and  $\tan\beta = \frac{1}{2m+1}$ . Find the possible values of  $\tan(\alpha + \beta)$

A. 2

B. 1

C. -1

D. 0

**Answer: B**

 [Watch Video Solution](#)

**22.**

If

$$\sin(A - B) = \frac{1}{\sqrt{10}}, \cos(A + B) = \frac{2}{\sqrt{29}}, f \in \text{dthe value of } 2A \text{ where } A \text{ and } B \text{ are acute angles}$$

 [Watch Video Solution](#)

**23.** Prove that  $\frac{\cos 100^\circ \sin 10^\circ}{\cos 10^\circ - \sin 10^\circ} = \tan 55^\circ$

 [Watch Video Solution](#)

**24.** Prove that  $\tan 70^\circ = \tan 20^\circ + 2 \tan 50^\circ$

 [Watch Video Solution](#)

25. Find the value of

$$\frac{\cot 25^\circ + \cot 55^\circ}{\tan 25^\circ + \tan 55^\circ} + \frac{\cot 55^\circ + \cot 100^\circ}{\tan 55^\circ + \tan 100^\circ} + \frac{\cot 100^\circ + \cot 25^\circ}{\tan 100^\circ + \tan 25^\circ}$$

 [Watch Video Solution](#)

26. Prove that  $(1 + \tan 1^\circ)(1 + \tan 2^\circ) \dots (1 + \tan 45^\circ) = 2^{23}$

 [Watch Video Solution](#)

27. If  $A = \frac{\pi}{5}$ , then find the value of  $\sum_{r=1}^8 \tan(rA)\tan((r+1)A)$ .

 [Watch Video Solution](#)

28. In  $ABC$ , if  $\angle A = \frac{\pi}{4}$ , then find all possible values of  $\tan B \tan C$ .

 [Watch Video Solution](#)



29. If  $\tan^3 A + \tan^3 B + \tan^3 C = 3 \tan A \tan B \tan C$ , then prove that triangle ABC is an equilateral triangle.

 [Watch Video Solution](#)

30. In  $\sin A = \sin B$  and  $\cos A = \cos B$ , then prove that  $\frac{\sin(A - B)}{2} = 0$

 [Watch Video Solution](#)

31. Prove that  $\cos 55^\circ + \cos 65^\circ + \cos 175^\circ = 0$

 [Watch Video Solution](#)

32. Prove that:  $\cos 18^\circ - \sin 18^\circ = \sqrt{2} \sin 27^\circ$

 [Watch Video Solution](#)

33. Prove that:  $\frac{\sin 5A - \sin 3A}{\cos 5A + \cos 3A} = \tan A \frac{\sin A - \sin 3A}{\cos A + \cos 3A} = \tan 2A$

 [Watch Video Solution](#)

34. Prove that  $\frac{\sin A + \sin 2A + \sin 4A + \sin 5A}{\cos A + \cos 2A + \cos 4A + \cos 5A} = \tan 3A$ .

 [Watch Video Solution](#)

35. Prove that

$$\cos \alpha + \cos \beta + \cos \gamma + \cos(\alpha + \beta + \gamma) = 4 \cos\left(\frac{\alpha + \beta}{2}\right) \cos\left(\frac{\beta + \gamma}{2}\right) \cos\left(\frac{\alpha + \gamma}{2}\right)$$

 [Watch Video Solution](#)

36. If  $n$  is an odd positive interger, then

$$\left(\frac{\cos A + \cos B}{\sin A - \sin B}\right)^n + \left(\frac{\sin A + \sin B}{\cos A - \cos B}\right)^n =$$

 [Watch Video Solution](#)

37. Find the Value of  $(\cos \alpha + \cos \beta)^2 + (\sin \alpha + \sin \beta)^2$

A.  $4 \cos^2\left(\frac{\alpha - \beta}{2}\right)$

B.  $4 \cos^2\left(\frac{\alpha + \beta}{2}\right)$

C.  $4 \cos^2(\alpha - \beta)$

D.  $\cos^2\left(\frac{\alpha + \beta}{2}\right)$

Answer: A

 [Watch Video Solution](#)

38. In quadrilateral  $ABCD$ , if

$$\sin\left(\frac{A+B}{2}\right)\cos\left(\frac{A-B}{2}\right) + \sin\left(\frac{C+D}{2}\right)\cos\left(\frac{C-D}{2}\right) = 2 \quad \text{then}$$

find the value of  $\frac{\sin A}{2} \frac{\sin B}{2} \frac{\sin C}{2} \frac{\sin D}{2}$ .

 [Watch Video Solution](#)

39. If  $ABC$ ,  $\sin C + \cos C + \sin(2B + C) - \cos(2B + C) = 2\sqrt{2}$ .

Prove that  $ABC$  is right-angled isosceles.

 [Watch Video Solution](#)

40. If  $\alpha$  and  $\beta$  are acute angles such that  $\alpha + \beta = \lambda$ , where constant, find the maximum possible value of the expression  $\sin \alpha + \sin \beta + \cos \alpha + \cos \beta$ .

 [Watch Video Solution](#)

41. Prove that

$$\sum_{r=1}^n \left( \frac{1}{\cos \theta + \cos(2r + 1)\theta} \right) = \frac{\sin n\theta}{2 \sin \theta \cos \theta \cos(n + 1)\theta}, \text{ (where } n \in \mathbb{N}\text{)}.$$

 [Watch Video Solution](#)

42. Prove that:

$$(a) \frac{\sin 2\theta}{1 + \cos 2\theta} = \tan \theta \quad (b) \frac{1 + \sin 2\theta + \cos 2\theta}{1 + \sin 2\theta - \cos 2\theta} = \cot \theta$$

 [Watch Video Solution](#)

43. Prove that  $\frac{1 + \sin 2\theta}{1 - \sin 2\theta} = \left( \frac{1 + \tan \theta}{1 - \tan \theta} \right)^2$

 [Watch Video Solution](#)

44. If  $\alpha + \beta = 90^\circ$ , find the maximum and minimum values of  $\sin \alpha \sin \beta$ .

 [Watch Video Solution](#)

45. If  $\sin A = \frac{3}{5}$ , where  $0^\circ < A < 90^\circ$ , then find the values of  $\sin 2A$ ,  $\cos 2A$ ,  $\tan 2A$  and  $\sin 4A$

 [Watch Video Solution](#)

46. Find the Value of  $\sqrt{2 + \sqrt{2 + 2 \cos 4\theta}}$

A.  $2 \cos \theta$

B.  $\cos \theta$

C.  $2 \cos \left( \frac{\theta}{2} \right)$

D.  $2 \cos \left( \frac{\theta}{4} \right)$

**Answer: A**



[Watch Video Solution](#)

47. Prove that  $\frac{\sec 8\theta - 1}{\sec 4\theta - 1} = \frac{\tan 8\theta}{\tan 2\theta}$



[Watch Video Solution](#)

48. Prove that  $\frac{\tan \pi}{16} + 2 \frac{\tan \pi}{8} + 4 = \frac{\cot \pi}{16}$ .



[Watch Video Solution](#)

49. Prove that:  $\frac{\cos^4 \pi}{8} + \frac{\cos^4(3\pi)}{8} + \frac{\cos^4(5\pi)}{8} + \frac{\cos^4(7\pi)}{8} = \frac{3}{2}$

 Watch Video Solution

50. If  $\pi$

 Watch Video Solution

51. If  $\sin \alpha + \sin \beta = a$  and  $\cos \alpha + \cos \beta = b$ , prove that

$$\frac{\tan(\alpha - \beta)}{2} = \pm \sqrt{\frac{4 - a^2 - b^2}{a^2 + b^2}}$$

 Watch Video Solution

52. Prove that  $\frac{1 - \tan^2\left(\frac{\pi}{4} - A\right)}{1 + \tan^2\left(\frac{\pi}{4} - A\right)} = \sin 2A$ .



 Watch Video Solution

53. If  $\frac{\tan \theta}{2} = \sqrt{\frac{2-b}{a+b}} \frac{\tan \phi}{2}$ , prove that  $\cos \alpha = \frac{1 \cos \phi + b}{a + b \cos \phi}$

 Watch Video Solution

54. If  $(\cos \theta = \cos \alpha \cos \beta)$ , prove that  $\tan \frac{\theta + \alpha}{2} \tan \frac{\theta - \alpha}{2} = \tan^2 \frac{\beta}{2}$ .

 Watch Video Solution

55. If  $\tan \beta = \frac{\tan \alpha + \tan \gamma}{1 + \tan \alpha \tan \gamma}$ , prove that  $\sin 2\beta = \frac{\sin 2\alpha + \sin 2\gamma}{1 + \sin 2\alpha \sin 2\gamma}$ .

 Watch Video Solution

56. Prove that  $(4 \cos^2 9^\circ - 3)(4 \cos^2 27^\circ - 3) = \tan 9^\circ$ .

 Watch Video Solution



57. Prove that  $\left(4 \frac{\cos(2\pi)}{7} \cdot \frac{\cos \pi}{7} - 1 = 2 \frac{\cos(2\pi)}{7}\right)$ .

 [Watch Video Solution](#)

58. Evaluate  $\cos \alpha \cos 2\alpha \cos 3\alpha \dots \cos 999\alpha$ , where  $\alpha = \frac{2\pi}{1999}$

 [Watch Video Solution](#)

59. prove that  $\sin \theta \sec 3\theta + \sin 3\theta, \sec 3^2\theta + \sin 3^2\theta \sec 3^3\theta + \dots \rightarrow n$   
terms  $= \frac{1}{2}[\tan 3^n\theta - \tan \theta]$

 [Watch Video Solution](#)

60. Let  $f(x) = 2 \cos ec2x + \sec x + \cos ecx$ . Then find the minimum value of  $f(x)$  or  $x \in \left(0, \frac{\pi}{2}\right)$ .

 [Watch Video Solution](#)

61. Find the maximum and minimum values of  $\cos^2 \theta - 6 \sin \theta \cos \theta + 3 \sin^2 \theta + 2$ .

 [Watch Video Solution](#)

62. If  $\tan \alpha = \frac{1}{7}$ ,  $\sin \beta = \frac{1}{\sqrt{10}}$ , prove that  $\alpha + 2\beta = \frac{\pi}{4}$ , where  $\alpha, \beta \in (0, \frac{\pi}{2})$ .

 [Watch Video Solution](#)

63. Prove that  $\frac{\tan \pi}{10}$  is a root of polynomial equation  $5x^4 - 10x^2 + 1 = 0$ .

 [Watch Video Solution](#)

64. If  $x + y + z = xyz$  prove that  $\frac{2x}{1-x^2} + \frac{2y}{1-y^2} + \frac{2z}{1-z^2} = \frac{2x}{1-x^2} \frac{2y}{1-y^2} \frac{2z}{1-z^2}$ .

 [Watch Video Solution](#)

65. Prove that  $1 + \cot^2 \theta = \csc^2 \theta$  for  $\theta \neq 0$

 [Watch Video Solution](#)

66. Find the angle  $\theta$  whose cosine is equal to its tangent.

 [Watch Video Solution](#)

67. Find the value of  $\cos 12^\circ + \cos 84^\circ + \cos 156^\circ + \cos 132^\circ$

 [Watch Video Solution](#)

68. Prove that  $\cos 36^\circ \cos 72^\circ \cos 108^\circ \cos 144^\circ = 1/16$ .

 [Watch Video Solution](#)

69. Show that  $4\sin 27^\circ = (5 + \sqrt{5})^{\frac{1}{2}} - (3 - \sqrt{5})^{\frac{1}{2}}$

 [Watch Video Solution](#)

70. Prove that:  $\tan \frac{\pi}{16} = \sqrt{4 + 2\sqrt{2}} - (\sqrt{2} + 1)$

 [Watch Video Solution](#)

71. Find the quadratic equation whose roots are  $\tan\left(\frac{\pi}{8}\right)$  and  $\tan\left(\frac{5\pi}{8}\right)$  ?

 [Watch Video Solution](#)

72. Prove that  $\cos 20^\circ \cos 40^\circ \cos 60^\circ \cos 80^\circ = \frac{1}{16}$

 [Watch Video Solution](#)

73. The value of  $\sin 10^\circ \sin 30^\circ \sin 50^\circ \sin 70^\circ$  is equal to .

A.  $\frac{1}{8}$

B.  $\frac{1}{32}$

C.  $\frac{1}{16}$

D.  $\frac{1}{12}$

**Answer: C**



**Watch Video Solution**

74. The value of  $\tan 20^\circ \tan 40^\circ \tan 80^\circ$  is equal to

A.  $\tan 60^\circ$

B.  $\cot 60^\circ$

C.  $\tan 45^\circ$

D.  $\tan 80^\circ$

**Answer: A**



**Watch Video Solution**

75. If  $\theta = \frac{\pi}{2^n + 1}$ , prove that:  $2^n \cos \theta \cos 2\theta \cos 2^2 \cos 2^{n-1} \theta = 1$ .



**Watch Video Solution**

76.  $\cos \frac{2\pi}{15} \cos \frac{4\pi}{15} \cos \frac{8\pi}{15} \cos \frac{16\pi}{15} = \frac{1}{16}$



**Watch Video Solution**

77. Prove that  $\sin 6^\circ \sin 42^\circ \sin 66^\circ \sin 78^\circ = \frac{1}{16}$



**Watch Video Solution**

78. Find the value of  $2 \frac{\cos^2 \pi}{7} - \frac{\cos^2 \pi}{7} - \frac{\cos \pi}{7}$

[Watch Video Solution](#)

79. Find the value of  $\frac{\cos(2\pi)}{7} + \frac{\cos(4\pi)}{7} + \frac{\cos(6\pi)}{7}$

[Watch Video Solution](#)

80. Prove that  $\sin \theta + \sin 3\theta + \sin 5\theta + \dots + \sin(2n - 1)\theta = \frac{\sin^2 n\theta}{\sin \theta}$ .

[Watch Video Solution](#)

81. Prove that

$$\begin{aligned} & \frac{\cos 3x}{\sin 2x \sin 4x} + \frac{\cos 5x}{\sin 4x \sin 6x} + \frac{\cos 7x}{\sin 6x \sin 8x} + \frac{\cos 9x}{\sin 8x \sin 10x} \\ &= \frac{1}{2} (\operatorname{cosec} x) [\operatorname{cosec} 2x - \operatorname{cosec} 10x] \end{aligned}$$

[Watch Video Solution](#)

82. Prove that

$$2\sin 2^\circ + 4\sin 4^\circ + 6\sin 6^\circ + \dots + 180\sin 180^\circ = 90\cot 10^\circ.$$

 [Watch Video Solution](#)

83. If  $A + B + C = 180$ , prove that

$$\cos^2 A + \cos^2 B + \cos^2 C = 1 - 2 \cos A \cos B \cos C.$$

 [Watch Video Solution](#)

84. If  $A + B + C = 180^\circ$ , prove that :

$$\cos^2 A + \cos^2 B - \cos^2 C = 1 - 2 \sin A \sin B \cos C$$

 [Watch Video Solution](#)

85. In triangle ABC, prove that

$$\sin(B + C - A) + \sin(C + A - B) + \sin(A + B - C) = 4 \sin A \sin B \sin C$$



 [Watch Video Solution](#)

86. If  $A + B + C = \pi$ , prove that

$$\frac{\sin^2 A}{2} + \frac{\sin^2 B}{2} - \frac{\sin^2 C}{2} = 1 - 2 \frac{\cos A}{2} \frac{\cos B}{2} \frac{\sin C}{2}.$$

 [Watch Video Solution](#)

87. Prove that, in triangle ABC

$$\sin^3 A \cos(B - C) + \sin^3 B \cos(C - A) + \sin^3 C \cos(A - B) = 3 \sin A \sin B \sin C.$$

 [Watch Video Solution](#)

88. If  $A + B + C = \pi$ , prove that

$$\cot A + \cot B + \cot C - \cos A \cos B \cos C = \cot A \cot B \cot C.$$

 [Watch Video Solution](#)

89. If  $A + B + C = \pi$ , prove that

$$\frac{\tan A}{\tan B \tan C} + \frac{\tan B}{\tan A \tan C} + \frac{\tan C}{\tan A \tan B} = \tan A + \tan B + \tan C - 2 \cot A$$

 [Watch Video Solution](#)

90. In triangle  $ABC$ , if  $\cot A \cot C = \frac{1}{2}$  and  $\cot B \cot C = \frac{1}{18}$ , then the value of  $\tan C$  is

 [Watch Video Solution](#)

91. If  $\cos(A + B + C) = \cos A \cos B \cos C$ , then find the value of

$$\frac{8 \sin(B + C) \sin(C + A) \sin(A + B)}{\sin 2A \sin 2B \sin 2C}$$

 [Watch Video Solution](#)

92. If  $x + y + z = \frac{\pi}{2}$ , then prove that

$$|\sin x \sin y \sin z \cos x \cos y \cos z \cos^3 x \cos^y y \cos^3 z| = 0$$

 [Watch Video Solution](#)

93. The product of the sines of the angles of a triangle is  $p$  and the product of their cosines is  $q$ . Show that the tangents of the angles are the roots of the equation  $qx^3 - px^2 + (1 + q)x - p = 0$ .

 [Watch Video Solution](#)

94. If  $x^2 + yx^2 = 4$  then find the maximum value of  $\frac{x^3 + y^3}{x + y}$

 [Watch Video Solution](#)

95. If  $\frac{x^2}{4} + \frac{y^2}{9} = 1$ , then find the range of  $2 + y$ .

 [Watch Video Solution](#)

96. If  $x^2 + y^2 = x^2y^2$  then find the range of  $\frac{5x + 12y + 7xy}{xy}$ .



Watch Video Solution

97. For all,  $x, y \in \mathbb{R}$ . find the range of  $\frac{(x+y)(1-xy)}{(1+x^2)(1+y^2)}$ .



Watch Video Solution

98. If  $x, y \in \mathbb{R}$  and  $x^2 + y^2 + xy = 1$ , then find the minimum value of  $x^3y + xy^3 + 4$ .



Watch Video Solution

99. Prove that in a  $ABC$ ,  $\sin^3 A + \sin^2 B + \sin^2 C \leq \frac{9}{4}$ .



Watch Video Solution

100. Prove that in  $\triangle ABC$ ,  $2 \cos A \cos B \cos C \leq \frac{1}{8}$ .



Watch Video Solution

101. In  $\triangle ABC$ , prove that  $\cos^2 A + \cos^2 B + \cos^2 C \geq \frac{3}{4}$ .

 [Watch Video Solution](#)

102. In  $\triangle ABC$ , prove that  $\sin\left(\frac{A}{2}\right) + \sin\left(\frac{B}{2}\right) + \sin\left(\frac{C}{2}\right) \leq \frac{3}{2}$ .

 [Watch Video Solution](#)

103. Find the least value of  $\sec A + \sec B + \sec C$  in an acute angled triangle.

 [Watch Video Solution](#)

104. If  $A + B + C = \pi$ , prove that  $\frac{\tan^2 A}{2} + \frac{\tan^2 B}{2} + \frac{\tan^2 C}{2} \geq 1$ .

 [Watch Video Solution](#)

105. In a  $ABC$ , if  $\frac{\tan A}{2}, \frac{\tan B}{2}, \frac{\tan C}{2}$  are in A.P.; then show that  $\cos A, \cos B, \cos C$  are in A.P.

 [Watch Video Solution](#)

106. If  $ABC$  is a triangle and  $\tan\left(\frac{A}{2}\right), \tan\left(\frac{B}{2}\right), \tan\left(\frac{C}{2}\right)$  are in H.P. Then find the minimum value of  $\cot\left(\frac{B}{2}\right)$

 [Watch Video Solution](#)

107. In  $ABC$ , if  $\sin^3 \theta = \sin(A - \theta)\sin(B - \theta)\sin(C - \theta)$ , then prove that  $\cot \theta = \cot A + \cot B + \cot C$ .

 [Watch Video Solution](#)

108. Find the sum of series  $\operatorname{cosec} \theta + \operatorname{cosec} 2\theta + \operatorname{cosec} 4\theta + \dots$  to  $n$  terms



Watch Video Solution

109. If  $\frac{x}{\tan(\theta + \alpha)} = \frac{y}{\tan(\theta + \beta)} = \frac{z}{\tan(\theta + \gamma)}$ , then show that

$$\sum \frac{x + y}{x - y} \sin^2(\alpha - \beta) = 0$$



Watch Video Solution

110. If  $\tan 6\theta = p/q$  find the value of  $\frac{1}{2}(p \csc 2\theta - q \sec 2\theta)$  terms of  $p$  and  $q$ .



Watch Video Solution

111. If  $\theta = 0$



Watch Video Solution

112. Let  $A, B, C$ , be three angles such that  $A = \frac{\pi}{4}$  and  $\tan B, \tan C = p$ . Find all possible values of  $p$  such that  $A, B, C$  are the angles of a triangle.

 [Watch Video Solution](#)

113. If  $\sin A = \frac{12}{13}$  and  $\sin B = \frac{4}{5}$ , where  $\theta \in \left(\frac{\pi}{2}, \pi\right)$

 [Watch Video Solution](#)

114. Prove that :

$$\frac{2\cos 2^n \theta + 1}{2\cos \theta + 1} = (2\cos \theta - 1)(2\cos 2\theta - 1)(2\cos 2^2\theta - 1) \dots (2\cos 2^{n-1}\theta - 1)$$

 [Watch Video Solution](#)

115. If  $\tan\left(\frac{\pi}{4} + y\right) = \tan^3\left(\frac{\pi}{4} + \frac{x}{2}\right)$ . Prove that  $\frac{\sin y}{\sin x} = \frac{3 + \sin^2 x}{1 + 3\sin^2 x}$ .

 [Watch Video Solution](#)



116. If  $(1 + \sin t)(1 + \cos t) = \frac{5}{4}$ . Find the value  $(1 - \sin t)(1 - \cos t)$

 [Watch Video Solution](#)

117. For all  $\theta$  in  $\left[0, \frac{\pi}{2}\right]$  show that the  $\cos(\sin \theta) \geq \sin(\cos \theta)$

 [Watch Video Solution](#)

118. Prove that  $\frac{\tan 3x}{\tan x}$  never lies between  $\frac{1}{3}$  and 3.

 [Watch Video Solution](#)

119. Prove that

$$\sum_{k=1}^{n-1} (n-k) \frac{\cos(2k\pi)}{n} = -\frac{n}{2}, \text{ where } n \geq 3 \text{ is an integer } \geq r$$

 [Watch Video Solution](#)

### Exercise 3.1

1. In  $\triangle ABC$ , if  $\cos A + \sin A - \frac{2}{\cos B + \sin B} = 0$  then prove that triangle is isosceles right angled.

 [Watch Video Solution](#)

2. If  $x$  is A. M. of  $\tan\left(\frac{\pi}{9}\right)$  and  $\tan\left(\frac{5\pi}{18}\right)$  and  $y$  is A. M. of  $\tan\left(\frac{\pi}{9}\right)$  and  $\tan\left(\frac{7\pi}{18}\right)$ , then

 [Watch Video Solution](#)

3. Find the value of

$$\cos \frac{\pi}{12} \left( \sin \frac{5\pi}{12} + \cos \frac{\pi}{4} \right) + \sin \frac{\pi}{12} \left( \cos \frac{5\pi}{12} - \sin \frac{\pi}{4} \right).$$

 [Watch Video Solution](#)

4. If  $\cos(\alpha + \beta) + \sin(\alpha - \beta) = 0$  and  $\tan \beta \neq 1$ , then find the value of  $\tan \alpha$ .

 [Watch Video Solution](#)

5. If  $\sin A + \cos 2A = 1/2$  and  $\cos A + \sin 2A = 1/3$ . Then find the value of  $\sin 3A$ .

 [Watch Video Solution](#)

6. If  $\sin x + \sin y + \sin z = 0 = \cos x + \cos y + \cos z$ , then find the value of  $\cos(\theta - x) + \cos(\theta - z)$

 [Watch Video Solution](#)

7. In a triangle ABC, if  $\sin A \sin(B - C) = \sin C \sin(A - B)$ , then prove that  $\cot A, \cot B, \cot C$  are  $\in AP$ .





Watch Video Solution

8. Find the value of

$$\frac{(\cos 1^\circ + \sin 1^\circ)(\cos 2^\circ + \sin 2^\circ)(\cos 3^\circ + \sin 3^\circ) \dots (\cos 45^\circ + \sin 45^\circ)}{\cos 1^\circ \cos 2^\circ \cos 3^\circ \dots \cos 45^\circ}$$



Watch Video Solution

9. Find the maximum value of  $\sqrt{3}\sin x + \cos x$  and  $x$  for which a maximum value occurs.



Watch Video Solution

10. The maximum value of  $1 + \sin\left(\frac{\pi}{4} + \theta\right) + 2\cos\left(\frac{\pi}{4} - \theta\right)$  for real values of  $\theta$  is



Watch Video Solution

11. show that  $2^{\sin x} + 2^{\cos x} \geq 2^{1 - \frac{1}{\sqrt{2}}}$



Watch Video Solution

### Exercise 3.2

1. If  $A + B = 225^\circ$ , then find the value of  $\frac{\cot A}{1 + \cot A} \times \frac{\cot B}{1 + \cot B}$



Watch Video Solution

2. If  $\tan A - \tan B = x$ , and  $\cot B - \cot A = y$ , then find the value of  $\cot(A - B)$ .



Watch Video Solution

3. Prove that  $\frac{\tan^2 2\theta - \tan^2 \theta}{1 - \tan^2 2\theta \tan^2 \theta} = \tan 3\theta \tan \theta$ .



Watch Video Solution

4. If  $A + B = 45^\circ$ , then  $(1 + \tan A)(1 + \tan B) = \underline{\hspace{2cm}}$

 [Watch Video Solution](#)

5. If  $\tan A = 1/2$ ,  $\tan B = 1/3$ , then prove that  $\cos 2A = \sin 2B$ .

 [Watch Video Solution](#)

6. If  $P + Q = \frac{7\pi}{6}$ , then find the value of  $(\sqrt{3} + \tan P) \times (\sqrt{3} + \tan Q)$ .

 [Watch Video Solution](#)

7. If  $\tan \beta = \frac{n \sin \alpha \cos \beta}{1 - n \sin^2 \alpha}$  then prove that  $\tan(\alpha - \beta) = (1 - n)\tan \alpha$ .

 [Watch Video Solution](#)

### Exercise 3.3

1. (a) Prove that  $\sin 65^\circ + \cos 65^\circ = \sqrt{2}\cos 20^\circ$

(b) Prove that  $\sin 47^\circ + \cos 77^\circ = \cos 17^\circ$

 [Watch Video Solution](#)

2. Prove that:  $\cos 80^\circ + \cos 40^\circ - \cos 20^\circ = 0$

 [Watch Video Solution](#)

3. Prove that  $\sin 10^\circ + \sin 20^\circ + \sin 40^\circ + \sin 50^\circ = \sin 70^\circ + \sin 80^\circ$

 [Watch Video Solution](#)

4.  $\frac{\cos \pi}{5} + \frac{\cos(2\pi)}{5} + \frac{\cos(6\pi)}{5} + \frac{\cos(7\pi)}{5} = 0$

 [Watch Video Solution](#)

5. If  $\sin \alpha - \sin \beta = \frac{1}{3}$  and  $\cos \beta - \cos \alpha = \frac{1}{2}$ , show that

$$\frac{\cot(\alpha + \beta)}{2} = \frac{2}{3}$$

 [Watch Video Solution](#)

6. If  $\cos ecA + \sec A = \cos ecB + \sec B$ , prove that:

$$\tan A \tan B = \frac{\cot(A + B)}{2}$$

 [Watch Video Solution](#)

7.  $\sin 25^\circ \cos 115^\circ = \frac{1}{2}(\sin 40^\circ - 1)$

 [Watch Video Solution](#)

8. If  $x \cos \theta = y \cos \left( \theta + \frac{2\pi}{3} \right) = z \cos \left( \theta + \frac{4\pi}{3} \right)$ , prove that

$$xy + yz + zx = 0.$$







Watch Video Solution

9. If  $y \sin \phi = x \sin(2\theta + \phi)$  show that

$$(x + y) \cot(\theta + \phi) = (y - x) \cot \theta.$$



Watch Video Solution

10. If  $\cos(A + B)\sin(C + D) = \cos(A - B)\sin(C - D)$ , prove that  $\cot A \cot B \cot C = \cot D$ .



Watch Video Solution

11. If  $\tan(A + B) = 3 \tan A$ , prove that (a)  $\sin(2A + B) = 2 \sin B$



Watch Video Solution

12. If  $\frac{x}{2} = \frac{\cos A}{\cos B}$  then prove that

$$\frac{x \tan A + y \tan B}{x + y} = \tan \frac{A + B}{2}$$



Watch Video Solution

13. If  $\frac{\cos 6x + 6 \cos 4x + 15 \cos 2x + 10}{\cos 5x + 5 \cos 3x + 10 \cos x} = 1$ , then find the smallest positive value of  $x$ .



Watch Video Solution

### Exercise 3.4

1. 
$$\frac{1 + \sin 2A - \cos 2A}{1 + \sin 2A + \cos 2A} = \tan A$$



Watch Video Solution

2. Prove that 
$$\frac{1 + \sin 2A}{\cos 2A} = \frac{\cos A + \sin A}{\cos A - \sin A} = \tan\left(\frac{\pi}{4} + A\right)$$



Watch Video Solution

3. Prove that  $\cot \theta - \tan \theta = 2 \cot 2\theta$ .

 [Watch Video Solution](#)

4. Prove that  $\frac{\cos \theta - \sin \theta}{\cos \theta + \sin \theta} = \sec 2\theta - \tan 2\theta$ .

 [Watch Video Solution](#)

5.  $\tan\left(\frac{\pi}{4} + \theta\right) - \tan\left(\frac{\pi}{4} - \theta\right) = 2 \tan 2\theta$

 [Watch Video Solution](#)

6. Prove that  $\cos ec A - 2 \cot 2A \cos A = 2 \sin A$ .

 [Watch Video Solution](#)

7. Prove that  $\cos^3 \theta \sin 3\theta + \sin^3 \theta \cos 3\theta = \frac{3}{4} \sin 4\theta$ .



Watch Video Solution

8.  $\frac{\sin^2 3A}{\sin^2 A} - \frac{\cos^2 3A}{\cos^2 A} =$



Watch Video Solution

9. Prove that  $(1 + \sec 2\theta)(1 + \sec 4\theta)(1 + \sec 8\theta) = \frac{\tan 8\theta}{\tan \theta}$



Watch Video Solution

10. If in an isosceles triangle with base 'a', vertical angle  $20^\circ$  and lateral side each of length 'b' is given then value of  $a^3 + b^3$  equals



Watch Video Solution

11. In  $\Delta ABC$ ,  $a = 3$ ,  $b = 4$  and  $c = 5$ , then value of  $\sin A + \sin 2B + \sin 3C$  is



Watch Video Solution

12. If  $\cos A = \frac{3}{4}$ , then  $32 \sin \frac{A}{2} \sin \frac{5A}{2} =$  (A)  $\sqrt{11}$  (B)  $-\sqrt{11}$  (C) 11  
(D) -11



Watch Video Solution

13. Find the value of  $(4 \cos^2 9^\circ - 1)(4 \cos^2 27^\circ - 1)$   
 $(4 \cos^2 81^\circ - 1)(4 \cos^2 243^\circ - 1)$ .



Watch Video Solution

14. If  $\theta$  is an acute angle and  $\sin\left(\frac{\theta}{2}\right) = \sqrt{\frac{x-1}{2x}}$ , then  $\tan \theta$  is equal to



Watch Video Solution

15. In a triangle ABC, if  $\sin A \sin(B - C) = \sin C \sin(A - B)$ , then prove that  $\cos 2A$ ,  $\cos 2B$  and  $\cos 2C$  are in AP.

 [Watch Video Solution](#)

16. Let  $a = \frac{\pi}{7}$ , then

(a) show that  $\sin^2 3a - \sin^2 a = \sin 2a \sin 3a$

(b) show that  $\operatorname{cosec} a = \operatorname{cosec} 2a + \operatorname{cosec} 4a$ .

(c) Prove that  $\cos a$  is a root of the equation  $8x^3 + 4x^2 - 4x + 1 = 0$ .

 [Watch Video Solution](#)

17. Show that  $\frac{1}{\sin 10^\circ} - \frac{\sqrt{3}}{\cos 10^\circ} = 4$

 [Watch Video Solution](#)

18.

Prove

that

$$2 \sin^2 \beta + 4 \cos(\alpha + \beta) \sin \alpha \sin \beta + \cos 2(\alpha + \beta) = \cos 2\alpha$$

[!\[\]\(9dfdaff1d86ba3c1f8353b4d1b61b8c5\_img.jpg\) Watch Video Solution](#)

19. If  $\tan x = \frac{a}{b}$  and  $\tan 2x = \frac{b}{a+b}$  find the smallest positive value of  $x$ .

[!\[\]\(8d0f0e0fe25b320c33272c52aec1fbca\_img.jpg\) Watch Video Solution](#)

$$20. \tan \theta + \tan(60^\circ + \theta) + \tan(120^\circ + \theta) = 3 \tan 3\theta$$

[!\[\]\(2b376d1a92330ab09dad2665d2f89bf5\_img.jpg\) Watch Video Solution](#)

21. If  $A = 110^\circ$ , then prove that  $\frac{1 + \sqrt{1 + \tan^2 2A}}{\tan 2A} = -\tan A$ .

[!\[\]\(d0262bbe9d2356661a2e89321dfcc781\_img.jpg\) Watch Video Solution](#)

22. If  $\alpha$  and  $\beta$  are the two different roots of equations  $a \cos \theta + b \sin \theta = c$ , prove that

$$(a) \tan(\alpha - \beta) = \frac{2ab}{a^2 - b^2} \quad (b) \cos(\alpha + \beta) = \frac{a^2 - b^2}{a^2 + b^2}$$

 [Watch Video Solution](#)

23. If  $\tan \beta = \cos \theta \tan \alpha$ , then prove that  $\tan^2 \frac{\theta}{2} = \frac{\sin(\alpha - \beta)}{\sin(\alpha + \beta)}$ .

 [Watch Video Solution](#)

24. If  $\cos \theta = \frac{a}{b+c}$ ,  $\cos \phi = \frac{b}{a+c}$  and  $\cos \psi = \frac{c}{a+b}$  where  $\theta, \phi, \psi \in (0, \pi)$  and  $a, b, c$  are sides of triangle  $ABC$  then  $\tan^2\left(\frac{\theta}{2}\right) + \tan^2\left(\frac{\phi}{2}\right) + \tan^2\left(\frac{\psi}{2}\right) =$

 [Watch Video Solution](#)

25. If  $\cos \theta = \frac{\cos \alpha - \cos \beta}{1 - \cos \alpha \cos \beta}$ , prove that  $\frac{\tan \theta}{2} = \pm \frac{\tan \alpha}{2} \frac{\cot \beta}{2}$ .





Watch Video Solution

26. If  $\tan \theta \tan \phi = \sqrt{\frac{a-b}{a+b}}$ , prove that

$a - b \cos 2\theta)(a - b \cos 2\phi)$  is independent of  $\theta$  and  $\phi$ .



Watch Video Solution

### Exercise 3.5

1. Find the value of  $(\cos^2 66^\circ - \sin^2 6^\circ)(\cos^2 48^\circ - \sin^2 12^\circ)$ .



Watch Video Solution

2.  $4(\sin 24^\circ + \cos 6^\circ) = \sqrt{3} + \sqrt{15}$



Watch Video Solution

3.  $\sin 47^\circ + \sin 61^\circ - \sin 11^\circ - \sin 25^\circ =$



Watch Video Solution

4. Find the values of following . (a)  $\frac{\tan^2 37\frac{1}{(7)^\circ} + 1}{\tan^2 37\frac{1}{(2)^\circ} - 1}$  (b)

$$\frac{3 \tan^2 5^\circ - 1}{3 \tan 5^\circ - \tan^3 5^\circ}$$



Watch Video Solution

5. If the value of

$$\cot\left(11\frac{1}{4}^\circ\right) + \tan\left(112\frac{1}{2}^\circ\right) - \cot\left(112\frac{1}{2}^\circ\right) - \tan\left(11\frac{1}{4}^\circ\right) = \sqrt{n}$$

where  $n \in N$  then find the value of  $n$



Watch Video Solution

6. Find the value of  $\frac{\tan 9^\circ + \cot 9^\circ}{\tan 27^\circ + \cot 27^\circ}$ .



 Watch Video Solution

### Exercise 3.6

1. Prove that:  $\sin 20^\circ \sin 40^\circ \sin 60^\circ \sin 80^\circ = \frac{3}{16}$

 Watch Video Solution

2. Prove that:  $\cos 10^\circ \cos 30^\circ \cos 50^\circ \cos 70^\circ = \frac{3}{16}$

 Watch Video Solution

3. Prove that  $\sin 12^\circ \sin 18^\circ \sin 42^\circ \sin 48^\circ \sin 72^\circ \sin 78^\circ = \frac{\cos 18^\circ}{32}$ .

 Watch Video Solution

4.  $\frac{\sin(9\pi)}{14} \frac{\sin(11\pi)}{14} \frac{\sin(13\pi)}{14}$  is equal to



 Watch Video Solution

5. The value of  $\frac{\sin \pi}{14} \frac{\sin(3\pi)}{14} \frac{\sin(5\pi)}{14} \frac{\sin(7\pi)}{14} \frac{\sin(9\pi)}{14} \frac{\sin(11\pi)}{14} \frac{\sin(13\pi)}{14}$  is equal to \_\_\_\_\_

 Watch Video Solution

### Exercise 3.7

1. The value of  $\cos \frac{\pi}{11} + \cos \frac{3\pi}{11} + \cos \frac{5\pi}{11} + \cos \frac{7\pi}{11} + \cos \frac{9\pi}{11}$ , is

 Watch Video Solution

2. The average value of  $\sin 2^\circ, \sin 4^\circ, \sin 6^\circ, \dots, \sin 180^\circ$  is  
(i)  $\frac{1}{90} \cos 1^\circ$  (ii)  $\frac{1}{90} \sin 1^\circ$  (iii)  $\frac{1}{90} \cot 1^\circ$  (iv) none of these

 Watch Video Solution

3.  $\sum_{r=0}^n \sin^2 \frac{r\pi}{n}$  is equal to

 [Watch Video Solution](#)

4. Sum the series:  $\sqrt{1 + \cos \alpha} + \sqrt{1 + \cos 2\alpha} + \sqrt{1 + \cos 3\alpha} + \dots$  to  $n$  terms, where  $0 < \alpha < \pi$

 [Watch Video Solution](#)

5. The value of  $(\cos^4 1^\circ + \cos^4 2^\circ + \dots + \cos^4 179^\circ) - (\sin^4 1^\circ + \sin^4 2^\circ + \dots + \sin^4 179^\circ)$  equals

 [Watch Video Solution](#)

### Exercise 3.8

1. If  $A + B + C = 180^\circ$ , prove that :

$$\cos^2 \frac{A}{2} + \cos^2 \frac{B}{2} - \cos^2 \frac{C}{2} = 2 \cos \frac{A}{2} \cos \frac{B}{2} \sin \frac{C}{2}$$

 [Watch Video Solution](#)

2. If  $A + B + C = \frac{\pi}{2}$ , show that :

$$\sin^2 A + \sin^2 B + \sin^2 C = 1 - 2 \sin A \sin B \sin C$$

 [Watch Video Solution](#)

3.  $\cos^2 A + \cos^2 B + \cos^2 C = 1 + 2 \cos A \cos B \cos C$ .

 [Watch Video Solution](#)

4. Prove that :

$$\cos^2(\beta - \gamma) + \cos^2(\gamma - \alpha) + \cos^2(\alpha - \beta) = 1 + 2 \cos(\beta - \gamma) \cos(\gamma - \alpha) \cos(\alpha - \beta)$$




Watch Video Solution

5. If  $A + B + C = \frac{\pi}{2}$ , show that :

$$\cot A + \cot B + \cot C = \cot A \cot B \cot C$$



Watch Video Solution

6. If  $A + B + C = \pi$ , prove that :

$$\cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2} = \cot \frac{A}{2} \cot \frac{B}{2} \cot \frac{C}{2}$$



Watch Video Solution

7. If  $A + B + C = \pi$ , prove that :

$$\frac{\cos A}{\sin B \sin C} + \frac{\cos B}{\sin C \sin A} + \frac{\cos C}{\sin A \sin B} = 2.$$



Watch Video Solution

8. In a triangle  $ABC$ ,  $\cos 3A + \cos 3B + \cos 3C = 1$  and  $\angle A + \angle B < \angle C$ , then find possible measure of  $\angle C$ .

 [Watch Video Solution](#)

9. In  $\triangle ABC$  if  $2\sin^2 C = 2 + \cos 2A + \cos 2B$ , then prove that triangle is right angled.

 [Watch Video Solution](#)

### Exercise 3.9

1. Let  $x, y \in R$ , then find the maximum and minimum values of expression  $\frac{x^2 + y^2}{x^2 + xy + 4y^2}$ .

 [Watch Video Solution](#)



2. Let  $a^2 + b^2 = \alpha^2 + \beta^2 = 2$ . Then show that the maximum value of  $S = (1 - \alpha)(a - b) + (1 - \alpha)(1 - \beta)$  is 8.

 [View Text Solution](#)

3. Find the maximum distance of any point on the curve  $x^2 + 2y^2 + 2xy = 1$  from the origin.

 [Watch Video Solution](#)

4. If  $\frac{x^2}{144} - \frac{y^2}{25} = 1$ . Find the range of  $\frac{144}{x} + \frac{25}{y}$ .

 [Watch Video Solution](#)

5. If  $x^2 + y^2 + 6x - 4y - 12 = 0$  then find the range of  $2x+y$

 [Watch Video Solution](#)

### Exercise 3.10

1. In  $\triangle ABC$  Prove that  $\frac{\cos^2 A}{2} + \frac{\cos^2 B}{2} + \frac{\cos^2 C}{2} \leq \frac{9}{4}$ . In  $\frac{\cos^2 A}{2} + \frac{\cos^2 B}{2} + \frac{\cos^2 C}{2} = y \left( x^2 + \frac{1}{x^2} \right)$  then find the maximum value of  $y$ .

 [Watch Video Solution](#)

2. Let  $\alpha, \beta, \gamma > 0$  and  $\alpha + \beta + \gamma = \frac{\pi}{2}$ . Then prove that  $\sqrt{\tan \alpha \tan \beta} + \sqrt{\tan \beta \tan \gamma} + \sqrt{\tan \alpha \tan \gamma} \leq \sqrt{3}$

 [Watch Video Solution](#)

3. In acute angled  $\triangle ABC$  prove that  $\tan^2 A + \tan^2 B + \tan^2 C \geq 9$ .

 [Watch Video Solution](#)

4. In a  $\triangle ABC$   $\sin A \sin B \sin C \leq \frac{3\sqrt{3}}{8}$



Watch Video Solution

5. In triangle ABC, prove that  $\sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} \leq \frac{1}{8}$  and hence, prove that  $\operatorname{cosec} \frac{A}{2} + \operatorname{cosec} \frac{B}{2} + \operatorname{cosec} \frac{C}{2} \geq 6$ .



Watch Video Solution

### Exercise (Single)

1. If  $\cos(A - B) = \frac{3}{5}$  and  $\tan A \tan B = 2$  then

A.  $\cos A \cos B = \frac{1}{5}$

B.  $\sin A \sin B = -\frac{2}{5}$

C.  $\cos A \cos B = -\frac{1}{5}$

D.  $\sin A \sin B = 1/5$

**Answer: A**



**Watch Video Solution**

2. If  $A = \sin 45^\circ + \cos 45^\circ$  and  $B = \sin 44^\circ$ , then  $A > B$  (b) A

**A.  $A > B$**

**B.  $A < B$**

**C.  $A = B$**

**D. none of these.**

**Answer: A**



**Watch Video Solution**

3.  $\tan 100^\circ + \tan 125^\circ + \tan 100^\circ \tan 125^\circ$  is equal to

**A. 0**

B.  $\frac{1}{2}$

C.  $-1$

D.  $1$

**Answer: D**

 [Watch Video Solution](#)

4. If  $\cos(\alpha + \beta) = 0$  then  $\sin(\alpha + 2\beta) =$

A.  $-\sin \alpha$

B.  $\sin \beta$

C.  $\cos \alpha$

D.  $\cos \beta$

**Answer: D**

 [Watch Video Solution](#)

5. In triangle  $ABC$ , if  $\sin A \cos B = \frac{1}{4}$  and  $3\tan A = \tan B$ , then  $\cot^2 A$  is equal to

A. 2

B. 3

C. 4

D. 5

Answer: B



Watch Video Solution

6. Let  $\frac{\sin(\theta - \alpha)}{\sin(\theta - \beta)} = \frac{a}{b}$  and  $\frac{\cos(\theta - \alpha)}{\cos(\theta - \beta)} = \frac{c}{d}$  then  $\frac{ac + bd}{ad + bc} =$

A.  $\cos(\alpha - \beta)$

B.  $\sin(\alpha - \beta)$

C.  $\cos(\alpha + \beta)$

D. none of these.

Answer: A



Watch Video Solution

7. If  $A, B, C$  are angles of a triangle, then

$$2 \frac{\sin A}{2} \cos ec \frac{B}{2} \frac{\sin C}{2} - \sin A \frac{\cot B}{2} - \cos A \text{ is independent of } A, B, C \text{ (b)}$$

function of  $A, B$  function of  $C$  (d) none of these

A. independent of  $A, B, C$

B. function of  $A, B$

C. function of  $C$

D. none of these.

Answer: A



Watch Video Solution

8. If  $a \leq 3 \cos x + 5 \sin \left( x - \frac{\pi}{6} \right) \leq b$  for all  $x$ , then  $(a, b)$  is

A.  $(-\sqrt{19}, \sqrt{19})$

B.  $(-17, 17)$

C.  $(-\sqrt{21}, \sqrt{21})$

D. none of these.

Answer: A

 [Watch Video Solution](#)

9. If  $\frac{x}{\cos \theta} = \frac{y}{\cos\left(\theta - \frac{2\pi}{3}\right)} = \frac{z}{\cos\left(\theta + \frac{2\pi}{3}\right)}$ , then  $x + y + z =$

A. 1

B. 0

C. -1

D. none of these.

Answer: B

 [Watch Video Solution](#)



10. Let  $x = \sin 1^\circ$  then find the value of the expression

$$\frac{1}{\cos 0^\circ \cos 1^\circ} + \frac{1}{\cos 1^\circ \cos 2^\circ} + \dots + \frac{1}{\cos 44^\circ \cos 45^\circ}$$

A.  $x$

B.  $1/x$

C.  $\sqrt{2}/x$

D.  $x/\sqrt{2}$

Answer: B



Watch Video Solution

11. If  $\theta$  is eliminated from the equations  $x = a \cos(\theta - \alpha)$  and  $y = b \cos(\theta - \beta)$ , then  $\sec^2(\alpha - \beta)$  (b)  $\cos ec^2(\alpha - \beta)$  (d)  $\sin^2(\alpha - \beta)$

A.  $\sec^2(\alpha - \beta)$

B.  $\operatorname{cosec}^2(\alpha - \beta)$

C.  $\cos^2(-\beta)$

D.  $\sin^2(\alpha - \beta)$

**Answer: D**



**Watch Video Solution**

12. The minimum vertical distance between the graphs of

$y = 2 + \sin x$  and  $y = \cos x$  is **(a) 2 (b) 1 (c)  $\sqrt{2}$  (d)  $2 - \sqrt{2}$**

A. 2

B. 1

C.  $\sqrt{2}$

D.  $2 - \sqrt{2}$

**Answer: D**



**Watch Video Solution**

13. If  $\frac{\tan^2(\pi - A)}{4} + \frac{\tan^2(\pi - B)}{4} + \frac{\tan^2(\pi - C)}{4} = 1$ , then  $ABC$  is equilateral (b) isosceles (c) scalene (d) none of these

A. equilateral

B. isosceles

C. scalene

D. none of these.

Answer: A



Watch Video Solution

14. if  $(1+\tan\alpha)(1+\tan4\alpha) = 2$  where  $\alpha \in (0, \frac{\pi}{16})$  then  $\alpha$  equal to

A.  $\frac{\pi}{20}$

B.  $\frac{\pi}{30}$

C.  $\frac{\pi}{40}$

D.  $\frac{\pi}{60}$

Answer: A



Watch Video Solution

15. If  $\cos 28^\circ + \sin 28^\circ = k^3$ , then  $\cos 17^\circ$  is equal to  $\frac{k^3}{\sqrt{2}}$  (b)  $-\frac{k^3}{\sqrt{2}}$  (c)  $\pm \frac{k^3}{\sqrt{2}}$  (d) none of these

A.  $\frac{k^3}{\sqrt{2}}$

B.  $-\frac{k^3}{\sqrt{2}}$

C.  $\pm \frac{k^3}{\sqrt{2}}$

D. none of these.

Answer: A



Watch Video Solution

16. Let  $f(\theta) = \frac{\cot \theta}{1 + \cot \theta}$  and  $\alpha + \beta = \frac{5\pi}{4}$  then the value  $f(\alpha)f(\beta)$  is

A.  $\frac{1}{2}$

B.  $-\frac{1}{2}$

C. 2

D. none of these.

Answer: A



Watch Video Solution

17. If  $y = (1 + \tan A)(1 - \tan B)$ , where  $A - B = \frac{\pi}{4}$  then  $(y + 1)^{y+1}$  is

equal to

A. 9

B. 4

C. 27

D. 81

**Answer: C**



**Watch Video Solution**

18. If  $\frac{\sin x}{\sin y} = \frac{1}{2}$ ,  $\frac{\cos x}{\cos y} = \frac{3}{2}$ , where  $x, y \in \left(0, \frac{\pi}{2}\right)$ , then the value of  $\tan(x + y)$  is equal to

A.  $\sqrt{13}$

B.  $\sqrt{14}$

C.  $\sqrt{17}$

D.  $\sqrt{15}$

**Answer: D**



**Watch Video Solution**

19. If  $\cot^2 x = \cot(x - y)\cot(x - z)$ , then  $\cot 2x$  is equal to  
(where  $x \neq \frac{\pi}{4}$ ) .  $\frac{1}{2}(\tan y + \tan z)$  (b)  $\frac{1}{2}(\cot y + \cot z)$

$\frac{1}{2}(\sin y + \sin z)$  (d) none of these

A.  $\frac{1}{2}(\tan y + \tan x)$

B.  $\frac{1}{2}(\cot y + \cot z)$

C.  $\frac{1}{2}(\sin y + \sin z)$

D. none of these.

Answer: B



Watch Video Solution

20. In a  $ABC$ , if  $\tan A : \tan B : \tan C = 3 : 4 : 5$ , then the value of

$\sin A \sin B \sin C$  is equal to (a)  $\frac{2}{\sqrt{5}}$  (b)  $\frac{2\sqrt{5}}{7}$  (c)  $\frac{2\sqrt{5}}{9}$  (d)  $\frac{2}{3\sqrt{5}}$

A.  $\frac{2}{\sqrt{5}}$

B.  $\frac{2\sqrt{5}}{7}$

C.  $\frac{2\sqrt{5}}{9}$

D.  $\frac{2}{3\sqrt{5}}$

Answer: B

 Watch Video Solution

21. find the value of  $\sin 27^\circ - \cos 27^\circ$  ?

A.  $-\frac{\sqrt{3 - \sqrt{3}}}{2}$

B.  $-\frac{\sqrt{5 - \sqrt{5}}}{2}$

C.  $-\frac{\sqrt{5} - 1}{2\sqrt{2}}$

D. none of these.

Answer: B

 Watch Video Solution

22. If  $\cos \theta_1 = 2 \cos \theta_2$ , then  $\frac{\tan(\theta_1 - \theta_2)}{2} \frac{\tan(\theta_1 + \theta_2)}{2}$  is equal to  $\frac{1}{3}$  (b)  $-\frac{1}{3}$

1 (d)  $-1$



A.  $\frac{1}{3}$

B.  $-\frac{1}{3}$

C. 1

D. -1

Answer: B

 Watch Video Solution

23. Let  $\alpha, \beta$  be such that it  $\pi < \alpha - \beta < 3\pi$ . if  $\sin \alpha + \sin \beta = -\frac{21}{65}$  and  $\cos \alpha + \cos \beta = -\frac{27}{65}$  then the value of  $\cos\left(\frac{\alpha - \beta}{2}\right)$  is

A.  $-\frac{3}{\sqrt{130}}$

B.  $\frac{3}{\sqrt{130}}$

C.  $\frac{6}{65}$

D.  $-\frac{6}{65}$

Answer: A



Watch Video Solution

24. If  $n = \frac{\pi}{4\alpha}$ , then  $\tan \alpha \tan 2\alpha \tan 3\alpha \dots \tan(2n - 1)\alpha$  is equal to

A. 1

B.  $1/2$

C. 2

D.  $1/3$

Answer: A



Watch Video Solution

25.  $\frac{\sin 3\theta + \sin 5\theta + \sin 7\theta + \sin 9\theta}{\cos 3\theta + \cos 5\theta + \cos 7\theta + \cos 9\theta}$  is equal to

A.  $\tan 3\theta$

B.  $\cot 3\theta$

C.  $\tan 6\theta$

D.  $\cot 6\theta$

Answer: C



Watch Video Solution

26. If  $x, y, z$  are in A.P., then  $\frac{\sin x - \sin z}{\cos z - \cos x}$  is equal to  $\tan y$  (b)  $\cot y$  (c)  $\sin y$  (d)  $\cot y$

A.  $\tan y$

B.  $\cot y$

C.  $\sin y$

D.  $\cos y$

Answer: B



Watch Video Solution

27. If  $\frac{\cos x}{a} = \frac{\cos(x + \theta)}{b} = \frac{\cos(x + 2\theta)}{c} = \frac{\cos(x + 3\theta)}{d}$  then  $\frac{a + c}{b + d}$  is equal to (A)  $\frac{a}{d}$  (B)  $\frac{c}{d}$  (C)  $\frac{b}{c}$  (D)  $\frac{d}{a}$

A.  $\frac{a}{d}$

B.  $\frac{C}{b}$

C.  $\frac{b}{c}$

D.  $\frac{d}{a}$

Answer: C



Watch Video Solution

28. If  $\cos \alpha + \cos \beta = 0 = \sin \alpha + \sin \beta$ , then  $\cos 2\alpha + \cos 2\beta =$

A.  $-2 \sin(\alpha + \beta)$

B.  $-2 \cos(\alpha + \beta)$

C.  $2 \sin(\alpha + \beta)$

D.  $2 \cos(\alpha + \beta)$

**Answer: B**



**Watch Video Solution**

29. Value of  $\frac{3 + \cot 80^\circ \cot 20^\circ}{\cot 80^\circ + \cot 20^\circ}$  is equal to

A.  $\cot 20^\circ$

B.  $\tan 50^\circ$

C.  $\cot 50^\circ$

D.  $\cot \sqrt{20^\circ}$

**Answer: B**



**Watch Video Solution**

30. If  $\tan \alpha$  is equal to the integral solution of the inequality  $4x^2 - 16x + 15 < 0$  and  $\cos \beta$  is equal to the slope of the bisector of the first quadrant, then  $\sin(\alpha + \beta)\sin(\alpha - \beta)$  is equal to  $\frac{3}{5}$  (b)  $\frac{3}{5}$  (c)  $\frac{2}{\sqrt{5}}$  (d)  $\frac{4}{5}$

A.  $\frac{3}{5}$

B.  $\frac{3}{5}$

C.  $\frac{2}{\sqrt{5}}$

D.  $\frac{4}{5}$

Answer: D



Watch Video Solution

31. Let  $f(n) = 2 \cos nx \forall n \in N$ , then  $f(1)f(n+1) - f(n)$  is equal to  $f(n+3)$  (b)  $f(n+2) f(n+1)f(2)$  (d)  $f(n+2)f(2)$

A.  $f(n+3)$

B.  $f(n + 2)$

C.  $f(n + 1)f(2)$

D.  $f(n + 2)f(2)$

**Answer: B**



**Watch Video Solution**

32. If  $\sin \theta_1 \sin \theta_2 - \cos \theta_1 \cos \theta_2 + 1 = 0$ , then the value of  $\tan\left(\frac{\theta_1}{2}\right) \cot\left(\frac{\theta_2}{2}\right)$  is equal to -1 (b) 1 (c) 2 (d) -2

A.  $a^2 + b^2 \geq 4$

B.  $a^2 + b^2 \leq 4$

C.  $a^2 + b^2 \geq 3$

D.  $a^2 + b^2 \leq 2$

**Answer: B**



**Watch Video Solution**

33.  $\frac{\sqrt{2} - \sin \alpha - \cos \alpha}{\sin \alpha - \cos \alpha}$  is equal to

A.  $\sec\left(\frac{\alpha}{2} - \frac{\pi}{8}\right)$

B.  $\cos\left(\frac{\pi}{8} - \frac{\alpha}{2}\right)$

C.  $\tan\left(\frac{\alpha}{2} - \frac{\pi}{8}\right)$

D.  $\cot\left(\frac{\alpha}{2} - \frac{\pi}{2}\right)$

Answer: C



Watch Video Solution

34. If  $x_1$  and  $x_2$  are two distinct roots of the equation

$a \cos x + b \sin x = c$ , then  $\tan \frac{x_1 + x_2}{2}$  is equal to

A.  $\frac{a}{b}$

B.  $\frac{b}{a}$

C.  $\frac{c}{a}$



D.  $\frac{a}{c}$

Answer: B



Watch Video Solution

35. If  $\sin(y + z - x)$ ,  $\sin(z + x - y)$ ,  $\sin(x + y - z)$  are in A.P., then  $\tan x$ ,  $\tan y$ ,  $\tan z$  are in A.P. (b) G.P. (c) H.P. (d) none of these

A. AP

B. GP

C. HP

D. none of these.

Answer: A



Watch Video Solution

36. If  $\frac{\tan(\alpha + \beta + \gamma)}{\tan(\alpha - \beta - \gamma)} = \frac{\tan \gamma}{\tan \beta}$ , ( $\beta \neq \gamma$ ) then

$$\sin 2\alpha + s \in 2\beta + s \in 2\gamma = \mathbf{0 (b) 1 (c) 2 (d)}$$

A. 0

B. 1

C. 2

D. 1/2

Answer: A



Watch Video Solution

37. If  $\sin \theta_1 \sin \theta_2 - \cos \theta_1 \cos \theta_2 + 1 = 0$ , then the value of  $\tan\left(\frac{\theta_1}{2}\right) \cot\left(\frac{\theta_2}{2}\right)$  is equal to  $-1 (b) 1 (c) 2 (d) -2$

A. -1

B. 1

C. 2

D.  $-2$

Answer: A

 Watch Video Solution

38. The value of the expression

$$\frac{2(\sin 1^\circ + \sin 2^\circ + \sin 3^\circ + \dots + \sin 89^\circ)}{2(\cos 1^\circ + \cos 2^\circ + \dots + \cos 44^\circ) + 1} \text{ equals}$$

A.  $\sqrt{2}$

B.  $\frac{1}{\sqrt{2}}$

C.  $\frac{1}{2}$

D. 0

Answer: A

 Watch Video Solution

39. If  $A, B, C$ , are the angles of a triangle such that  $\frac{\cot A}{2} = 3\frac{\tan C}{2}$ , then  $\sin A, \sin B, \sin C$  are in  $AP$ . (b)  $GP$ . (c)  $HP$ . (d) none of these

A. AP

B. GP

C. HP

D. none of these.

Answer: A



Watch Video Solution

40. If  $2 \sec 2\theta = \tan \phi + \cot \phi$ , then one of the values of  $\theta + \phi$  is

A.  $\frac{\pi}{2}$

B.  $\frac{\pi}{4}$

C.  $\frac{\pi}{3}$

D. none of these

**Answer: B**



**Watch Video Solution**

**41. The roots of the equation  $4x^2 - 2\sqrt{5}x + 1 = 0$  are .**

**A.**  $\sin 36^\circ, \sin 18^\circ$

**B.**  $\sin 18^\circ, \cos 36^\circ$

**C.**  $\sin 36^\circ, \cos 18^\circ$

**D.**  $\cos 18^\circ, \cos 36^\circ$

**Answer: B**



**Watch Video Solution**

**42. If  $A$  and  $B$  are acute positive angles satisfying the equations**

**$3 \sin^2 A + 2 \sin^2 B = 1$  and  $3 \sin 2A - 2 \sin 2B = 0$ , then  $A + 2B$  is**

**equal to  $\pi$  (b)  $\frac{\pi}{2}$  (c)  $\frac{\pi}{4}$  (d)  $\frac{\pi}{6}$**

A.  $\pi$

B.  $\frac{\pi}{2}$

C.  $\frac{\pi}{4}$

D.  $\frac{\pi}{6}$

Answer: B

 Watch Video Solution

43. If  $\cos 25^\circ + \sin 25^\circ = p$ , then  $\cos 50^\circ$  is equal to

A.  $\sqrt{2 - p^2}$

B.  $-\sqrt{2 - p^2}$

C.  $p\sqrt{2 - p^2}$

D.  $-p\sqrt{2 - p^2}$

Answer: C

 Watch Video Solution

44. The value of  $\frac{\cot(7\pi)}{16} + 2\frac{\cot(3\pi)}{8} + \frac{\cot(15\pi)}{16}$  is 4 (b) 2 (c) -2 (d)

-4

A. 4

B. 2

C. -2

D. -4

Answer: D



Watch Video Solution

45. If  $\alpha, \beta, \gamma, \delta$  are the smallest positive angles in ascending order of magnitude which have their sines equal to the positive quantity  $k$ , then

the value of  $4\frac{\sin \alpha}{2} + 3\frac{\sin \beta}{2} + 2\frac{\sin \gamma}{2} + \frac{\sin \delta}{2}$  is equal to  $2\sqrt{1-k}$  (b)

$2\sqrt{1+k} \frac{\sqrt{1-k}}{2}$  (d) none of these

A.  $2\sqrt{1 - k}$

B.  $2\sqrt{1 + k}$

C.  $\frac{\sqrt{1 + k}}{2}$

D. none of these.

Answer: B



Watch Video Solution

46.  $\frac{\sin^2 A - \sin^2 B}{\sin A \cos A - \sin B \cos B}$  is equal to  $\tan(A - B)$  (b)  $\tan(A + B)$   
 $\cot(A - B)$  (d)  $\cot(A + B)$

A.  $\tan(A - B)$

B.  $\tan(A + B)$

C.  $\cot(A - B)$

D.  $\cot(A + B)$

Answer: B





Watch Video Solution

47. If  $\cos(\alpha - \beta) = 3\sin(\alpha + \beta)$ , then  $\frac{1}{1 - 3\sin 2\alpha} + \frac{1}{1 - 3\sin 2\beta} =$   
 $\frac{1}{2}$  (b)  $\frac{-1}{2}$  (c)  $\frac{1}{4}$  (d)  $\frac{-1}{4}$

A.  $\frac{1}{2}$

B.  $\frac{-1}{2}$

C.  $\frac{1}{4}$

D.  $\frac{-1}{4}$

Answer: D



Watch Video Solution

48. The value of  $\cos^2 10^\circ - \cos 10^\circ \cos 50^\circ + \cos^2 50^\circ$  is equal to  $\frac{4}{3}$  (b)  $\frac{1}{3}$   
(c)  $\frac{3}{4}$  (d) 3

A.  $\frac{4}{3}$

B.  $\frac{1}{3}$

C.  $\frac{3}{4}$

D. 3

Answer: C



Watch Video Solution

49. If  $\tan^2\theta = 2\tan^2\varphi + 1$ , prove that  $\cos 2\theta + \sin^2\varphi = 0$ .

A. -1

B. 0

C. 1

D. none of these.

Answer: B



Watch Video Solution

50. If  $\sin x + \operatorname{cosec} x + \tan y = 4$  where  $x$  and  $y \in \left[0, \frac{\pi}{2}\right]$

then  $\tan \frac{y}{2}$  is a root of the equation.

A.  $a^2 + 2a + 1 = 0$

B.  $a^2 + 2a = 0$

C.  $2a^2 - 2a - 1 = 0$

D.  $a^2 - a - 1 = 0$

Answer: D



[View Text Solution](#)

51. If  $2 \sin 2\alpha = \tan \beta$ ,  $\alpha, \beta \in \left(\frac{\pi}{2}, \pi\right)$ , then the value of  $\alpha + \beta$  is

A.  $\frac{3\pi}{4}$

B.  $\pi$

C.  $\frac{3\pi}{2}$

D.  $\frac{5\pi}{4}$

**Answer: B**



**Watch Video Solution**

52. If  $2|\sin 2\alpha| = \tan \beta + \cot \beta$ ,  $\alpha, \beta \in \left(\frac{\pi}{2}, \pi\right)$ , then the value of  $\alpha + \beta$  is **(a)**  $\frac{3\pi}{4}$  **(b)**  $\pi$  **(c)**  $\frac{3\pi}{2}$  **(d)**  $\frac{5\pi}{4}$

**A.**  $\frac{3\pi}{4}$

**B.**  $\pi$

**C.**  $\frac{3\pi}{2}$

**D.**  $\frac{5\pi}{4}$

**Answer: C**



**Watch Video Solution**

53. The value of  $\sin^3 10^\circ + \sin^3 50^\circ - \sin^3 70^\circ$  is equal to

A.  $-\frac{3}{2}$

B.  $\frac{3}{4}$

C.  $-\frac{3}{4}$

D.  $-\frac{3}{8}$

Answer: D



Watch Video Solution

54. Let  $P(x) = \left( \frac{1 - \cos 2x + \sin 2x}{1 + \cos 2x + \sin 2x} \right)^2 + \left( \frac{1 + \cot x + \cot^2 x}{1 + \tan x + \tan^2 x} \right)$ ,

then the minimum value of  $P(x)$  equal 1 (b) 2 (c) 4 (d) 16

A. 1

B. 2

C. 4

D. 16

Answer: B



Watch Video Solution

55. If  $\frac{3 - \tan^2 \frac{\pi}{7}}{1 - \tan^2 \frac{\pi}{7}} = k \cos \frac{\pi}{7}$  then the value of k is

A. 1

B. 2

C. 3

D. 4

Answer: D



Watch Video Solution

56.  $\cos ec \frac{360^0}{7} + \cos ec \frac{540^0}{7} = \cos ec \frac{180^0}{7}$  (b)  $\cos ec \frac{90^0}{7} \frac{\sec(180^0)}{7}$   
(d)  $\frac{\sec(90^0)}{7}$

A.  $\cos ec \left( \frac{180^0}{7} \right)$

B.  $\operatorname{cosec}\left(\frac{90^\circ}{7}\right)$

C.  $\sec\left(\frac{180^\circ}{7}\right)$

D.  $\sec\left(\frac{90^\circ}{7}\right)$

**Answer: A**



**Watch Video Solution**

**57. If  $\theta_1$  and  $\theta_2$  are two values lying in  $[2, 2\pi]$  for which  $\tan\theta = \lambda$ , then**

**$\frac{\tan(\theta_1)}{2} \frac{\tan(\theta_2)}{2}$  is equal to 0 (b)  $-1$  (c)  $2$  (d)  $1$**

**A. 0**

**B.  $-1$**

**C. 2**

**D. 1**

**Answer: B**



**Watch Video Solution**

58. If  $\tan \theta = \sqrt{n}$ , where  $n \in N, \geq 2$ , then  $\sec 2\theta$  is always a rational number (b) an irrational number a positive integer (d) a negative integer

A. a rational number

B. an irrational number

C. a positive integer

D. a negative integer.

Answer: A



Watch Video Solution

59. If  $\sin 2\theta = \cos 3\theta$  and  $\theta$  is an acute angle, then  $\sin \theta$  is equal to

A.  $\frac{\sqrt{5} - 1}{4}$

B.  $-\left(\frac{\sqrt{5} - 1}{4}\right)$

C.  $\frac{\sqrt{5} + 1}{4}$



D.  $\frac{-\sqrt{5}-1}{4}$

Answer: A



Watch Video Solution

60. If  $\cos x = \tan y$ ,  $\cos y = \tan z$  and  $\cos z = \tan x$ , prove that  $\sin x = \sin y = \sin z = \sin 18^\circ$

A.  $2\cos 18^\circ$

B.  $\cos 18^\circ$

C.  $\sin 18^\circ$

D.  $2\sin 18^\circ$

Answer: D



Watch Video Solution

61. The value of  $\cot 70^\circ + 4\cos 70^\circ$  is

A.  $\frac{1}{\sqrt{3}}$

B.  $\sqrt{3}$

C.  $2\sqrt{3}$

D.  $\frac{1}{2}$

Answer: B



Watch Video Solution

62. If  $\sin x + \cos x = \frac{\sqrt{7}}{2}$  where  $x \in \left[0, \frac{\pi}{4}\right]$  then  $\tan\left(\frac{x}{2}\right)$  is equal to

A.  $\frac{3 - \sqrt{7}}{3}$

B.  $\frac{\sqrt{7} - 2}{3}$

C.  $\frac{4 - \sqrt{7}}{4}$

D. none of these.

Answer: B

 Watch Video Solution

63. If  $\frac{\tan 3A}{\tan A} = k (k \neq 1)$  then which of the following is not true?

$\frac{\cos A}{\cos 3A} = \frac{k-1}{2}$  (b)  $\frac{\sin 3A}{\sin A} = \frac{2k}{k-1}$   $\frac{\cot 3A}{\cot A} = \frac{1}{k}$  (d) none of these

A.  $\frac{\cos A}{\cos 3A} = \frac{k-1}{2}$

B.  $\frac{\sin 2A}{\sin A} = \frac{2k}{k-1}$

C.  $\frac{\cos 3A}{\cot A} = \frac{1}{k}$

D. none of these.

Answer: D

 Watch Video Solution

64. If  $x \in \left(\pi, \frac{3\pi}{2}\right)$ , then  $4\cos^2\left(\frac{\pi}{4} - \frac{x}{2}\right) + \sqrt{4\sin^4 x + \sin^2 2x}$  is always equal to 1 (b) 2 (c) -2 (d) none of these

A. 1

B. 2

C. -2

D. none of these.

Answer: B



Watch Video Solution

65. If  $\cos x = \frac{2 \cos y - 1}{2 - \cos y}$ , where  $x, y \in (0, \pi)$  then  $\frac{\tan x}{2} \frac{\cot y}{2}$  is equal to  $\sqrt{2}$  (b)  $\sqrt{3}$  (c)  $\frac{1}{\sqrt{2}}$  (d)  $\frac{1}{\sqrt{3}}$

A.  $\sqrt{2}$

B.  $\sqrt{3}$

C.  $\frac{1}{\sqrt{2}}$

D.  $\frac{1}{\sqrt{3}}$

Answer: B



Watch Video Solution

66.  $\cot 16^\circ \cot 44^\circ + \cot 44^\circ \cot 76^\circ - \cot 76^\circ \cot 16^\circ = 1$  (b) 2 (c) 3 (d) 4

A. 1

B. 2

C. 3

D. 4

Answer: C



Watch Video Solution

67. If  $\tan x = \frac{b}{a}$  then  $\sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}}$  is equal to

A.  $2 \sin x / \sqrt{2 \sin 2x}$

B.  $2 \cos x / \sqrt{\cos 2x}$

C.  $2 \cos x / \sqrt{\sin 2x}$

D.  $2 \sin x / \sqrt{\cos 2x}$

**Answer: B**



**Watch Video Solution**

**68. Given that  $(1 + \sqrt{1+x})\tan y = 1 + \sqrt{1-x}$ . Then  $\sin 4y$  is equal to  $4x$  (b)  $2x$  (c)  $x$  (d) none of these**

**A.  $4x$**

**B.  $2x$**

**C.  $x$**

**D. none of these.**

**Answer: C**



**Watch Video Solution**

69. If  $\cos 2B = \frac{\cos(A + C)}{\cos(A - C)}$ , then  $\tan A, \tan B, \tan C$  are in

A. AP

B. GP

C. HP

D. none of these.

Answer: B



Watch Video Solution

70. If  $\frac{\cos(x - y)}{\cos(+y)} + \frac{\cos(z + t)}{\cos(z - t)} = 0$ , then the value of expression

$\tan x \tan y \tan z \tan t$  is equal to 1 (b)  $-1$  (c)  $2$  (d)  $-2$

A. 1

B.  $-1$

C. 2

D.  $-2$

**Answer: B**



**Watch Video Solution**

**71. If  $\tan \beta = 2 \sin \alpha \sin \gamma \csc(\alpha + \gamma)$ , then  $\cot \alpha, \cot \beta, \cot \gamma$  are in**

**A. AP**

**B. GP**

**C. HP**

**D. none of these.**

**Answer: A**



**Watch Video Solution**

**72. the value of  $\tan 9^\circ - \tan 27^\circ - \tan 63^\circ + \tan 81^\circ$  is equal to**

**A. 2**



B. 3

C. 4

D. none of these.

Answer: C



Watch Video Solution

73. If  $\cos^3 x \sin 2x = \sum_{r=0}^n a_x \sin(rx)$ ,  $\forall x \in R$  then

A.  $n = 5, a_1 = 1/2$

B.  $n = 5, a_1 = 1/4$

C.  $n = 5, a_2 = 1/8$

D.  $n = 5, a_2 = 1/4$

Answer: B



Watch Video Solution

74.  $\frac{\cos^6 \pi}{9} - 33\frac{\tan^4 \pi}{9} + 27\frac{\tan^2 \pi}{9}$  is equal to 0 (b)  $\sqrt{3}$  (c) 3 (d) 9

A. 0

B.  $\sqrt{3}$

C. 3

D. 9

Answer: C



Watch Video Solution

75. Given that  $a, b, c,$  are the side of a  $ABC$  which is right angled at  $C,$  then the minimum value of  $\left(\frac{c}{a} + \frac{c}{b}\right)^2$  is 0 (b) 4 (c) 6 (d) 8

A. 0

B. 4

C. 6

D. 8

Answer: D



Watch Video Solution

76. If  $\theta = 3\alpha$  and  $\sin \theta = \frac{a}{\sqrt{a^2 + b^2}}$ , the value of the expression  $a \cos \alpha - b \sec \alpha$  is

A.  $\frac{a}{\sqrt{a^2 + b^2}}$

B.  $2\sqrt{a^2 + b^2}$

C.  $a+b$

D. none of these.

Answer: B



Watch Video Solution

77. The value of  $\tan 6^\circ \tan 42^\circ \tan 66^\circ \tan 78^\circ$  is

A. 1

B.  $1/2$

C.  $1/4$

D.  $1/8$

Answer: A



Watch Video Solution

78. In triangle  $ABC$ , if angle is  $90^\circ$  and the area of triangle is  $30^0$  sq. units, then the minimum possible value of the hypotenuse  $c$  is equal to  $30\sqrt{2}$  (b)  $60\sqrt{2}$  (c)  $120\sqrt{2}$  (d)  $2\sqrt{30}$

A.  $30\sqrt{2}$

B.  $60\sqrt{2}$

C.  $120\sqrt{2}$

D.  $2\sqrt{30}$

Answer: D



Watch Video Solution

79. If  $\sqrt{2} \cos A = \cos B + \cos^3 B$ , and  $\sqrt{2} \sin A = \sin B - \sin^3 B$  then

$\sin(A - B) =$

A.  $\pm 1$

B.  $\pm \frac{1}{2}$

C.  $\pm \frac{1}{3}$

D.  $\pm \frac{1}{4}$

Answer: C



Watch Video Solution

80. In a right angled triangle the hypotenuse is  $2\sqrt{2}$  times the perpendicular distance drawn from the opposite vertex. then the other acute angles of the triangle are

A.  $\frac{\pi}{3}$  and  $\frac{\pi}{6}$

B.  $\frac{\pi}{8}$  and  $\frac{3\pi}{8}$

C.  $\frac{\pi}{4}$  and  $\frac{\pi}{4}$

D.  $\frac{\pi}{5}$  and  $\frac{3\pi}{10}$

Answer: B



Watch Video Solution

81. A circular ring of radius 3cm hangs horizontally from a point 4cm vertically above the centre by 4 strings attached at equal intervals to its circumference. If the angle between two consecutive strings be  $\theta$ , then

$\cos \theta$  is equal to  $\frac{4}{5}$  (b)  $\frac{4}{25}$  (d)  $\frac{16}{25}$  (d) none of these

A.  $\frac{4}{5}$

B.  $\frac{4}{25}$

C.  $\frac{16}{25}$

D. none of these.

Answer: C

 [Watch Video Solution](#)

82. The distance between the two parallel lines is 1 unit. A point A is chosen to lie between the lines at a distance 'd' from one of them. Triangle ABC is equilateral with B on one line and C on the other parallel line. The length of the side of the equilateral triangle is

A.  $\frac{2}{3}\sqrt{d^2 + d + 1}$

B.  $2\sqrt{\frac{d^2 - d + 1}{3}}$

C.  $2\sqrt{d^2 - d + 1}$

D.  $\sqrt{d^2 - d + 1}$

Answer: B



Watch Video Solution

83. If  $\sin^{-1} a + \sin^{-1} b + \sin^{-1} c = \pi$ , then  $a\sqrt{1-a^2} + b\sqrt{1-b^2} + c\sqrt{1-c^2}$  is equal to  $a + b + c$  (b)  $a^2b^2c^2$   $2abc$  (d)  $4abc$

A.  $a+b+c$

B.  $a^2b^2c^2$

C.  $2abc$

D.  $4abc$

Answer: C



Watch Video Solution

84. If  $A + B + C = 3\frac{\pi}{2}$ . Then  $\cos 2A + \cos 2B + \cos 2C$  is equal to



A.  $1 - 4 \cos A \cos B \cos C$

B.  $4 \sin A \sin B \sin C$

C.  $1 + 2 \cos A \cos B \cos C$

D.  $1 - 4 \sin A \sin B \sin C$

**Answer: D**

 [Watch Video Solution](#)

85. If  $\tan(\alpha - \beta) = \frac{\sin 2\beta}{3 - \cos 2\beta}$ , then

A.  $\tan \alpha = 2 \tan \beta$

B.  $\tan \beta = 2 \tan \alpha$

C.  $2 \tan \alpha = 3 \tan \beta$

D.  $3 \tan \alpha = 2 \tan \beta$

**Answer: A**

 [Watch Video Solution](#)

86. In any triangle  $ABC$ ,  $\sin^2 A - \sin^2 B + \sin^2 C$  is always equal to  
 $2 \sin A \sin B \cos C$    (b)    $2 \sin A \cos B \sin C$     $2 \sin A \cos B \cos C$    (d)  
 $2 \sin A \sin B \sin C$

A.  $2 \sin A \sin B \cos C$

B.  $2 \sin A \cos B \sin C$

C.  $2 \sin A \cos B \cos C$

D.  $2 \sin A \sin B \sin C$

Answer: B

 [Watch Video Solution](#)

87. The value of  $\sum_{r=0}^{10} \cos^3\left(\frac{r\pi}{3}\right)$  is

A.  $1/4$

B.  $1/8$

C.  $-1/4$

D.  $-1/8$

**Answer: D**



**Watch Video Solution**

88. In triangle ABC,  $\frac{\sin A + \sin B + \sin C}{\sin A + \sin B - \sin C}$  is equal to

A.  $\tan\left(\frac{A}{2}\right) \cot\left(\frac{B}{2}\right)$

B.  $\cot\left(\frac{A}{2}\right) \tan\left(\frac{B}{2}\right)$

C.  $\cot\left(\frac{A}{2}\right) \cot\left(\frac{B}{2}\right)$

D.  $\tan\left(\frac{A}{2}\right) \tan\left(\frac{B}{2}\right)$

**Answer: C**



**Watch Video Solution**

89.  $\frac{\sin 2A + \sin 2B + \sin 2C}{\sin A + \sin B + \sin C}$  is equal to

A.  $8 \sin\left(\frac{A}{2}\right) \sin\left(\frac{B}{2}\right) \sin\left(\frac{C}{2}\right)$

B.  $8 \cos\left(\frac{A}{2}\right) \cos\left(\frac{B}{2}\right) \cos\left(\frac{C}{2}\right)$

C.  $8 \tan\left(\frac{A}{2}\right) \tan\left(\frac{B}{2}\right) \tan\left(\frac{C}{2}\right)$

D.  $8 \cot\left(\frac{A}{2}\right) \cot\left(\frac{B}{2}\right) \cot\left(\frac{C}{2}\right)$

Answer: A



Watch Video Solution

90. If  $\cos^2 A + \cos^2 B + \cos^2 C = 1$ , then  $\triangle ABC$  is

A. equilateral

B. isosceles

C. right angled

D. none of these.

**Answer: C**



**Watch Video Solution**

**91. In a triangle  $\tan A + \tan B + \tan C = 6$  and  $\tan A \tan B = 2$ , then the values of  $\tan A$ ,  $\tan B$  and  $\tan C$  are**

**A. 1,2,3**

**B. 3,  $2/3$ ,  $7/3$**

**C. 4,  $1/2$ ,  $3/2$**

**D. none of these.**

**Answer: A**



**Watch Video Solution**

**92. If  $\cos x + \cos y - \cos(x + y) = \frac{3}{2}$ , then**

A.  $x + y = 0$

B.  $x=2y$

C.  $x=y$

D.  $2x=y$

Answer: B



Watch Video Solution

93. If  $a \sin x + b \cos(x + \theta) + b \cos(x - \theta) = d$ , then the minimum value of  $|\cos \theta|$  is equal to

A.  $\frac{1}{2|b|} \sqrt{d^2 - a^2}$

B.  $\frac{1}{2|a|} \sqrt{d^2 - a^2}$

C.  $\frac{1}{2|d|} \sqrt{d^2 - a^2}$

D. none of these.

Answer: A



Watch Video Solution

94. If  $u = \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} + \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta}$ , then the difference between maximum and minimum values of  $u^2$  is

A.  $2(a^2 + b^2)$

B.  $2\sqrt{a^2 + b^2}$

C.  $(a + b)^2$

D.  $(a - b)^2$

Answer: D



Watch Video Solution

95. If  $\tan x = n \tan y$ ,  $n \in R^+$  then the maximum value of  $\sec^2(x - y)$  is

A.  $\frac{(n + 1)^2}{2n}$

B.  $\frac{(n + 1)^2}{n}$

C.  $\frac{(n+1)^2}{2}$

D.  $\frac{(n+1)^2}{4n}$

Answer: D

 [Watch Video Solution](#)

96. If  $0 \leq x \leq \frac{\pi}{3}$  then range of  $f(x) = \sec\left(\frac{\pi}{6} - x\right) + \sec\left(\frac{\pi}{6} + x\right)$  is

$\left(\frac{4}{\sqrt{3}}, \infty\right)$  (b)  $\left(\frac{4}{\sqrt{3}}, \infty\right)$   $\left(0, \frac{4}{\sqrt{3}}\right)$  (d)  $\left(0, \frac{4}{\sqrt{3}}\right)$

A.  $\left(\frac{4}{\sqrt{3}}, \infty\right)$

B.  $\left[\frac{4}{\sqrt{3}}, \infty\right)$

C.  $\left[0, \frac{4}{\sqrt{3}}\right]$

D.  $\left(0, \frac{4}{\sqrt{3}}\right)$

Answer: B

 [Watch Video Solution](#)



97. The maximum value of  $\cos x \sin x + \sqrt{\sin^2 x + \sin^2\left(\frac{\pi}{6}\right)}$  is

A.  $\frac{\sqrt{5}}{3}$

B.  $\sqrt{\frac{3}{2}}$

C.  $\sqrt{\frac{5}{2}}$

D.  $\frac{\sqrt{5}}{2}$

Answer: D



Watch Video Solution

98. If  $\alpha, \beta, \gamma$  are acute angles and

$$\cos \theta = \sin \beta / \sin \alpha, \cos \varphi = \sin \gamma \sin \alpha \text{ and } \cos(\theta - \varphi) = \sin \beta \sin \gamma,$$

then the value of  $\tan^2 \alpha - \tan^2 \beta - \tan^2 \gamma$  is equal to (a)  $-1$  (b)  $0$  (c)  $1$  (d)  $2$

A.  $-1$

B.  $0$

C.  $1$

Answer: B

 Watch Video Solution

$$99. \sum_{n=1}^{\infty} \frac{\tan\left(\frac{\theta}{2^n}\right)}{2^{n-1} \cos\left(\frac{\theta}{2^{n-1}}\right)}$$

A.  $\frac{2}{\sin 2\theta} - \frac{1}{\theta}$

B.  $\frac{2}{\sin 2\theta} + \frac{1}{\theta}$

C.  $\frac{1}{\sin 2\theta} - \frac{1}{\theta}$

D.  $\frac{1}{\sin \theta} - \frac{1}{\theta}$

Answer: A

 Watch Video Solution

100. If  $x \sin a + y \sin 2a + z \sin 3a = \sin 4a$   
 $x \sin b + y \sin 2b + z \sin 3b = \sin 4b$   $x \sin c + y \sin 2c + z \sin 3c = \sin 4c$

then the roots of the equation

$$t^3 - \left(\frac{z}{2}\right)t^2 - \left(\frac{y+2}{4}\right)t + \left(\frac{z-x}{8}\right) = 0, \quad a, b, c, \neq n\pi, \quad \text{are}$$

$\sin a, \sin b, \sin c$  (b)  $\cos a, \cos b, \cos c$   $\sin 2a, \sin 2b, \sin 2c$  (d)

$\cos 2a, \cos 2b \cos 2c$

A.  $\sin a, \sin b, \sin c$

B.  $\cos a, \cos b, \cos c$

C.  $\sin 2a, \sin 2b, \sin 2c$

D.  $\cos 2a, \cos 2b \cos 2c$

Answer: B



Watch Video Solution

Exercise (Multiple)

1. If  $\cos \beta$  is the geometric mean between  $\sin \alpha$  and  $\cos \alpha$ , where  $0 < \alpha, \beta < \pi/2$ . Then  $\cos 2\beta$  is equal to

A.  $-2 \sin^2\left(\frac{\pi}{4} - \alpha\right)$

B.  $-2 \cos^2\left(\frac{\pi}{4} + \alpha\right)$

C.  $2 \sin^2\left(\frac{\pi}{4} + \alpha\right)$

D.  $2 \cos^2\left(\frac{\pi}{4} - \alpha\right)$

Answer: A:B



Watch Video Solution

2. Which of the following statements are always correct (where  $Q$  denotes

the set of rationals)?  $\cos 2\theta \in Q$  and  $\sin 2\theta \in Q$   $\tan \theta \in Q$

( if  $\sin \theta \in Q$  )  $\tan \theta \in Q$   $\sin 2\theta, \cos 2\theta$  and  $\tan 2\theta \in Q$  ( if  $\sin \theta \in Q$  )

if  $\sin \theta \in Q$  and  $\cos \theta \in Q$   $\tan 3\theta \in Q$  ( if  $\sin \theta \in Q$  ) **fi** **sin**

$\theta \in Q$   $\cos 3\theta \in Q$

A.  $\cos 2\theta \in Q$  and  $\sin 2\theta \in Q \Rightarrow \tan \theta \in Q$  (if defined)

B.  $\tan \theta \in Q \Rightarrow \sin 2\theta, \cos 2\theta$  and  $\tan 2\theta \in Q$  (if denied)

C. if  $\sin \theta \in Q$  and  $\cos \theta \in Q \Rightarrow \tan 3\theta \in Q$  (if denied)

D. if  $\sin \theta \in Q \Rightarrow \cos 3\theta \in Q$

Answer: A::B::C

 Watch Video Solution

3. Which of the following quantities are rational?  $\sin\left(\frac{11\pi}{12}\right)\sin\left(\frac{5\pi}{12}\right)$

(b)  $\operatorname{cosec}\left(\frac{9\pi}{10}\right)\sec\left(\frac{4\pi}{5}\right)$   $\sin^4\left(\frac{\pi}{8}\right) + \cos^4\left(\frac{\pi}{8}\right)$   
 $\left(1 + \frac{\cos(2\pi)}{9}\right)\left(1 + \frac{\cos(4\pi)}{9}\right)\left(1 + \frac{\cos(8\pi)}{9}\right)$

A.  $\sin\left(\frac{11\pi}{12}\right)\sin\left(\frac{5\pi}{12}\right)$

B.  $\operatorname{cosec}\left(\frac{9\pi}{10}\right)\sec\left(\frac{4\pi}{5}\right)$

C.  $\sin^4\left(\frac{\pi}{8}\right) + \cos^4\left(\frac{\pi}{8}\right)$

D.  $\left(1 + \cos\left(\frac{2\pi}{9}\right)\right)\left(1 + \cos\left(\frac{4\pi}{9}\right)\right)\left(1 + \cos\left(\frac{8\pi}{9}\right)\right)$

Answer: A::B::C::D



Watch Video Solution

4. In which of the following sets the inequality  $\sin^6 x + \cos^6 x > \frac{5}{8}$  holds good? (a)  $\left(-\frac{\pi}{3}, \frac{\pi}{8}\right)$  (b)  $\left(\frac{3\pi}{8}, \frac{5\pi}{8}\right)$  (c)  $\left(\frac{\pi}{4}, \frac{3\pi}{4}\right)$  (d)  $\left(\frac{7\pi}{8}, \frac{9\pi}{8}\right)$

A.  $\left(-\pi/8, \pi/8\right)$

B.  $\left(3\pi/8, 5\pi/8\right)$

C.  $\left(\pi/4, 3\pi/4\right)$

D.  $\left(7\pi/8, 9\pi/8\right)$

Answer: A::B::D



Watch Video Solution

5. Let  $f(x) = x^2 - 2\sqrt{(\sin \sqrt{3} - \sin \sqrt{2})}x - (\cos \sqrt{3} - \cos \sqrt{2})$

A.  $f(x)$  is positive  $\forall x \in R$

B.  $f(x)$  assumes both positive and negative values

C.  $f(x)=0$  has no real roots

D.  $y=f(x)$  touches the line  $y=0$ .

Answer: A::C

 [Watch Video Solution](#)

6. For  $\alpha = \frac{\pi}{7}$  which of the following hold (s) good?

A.  $\tan \alpha \tan 2\alpha \tan 3\alpha = \tan 3\alpha - \tan 2\alpha - \tan \alpha$

B.  $\operatorname{cosec} \alpha = \operatorname{cosec} 2\alpha + \operatorname{cosec} 4\alpha$ .

C.  $\cos \alpha - \cos 2\alpha + \cos 3\alpha = 1/2$

D.  $8 \cos \alpha \cos 2\alpha \cos 4\alpha = 1$

Answer: A::B::C

 [Watch Video Solution](#)

7. Which of the following identities, wherever defined, hold(s) good?

A.  $\cot \alpha - \tan \alpha = 2 \cot 2\alpha$

B.  $\tan(45^\circ + \alpha) - \tan(45^\circ - \alpha) = 2 \cot 2\alpha$

C.  $\tan(45^\circ + \alpha) + \tan(45^\circ - \alpha) = 2 \sec 2\alpha$

D.  $\tan \alpha + \cot \alpha = 2 \tan 2\alpha$ .

Answer: A::C



Watch Video Solution

8. The expression  $(\tan^4 x + 2 \tan^2 x + 1) \cos^2 x$ , when  $x = \frac{\pi}{12}$ , can be

equal to  $4(2 - \sqrt{3})$  (b)  $4(\sqrt{2} + 1)$   $16 \frac{\cos^2 \pi}{12}$  (d)  $16 \frac{\sin^2 \pi}{12}$

A.  $4(2 - \sqrt{3})$

B.  $4(\sqrt{2} + 1)$

C.  $16 \cos^2 \pi / 12$



D.  $16 \sin^2 \pi / 12$

Answer: A::D

 [Watch Video Solution](#)

9. Let  $\alpha, \beta$  and  $\gamma$  be some angles in the first quadrant satisfying

$\tan(\alpha + \beta) = \frac{15}{8}$  and  $\operatorname{cosec} \gamma = \frac{17}{8}$ , then which of the following

hold(s) good?  $\alpha + \beta + \gamma = \pi$   $\cot \alpha + \tan \beta + \tan \gamma = \tan \alpha \tan \beta \tan \gamma$

$$\tan \alpha + \tan \beta + \tan \gamma = \tan \alpha \tan \beta \tan \gamma$$

$$\tan \alpha \tan \beta + \tan \beta \tan \gamma + \tan \gamma \tan \alpha = 1$$

A.  $\alpha + \beta + \gamma = \pi$

B.  $\cot \alpha \cot \beta \cot \gamma = \cot \alpha + \cot \beta + \cot \gamma$

C.  $\tan \alpha + \tan \beta + \tan \gamma = \tan \alpha \tan \beta \tan \gamma$

D.  $\tan \alpha \tan \beta + \tan \beta \tan \gamma + \tan \gamma \tan \alpha = 1$

Answer: B::D

 [Watch Video Solution](#)

10. Let  $f_n(\theta) = \frac{\frac{\cos \theta}{2} + \cos 2\theta + \frac{\cos(7\theta)}{2} + \dots + \cos(3n-2)\left(\frac{\theta}{2}\right)}{\frac{\sin \theta}{2} + \sin 2\theta + \frac{\sin(7\theta)}{2} + \dots + \sin(3n-2)\left(\frac{\theta}{2}\right)}$  then  $f_3\left(\frac{3\pi}{16}\right)$

A.  $f_3\left(\frac{3\pi}{16}\right) = \sqrt{2} - 1$

B.  $f_5\left(\frac{\pi}{28}\right) = \sqrt{2} + 1$

C.  $f_7\left(\frac{\pi}{60}\right) = (2 + \sqrt{3})$

D. none of these.

Answer: A::B::C



Watch Video Solution

11. If  $\sin(x + 20^\circ) = 2 \sin x \cos 40^\circ$ , where  $x \in \left(0, \frac{\pi}{2}\right)$ , then which of the following hold(s) good?  $\cos 2x = \frac{1}{2}$  (b)  $\cos ec 4x = 2 \frac{\sec x}{2} = \sqrt{6} - \sqrt{2}$

(d)  $\frac{\tan x}{2} = (2 - \sqrt{3})$

A.  $\cos 2x = 1/2$

B.  $\operatorname{cosec} 4x = 2$

C.  $\sec\left(\frac{x}{2}\right) = \sqrt{6} - \sqrt{2}$

D.  $\tan\left(\frac{x}{2}\right) = (2 - \sqrt{3})$

Answer: A::C::D

 [Watch Video Solution](#)

12. The expression  $\cos^2(\alpha + \beta) + \cos^2(\alpha - \beta) - \cos 2\alpha \cos 2\beta$  is

A. independent of  $\alpha$

B. independent of  $\beta$

C. independent of  $\alpha$  and  $\beta$

D. dependent on  $\alpha$  and  $\beta$ .

Answer: A::B::C

 [Watch Video Solution](#)

13. If  $\cot^3 \alpha + \cot^2 \alpha + \cot \alpha = 1$  then which of the following is/are correct

A.  $\cos 2\alpha \cdot \tan \alpha = -1$

B.  $\cos 2\alpha \cdot \tan \alpha = 1$

C.  $\cos 2\alpha - \tan 2\alpha = 1$

D.  $\cos 2\alpha - \tan 2\alpha = 1.$

Answer: A::C



Watch Video Solution

14.

If

$$p = \sin(A - B)\sin(C - D), q = \sin(B - C)\sin(A - D), r = \sin(C - A)\sin(B - D)$$

then  $p + q - r = 0$  (b)  $p + q + r = 0$   $p - q + r = 0$  (d)

$$p^3 + q^3 + r^3 = 3pqr$$

A.  $p + q - r = 0$

B.  $p + q + r = 0$

C.  $p - q + r = 0$

D.  $p^3 + q^3 + r^3 = 3pqr$

Answer: B::D

 [Watch Video Solution](#)

15. If  $\cos x - \sin \alpha \cot \beta \sin x = \cos a$ , then the value of  $\tan\left(\frac{x}{2}\right)$  is  $-\tan\left(\frac{\alpha}{2}\right)\cot\left(\frac{\beta}{2}\right)$  (b)  $\tan\left(\frac{\alpha}{2}\right)\tan\left(\frac{\beta}{2}\right) - \cot\left(\frac{\alpha\beta}{2}\right)\tan\left(\frac{\beta}{2}\right)$  (d)  $\cot\left(\frac{\alpha}{2}\right)\cot\left(\frac{\beta}{2}\right)$

A.  $-\tan(\alpha/2)\cot(\beta/2)$

B.  $\tan(\alpha/2)\tan(\beta/2)$

C.  $-\cot(\alpha/2)\tan(\beta/2)$

D.  $\cot(\alpha/2)\cot(\beta/2)$

Answer: A::B



Watch Video Solution

16. Let  $f(x) = ab \sin x + b\sqrt{1 - a^2} \cos x + c$ , where  $|a| < 1, b > 0$  then

A. maximum value of  $f(x) = -\cos^{-1} a$

B.  $f(x) = c$  if  $x = \cos^{-1} a$

C.  $f(x) = c$  if  $x = -\cos^{-1} a$

D.  $f(x) = c$  if  $x = \cos^{-1} a$ .

Answer: A::B::C



Watch Video Solution

17. Let  $P(k) = \left(1 + \frac{\cos \pi}{4k}\right) \left(1 + \frac{\cos((2k-1)\pi)}{4k}\right)$   
 $\left(1 + \frac{\cos((2k+1)\pi)}{4k}\right) \left(1 + \frac{\cos((4k-1)\pi)}{4k}\right)$  Then  $P(3) = \frac{1}{16}$  (b)  
 $P(4) = \frac{2 - \sqrt{2}}{16}$   $P(5) = \frac{3 - \sqrt{5}}{32}$  (d)  $P(6) = \frac{2 - \sqrt{3}}{16}$

A.  $P(3) = \frac{1}{16}$

B.  $P(4) = \frac{2 - \sqrt{2}}{16}$

C.  $P(5) = \frac{3 - \sqrt{5}}{32}$

D.  $P(6) = \frac{2 - \sqrt{3}}{16}$

Answer: A::B::C::D

 [Watch Video Solution](#)

18. If  $3 \sin \beta = \sin(2\alpha + \beta)$ , then  $\tan(\alpha + \beta) - 2 \tan \alpha$  is

A. independent of  $\alpha$

B. independent of  $\beta$

C. dependent of both  $\alpha$  and  $\beta$ .

D. independent of both  $\alpha$  and  $\beta$ .

Answer: A::B::D

 [Watch Video Solution](#)

19.  $x = \sqrt{a^2 \cos^2 \alpha + b^2 \sin^2 \alpha} + \sqrt{a^2 \sin^2 \alpha + b^2 \cos^2 \alpha}$  then

$x^2 = a^2 + b^2 + 2\sqrt{p(a^2 + b^2) - p^2}$ , where  $p$  can be is equal to

A.  $a^2 \cos^2 \alpha + b^2 \sin^2 \alpha$

B.  $a^2 \sin^2 \alpha + b^2 \cos^2 \alpha$

C.  $\frac{1}{2} [a^2 + b^2 + (a^2 - b^2) \cos 2\alpha]$

D.  $\frac{1}{2} [a^2 + b^2 - (a^2 - b^2) \cos 2\alpha]$

Answer: A::B::C::D



Watch Video Solution

20. If

$$(x - a) \cos \theta + y \sin \theta = (x - a) \cos \varphi + y \sin \theta = a \text{ and } \tan\left(\frac{\theta}{2}\right) = \tan\left(\frac{\varphi}{2}\right)$$

$$y^2 = 2ax - (1 - b^2)x^2 \quad \frac{\tan \theta}{2} = \frac{1}{x}(y + bx) \quad y^2 = 2bx - (1 - a^2)x^2$$

$$\frac{\tan \varphi}{2} = \frac{1}{x}(y - bx)$$



A.  $y^2 = 2ax - (1 - b^2)x^2$

B.  $\tan\left(\frac{\theta}{2}\right) = \frac{1}{x}(y + bx)$

C.  $y^2 = 2bx - (1 - a^2)x^2$

D.  $\tan\left(\frac{\phi}{2}\right) = \frac{1}{x}(y - bx)$

Answer: A::B::D



Watch Video Solution

21. If  $\cos(x - y)$ ,  $\cos x$  and  $\cos(x + y)$  are in H.P., then

$\cos x \cdot \sec\left(\frac{y}{2}\right) =$

A.  $-\sqrt{3}$

B.  $-\sqrt{2}$

C.  $\sqrt{2}$

D.  $\sqrt{3}$

Answer: B::C



[Watch Video Solution](#)

22. Difference between maximum and minimum values of  $(60 \sin \alpha + p \cos \alpha)$  is 122 then p can be

A. 61

B. 11

C. - 61

D. - 11

Answer: B::D



[Watch Video Solution](#)

### Exercise (Comprehension)

1. If  $\sin \alpha = A \sin(\alpha + \beta)$ ,  $A \neq 0$ , then

The value of  $\tan \alpha$  is

A.  $\frac{A \sin \beta}{1 - A \cos \beta}$

B.  $\frac{A \sin \beta}{1 + A \cos \beta}$

C.  $\frac{A \cos \beta}{1 - A \sin \beta}$

D.  $\frac{A \sin \beta}{1 + A \cos \beta}$

**Answer: A**



**Watch Video Solution**

**2. If  $\sin \alpha = A \sin(\alpha + \beta)$ ,  $A \neq 0$ , then**

**The value of  $\tan \beta$  is**

A.  $\frac{\sin \alpha(1 + A \cos \beta)}{A \cos \alpha \cos \beta}$

B.  $\frac{\sin \alpha(1 - A \cos \beta)}{A \cos \alpha \cos \beta}$

C.  $\frac{\cos(1 - A \sin \beta)}{A \cos \alpha \cos \beta}$

D.  $\frac{\cos \alpha(1 + A \sin \beta)}{A \cos \alpha \cos \beta}$

**Answer: B**

 [Watch Video Solution](#)

3. If  $\sin \alpha = A \sin(\alpha + \beta)$ ,  $A \neq 0$ , then

Which of the following is not the value of  $\tan(\alpha + \beta)$ ?

A.  $\frac{\sin \beta}{\cos \beta - A}$

B.  $\frac{\sin \alpha \cos \alpha}{A \cos \beta - \sin^2 \alpha}$

C.  $\frac{\sin \alpha \cos \alpha}{A \cos \beta + \sin^2 \alpha}$

D. none of these.

Answer: C

 [Watch Video Solution](#)

4. If  $\alpha, \beta, \gamma, \delta$  are the four solutions of the equation

$\tan\left(\theta + \frac{\pi}{4}\right) = 3 \tan 3\theta$ . No two of which have equal tangents, then the

value of  $\tan \alpha + \tan \beta + \tan \gamma + \tan \delta =$

A.  $1/3$

B.  $8/3$

C.  $-8/3$

D. 0

Answer: D



Watch Video Solution

5. If  $\alpha, \beta, \gamma, \delta$  are the four solutions of the equation  $\tan\left(\theta + \frac{\pi}{4}\right) = 3 \tan 3\theta$ . No two of which have equal tangents, then the value of  $\tan \alpha + \tan \beta + \tan \gamma + \tan \delta =$

A.  $-1/3$

B.  $-2$

C. 0

D. none of these

**Answer: A**



**Watch Video Solution**

**6. If  $\alpha, \beta, \gamma$  are the solutions of the equation  $\tan\left(\theta + \frac{\pi}{4}\right) = 3 \tan 3\theta$ , no two of which have equal tangents.**

**The value of  $\frac{1}{\tan \alpha} + \frac{1}{\tan \beta} + \frac{1}{\tan \gamma} + \frac{1}{\tan \delta}$  is**

**A.  $-8$**

**B.  $8$**

**C.  $2/3$**

**D.  $1/3$**

**Answer: B**



**Watch Video Solution**

**7.  $\sin \alpha + \sin \beta = \frac{1}{4}$  and  $\cos \alpha + \cos \beta = \frac{1}{3}$  the value of  $\sin(\alpha + \beta)$**

A.  $\frac{24}{25}$

B.  $\frac{13}{25}$

C.  $\frac{12}{13}$

D. none of these.

Answer: A



Watch Video Solution

8.  $\sin \alpha + \sin \beta = \frac{1}{4}$  and  $\cos \alpha + \cos \beta = \frac{1}{3}$

The value of  $\cos(\alpha + \beta)$  is

A.  $\frac{12}{25}$

B.  $\frac{7}{25}$

C.  $\frac{12}{13}$

D. none of these.

Answer: B

 [Watch Video Solution](#)

9.  $\sin \alpha + \sin \beta = \frac{1}{4}$  and  $\cos \alpha + \cos \beta = \frac{1}{3}$

The value of  $\tan(\alpha + \beta)$  is

A.  $\frac{25}{7}$

B.  $\frac{25}{12}$

C.  $\frac{25}{13}$

D.  $\frac{24}{7}$

Answer: D

 [Watch Video Solution](#)

10. To find the sum  $\sin^2 \frac{2\pi}{7} + \sin^2 \frac{4\pi}{7} + \sin^2 \frac{8\pi}{7}$ , we follow the following method.

Put  $7\theta = 2n\pi$ , where  $n$  is any integer. Then

$$\sin 4\theta = \sin(2n\pi - 3\theta) = -\sin 3\theta$$



**This means that  $\sin \theta$  takes the values**

$0, \pm \sin(2\pi/7), \pm \sin(4\pi/7), \text{ and } \pm \sin(8\pi/7).$

**From Eq. (i), we now get**

$$2 \sin 2\theta \cos 2\theta = 4 \sin^3 \theta - 3 \sin \theta$$

$$\text{or } 4 \sin \theta \cos \theta (1 - 2 \sin^2 \theta) = \sin \theta (4 \sin^2 \theta - 3)$$

**Rejecting the value  $\sin \theta = 0$ , we get**

$$4 \cos \theta (1 - 2 \sin^2 \theta) = 4 \sin^2 \theta - 3$$

$$\text{or } 16 \cos^2 \theta (1 - 2 \sin^2 \theta)^2 = (4 \sin^2 \theta - 3)^2$$

$$\text{or } 16(1 - \sin^2 \theta)(1 - 4 \sin^2 \theta + 4 \sin^4 \theta)$$

$$= 16 \sin^4 \theta - 24 \sin^2 \theta + 9$$

$$\text{or } 64 \sin^6 \theta - 112 \sin^4 \theta - 56 \sin^2 \theta - 7 = 0$$

**This is cubic in  $\sin^2 \theta$  with the roots**

$\sin^2(2\pi/7), \sin^2(4\pi/7), \text{ and } \sin^2(8\pi/7).$

**The sum of these roots is**

$$\sin^2 \frac{2\pi}{7} + \sin^2 \frac{4\pi}{7} + \sin^2 \frac{8\pi}{7} = \frac{112}{64} = \frac{7}{4}.$$

**The value of**

$$\left( \tan^2 \frac{\pi}{7} + \tan^2 \frac{2\pi}{7} + \tan^2 \frac{3\pi}{7} \right) \times \left( \cot^2 \frac{\pi}{7} + \cot^2 \frac{2\pi}{7} + \cot^2 \frac{3\pi}{7} \right) \text{ is}$$

B. 35

C. 210

D. none of these.

Answer: A



Watch Video Solution

11. To find the sum  $\sin^2 \frac{2\pi}{7} + \sin^2 \frac{4\pi}{7} + \sin^2 \frac{8\pi}{7}$ , we follow the following method.

Put  $7\theta = 2n\pi$ , where  $n$  is any integer. Then

$$\sin 4\theta = \sin(2n\pi - 3\theta) = -\sin 3\theta$$

This means that  $\sin \theta$  takes the values

$0, \pm \sin(2\pi/7), \pm \sin(4\pi/7), \pm \sin(6\pi/7),$  and  $\pm \sin(8\pi/7).$

From Eq. (i), we now get

$$2 \sin 2\theta \cos 2\theta = 4 \sin^3 \theta - 3 \sin \theta$$

$$\text{or } 4 \sin \theta \cos \theta (1 - 2 \sin^2 \theta) = \sin \theta (4 \sin^2 \theta - 3)$$

Rejecting the value  $\sin \theta = 0$ , we get

$$4 \cos \theta(1 - 2 \sin^2 \theta) = 4 \sin^2 \theta - 3$$

$$\text{or } 16 \cos^2 \theta(1 - 2 \sin^2 \theta)^2 = (4 \sin^2 \theta - 3)^2$$

$$\text{or } 16(1 - \sin^2 \theta)(1 - 4 \sin^2 \theta + 4 \sin^4 \theta)$$

$$= 16 \sin^4 \theta - 24 \sin^2 \theta + 9$$

$$\text{or } 64 \sin^6 \theta - 112 \sin^4 \theta - 56 \sin^2 \theta - 7 = 0$$

This is cubic in  $\sin^2 \theta$  with the roots

$$\sin^2(2\pi/7), \sin^2(4\pi/7), \text{ and } \sin^2(8\pi/7).$$

The sum of these roots is

$$\sin^2 \frac{2\pi}{7} + \sin^2 \frac{4\pi}{7} + \sin^2 \frac{8\pi}{7} = \frac{112}{64} = \frac{7}{4}.$$

The value of  $\frac{\tan^2 \frac{\pi}{7} + \tan^2 \frac{2\pi}{7} + \tan^2 \frac{3\pi}{7}}{\cot^2 \frac{\pi}{7} + \cot^2 \frac{2\pi}{7} + \cot^2 \frac{3\pi}{7}}$  is

A. 7

B.  $35/3$

C.  $21/5$

D. none of these

Answer: C



Watch Video Solution

12. To find the sum  $\sin^2 \frac{2\pi}{7} + \sin^2 \frac{4\pi}{7} + \sin^2 \frac{8\pi}{7}$ , we follow the following method.

Put  $7\theta = 2n\pi$ , where  $n$  is any integer. Then

$$\sin 4\theta = \sin(2n\pi - 3\theta) = -\sin 3\theta$$

This means that  $\sin \theta$  takes the values

$0, \pm \sin(2\pi/7), \pm \sin(4\pi/7), \pm \sin(6\pi/7),$  and  $\pm \sin(8\pi/7).$

From Eq. (i), we now get

$$2 \sin 2\theta \cos 2\theta = 4 \sin^3 \theta - 3 \sin \theta$$

or  $4 \sin \theta \cos \theta (1 - 2 \sin^2 \theta) = \sin \theta (4 \sin^2 \theta - 3)$

Rejecting the value  $\sin \theta = 0$ , we get

$$4 \cos \theta (1 - 2 \sin^2 \theta) = 4 \sin^2 \theta - 3$$

or  $16 \cos^2 \theta (1 - 2 \sin^2 \theta)^2 = (4 \sin^2 \theta - 3)^2$

or  $16(1 - \sin^2 \theta)(1 - 4 \sin^2 \theta + 4 \sin^4 \theta)$

$$= 16 \sin^4 \theta - 24 \sin^2 \theta + 9$$

or  $64 \sin^6 \theta - 112 \sin^4 \theta - 56 \sin^2 \theta - 7 = 0$

This is cubic in  $\sin^2 \theta$  with the roots

$\sin^2(2\pi/7), \sin^2(4\pi/7),$  and  $\sin^2(8\pi/7).$

The sum of these roots is

$$\sin^2 \frac{2\pi}{7} + \sin^2 \frac{4\pi}{7} + \sin^2 \frac{8\pi}{7} = \frac{112}{64} = \frac{7}{4}.$$

The value of  $\tan^2 \frac{\pi}{7} \tan^2 \frac{2\pi}{7} \tan^2 \frac{3\pi}{7}$  is

- A.  $-3$
- B.  $7$
- C.  $-5$
- D. none of these

Answer: B



Watch Video Solution

13. In a  $\triangle ABC$ , if

$$\cos A \cos B \cos C = \frac{\sqrt{3} - 1}{8} \text{ and } \sin A \sin B \sin C = \frac{3 + \sqrt{3}}{8}, \text{ then}$$

The value of  $\tan A + \tan B + \tan C$  is

- A.  $\frac{3 + \sqrt{3}}{\sqrt{3} - 1}$
- B.  $\frac{\sqrt{3} + 4}{\sqrt{3} - 1}$

C.  $\frac{6 - \sqrt{3}}{\sqrt{3} - 1}$

D.  $\frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} - 1}$

**Answer: A**



**Watch Video Solution**

14. In a  $\triangle ABC$ , if  $\cos A \cos B \cos C = \frac{\sqrt{3} - 1}{8}$  and  $\sin A \sin B \sin C = \frac{3 + \sqrt{3}}{8}$ , then

The value of  $\tan A \tan B \tan C$  is

A.  $5 - 4\sqrt{3}$

B.  $5 + 4\sqrt{3}$

C.  $6 + \sqrt{3}$

D.  $6 - \sqrt{3}$

**Answer: B**



**Watch Video Solution**

15. In a  $\triangle ABC$ , if

$$\cos A \cos B \cos C = \frac{\sqrt{3} - 1}{8} \text{ and } \sin A \sin B \sin C = \frac{3 + \sqrt{3}}{8}, \text{ then}$$

the respective values of  $\tan A$ ,  $\tan B$  and  $\tan C$  are

A.  $1, \sqrt{3}, \sqrt{2}$

B.  $1, \sqrt{3}, 2$

C.  $1, 2, \sqrt{3}$

D.  $1, \sqrt{3}, 2 + \sqrt{3}$

Answer: D



Watch Video Solution

16. If the angles  $\alpha, \beta, \gamma$  of a triangle satisfy the relation,

$$\sin\left(\frac{\alpha - \beta}{2}\right) + \sin\left(\frac{\alpha - \gamma}{2}\right) + \sin\left(\frac{3\alpha}{2}\right) = \frac{3}{2}, \text{ then}$$

The measure of the smallest angle of the triangle is

A.  $30^\circ$

B.  $40^\circ$

C.  $45^\circ$

D.  $50^\circ$

Answer: B



Watch Video Solution

17. If the angles  $\alpha, \beta, \gamma$  of a triangle satisfy the relation,

$$\sin\left(\frac{\alpha - \beta}{2}\right) + \sin\left(\frac{\alpha - \gamma}{2}\right) + \sin\left(\frac{3\alpha}{2}\right) = \frac{3}{2}, \text{ then}$$

Triangle is

A. acute angled

B. right angled but not isosceles

C. isosceles

D. isosceles right angled



Answer: C



Watch Video Solution

18. A line OA of length  $r$  starts from its initial position OX and traces an angle  $\text{AOB} = \alpha$  in the anticlockwise direction. It then traces back in the clockwise direction an angle  $\text{BOC} = 3\theta$  ( where  $\alpha > 3\theta$ ). L is the foot of the

perpendicular from C on OA. Also,  $\frac{\sin^3 \theta}{CL} = \frac{\cos^3 \theta}{OL} = 1$

$\frac{1 - r \cos \alpha}{r \sin \alpha}$  is equal to

A.  $\tan 2\theta$

B.  $\cot 2\theta$

C.  $\sin 2\theta$

D.  $\sin 2\theta$

Answer: A



Watch Video Solution

19. A line OA of length  $r$  starts from its initial position OX and traces an angle  $\text{AOB} = \alpha$  in the anticlockwise direction. It then traces back in the clockwise direction an angle  $\text{BOC} = 3\theta$  ( where  $\alpha > 3\theta$ ). L is the foot of the perpendicular from C on OA. Also,  $\frac{\sin^3 \theta}{CL} = \frac{\cos^3 \theta}{OL} = 1$

$\frac{2r \sin \alpha}{1 + 2r \cos \alpha}$  is equal to

A.  $\tan^2 \theta$

B.  $\cot^2 \theta$

C.  $\cot 2\theta$

D.  $\tan 2\theta$

Answer: D



Watch Video Solution

20. A line OA of length  $r$  starts from its initial position OX and traces an angle  $\text{AOB} = \alpha$  in the anticlockwise direction. It then traces back in the clockwise direction an angle  $\text{BOC} = 3\theta$  ( where  $\alpha > 3\theta$ ). L is the foot of the

perpendicular from C on OA. Also,  $\frac{\sin^3 \theta}{CL} = \frac{\cos^3 \theta}{OL} = 1$

$\frac{2r^2 - 1}{r}$  is equal to

A.  $\sin \alpha$

B.  $\cos \alpha$

C.  $\sin \theta$

D.  $\cos \theta$

**Answer: B**



[Watch Video Solution](#)

### Exercise (Matrix Match Type )

1. If  $\cos \theta - \sin \theta = \frac{1}{5}$ , where  $0 < \theta < \frac{\pi}{4}$ , then



[View Text Solution](#)

2. If  $\cos \alpha + \cos \beta = 1/2$  and  $\sin \alpha + \sin \beta = 1/3$ , then



[View Text Solution](#)

3. 



[View Text Solution](#)

4. 



[View Text Solution](#)

5. 



[View Text Solution](#)

6. 



[View Text Solution](#)

7. 



[View Text Solution](#)

8. 

- A.  $\begin{matrix} a & b & c & d \\ (1) & s & q & r & p \end{matrix}$
- B.  $\begin{matrix} a & b & c & d \\ (2) & r & s & q & p \end{matrix}$
- C.  $\begin{matrix} a & b & c & d \\ (3) & p & r & s & q \end{matrix}$
- D.  $\begin{matrix} a & b & c & d \\ (4) & q & r & s & p \end{matrix}$

**Answer: A**



[View Text Solution](#)

9. 

- A.  $\begin{matrix} a & b & c & d \\ (1) & s & q & r & p \end{matrix}$
- B.  $\begin{matrix} a & b & c & d \\ (2) & r & s & q & p \end{matrix}$
- C.  $\begin{matrix} a & b & c & d \\ (3) & q & s & p & r \end{matrix}$
- D.  $\begin{matrix} a & b & c & d \\ (4) & q & r & s & p \end{matrix}$

Answer: C



[View Text Solution](#)

## Exercise (Numerical)

1. If  $f(\theta) = \frac{1 - \sin 2\theta + \cos 2\theta}{2 \sin 2\theta}$  then value of  $f(11^\circ)$ .  $f(34^\circ)$  is



[Watch Video Solution](#)

2. If  $f(x) = 2(7 \cos x + 24 \sin x)(7 \sin x - 24 \cos x)$ , for even  $x \in R$  then maximum value of  $f(x)$  is \_\_\_\_\_

 [Watch Video Solution](#)

3. In a triangle  $ABC$ ,  $\angle C = \frac{\pi}{2}$ . If  $\tan\left(\frac{A}{2}\right)$  and  $\tan\left(\frac{B}{2}\right)$  are the roots of the equation  $ax^2 + bx + c = 0$ , ( $a \neq 0$ ), then the value of  $\frac{a+b}{c}$  (where  $a, b, c$ , are sides of opposite to angles  $A, B, C$ , respectively) is

 [Watch Video Solution](#)

4. If  $x, y \in R$  satisfies  $(x + 5)^2 + (y - 12)^2 = (14)^2$ , then the minimum value of  $\sqrt{x^2 + y^2}$  is \_\_\_\_\_

 [Watch Video Solution](#)

5. Suppose  $x$  and  $y$  real number such that  $\tan x \tan y = 42$  and  $\cot x + \cot y = 49$  the value of  $\tan(x + y)$  is



[Watch Video Solution](#)

6. Let  $0 \leq a, b, c, d \leq \pi$ , where  $a, b, c, d$  are not complementary, such that

$$2 \cos a + 6 \cos b + 7 \cos c + 9 \cos d = 0 \text{ and } 2 \sin a - 6 \sin b + 7 \sin c - 9 \sin d = 0$$

then the value of  $3 \frac{\cos(a+d)}{\cos(b+c)}$  is \_\_\_\_\_



[Watch Video Solution](#)

7. Suppose  $A$  and  $B$  are two angles such that  $A, B \in (0, \pi)$  and satisfy

$$\sin A + \sin B = 1 \quad \text{and} \quad \cos A + \cos B = 0. \quad \text{Then the value of}$$

$$12 \cos 2A + 4 \cos 2B \text{ is } \underline{\hspace{2cm}}$$



[Watch Video Solution](#)

8.  $\alpha$  and  $\beta$  are the positive acute angles and satisfying equation

$$5 \sin 2\beta = 3 \sin 2\alpha \text{ and } \tan \beta = 3 \tan \alpha \text{ simultaneously. Then the value of}$$



$\tan \alpha + \tan \beta$  is \_\_\_\_\_

 [Watch Video Solution](#)

9. The absolute value of the expression

$$\frac{\tan \pi}{16} + \frac{\tan(5\pi)}{16} + \frac{\tan(9\pi)}{16} + \frac{\tan(13\pi)}{16} \text{ is } \underline{\hspace{2cm}}$$

 [Watch Video Solution](#)

10. The greatest integer less than or equal to  $\frac{1}{\cos 290^\circ} + \frac{1}{\sqrt{3}\sin 250^\circ}$  is

\_\_\_\_\_

 [Watch Video Solution](#)

11. The maximum value of  $y = \frac{1}{\sin^6 x + \cos^6 x}$  is \_\_\_\_\_

 [Watch Video Solution](#)

12. The maximum value of  $\cos^2(45^\circ + x) + (\sin x - \cos x)^2$  is \_\_\_\_\_



Watch Video Solution

13. Find the exact value of  $\operatorname{cosec} 10^\circ + \operatorname{cosec} 50^\circ - \operatorname{cosec} 70^\circ$



Watch Video Solution

14. Number of triangles  $ABC$  if

$\tan A = x$ ,  $\tan B = x + 1$ , and  $\tan C = 1 - x$  is \_\_\_\_\_



Watch Video Solution

15. If

$$\log_{10} \sin x + \log_{10} \cos x = -1 \text{ and } \log_{10}(\sin x + \cos x) = \frac{(\log_{10} n)n - 1}{2}$$

then the value of  $n$  is \_\_\_\_\_



Watch Video Solution

16. The value of  $\frac{\sin 1^\circ + \sin 3^\circ + \sin 5^\circ + \sin 7^\circ}{\cos 1^\circ \cdot \cos 2^\circ \sin 4^\circ}$  is \_\_\_\_\_



Watch Video Solution

17. In a triangle  $ABC$ , if  $A - B = 120^\circ$  and  $\frac{\sin A}{2} \frac{\sin B}{2} \frac{\sin C}{2} = \frac{1}{32}$ , then the value of  $8 \cos C$  is \_\_\_\_\_



Watch Video Solution

18. If  
 $\frac{\tan x}{2} = \frac{\tan y}{3} = \frac{\tan z}{5}$ ,  $x + y + z = \pi$  and  $\tan^2 x + \tan^2 y + \tan^2 z = \frac{38}{K}$



Watch Video Solution

19. If  $\sin^3 x \cos 3x + \cos^3 x \sin 3x = 3/8$ , then the value of  $\sin 4x$  is \_\_\_\_\_



Watch Video Solution

20. The value of  $\operatorname{cosec} \frac{\pi}{18} - 4 \sin \frac{7\pi}{18}$  is \_\_\_\_\_



Watch Video Solution

21. If  $\tan x + \tan 2x + \tan 3x = \tan x \tan 2x \tan 3x$  then value of  $|\sin 3x + \cos 3x|$  is \_\_\_\_\_



Watch Video Solution

22.

$$16 \left( \cos \theta - \frac{\cos \pi}{8} \right) \left( \cos \theta - \frac{\cos(3\pi)}{8} \right) \left( \cos \theta - \frac{\cos(5\pi)}{8} \right) \left( \cos \theta - \frac{\cos(7\pi)}{8} \right)$$

then the value of  $\lambda$  is \_\_\_\_.



Watch Video Solution

23. If  $\frac{\tan(6n)\tan(2n)\tan(3n)}{\tan(6n) - \tan(2n) - \tan(3n)} = k$ , then the value of  $k$  is \_\_\_\_\_

 [Watch Video Solution](#)

24. If  $\cot(\theta - \alpha)$ ,  $3 \cot \theta$ ,  $\cot(\theta + \alpha)$  are in A.P. and  $\theta$  is not an integral multiple of  $\frac{\pi}{2}$ , then the value of  $\frac{4 \sin^2 \theta}{3 \sin^2 \alpha} =$  \_\_\_\_\_

 [Watch Video Solution](#)

25. The value of  $\frac{2 \sin x}{\sin 3x} + \frac{\tan x}{\tan 3x}$  \_\_\_\_\_

 [Watch Video Solution](#)

26. If  $\cot^2 A \cot^2 B = 3$ , then the value of  $(2 - \cos 2A)(2 - \cos 2B)$  is \_\_\_\_\_

 [Watch Video Solution](#)

27. The value of  $f(x) = x^4 + 4x^3 + 2x^2 - 4x + 7$ , when  $x = \frac{\cot(11\pi)}{8}$  is \_\_\_\_\_

 Watch Video Solution

28. The value of  $\sin^2 12^\circ + \sin^2 21^\circ + \sin^2 39^\circ + \sin^2 48^\circ - \sin^2 9^\circ - \sin^2 18^\circ$  is \_\_\_\_\_

 Watch Video Solution

29. Given that  $f(n\theta) = \frac{2 \sin 2\theta}{\cos 2\theta - \cos 4n\theta}$ , and  $f(\theta) + f(2\theta) + f(3\theta) + \dots + f(n\theta) = \frac{\sin \lambda\theta}{\sin \theta \sin \mu\theta}$ , then the value of  $\mu - \lambda$  is \_\_\_\_\_

 Watch Video Solution

30. Suppose  $\sin^3 x \sin 3x = \sum_{m=0}^n C_m \cos mx$  is an identity in  $x$ , where

$C_0, \dots, C_n$  are constant and  $C_n \neq 0$  then the value of  $n$  is \_\_\_\_\_

 [Watch Video Solution](#)

31. If  $\sec \alpha$  is the average of  $\sec(\alpha - 2\beta)$  and  $\sec(\alpha + 2\beta)$  then the value of  $(2\sin^2 \beta - \sin^2 \alpha)$  where  $\beta \neq n\pi$  is

 [Watch Video Solution](#)

32. If  $A, B$  and  $C$  are three values lying in  $[0, 2\pi]$  for which  $\tan \theta = K$

then  $\tan \frac{A}{3} \tan \frac{B}{3} + \tan \frac{B}{3} \tan \frac{C}{3} + \tan \frac{C}{3} \tan \frac{A}{3}$  is equal to

\_\_\_\_\_.

 [Watch Video Solution](#)

33. The value of  $\left[ \left( \sin \frac{\pi}{9} \right) \left( 4 + \sec \frac{\pi}{9} \right) \right]^2$  is \_\_\_\_\_.

 [Watch Video Solution](#)

34.  $\left(\frac{\sin 33^\circ}{\sin 11^\circ \sin 49^\circ \sin 71^\circ}\right)^2 + \left(\frac{\cos 33^\circ}{\cos 11^\circ \cos 49^\circ \cos 71^\circ}\right)^2$  is equal to \_\_\_\_\_.

 [Watch Video Solution](#)

35. If  $f(\theta) = \sin^3 \theta + \sin^3\left(\theta + \frac{2\pi}{3}\right) + \sin^3\left(\theta + \frac{4\pi}{3}\right)$  then the value of  $f\left(\frac{\pi}{18}\right) + f\left(\frac{7\pi}{18}\right)$  is \_\_\_\_\_.

 [Watch Video Solution](#)

36. The expression  $\frac{1 + \sin 22^\circ \sin 33^\circ \sin 35^\circ}{\cos^2 22^\circ + \cos^2 33^\circ + \cos^2 35^\circ}$  simplifies to

 [Watch Video Solution](#)

37. If  $A > 0$ ,  $B > 0$ , and  $A + B = \frac{\pi}{3}$  then the maximum value of  $\tan A \tan B$  is





Watch Video Solution

38. If  $\frac{\sin^3 \theta}{\sin(2\theta + \alpha)} = \frac{\cos^3 \theta}{\cos(2\theta + \alpha)}$  and  $\tan 2\theta = \lambda \tan(3\theta + \alpha)$  then the value of  $\lambda$  is \_\_\_\_\_.



Watch Video Solution

JEE Main Previous Year

1. Let  $A$  and  $B$  denote the statements

$$A: \cos \alpha + \cos \beta + \cos \gamma = 0$$

$$B: \sin \alpha + \sin \beta + \sin \gamma = 0$$

$$\text{If } \cos(\beta - \gamma) + \cos(\gamma - \alpha) + \cos(\alpha - \beta) = -\frac{3}{2},$$

then

A.  $A$  is true and  $B$  is false.

B.  $A$  is false and  $B$  is true.

C. Both  $A$  and  $B$  are true.

D. Both  $A$  and  $B$  are false.

Answer: C



Watch Video Solution

2. Let  $\cos(\alpha + \beta) = \frac{4}{5}$  and  $\sin(\alpha - \beta) = \frac{5}{13}$  where  $0 \leq \alpha, \beta \leq \frac{\pi}{4}$

then find  $\tan(2\alpha)$

A.  $\frac{20}{7}$

B.  $\frac{25}{16}$

C.  $\frac{56}{33}$

D.  $\frac{19}{12}$

Answer: C



Watch Video Solution

3. If  $A = \sin^2 x + \cos^4 x$ , then for all real  $x$ :

A.  $\frac{3}{4} \leq A \leq \frac{13}{16}$

B.  $\frac{3}{4} \leq A \leq 1$

C.  $\frac{13}{16} \leq A \leq 1$

D.  $1 \leq A \leq 2$

Answer: B



Watch Video Solution

4. In a  $\Delta PQR$ . if  $3 \sin P + 4 \cos Q = 6$  and  $4 \sin Q + 3 \cos P = 1$ , then the angle  $R$  is equal to:

A.  $\frac{5\pi}{6}$

B.  $\frac{\pi}{6}$

C.  $\frac{\pi}{4}$

D.  $\frac{3\pi}{4}$

**Answer: B**



**Watch Video Solution**

5. If  $5(\tan^2 x - \cos^2 x) = 2 \cos 2x + 9$ , then the value of  $\cos 4x$  is

A.  $-\frac{7}{9}$

B.  $-\frac{3}{5}$

C.  $\frac{1}{3}$

D.  $\frac{2}{9}$

**Answer: A**



**Watch Video Solution**

1. The value of  $\sum_{k=1}^{13} \frac{1}{\sin\left(\frac{\pi}{4} + \frac{(k-1)\pi}{6}\right)\sin\left(\frac{\pi}{4} + \frac{k\pi}{6}\right)}$  is equal to

A.  $3 - \sqrt{3}$

B.  $2(3 - \sqrt{3})$

C.  $2(\sqrt{3} - 1)$

D.  $2(2 + \sqrt{3})$

Answer: C



Watch Video Solution

2. Let  $f: (-1, 1) \rightarrow R$  be such that  $f(\cos 4\theta) = \frac{2}{2 - \sec^2 \theta}$  for  $\theta \in \left(0, \frac{\pi}{4}\right) \cup \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$ . Then the value(s) of  $f\left(\frac{1}{3}\right)$  is/are

A.  $1 - \sqrt{\frac{3}{2}}$

B.  $1 + \sqrt{\frac{3}{2}}$

C.  $1 - \sqrt{\frac{2}{3}}$

D.  $1 + \sqrt{\frac{2}{3}}$

Answer: A::B



Watch Video Solution

3. If  $\alpha$  and  $\beta$  are non-zero real number such that

$2(\cos \beta - \cos \alpha) + \cos \alpha \cos \beta = 1$ . Then which of the following is true?

A.  $\tan\left(\frac{\alpha}{2}\right) + \sqrt{3}\tan\left(\frac{\beta}{2}\right) = 0$

B.  $\sqrt{3}\tan\left(\frac{\alpha}{2}\right) + \tan\left(\frac{\beta}{2}\right) = 0$

C.  $\tan\left(\frac{\alpha}{2}\right) - \sqrt{3}\tan\left(\frac{\beta}{2}\right) = 0$

D.  $\sqrt{3}\tan\left(\frac{\alpha}{2}\right) - \tan\left(\frac{\beta}{2}\right) = 0$

Answer: A::C



Watch Video Solution

4. The maximum value of the expression  $\frac{1}{\sin^2 \theta + 3 \sin \theta \cos \theta + 5 \cos^2 \theta}$  is.....

 [Watch Video Solution](#)

5. The positive integer value of  $n > 3$  satisfying the equation

$$\frac{1}{\sin\left(\frac{\pi}{n}\right)} = \frac{1}{\sin\left(\frac{2\pi}{n}\right)} + \frac{1}{\sin\left(\frac{3\pi}{n}\right)}$$
 is

 [Watch Video Solution](#)

## Matrix Match Type

1. Match List I with List II and select the correct answer using the codes given below the lists :



A. 

(P)	(Q)	(R)	(S)
IV	III	I	II

- B.  $(P)$   $(Q)$   $(R)$   $(S)$   
 $IV$   $III$   $II$   $I$
- C.  $(P)$   $(Q)$   $(R)$   $(S)$   
 $III$   $IV$   $II$   $I$
- D.  $(P)$   $(Q)$   $(R)$   $(S)$   
 $III$   $IV$   $I$   $II$

**Answer: 2**



**View Text Solution**