



BIOLOGY

BOOKS - S DINESH & CO BIOLOGY (HINGLISH)

CHORDATA- PROTOCHORDATA

Mcq

1. Chordates differ from other animals in having

A. Notochord

B. Ventral nerve cord

C. Dorsal heart

D. Schizocoel

Answer: A



Watch Video Solution

2. Notochord occurs in the adult animal of

A. Osteichthyes

B. Chondrichthyes

C. Protochordata

D. Amphibia

Answer: C



Watch Video Solution

3. Which one of the following is a protochordate

A. Amphioxus

B. Lamprey

C. Labeo

D. Exocoetus

Answer: A



Watch Video Solution

4. Which group of protochordates contains

Balanoglossus

A. Urochordata

B. Tunicata

C. Tunicata

D. Hemichordata

Answer: D



Watch Video Solution

5. Only the tail portion contains notochord in

A. Chordata

B. Cephalochordata

C. Hemichordata

D. Urochordata

Answer: D



Watch Video Solution

6. A free swimming urochordate is

A. Herdmania

B. Botryllus

C. Salpa

D. Ciona

Answer: C



Watch Video Solution

7. A member of cephalochordata is

A. Herdmania

B. Ciona

C. Amphioxus

D. Balanoglossus

Answer: C



Watch Video Solution

8. The most important character of chordata is

A. Dorsal hollow nervous system

B. Vertebral column

C. Kidneys

D. None of the above

Answer: A



Watch Video Solution

9. A current of water is produced in Branchiostoma by

A. Cilia

B. Cirri

C. Wheel organ

D. Rostrum

Answer: C



 [Watch Video Solution](#)

10. Which one is absent in Branchiostoma

A. Head

B. Gill slits

C. Notochord

D. Dorsal hollow nerve cord

Answer: A



[Watch Video Solution](#)

11. The anterior part of Amphioxus is

A. Oral hood

B. Rostrum

C. Mouth

D. Both A and B

Answer: B



Watch Video Solution

12. Cephalochordate Branchiostoma possesses

A. Laterally compressed spindle-shaped body

B. Caudal fin

C. Dorsal and ventral fins

D. All the above

Answer: D



Watch Video Solution

13. Hepatoc portal system has evolved in protochordates

- A. Hemichordates
- B. Urochordates
- C. Cephalochordates
- D. All the above

Answer: C



Watch Video Solution

14. In Amphioxus, circulatory system is

A. Closed

B. Without heart

C. Without respiratory pigment

D. All the above

Answer: D



Watch Video Solution

15. Branchiostoma is

A. Filter feeder

B. Carnivorous

C. Omnivorous

D. Herbivorous

Answer: A



Watch Video Solution

16. In urochordates the chordates characters are

A. Well developed

B. Present only in larva

C. Present in adult

D. Present both in adult and larva

Answer: B



Watch Video Solution

17. The larva of urochordates contains notochord

A. Throughout

B. Head region

C. Tail region

D. Trunk region

Answer: C



Watch Video Solution

18. An ascidian is

A. Balanoglossus

B. Herdmania

C. Branchiostoma

D. None of the above

Answer: B



Watch Video Solution

19. Tunic is

- A. Sac- like covering of urochordates
- B. A string inside cephalochordates
- C. Collar of hemichordates
- D. None of the above

Answer: A



Watch Video Solution

20. Tunicin is

- A. Lipopolysaccharide
- B. Cellulose like polysaccharide
- C. Non - cellulosic polysaccharide
- D. Proteinaceous

Answer: B



Watch Video Solution

21. Retrogressive metamorphosis is peculiar to

- A. Urochordates
- B. Hemichordates
- C. Cephalochordates
- D. All the above

Answer: A



Watch Video Solution

22. Mouth of Herdmania is

- A. Atrial aperture
- B. Excurrent aperture
- C. Branchial aperture
- D. Both A and B

Answer: C



Watch Video Solution

23. Atrial and branchial apertures of Herdmania are surrounded by

- A. Two lips
- B. Three lips
- C. Four lips
- D. Five lips

Answer: C



View Text Solution

24. Water current is maintained in Herdmania by

A. Rhythmic contractions of tunic

B. Opening and closing of atrial and
branchial apertures

C. Constriction of incurrent siphon

D. Cilia bordering gill slits

Answer: D



Watch Video Solution

25. Lancelet is

A. Herdmania

B. Salpa

C. Branchiostoma

D. Doliolum

Answer: C



Watch Video Solution

26. In protochordates the epidermis is

- A. Absent
- B. Multilayered
- C. A few layered
- D. Single layered

Answer: D



Watch Video Solution

27. Part of Herdmania embedded in sand is

A. Test

B. Foot

C. Tunic

D. Both A and B

Answer: B



Watch Video Solution

28. Herdmania is

A. Sea purse

B. Sea Squirt

C. Sea potato

D. Both B and C

Answer: D



Watch Video Solution

29. Which one emits jet of water when disturbed

A. Balanoglossus

B. Herdmania

C. Salpa

D. Branchiostoma

Answer: B



Watch Video Solution

30. In adult urochordate the dorsal nerve cord of larva is changed into

- A. Ganglion
- B. Brain and spinal cord
- C. Brain and nerve cord
- D. Remains as such

Answer: A



Watch Video Solution

31. In cephalochordates the coelom is

A. Reduced

B. Absent

C. Schizocoel

D. Lined by ectoderm

Answer: A



Watch Video Solution

32. Coelom of urochordate is

A. Schizocoel

B. Absent

C. Enterocoel

D. Segmented

Answer: B



Watch Video Solution

33. Acorn Worm is

A. Doliolum

B. Salpa

C. Balanoglossus

D. Rhabdopleura

Answer: C



Watch Video Solution

34. The body of Tongue Worm *Balanoglossus* is differentiated into

- A. Head, trunk and tail
- B. Trunk, and tail
- C. Proboscis, collar and trunk
- D. Proboscis and collar

Answer: C



Watch Video Solution

35. Which one swallows mud

- A. Amphioxus
- B. Sea Squirt
- C. Tongue Worm
- D. Asymmetron

Answer: C



Watch Video Solution

36. In hemichordates the dorsal nerve cord is

A. Absent

B. Present throughout

C. Restricted to tongue region

D. Restricted to collar region

Answer: D



Watch Video Solution

37. Coelom of hemichordates is

A. Schizocoel

B. Enterocoel

C. Enterocoel with differentiation of three regions

D. Absent

Answer: C



View Text Solution

38. Heart of hemichordates is

A. Ventral

B. Dorsal

C. Lateral

D. Absent

Answer: B



Watch Video Solution

39. Who raised the status of hemichordates from subphylum to phylum

A. Linnaeus

B. Lamarck

C. Horst

D. Belfour

Answer: D



Watch Video Solution

40. Notochord like structure is present in hemichordates in the region

A. Proboscis

B. Collar

C. Trunk

D. Tail

Answer: A



Watch Video Solution

41. Notochord like structure of hemichordates

is

A. Protochord

B. Stomochord

C. Pallium

D. Glomerulus

Answer: B



Watch Video Solution

42. Pharyngeal gill slits are dorsal in

A. Cephalochordates

B. Urochordates

C. Hemichordates

D. Euchordates

Answer: C



Watch Video Solution

43. Protochordates are

A. Marine

B. Fresh water

C. Terrestrial

D. All the above

Answer: A



Watch Video Solution

44. Protochordates are

A. Non - vertebrates

B. Acraniates

C. Prevertebrates

D. All the above

Answer: D



Watch Video Solution

45. Chorda dorsalis is

- A. Dorsal solid nerve cord
- B. Dorsal hollow nerve cord
- C. Notochord
- D. Stomochord

Answer: C



Watch Video Solution

46. Amongst chordates, the paired appendages are absent in

- A. Hemichordates
- B. Urochordates
- C. Cephalochordates
- D. All the above

Answer: D



47. Notochord is

A. Solid

B. Hollow

C. Lacunate

D. Partially solid and partially hollow

Answer: A



48. Notochord is used in

- A. Attachment of muscles
- B. Development of dorsal nerve cord
- C. Formation of gill slits
- D. Development of kidneys

Answer: A



Watch Video Solution

49. Notochord is

A. Flat plate

B. Rod-like

C. Made of turgid vacuolated cells

D. Both B and C

Answer: D



Watch Video Solution

50. Epiderms of chordates is

A. Single layered

B. Transitional

C. Stratified

D. Absent

Answer: C



Watch Video Solution

51. Gill slits of chordates are

A. Paired

B. Lateral perforations

C. Pharyngeal

D. All the above

Answer: D



Watch Video Solution

52. Nerve cord of chordates is

A. Dorsal

B. Hollow

C. Ventral

D. Both A and B

Answer: D



Watch Video Solution

53. Notochord has been replaced in vertebrates by

- A. Brain and spinal cord
- B. Cranium and vertebral column
- C. Post anal tail
- D. Both A and B

Answer: B



Watch Video Solution

54. Brain and spinal cord of vertebrates have been derived from

- A. Dorsal hollow nerve cord
- B. Ventral solid nerve cord
- C. Lateral nerve cord
- D. Dorsal solid nerve cord

Answer: A



Watch Video Solution

55. Type of coelom present in chordates is

A. Psuedocoelom

B. Schizocoelom

C. Enterocoelom

D. Both A and C

Answer: C



Watch Video Solution

56. In higher chordates, pharyngeal gill slits are

A. Absent

B. Present in the embryo

C. Replaced by abdominal gill slits

D. Replaced by thoracic gill slits

Answer: B



Watch Video Solution

1. Larva of Balanoglossus is

A. Muller's larva

B. Tadpole

C. Tornaria

D. Kentrogen larva

Answer: C



Watch Video Solution

2. Which is not a vertebrate

A. Fish

B. Amphioxus

C. Oryctolagus

D. Sparrow

Answer: B



Watch Video Solution

3. All chordates possess

A. Exoskeleton

B. Limbs

C. Skull

D. Axial skeletal rod of notochord

Answer: D



Watch Video Solution

4. In which group, the notochord is limited to only anterior part or proboscis

A. Hemichordata

B. Urochordata

C. Cephalochordata

D. Mammalia

Answer: A



Watch Video Solution

5. Amphioxus belongs to

A. Craniata

B. Hemichordata

C. Urochordata

D. Cephalochordata

Answer: D



Watch Video Solution

6. Which is an exclusive chordata character

- A. True coelom
- B. Pharyngeal gill slits
- C. Bilateral symmetry
- D. Triploblastic

Answer: B



Watch Video Solution

7. Which one is a link between chordates and nonchordates?

A. Tadpole larva

B. Sphenodon

C. Balanoglossus

D. Crocodilia

Answer: C



Watch Video Solution

8. Notochord occurs only in the larva of

A. Balanoglossus

B. Amphixus

C. Herdmania

D. Cephalodiscus

Answer: C



Watch Video Solution

9. Phosphorescence occurs in

A. Salpa

B. Pyrosoma

C. Petromyzon

D. Rana

Answer: B



Watch Video Solution

10. Group where adults are degenerated while larvae are well developed is

A. Tunicata

B. Agnatha

C. Amphibia

D. Cephalochordata

Answer: A



Watch Video Solution

11. Which is not characteristic of all chordates

?

A. Dorsal nerve cord

B. Diaphragm between thorax and abdomen

C. Coelom

D. Pharyngeal gill slits

Answer: B



Watch Video Solution

12. A primary character of chordates is

A. Paired nerve cord

B. Ganglionated nerve cord

C. Dorsal tubular nerve cord

D. Solid ventral nerve cord

Answer: C



Watch Video Solution

13. Notochord occurs all through the length of body and throughout life in

A. Hemichordata

B. Urochordata

C. Cephalochordata

D. Vertebrata

Answer: C



Watch Video Solution

14. mode of feeding in tunicates is

A. Mixotrophic

B. Parasitic

C. Macrophagous

D. Ciliary feeders

Answer: D



Watch Video Solution

15. Which one belongs to hemichordata

A. Salpa

B. Balanoglossus

C. Petromyzon

D. Myxine

Answer: B



Watch Video Solution

16. Which is true ?

A. Chordates have ventral hollow nerve cord

B. All vertebrates have open circulatory system

C. All chordates are vertebrates

D. All vertebrates are chordates

Answer: D



Watch Video Solution

17. Notochord occurs in the embryonic stage

in

A. Some chordates

B. All chordates

C. All vertebrates

D. Some vertebrates

Answer: B



Watch Video Solution

18. All chordates at one or the other stage possess

A. Two pairs of pentadactyl limbs

B. Movable jaw

C. Vertebral column

D. Pharyngeal gill slits

Answer: D



Watch Video Solution

19. Acorn Worm is

A. Cestoda

B. Trematoda

C. Hemichordata

D. Echinodermata

Answer: C



Watch Video Solution

20. which of the following is a distinct character

A. Chorda dorsalis

B. Cephalization

C. Claws

D. Pharyngostomy

Answer: A



Watch Video Solution

21. The animal who possess notochord throughout life is

Or

Which of the following animals is not a vertebrate

A. Fish

B. Amphixus

C. Snakes

D. Birds

Answer: B



Watch Video Solution

22. Retrogressive metamorphosis is found in

A. Cephalochordata

B. Urochordata

C. Fishes

D. Amphibia

Answer: B



Watch Video Solution

23. Which of the following statements is/are not true

A. In Urochordata, notochord is present only in larval tail.

B. In Cephalochordata, notochord extends

from head to tail region.

C. Branchiosoma belongs to Hemichordata

D. Only one Class of living members, Class Cyclostomata represents the Super Class Agnatha.

A. a, b and d only

B. c, d and a only

C. c only

D. a and d only

Answer: C



24. Three fundamental characters of all chordates are

A. Dorsal tubular nerve cord, diaphragm

notochord

B. Notochord, diaphragm gill slits

C. Gill slits, diaphragm dorsal tubular nerve

cord

D. Notochord, gill slits dorsal tubular nerve
cord

Answer: D



Watch Video Solution

25. In which of the following a notochord is
absent

A. Adult tunicate

B. Amphioxus

C. Myxine

D. Larval tunicate

Answer: A



Watch Video Solution

26. Post anal tail is present in

A. Vertebrates

B. Invertebrates

C. Chordates

D. All the above

Answer: C



Watch Video Solution

27. Crocodile and Penguin are similar to whale and Dog fish in which one of the following characters

A. Possess bony skeleton

B. Possess a solid single stranded central nervous system

C. Have gill slits at some stage

D. Lay eggs and guard them till they hatch

Answer: C



Watch Video Solution

28. stomochord is found in

A. Hemichordata

B. Cephalochordata

C. Urochordata

D. Both B and C

Answer: A



Watch Video Solution

29. Proboscis gland in Balanoglossus is associated with

A. Digestion

B. Respiration

C. Excretion

D. Reproduction

Answer: C



Watch Video Solution

30. All chordates possess the following features except

A. Dorsal hollow nerve cord

B. Pharyngeal gill slits

C. Notochord

D. Amnion

Answer: D



Watch Video Solution

31. Which one of the following statements is incorrect about the occurrence of notochord ?

A. It is present only in larval tail of
ascidians

B. It is absent throughout life in humans
from the very beginning

C. it is replaced by vertebral column in
adult frog

D. It is present throughout life in
Amphioxus

Answer: B



Watch Video Solution

32. Chordates differ from non - chordates in having

A. Notochord

B. Dorsal hollow nerve cord

C. Pharyngeal gill slits

D. All the above

Answer: D



Watch Video Solution

33. Paedomorphosis occurs in

A. Oikopleura

B. Salpa

C. Balanoglossus

D. Branchiostoma

Answer: A



Watch Video Solution

34. Ascidia belongs to subphylum

A. Cephalochordata

B. Craniata

C. Urochordata

D. None of these

Answer: C



Watch Video Solution

35. Which is not a characteristic of chordates

A. Pharyngeal gill slits

B. Dorsal nerve cord

C. Notochord

D. Diploblastic

Answer: D



Watch Video Solution

36. Statement (S) Lancelets are jawless primitive fish like vertebrates

Statement (R) In lancelets notochord tubular nerve cord and pharyngeal gill slits are present throughout their life

A. S is correct but R is wrong

B. S is wrong but R is correct

C. Both S and R are correct and R is correct

explanation of S

D. Both S and R are correct but R is not correct explanation of S

Answer: B



Watch Video Solution

37. Assertion (A) It is necessary to study life history of an ascidian to consider it as chordate

Reason (R) Tadpole larva of ascidians have a notochord that disappears in the adult

- A. Both A and R are true and R is the correct explanation of A
- B. Both A and R are true but R is not the correct explanation of A
- C. A is true but R is false
- D. A is false but R is true

Answer: A



Watch Video Solution

38. Select the animal which exhibits retrogressive metamorphosis

A. Bufo

B. Amphioxus

C. Limulus

D. Herdmania

Answer: D



Watch Video Solution

39. Assertion Branchiostoma and Balanoglossus are bilaterally symmetrical and triploblastic animals

Reason They are exclusively marine and possess notochord



Watch Video Solution

40. Velum is found in

A. Herdmania

B. Amphioxus

C. Branchiostoma

D. Both B and C

Answer: D



Watch Video Solution

41. One of these is not a feature of nonchordates

A. Absence of postanal tail

B. Ventrally located central nervous system

C. Absence of notochord

D. Ventrally of located heart

Answer: D



Watch Video Solution

42. Hemichordates have close affinities with which of the following phylum

A. Annelida

B. Arthropoda

C. Echinodermata

D. Mollusca

Answer: C



Watch Video Solution

43. Which one of the following features is common is Earthworm, Butterfly Spider and Prawn

A. Setae

B. Antennae

C. Ventral nerve cord

D. Nephridia

Answer: C



Watch Video Solution

44. Retrogressive metamorphosis is peculiar to

A. Hemichordata

B. Cephalochordata

C. Urochordata

D. Vertebrata

Answer: C



Watch Video Solution

45. An important characteristic that hemichordates share with chordates is

A. Absence of notochord

B. Ventral tubular nerve cord

C. Pharynx with gill slits

D. Pharynx without gill slits

Answer: C



Watch Video Solution

Cyg

1. Hemichordates constitute a connecting link between

A. Protochordates and Chordates

B. Echinoderms and Chordates

C. Molluscs and Echinoderms

D. Molluscs and Chordates

Answer:



Watch Video Solution

2. Valveless single chambered heart is found in

A. Herdmania

B. Scoliodon

C. Catla

D. Amphioxus

Answer:



Watch Video Solution

3. Vanadocytes occur in

A. Cephalochordates

B. Urochordates

C. Hemichordates

D. All the above

Answer:



Watch Video Solution

4. Sea Potato is the common name of

A. Branchiostoma

B. Doliolum

C. Herdmania

D. Balanoglossus

Answer:



Watch Video Solution

5. Stomochord is not similar to notochord because it is

A. Hollow

B. Outgrowth of gut

C. Outgrowth of nerve cord

D. Ingrowth of body wall

Answer:



Watch Video Solution

6. Notochord occurs between

A. Body wall and dorsal nerve cord

B. Dorsal nerve cord and alimentary canal

C. Ventral nerve cord and alimentary canal

D. Ventral nerve cord and body wall

Answer:



Watch Video Solution

7. In *Balanoglossus* excretion occurs with the help of

A. Protonephridium

B. Kidney

C. Proboscis gland

D. Both A and C

Answer:



Watch Video Solution