



# BIOLOGY

## BOOKS - S DINESH & CO BIOLOGY (HINGLISH)

### LEAF

#### Mcq

1. Part of the leaf modified into tendril in Clematic is

A. Petiole

B. Rachis

C. Petiolules

D. All the above.

**Answer: D**



**Watch Video Solution**

**2. Petiole is modified into tendril in**

A. Gloriosa

B. Passiflora

C. Nepenthes

D. Luffa.

**Answer: C**



**Watch Video Solution**

**3.** In *Opuntia*, the leaves of areoles are modified into

A. spines

B. Glochidia

C. Scales

D. Tendrils.

**Answer: A**



**Watch Video Solution**

**4. Which one of the following has the largest floating leaves**

**A. Banana**



B. Victoria

C. Nelumbo

D. Data Palm.

**Answer: B**



**View Text Solution**

**5. Ptyxis is**

A. Arrangement of leaves on the stem

B. Arrangement of leaves in the bud

C. Folding of lamina in bud

D. Both B and C.

**Answer: C**



**Watch Video Solution**

6. Arrangement of leaves on the stem branches is called.

A. Ptyxis

B. Vernation

C. Prefoliation

D. Phyllotaxy.

**Answer: D**



**Watch Video Solution**

7. Opposite phyllotaxy is present in

A. Banana

B. Calotropis

C. Grass

D. China Rose.

**Answer: B**



**Watch Video Solution**

**8. A swollen leaf base is called**

A. Pulvinus

B. Amplexicaul

C. Windge

D. Sheathing leaf base.

**Answer: A**



**Watch Video Solution**

**9. Petiole modified for photosynthesis is**

A. Cladode

B. Phylloclade

C. Phyllode

D. Tuber.

**Answer: C**



[Watch Video Solution](#)

**10.** In Banyan, bud scales are

- A. Young leaves
- B. Modified leaves
- C. Trichomes
- D. Stipules.

**Answer: D**



[Watch Video Solution](#)

**11. Main function of leaf is**

A. Transpiration

B. Exchange of gases

C. Photosynthesis

D. Cooling.

**Answer: C**



**Watch Video Solution**

12. Leaf margin is spiny in

A. Poplar

B. Argemone

C. Agave

D. Bryophillum.

**Answer: B**



**View Text Solution**



**13.** Occurrence of more than one type of leaves is

A. Decussate phyllotaxy

B. Heterophylly

C. Aestivation

D. Anisophylly.

**Answer: B**



**View Text Solution**

**14.** Vernation is

- A. Arrangement of veins on the lamina
- B. Arrangement of leaves on the stem
- C. Folding of leaves in bud
- D. Arrangement of leaves in bud.

**Answer: D**



**View Text Solution**

**15.** Compound leaves occur in

A. Mustard

B. Syzygium

C. Wheat

D. Sweet Pea.

**Answer: D**



**Watch Video Solution**

**16.** In *Pisum sativum*, tendrils are

A. Modified leaves

B. Modified upper leaflets

C. Modified lower leaflets

D. Modified stipules.

**Answer: B**



**Watch Video Solution**

**17. Stipules are modified into spines in**

A. Shoe Flower

B. Cotton

C. Zizyphus

D. Dalbergia.

**Answer: C**



**Watch Video Solution**

**18.** A plant showing reticulate venation is

A. Ficus

B. Canna

C. Musa

D. Zea.

**Answer: A**



**Watch Video Solution**

**19.** Parallel venation is found in

A. Mentha

B. Banana

C. Dalbergia

D. Syzygium.

**Answer: B**



**Watch Video Solution**

**20. Whole leaf is modified into tendril in**

- A. *Lathyrus aphaca*
- B. *Lathyrus odoratus*
- C. *Pisum sativum*
- D. *Lens culinaris.*

**Answer: A**



Watch Video Solution

21. A monocot showing reticulate venation is

A. Zea

B. Plum

C. Bambusa

D. Smilax

**Answer: D**



Watch Video Solution



22. In Musa, the venation is

A. Unicostate reticulate

B. Unicostate parallel

C. Multicostate reticulate

D. Multicostate parallel.

**Answer: B**



**Watch Video Solution**

23. Which one of the following possesses winged petiole

A. Citrus

B. Bombax

C. Acacia

D. Asparagus.

**Answer: A**



**Watch Video Solution**

24. A plane with unifoliate leaves is

A. Balanites

B. Citrus

C. Aegle

D. Paris.

**Answer: B**



**Watch Video Solution**

25. The leaf of *Mimosa pudica* is

A. Simple

B. Bifoliate

C. Bipinnate

D. Trifoliate

**Answer: C**



**Watch Video Solution**

**26. Multifoliate leaves are found in**

A. Aegle

B. Paris

C. Oleander

D. Bombax

**Answer: D**



**Watch Video Solution**

**27.** Rachis is modified into a leafy structure called

A. Phyllode

B. Ochrea

C. Phylloclade

D. Phyllome.

**Answer: A**



**Watch Video Solution**

**28.** Lamina is reduced in

A. Xerophytes

B. Mesophytes

C. Hydrophytes

D. Climbers.

**Answer: A**



**Watch Video Solution**

**29. Heterophylly of Limnophila is**

A. Environmental

B. Developmental

C. Habitual

D. Both B and C.

**Answer: A**



**Watch Video Solution**

**30. Heterophylly found in Eucalytus is**

A. Habitual

B. Developmental

C. Adaptive

D. Environmental.



**Answer: B**



**Watch Video Solution**

**31. Phyllotaxy is meant for**

A. Protection of leaves against sunlight

B. Exposure of all the leaves equally to  
sunlight

C. Minimising the number of leaves on a  
branch

D. Maximising the number of leaves on a branch.

**Answer: B**



**Watch Video Solution**

**32. Leaf is imparipinnate in**

A. Rose

B. Cassia

C. Quissqualis

D. Guava.

**Answer: A**



**Watch Video Solution**

**33.** Phyllode is an adaptation to

A. Aquatic environment

B. Halophytic environment

C. Mesophytic environment

D. Xerophytic environment.

**Answer: D**



**Watch Video Solution**

**34. Ochreate stipules occur in**

A. Rose

B. Polygonum

C. Lathyrus aphaca

D. Smilax.

**Answer: B**



Watch Video Solution

**35.** In *Nepenthes* the pitcher is modified

A. Leaf apex

B. Leaf base

C. Lamina

D. Leaf stalk.

**Answer: C**



Watch Video Solution

**36.** Leaves borne on the main stem are called

A. Radical

B. Ramal

C. Petiole

D. Cauline.

**Answer: D**



**Watch Video Solution**

37. In Calotropis the phyllotaxy is

A. Alternate

B. Verticillate

C. Opposite and superposed

D. Opposite and decussate.

**Answer: D**



**Watch Video Solution**

**38.** Leaves are modified into hygroscopic appendages in

A. Tamarix

B. Tamarindus

C. Albizzia

D. Butea.

**Answer: A**



**Watch Video Solution**



**39.** Plants without leaves during one season are

A. Caducous

B. Deciduous

C. Evergreen

D. Semigreen.

**Answer: B**



**Watch Video Solution**

**40.** In spiral phyllotaxy, the number of leaves at each node is

A. One

B. Two

C. Many

D. Three

**Answer: A**



**Watch Video Solution**

41. Unicostate venation is called as

A. Palmate

B. Pinnate

C. Reticulate

D. Parallel.

**Answer: B**



**Watch Video Solution**

**42.** Palmate compound leaf is the one in which the leaflets develop from

- A. Rachis
- B. Tip of petiole
- C. Branch of rachis
- D. Node.

**Answer: B**



**Watch Video Solution**

**43.** A compound leaf with more than thrice pinnate nature is

- A. Multifoliate
- B. Decompound
- C. Quadrifoliate
- D. Tripinnate

**Answer: B**



**Watch Video Solution**

**44.** A unifoliate compound leaf can be differentiated from simple leaf in having

A. Joint

B. Stalk

C. Unicostate reticulate venation

D. Multicostate reticulate venation.

**Answer: A**



**Watch Video Solution**

**45.** The leaves which fall down very soon after their formation are

A. Deciduous

B. Caducous

C. Ramal

D. Cauline.

**Answer: B**



**Watch Video Solution**

**46.** Leaves developing from discoid reduced stem of Radish are

A. Ramal

B. Radical

C. Cauline

D. Deciduous

**Answer: B**



**Watch Video Solution**



47. Leaves arising from stem branches are

A. Radical

B. Cauline

C. Ramal

D. Pelate

**Answer: C**



**Watch Video Solution**

48. A leaf without petiole is

A. Sessile

B. Subsessile

C. Subpetiolate

D. Simple.

**Answer: A**



**Watch Video Solution**

**49.** A sensitive thread-like structure which can coil around a support is

A. Spring

B. Tendril

C. Rachis

D. Twiner.

**Answer: B**



**Watch Video Solution**

**50. Phyllode is a modification of**

A. Petiole

B. Stem

C. Inflorescence

D. Root.

**Answer: A**



**Watch Video Solution**

**51.** Tendrillar stipules occur in

A. Dolichos lablab

B. Acacia

C. Smilax

D. Mango.

**Answer: C**



**Watch Video Solution**

52. Petiole is modified into green leafy structure called

A. Phyllode

B. Phylloclade

C. Cladode

D. Foliaceous petiole.

**Answer: A**



**Watch Video Solution**

**53.** A leaf without petiole is

A. Subpetiolate

B. Sessile

C. Subsessile

D. All the above.

**Answer: B**



**Watch Video Solution**

**54.** Arrangement of leaves on a stem branch is

A. Venation

B. Vernation

C. Ptyxis

D. Phyllotaxy.

**Answer: D**



**Watch Video Solution**

**55.** Occurrence of more than one type of leaves on the same plant is

A. Heterophylly

B. Phyllotaxy

C. Venation

D. Vernation.



**Answer: A**



**Watch Video Solution**

**56.** Spiny leaf margins are found in

A. Opuntia

B. Papaver

C. Argemone

D. Polyalthia.

**Answer: C**



[Watch Video Solution](#)

57. A modification of leaf is

A. Phyllode

B. Phylloclade

C. Caldode

D. Corm.

**Answer: A**



[Watch Video Solution](#)

**58.** Free lateral stipules occur in

A. Mango/Mangifera

B. Maize/Zea

C. Rice/Oryza

D. China Rose/Hibiscus.

**Answer: D**



**Watch Video Solution**

59. Phyllotaxy is arrangement of

A. Leaflets

B. Leaves

C. Stipules

D. Branches.

**Answer: B**



**Watch Video Solution**

**60.** In Acacia species, the first few leaves are pinnately compound. Then there are leaves with flattened petiole and fewer pinnae. The leaf of adult plant has parallel veined flattened petiole and no pinnae. It shows that

A. Leaves of adult plant are reduced to phyllodes while those of the seedling are unreduced.

B. The parallel-veined green structures of the adult plant are phylloclades

C. The plant shows developmental heterophylly, compound in seedling and simple in adult plant

D. The leaves of adult plant are unreduced while they are reduced in the seedling stage.

**Answer: A**



**Watch Video Solution**

61. Finely dissected leaves occur in

A. Free floating plants

B. Rooted floating leaved plants

C. Submerged plants

D. Emerged plants.

**Answer: C**



**Watch Video Solution**

62. Imparipinnate leaf is the one where

A. Leaflets are borne in pairs

B. Leaflets are small

C. Leaflets are large

D. Rachis is terminated by an odd leaflet.

**Answer: D**



**Watch Video Solution**



**63.** In Tamaring (Imli) the pinnate leaf is

A. Tripinnate

B. Bipinnate

C. Pripinnate

D. Imparipinnate.

**Answer: C**



**Watch Video Solution**

**64.** The largest leaf belongs to

A. Nerium

B. Tobacco

C. Victoria/Musa

D. Rafflesia.

**Answer: C**



**Watch Video Solution**

**65.** Leaf apex is modified into tendrial in

A. Smilax

B. Gloriosa

C. Australian Acacia

D. All the above.

**Answer: B**



**Watch Video Solution**

**66.** Stem modified for photosynthetic function by appearing like leaves are known as

Or

Leaves are changed into spines in xerophytic structures Called

A. Phyllode

B. Cladode

C. Phylloclade

D. All the above.

**Answer: C**



Watch Video Solution

67. Phyllotaxis is

- A. Mode of leaf arrangement on stem
- B. Types of roots
- C. Arrangement of sepals and petals in a flower
- D. Type of ovary.

**Answer: A**



Watch Video Solution

68. Approximate diameter of Victoria leaf is

A. 1 m

B. 1.3 m

C. 2 m

D. 3 m.

**Answer: B**



Watch Video Solution

**69.** Petiole is modified into tendril in

A. Passiflora

B. Gloriosa

C. Pisum

D. Clematis

**Answer: D**



**Watch Video Solution**

70. Swollen spongy petiole is present in

A. Hydrilla

B. Eichhornia

C. Ruppia

D. Pistia.

**Answer: b**



**Watch Video Solution**



71. Presence of sheathing leaf base and ligule are characteristic of

- A. Cycas leaf
- B. Fern leaf
- C. Banana leaf
- D. Grass leaf.

**Answer: D**



**Watch Video Solution**

72. In *Lathyrus aphaca*, the leaves are modified into

- A. Spines
- B. Tendrils
- C. Scales
- D. Stem-like structures.

**Answer: B**



**Watch Video Solution**

73. Arrangement of leaves in the bud condition is called

A. Phyllotaxy

B. Ptyxis

C. Vernation

D. Venation.

**Answer: C**



**Watch Video Solution**

74. Occurrence of different types of leaves in Limnophylla is called

- A. Heterophylly
- B. Pseudophylly
- C. Heterophily
- D. Heterotrophy.

**Answer: A**



**Watch Video Solution**

75. Tripinnate leaves occur in

A. Acacia

B. Oxalis

C. Moringa

D. Gynandropsis.

**Answer: C**



**Watch Video Solution**

76. A leaf is identified from

- A. Flat green lamina
- B. Presence of leaf blade and petiole
- C. Presence of axillary bud
- D. Occurrence of chlorophyll.

**Answer: C**



**Watch Video Solution**

77. Swollen lower end of leaf stalk is

A. Petiole

B. Pulvinus

C. Thalamus

D. Disc.

**Answer: B**



**Watch Video Solution**

78. In sweet pea, the tendrils are modified

A. Stem branches

B. Leaflets

C. Leaves

D. Stipules.

**Answer: B**



**Watch Video Solution**



**79.** Parallel venation occurs in

A. Monocots

B. Dicots

C. All angiosperms

D. Ferns.

**Answer: A**



**Watch Video Solution**

**80.** Bombax leaf is

A. Tripinnate

B. Unipinnate

C. Multifoliate

D. Quadrifoliate

**Answer: C**



**Watch Video Solution**

**81.** Name the plant having reticulate venation

A. Musa

B. Mangifera

C. Oryza

D. Canna

**Answer: B**



**Watch Video Solution**

**82.** In *Nepenthes* the pitcher is modified

A. Whole leaf

B. Leaf apex

C. Lamina

D. Petiole

**Answer: C**



**Watch Video Solution**

**83.** Pinnately parrallel venation is found in

A. Canna

B. Grass

C. Zizyphus

D. Castor.

**Answer: A**



**Watch Video Solution**

**84.** Ochreate stipules occur in leafy vegetable

A. Amaranthus

B. Mentha

C. Platanus

D. Rumex.

**Answer: D**



**Watch Video Solution**

**85.** Spiral phyllotaxy in which sixth leaf lies above the first one after completing two circles is

- A. Destichous
- B. Tristichous
- C. Pentastichous
- D. Octastichous.

**Answer: C**



**Watch Video Solution**

**86. Match the columns**

- |                  |            |
|------------------|------------|
| (i) Acicular     | (1) Grass  |
| (ii) Linear      | (2) Nerium |
| (iii) Lanceolate | (3) Banana |
| (iv) Oblong      | (4) Pine   |

**Options**

A. (i) d (ii) a (iii) b (iv) c

B. (i) d (ii) a (iii) c (iv) b

C. (i) d (ii) b (iii) c (iv) a

D. (i) d (ii) c (iii) b (iv) a

**Answer: A**







Watch Video Solution

87. Leaves develop from

- A. Nodes
- B. Internodes
- C. Epidermis
- D. Endodermis

**Answer: A**



Watch Video Solution

**88.** Phyllode is found in

A. Clematis

B. Gloriosa

C. Australian Acacia

D. Dischidia.

**Answer: C**



**Watch Video Solution**

**89.** Pitcher is found in

A. Dionaea

B. Drosera

C. Nepenthes

D. Viscum.

**Answer: C**



**Watch Video Solution**

90. In sarracenia, the insect trapping structure is modified

A. Leaf

B. Leaf base

C. Stipule

D. Axillary shoot

**Answer: A**



**Watch Video Solution**

91. Which one is modified leaf

A. Pitcher of *Nepenthes*

B. Tendril of *Pisum sativum*

C. Spine of Cactus

D. All the above.

**Answer: D**



**Watch Video Solution**

92. Pick up the leaf modification

A. Cladode

B. Phyllode

C. Corm

D. Phylloclade

**Answer: B**



**Watch Video Solution**

**93.** Bladders of *Utricularia* and pitcher of *Nepenthes* are modifications of

A. Leaves

B. Stem

C. Roots

D. Flowers.

**Answer: A**



**Watch Video Solution**

**94.** Smallest leaf occurs in

A. Victoria

B. Wolffia

C. Cycas

D. Spiraea.

**Answer: B**



**Watch Video Solution**

**95.** Identify in order, the plants showing alternate, opposite and whorled phyllotaxy

A. Chine Rose, Calotropis and Nerium



B. Chine Rose, Nerium and Calotropic

C. Nerium, Chine Rose and Calotropis

D. Nerium, Calotropis and Chine Rose

**Answer: A**



**Watch Video Solution**

**96.** Pulvinus is found in

A. Calotropis

B. Ocimum

C. Legume plants

D. Alstonia

**Answer: C**



**Watch Video Solution**

**97. Venation is generally reticulate in**

A. Monocot plants

B. Bryophytes

C. Thallophytes

D. Dicot plants.

**Answer: D**



**Watch Video Solution**

**98.** Venation in monocots is

A. Pinnate reticulate

B. Palmate reticulate

C. Pinnate parallel

D. Parallel.

**Answer: D**



**Watch Video Solution**

**99.** Stipules are modified into spines in

- A. Citrus and Euphorbia
- B. Euphorbia and Zizyphus
- C. Zizyphus and Bougainvillea
- D. Citrus and Bougainvilla.

**Answer: B**



Watch Video Solution

**100.** A compound leaf which appears simple due to suppression of 1-2 lateral leaflets is found in

- A. Hardwickia
- B. Parkinsonia
- C. Citrus
- D. Coriandrum.

**Answer: C**



Watch Video Solution

**101.** In *Utricularis*, the leaves are modified to form

A. Bladders

B. Tendrils

C. Hooks

D. Pitchers

**Answer: A**



**102.** In *Opuntia*, spines are modification of

- A. Epidermal hair
- B. Stem
- C. Flowers
- D. Leaves of axillary bud.

**Answer: D**



**103.** The leaves are modified into tendrils, hooks, pitcher and bladder in the following plants respectively

A. Sweet Pea, Cat's Nail, Nepenthes,  
Utricularia

B. Sweet Pea, Cat's Nail, Utricularia,  
Nepenthes

C. Nepenthes, Sweet Pea, Cat's Nail,  
Utricularia



D. Utricularia, Nepenthes, Cat's Nail, Sweet

Pea.

**Answer: A**



**Watch Video Solution**

**104.** In distichous condition

A. First leaf stands over the second

B. Second leaf stands over the first

C. Third leaf stands over the first

D. Fourth leaf stands over the first.

**Answer: C**



**Watch Video Solution**

**105.** Leaves of Nelumbo plant are

- A. Epistomatic
- B. Hypostomatic
- C. Amphistomatic
- D. None of the above.

**Answer: A**



**Watch Video Solution**

**106.** Phyllotaxy is decussate in

- A. Nerium indicum
- B. Pisum sativum
- C. Hibiscus rosa-sinensis
- D. Catharanthus roseus.

**Answer: D**



[Watch Video Solution](#)

**107.** Which ones show stipular modifications

- (a) Spines of Zizyphus
- (b) Tendrils in Smilax
- (c) Tendrils in Nepenthes
- (d) Spines in Argemone
- (e) Thorn in B



[Watch Video Solution](#)

**108.** Phyllotaxy in Calotropis is

A. Alternate

B. Opposite

C. Whorled

D. None of the above.

**Answer: B**



**Watch Video Solution**

**109.** Multicostate parallel venation occurs in

A. Banana and Canna

B. Mango and Peepal

C. Grasses and Palms

D. Caster and Tapioca

**Answer: C**



**Watch Video Solution**

**110.** Leaves are modified into spines in

A. Nepenthes

B. Australian Acacia

C. Opuntia

D. Utricularia

**Answer: C**



**Watch Video Solution**

# 111. Find the correct combination

I		II	
<i>a</i>	Entire leaf with stipules modified into spines	<i>i</i>	<i>Clematis</i>
<i>b</i>	Leaf except stipules modified into tendrils	<i>ii</i>	<i>Citrus</i>
<i>c</i>	Stipules modified into tendrils	<i>iii</i>	<i>Euphorbia</i>
<i>d</i>	First leaf of axillary bud modified into spine	<i>iv</i>	<i>Lathyrus</i>
		<i>v</i>	<i>Smilax</i>

- A. a-iii, b-i, c-iv, d-ii
- B. a-ii, b-iii, c-i, d-v
- C. a-v, b-ii, c-i, d-iii
- D. a-iii, b-iv, c-v, d-ii.

**Answer: D**





**112.** Find the correct match

- A. Mustard-Leaves opposite
- B. Mustard-Leaves alternate
- C. Guava-Leaves alternate
- D. Guava-Leaves whorled.

**Answer: B**



Watch Video Solution

**113.** Whorled phyllotaxy with simple reticulate leaves occurs in

A. Alstonia

B. Guava

C. Calotropis

D. Mustard

**Answer: A**



**Watch Video Solution**

114. Bladderwort is

A. Drosera

B. Nepenthes

C. Dionaea

D. Utricularia.

**Answer: D**



**Watch Video Solution**

115.  $120^\circ$  phyllotaxy is found in

A. Distichous

B. Tristichous

C. Pentastichous

D. Octastichous.

**Answer: B**



**Watch Video Solution**

**116.** Study the following statements and select the correct option

(A) Buds are present in the axil of leaflets of

the compound leaf

(B) Pulvinus leaf-base is present in some leguminous plants

(C) In Alstonia, the petioles expand, become green and synthesize food

(D) Opposite phyllotaxy is seen in guava.

A. b and d

B. a and c

C. a and d

D. b, c and d

**Answer: A**



[Watch Video Solution](#)

**117.** In the leaves, veins are useful for

- A. Transport of water and minerals
- B. Mechanical support
- C. Transport of organic nutrients
- D. All the above.

**Answer: D**



[Watch Video Solution](#)

**118.** Angle of divergence in tristichous phyllotaxy is  $140^\circ$

A.  $140^\circ$

B.  $135^\circ$

C.  $180^\circ$

D.  $120^\circ$

**Answer: D**



**Watch Video Solution**

**119.** Petiole part of the leaf is known as

A. Epipodium

B. Mesopodium

C. Hypopodium

D. None of the above.

**Answer: B**



**Watch Video Solution**



**120.** Pulvinate leaf base is found in

A. *Lycopersicum*

B. *Trifolium*

C. *Nicotiana*

D. *Petunia*.

**Answer: B**



**Watch Video Solution**

**121.** Identify the correct pair of plants with odd number of leaflets in compound leaf

A. Hardwickia, Gynandropsis

B. Citrus, Aegle marmelos

C. Marsilea, Gynandropsis

D. Aegle marmelos, Hardwickia

**Answer: B**



**Watch Video Solution**

**122. Identify the wrong statement**

- A. Stipules of Lathyrus are persistent
- B. Phyllotaxy in Trillium is alternate
- C. Venation in Calophyllum is parallel
- D. Cauline leaves are found in Cocos.

**Answer: B**



**Watch Video Solution**

**123.** How many plants among China rose, Ocimum, sunflower, mustard, Alstonia, guava, Calotropis and Nerium (Olender) have opposite phyllotaxy

A. Four

B. Five

C. Two

D. Three

**Answer: D**



Watch Video Solution

124. I. Pulvinus leaf base is present in some leguminous plants. II. Whorled phyllotaxy is seen in Calotropis. III. In Australian Acacia, the petioles expand, become green and synthesise food, IV. A bud is present in the axils of leaflets of a compound leaf.

A. I and IV correct, II and III wrong

B. II and III correct, I and IV wrong

C. I and III correct, II and IV wrong

D. III and IV correct, I and II wrong

Answer: C



Watch Video Solution

125. Match the lists

I		II	
(a) <i>Alstonia</i>	I.	I.	Roots at lower nodes of stem
(b) <i>Ananas sativus</i>	II.	II.	Leaflets are attached at common point in leaf
(c) Sugarcane	III.	III.	Swollen placenta
(d) <i>Bombax ceiba</i>	IV.	IV.	More than two leaves at every node
	V.	V.	Underground lateral branches producing aerial leafy shoots.

The correct match is

A. a-IV, b-V, c-I, d-II

B. a-V, B-III, c-I, d-II

C. a-V, b-III, c-II, d-IV

D. a-IV, b-II, c-V, d-I

**Answer: A**



**Watch Video Solution**

**126.** Multicostate divergent reticulate venation  
is seen in ....leaf

A. Zizyphus

B. Bamboo

C. Castor

D. Mango.

**Answer: C**



**Watch Video Solution**

**127.** Leaf tendrils are found in

A. Peas



B. Cucumber

C. Grape vine

D. All the above.

**Answer: A**



**Watch Video Solution**

**128.** Consider the following statements :

(a) In leguminous plants, leaf base becomes swollen, called pulvinus

(b) The fleshy leaves of Onion and Garlic store

food

(c) The buds in Australian acacia become green and synthesise food

(d) In Alstonia, leaves show alternate phyllotaxy.

Of the above statements

A. (b) and (d) are correct

B. (a) and (c) are correct

C. (a) and (b) are correct

D. (a) and (d) are correct

**Answer: C**



Watch Video Solution

**129.** Match the columns and choose the right option

1. Pneumatophores	(a) Axillary buds
2. Tendrils in pea	(b) Roots
3. Thorns in <i>Citrus</i>	(c) Leaves

A. 1-(b), 2-(a), 3-(c)

B. 1-(c), 2-(a), 3-(b)

C. 1-(b), 2-(c), 3-(a)

D. 1-(a), 2-(b), 3-(c).

**Answer: D**



**Watch Video Solution**

**130.** Identify the correct combination

A. Neem - Absence of buds in the axile of  
leaflets-Pinnately compound leaf

B. Sunflower - Flowers brought to same  
height due to varied lengths of pedicels -  
Involucre of bracts

C. Carrot - Flowers brought to same height

due to varied lengths of pedicels -

Involucre of bracts

D. Pistia - Discoid stem - Lateral branch with

many internodes.

**Answer: A and C**



**Watch Video Solution**

**131.** It is an example of pinnate type venation of leaf blade

A. Cosmos

B. Castor

C. Tapioca

D. Semul.

**Answer: A**



**Watch Video Solution**

**132.** This has glandular hair

A. Calotropis

B. Castor

C. Lemon

D. Yucca.

**Answer: B**



**Watch Video Solution**

**133.** Which of these plants has pinnately compound leaf

A. Alstonia

B. Calotropis

C. Guava

D. neem

**Answer:**



**Watch Video Solution**



**134.** Match the columns and find the correct option

I	II
(i) <i>Dahlia</i>	(a) Eyes
(ii) <i>Solanum tuberosum</i>	(b) Runner
(iii) <i>Begonia</i>	(c) Fasciculated tuberous roots
(iv) <i>Cycotodon</i>	(d) Epiphyllous buds

A. i-c, ii-a, iii-b, iv-d

B. i-d, ii-a, iii-b, iv-c

C. i-c, ii-a, iii-d, iv-b

D. i-b, ii-c, iii-a, iv-d.

**Answer: C**



**Watch Video Solution**

**135.** Identify the wrong statements from the following : (i) Branches of limited growth in *Asparagus* perform photosynthesis (ii) Petiole of Australian *Acacia* helps in climbing (iii) Floral buds of *Agave* store food materials (iv) Aerial roots of *Taeniophyllum* help in vegetative propagation

A. *I, iv*

B. *I, ii*

C. *iii, iv*

D. *ii, iv*.

**Answer: D**



**Watch Video Solution**

**136.** Assertion : In opposite phyllotaxy, the leaves are borne on the opposite sides of a single node

Reason : Opposite phyllotaxy is seen in China

Rose and Oleander

A. both true but reason is not correct  
explanation

B. assertion true but reason is wrong

C. both are wrong

D. both are wrong

**Answer: C**



**Watch Video Solution**

1. In grasses the ligule occurs

A. At leaf base

B. Between leaf base and lamina

C. Between leaf base and petiole

D. Between petiole and lamina.

**Answer:**



**Watch Video Solution**

2. In *Nelumbium*/*Victoria*/*Tropaeolum* the leaf blade is

A. Petate

B. Centric

C. Unifacial

D. Ligulate

**Answer: A**



**Watch Video Solution**

3. In examples of interpetiolar stipules, the number of leaves and stipules at each node are

A. One and two

B. Two and two

C. Two and one

D. One and one.

**Answer:**



**Watch Video Solution**

4. Arrangement of leaves in the bud condition is called

A. Phyllotaxy

B. Insertion

C. Ptyxis

D. Vernation.

**Answer:**



**Watch Video Solution**



5. Number of orthostichies present in opposite and decussate phyllotaxy is

A. Two

B. Four

C. One

D. Three.

**Answer: B**



**Watch Video Solution**

6. A monocot leaf having reticulate venation is

A. Dioscorea

B. Alocasia

C. Amilax

D. All the above.

**Answer:**



**Watch Video Solution**

7. A dicot leaf having parallel venation is

A. Eryngium

B. Zizyphus

C. Ricinus

D. Cassia.

**Answer:**



**Watch Video Solution**

**8. A simple pinnate compound leaf is**

A. Decomound

B. Unipinnate

C. Bipinnate and tripinnate

D. Both B and C.

**Answer: b**



**Watch Video Solution**

9. In common weed Parthenium the leaf is compound

A. Multifoliate

B. Tripinnate

C. Decompound

D. Bipinnate.

**Answer:**



**Watch Video Solution**

**10. Branched leaf spines occur in**

A. Acacia

B. Opuntia

C. Barberry

D. Asparagus.

**Answer:**



**Watch Video Solution**