



CHEMISTRY

BOOKS - NCERT CHEMISTRY (HINGLISH)

HYDROCARBONS

Multiple Choice Questions Mcqs

1. Arrange the following in decreasing order of their boiling points.

- (A). N-butane
- (B). 2-methylbutane
- (C). N-pentane
- (D). 2,2-dimethylpropane

A. $A > B > C > D$

B. $B > C > D > A$

C. $D > C > B > A$

D. $C > B > D > A$

Answer: D




Watch Video Solution

2. Arrange the halogens F_2 , Cl_2 , Br_2 , I_2 , in order of their increasing reactivity with alkanes.

A. $I_2 < Br_2 < Cl_2 < F_2$

B. $Br_2 < Cl_2 < F_2 < I_2$





C. $F_2 < Cl_2 < Br_2 < I_2$



Answer: A

 [Watch Video Solution](#)


3. The increasing order of reduction of alkyl halides with zinc and dilute HCl is



Answer: B

 [Watch Video Solution](#)

4. The correct IUPAC name of the following alkane is



- A. 3, 6-diethyl-2-methyloctane
- B. 5-isopropyl -3-ethyloctane
- C. 3-ethyl-5-isopropyloctane
- D. 3-isopropyl-6-ethyloctane


Answer: A



Watch Video Solution

5. The addition of HBr of 1-butene gives a mixture of products

A, B and C



(C) $CH_3 - CH_2 - CH_2 - CH_2 - Br$

The mixture consists of





- A. A and B as major and C as minor products
- B. B as major, A and C as major products
- C. B as minor, A and C as major products
- D. A and B as minor and C as major products

Answer: A



Watch Video Solution

6. Which of the following will not show geometrical isomerism ?




Answer: D



Watch Video Solution

7. Arrange the following hydrogen halides in order of their decreasing reactivity with propene.







Answer: C



Watch Video Solution

8. Arrange the carbanions, $(CH_3)_3\bar{C}$, $\bar{C}Cl_3$, $(CH_3)_2\bar{C}H$, $C_6H_5\bar{C}H_2$, in order of their decreasing stability









Answer: B

 [Watch Video Solution](#)

9. Arrange the following alkyl halides in decreasing order of the rate of β -elimination reaction with alcoholic KOH.



$$B. C > B > A$$




$$C. B > C > A$$

$$D. A > C > B$$

Answer: D

 [Watch Video Solution](#)

10. Which of the following reactions of methane is incomplete combustion:








Answer: C



Watch Video Solution

Multiple Choice Questions More Than One Options

1. Some oxidation reactions of methane are given below. Which of them is/are controlled oxidation reactions?







Answer: C::D



Watch Video Solution

2. Which of the following alkenes on ozonolysis give a mixture of ketones only?



Answer: C



Watch Video Solution

3. Which are the correct IUPAC names of the following compound?




- A. 5-Butyl-4-isopropyldecane
- B. 5-Ethyl-4-propyldecane
- C. 5-sec-Butyl-4-iso-propyldecane
- D. 4-(1-methylentyl)-5-(1-methylpropyl)-decane

Answer: C::D



Watch Video Solution



4.

Which are the correct IUPAC names of the following compound?

- A. 5-(2', 2'-Dimethylpropyl)-decane
- B. 4-Butyl-2,2-dimethylnonane
- C. 2,2-Dimethyl-4-pentyl-octane
- D. 5-neo-Pentyldecane

Answer: A::D



Watch Video Solution

5. For an electrophilic substitution reaction , the presence of a halogen atom in the benzene ring

A. deactivates the ring by inductive effect

B. deactivates the ring by resonance

C. increases the charge density at ortho and para position relative to meta position by resonance.

D. directs the incoming electrophile to meta position by increasing the charge density relative to ortho and para position.

Answer: A::C



Watch Video Solution

6. In an electrophilic substitution reaction of nitrobenzene, the presence of nitro group.....

A. deactivates the ring by inductive effect

B. activates the ring by inductive effect

C. decreases the charge density at ortho and para position of the ring relative to meta position by resonance

D. increases the charge density at meta position relative to the ortho and para positions of the ring by resonance

Answer: A::C



[Watch Video Solution](#)

7. Which of the following are correct ?

A. $CH_3 - O - CH_2^{\oplus}$ is more stable than $CH_3 - CH_2^{\oplus}$

B. $(CH_3)_2CH^{\oplus}$ is less stable than $CH_3 - CH_2 - CH_2^{\oplus}$

C.


$C_2 = CH - CH_2^{\oplus}$ is more stable than $CH_3 - CH_2 - CH_2^{\oplus}$


D. $CH_2 = CH^{\oplus}$ is more stable than $CH_3 - CH_2^{\oplus}$

Answer: A::C


 [Watch Video Solution](#)

8. Four structures are given in options (a) to (d) . Examine them and select the aromatic structures.






B.



C.



D.

Answer: A::C



Watch Video Solution

9. The molecules having dipole moment are :

A. 2,2-Dimethylpropane

B. trans-Pent-2-ene

C. cis-Hex-3-ene

D. 2,2,3,3-Tetramethylbutane

Answer: B::C

 [Watch Video Solution](#)

Short Answer Type Questions

1. Why do alkenes prefer to undergo electrophilec addition reaction while arenes prefer electrophilic substitution reactions ? Explain.

 [Watch Video Solution](#)

2. Alkynes on reduction with sodium in liquid ammonia form trans alkenes. Will the butene thus formed on reduction of 2-butyne show the geometrical isomerism ?

 [Watch Video Solution](#)

3. Rotation around carbon-carbon single bond of ethane is not completely free. Justify the statement

 [Watch Video Solution](#)


4. Draw Newman and sawhorse projections for the eclipsed and staggered conformations of ethane. Which of these conformations is more stable and why ?

 [Watch Video Solution](#)

5. The intermediate carbocation formed in the reactions of HI, HBr, and HCl with propene is the same and the bond energy of HCl, HBr, and HI is 430.5 KJmol^{-1} , 363.7 KJmol^{-1} and 296.8 KJmol^{-1} respectively. What will be the order of reactivity of these halogen acids?

 [Watch Video Solution](#)

6. What will be the product obtain as a result of the following reaction?



 [Watch Video Solution](#)

7. How will you convert benzene into (a) p-nitrobromobenzene
(b) m-nitrobromobenzene

 [Watch Video Solution](#)

8. Arrange the following set of compounds in the order of their decreasing relative reactivity with an electrophile . Give reason.

Arrange the following in
following order



Anisole

>



Chlorobenzene

>



Nitrobenzene

 [Watch Video Solution](#)

9. Despite their-I effect, halogens are o- and p- directing in haloarenes. Explain .



[Watch Video Solution](#)

10. Why does presence of a nifro group make the benzene ring less reactive in comparison to the unsubstituted benzene ring . Explain .




[Watch Video Solution](#)

11. Suggest a route for the preparation of nitrobenzene starting from acetylene ?






[Watch Video Solution](#)




12. Predict the major product(S) of the following reactions and explain their formation .



 [Watch Video Solution](#)

13. Nucleophiles and electrophiles are reaction intermediates having electron rich and electron deficient centres respectively . Hence, they tend to attack electron deficient and electron rich centres respectively. Classify the following species as electrophiles and uncheophiles.





 [Watch Video Solution](#)

14. The relative reactivity of 1° , 2° and 3° hydrogen's towards chlorination is 1:3.8:5. Calculate the percentages of all monochlorinated products obtained from 2-methylbutane.

 [Watch Video Solution](#)

15. Write the structures and names of products obtained in the reactions of sodium with a mixture of 1-iodo-2-methylpropane and 2-iodopropane.

 [Watch Video Solution](#)

16. Write hydrocarbon radicals that can be formed as intermediates during monochlorination of 2-methylpropane ?

Which of them is more stable? Give reasons.

 [Watch Video Solution](#)

17. An alkane C_8H_{18} is obtained as the only product on subjecting a primary alkyl halide to Wurtz reaction. On monobromination this alkane yields a single isomer of a tertiary bromide. Write the structure of alkane and the tertiary bromide.

 [Watch Video Solution](#)

18. The ring systems having following characteristics are aromatic.

(i) Planar ring containing conjugated π bonds .


(ii) complete delocalisation of the π -electron in ring system i.e.

, each atom in the ring has unhybridised p-orbital , and

(iii) Presence of $(4n + 2)\pi$ -electrons in the ring where n is an integer($n = 0, 1, 2, \dots$) [Huckel rule]. Using this information







classify the following compounds as aromatic/non-aromatic.

QUESTION



Watch Video Solution

19. Which of the following compounds are aromatic according to Huckel's rule?



A. A,B,C

B. B,D,E,F

C. B,D,E

D. All are correct

Answer: C



Watch Video Solution

20. Suggest a route to prepare ethyl hydrogensulphate ($CH_3 - CH_2 - OSO_2 - OH$) starting from ethanol (C_2H_5OH).

 Watch Video Solution

Matching The Columns

1. Match the reagent from Column I which on reaction with $CH_3 - CH = CH_2$ gives some product given in Column II as per the codes given below

Column I	Column II
A. $O_3 / Zn + H_2O$	1. Acetic acid and CO_2
B. $KMnO_4 / H^+$	2. Propan-1-ol
C. $KMnO_4 / OH^-$	3. Propan-2-ol
D. H_2O / H^+	4. Acetaldehyde and formaldehyde
E. $B_2H_6 / NaOH^+$ and H_2O_2	5. Propane-1, 2-diol



2. Match the hydrocarbons in column I with the boiling points given in column II.

Column I

Column II

(i). n-Pentane

(a). 282.5 K

(ii). iso-Pentane

(b). 309 K

(iii). neo-Pentane

(c). 301 K



Watch Video Solution

3. Match the following reactants in Column I with the corresponding reaction products in Column II.

Column I	Column II
A. Benzene + $\text{Cl}_2 \xrightarrow{\text{AlCl}_3}$	1. Benzoic acid
B. Benzene + $\text{CH}_3\text{Cl} \xrightarrow{\text{AlCl}_3}$	2. Methyl phenyl ketone
C. Benzene + $\text{CH}_3\text{COCl} \xrightarrow{\text{AlCl}_3}$	3. Toluene
D. Toluene $\xrightarrow{\text{KMnO}_4 / \text{NaOH}}$	4. Chlorobenzene
	5. Benzene hexachloride



Watch Video Solution

4. Match the reactions given in Column I with the reaction types in Column II.

Column I	Column II
A. $\text{CH}_2 = \text{CH}_2 + \text{H}_2\text{O} \xrightarrow{\text{H}^+} \text{CH}_3\text{CH}_2\text{OH}$	1. Hydrogenation
B. $\text{CH}_2 = \text{CH}_2 + \text{H}_2 \xrightarrow{\text{Pd}} \text{CH}_3 - \text{CH}_3$	2. Halogenation
C. $\text{CH}_2 = \text{CH}_2 + \text{Cl}_2 \longrightarrow \text{Cl} - \text{CH}_2 - \text{CH}_2 - \text{Cl}$	3. Polymerisation
D. $3\text{CH} \equiv \text{CH} \xrightarrow[\text{Heat}]{\text{Cu tube}} \text{C}_6\text{H}_6$	4. Hydration
	5. Condensation



Watch Video Solution

Assertion And Reason

1. Assertion (A) The compound tetraene has the following structural formul.



It is cyclic and has conjugated 8π -electron system but it is not an aromatic compound.

Reason (R) $(4n + 2)\pi$ electrons rule does not hold good and ring is not planar.

- A. Both A and B are correct and R is the correct explanation of A
- B. Both A and R are not correct
- C. Both A and R are not correct
- D. A is not correct but R is correct

Answer: a



Watch Video Solution

2. Assertion (A) Toluene on Friedal Crafts methylation gives o - and p-xylene.

Reason (R) CH_3 -group bonded to benzene ring increases density at o - and p- position.

- A. Both A and B are correct and R is the correct explanation of A
- B. Both A and R are not correct
- C. Both A and R are not correct
- D. A is not correct but R is correct

Answer: a



Watch Video Solution

3. S-I: Nitration of benzene with nitric acid requires the use of concentrated sulphuric acid

S-II: The mixture of concentrated sulphuric acid and concentrated nitric acid produces the electrophile, nitronium ion.

- A. Both A and B are correct and R is the correct explanation of A
- B. Both A and R are not correct
- C. Both A and R are not correct
- D. A is not correct but R is correct

Answer: a



Watch Video Solution

4. Assertion (A) Among isomeric pentanes, 2, 2-dimethylpentane has highest boiling point.

Reason (R) Branching does not affect the boiling point.

- A. Both A and R are correct and R is the correct explanation of A
- B. Both A and R are not correct
- C. Both A and R are not correct
- D. A is not correct but R is correct

Answer: c



Watch Video Solution

Long Answer Type Questions

1. An alkyl halide C_5H_{11} (A) reacts with ethanolic KOH to give an alkene 'B' which reacts with Br_2 to give a compound 'C' which on dehydrohalogenation gives an alkyne 'D'. On treatment with sodium metal in liquid ammonia one mole of 'D' gives one mole of the sodium salt of 'D' and half a mole of hydrogen gas. Complete hydrogenation of 'D' yields a straight chain alkane. Identify A, B, C and D. Give the reactions involved.



[Watch Video Solution](#)

2. 448 mL of a hydrocarbon (A) having C (87.80%), H (12.19%) weight 1.64 g at NTP. On hydrogenation it gives 2 methyl pentane. Treatment of (A) with acidic $HgSO_4$ gives a new compound (B) of molecular weight $C_6H_{12}O$. Compound (A) does not react with ammoniacal $AgNO_3$. What is the structure of (A)?

 [Watch Video Solution](#)

3. An unsaturated hydrocarbon 'A' adds two molecules of H_2 and on reductive ozonolysis gives butane-1, 4-dial, ethanal and propanone. Give the structure of 'A', write its IUPAC name and explain the reactions involved.

 [Watch Video Solution](#)

4. In the presence of peroxide addition of HBr to propene takes place according to anti Markownikoff's rule but peroxide effect is not seen in the case of HCl and HI. Explain.

 [Watch Video Solution](#)