



## PHYSICS

### BOOKS - NCERT PHYSICS (HINGLISH)

#### ELECTROMAGNETIC WAVES

##### Electromagnetic Waves

1. One requires  $11eV$  of energy to dissociate a carbon monoxide molecule into carbon and oxygen atoms. The minimum frequency of the appropriate electromagnetic radiation to achieve the dissociation lies in.

- A. visible region
- B. infrared region
- C. ultraviolet region
- D. microwave region

**Answer: C**



**Watch Video Solution**

2. A linearly polarised electromagnetic wave given as  $E = E_0 \hat{i} \cos(kz - \omega t)$  is incident normally on a perfectly reflecting wall  $z = a$ . Assuming that the material of the wall is optically inactive, the reflected wave will be given as

A.  $E_r = E_0 \hat{i} (kz - \omega t)$

B.  $E_r = E_0 \hat{i} \cos(kz + \omega t)$

C.  $E_r = -E_0 \hat{i} \cos(kz + \omega t)$

D.  $E_r = E_0 \hat{i} \sin(kz - \omega t)$

**Answer: B**



**Watch Video Solution**

3. Light with an energy flux  $20 \text{ W/cm}^2$  falls on a non-reflecting surface at normal incidence. If the surface has an area of  $30 \text{ cm}^2$ , the total momentum delivered (for complete absorption) during 30 minutes is

A.  $36 \times 10^{-5} \text{ kg} - \text{m} / \text{s}$

B.  $36 \times 10^{-4} \text{ kg} - \text{m} / \text{s}$

C.  $108 \times 10^4 \text{ kg} - \text{m} / \text{s}$

D.  $1.08 \times 10^7 \text{ kg} - \text{m} / \text{s}$

**Answer: B**



**Watch Video Solution**

4. The electric field intensity produced by the radiations coming from 100W bulbs at a 3m distance is

E. The electric field intensity produced by the radiations coming from 50W bulb at the same distance is

A.  $\frac{E}{2}$

B.  $2E$

C.  $\frac{E}{\sqrt{2}}$

D.  $\sqrt{2E}$

**Answer: C**



**Watch Video Solution**

5. If  $E$  and  $B$  represent electric and magnetic field vectors of the electromagnetic wave, the direction of propagation of electromagnetic wave is along.

A.  $E$

B.  $B$

C.  $B \times E$

D.  $E \times B$

**Answer: D**



**Watch Video Solution**

6. The ratio of contributions made by the electric field and magnetic field components to the intensity of an  $EM$  wave is.

A.  $C : 1$

B.  $C^2 : 1$

C. 1:1

D.  $\sqrt{C}:1$

**Answer: C**



**Watch Video Solution**

7. An  $EM$  wave radiates out wards from a dipole antenna with  $E_0$  as the amplitude of its electric field vector. The electric field  $E_0$  which transports significant energy from the source falls off as

A.  $\frac{1}{r^3}$

B.  $\frac{1}{r^2}$

C.  $\frac{1}{r}$

D. remains constant

**Answer: C**



**Watch Video Solution**

**8.** An EM wave travels in vacuum along z direction:

$$\vec{E} = (E_1 \hat{i} + E_2 \hat{j}) \cos(kz - \omega t). \text{ Choose the correct option from the following :}$$

A. The associated magnetic field is given as

$$B = \frac{1}{C} (E_1 \hat{i} - E_2 \hat{j}) \cos(kz - \omega t)$$



B. The associated magnetic field is given as

$$B = \frac{1}{c} (E_1 \hat{i} - E_2 \hat{j}) \cos (kz - \omega t)$$

C. The given electromagnetic field is circularly

polarised

D. The given electromagnetic wave is plane

polarised

**Answer: D**

 [Watch Video Solution](#)

9. An electromagnetic wave travelling along z-axis is given as  $E = E_0 \cos (kz - \omega t)$ . Choose the correct

options from the following

A. The associated magnetic field is given as

$$B = \frac{1}{c} \hat{k} \times E = \frac{1}{\omega} (\hat{k} \times E)$$

B. The electromagnetic field can be written in terms

of the associated magnetic field as  $E = c(B \times \hat{k})$

C.  $\hat{k} \cdot E = 0, \hat{k} \cdot B = 0$

D.  $\hat{k} \times E = 0, \hat{k} \times B = 0$

**Answer: A::B::C**



**Watch Video Solution**

10. A plane electromagnetic wave propagating along x-direction can have the following pairs of  $E$  and  $B$ .

A.  $E_x, B_y$

B.  $E_y, B_z$

C.  $b_x, E_y$

D.  $E_z, B_y$

**Answer: B::D**



**Watch Video Solution**

11. A charged particle oscillates about its mean equilibrium position with a frequency of  $10^9 \text{ Hz}$ . The electromagnetic waves produced.

- A. will have frequency of  $10^9 \text{ Hz}$
- B. will have frequency of  $2 \times 10^9 \text{ Hz}$
- C. will have wavelength of  $0.3 \text{ m}$
- D. fall in the region of radiowaves

**Answer: A::B::C**



**Watch Video Solution**

12. The source of electromagnetic waves can be a charge.

A. moving with a constant velocity

B. moving in a circular orbit

C. at rest

D. falling in an electric field

**Answer: B::D**



**Watch Video Solution**

**13.** An electromagnetic wave of intensity  $I$  falls on a surface kept in vacuum and exerts radiation pressure  $p$  on it. Which of the following are true ?

A. Radiation pressure is  $\frac{I}{C}$  if the wave is totally absorbed

B. Radiation pressure is  $\frac{I}{C}$  if the wave is totally reflected

C. Radiation pressure is  $\frac{2I}{C}$  if the wave is totally reflected

D. Radiation pressure is in the range  $\frac{I}{C} < p < \frac{2I}{C}$  for real surfaces

**Answer: A::C::D**

 [Watch Video Solution](#)

**14.** Why is the orientation of the portable radio with respect to broadcasting station important?

 [Watch Video Solution](#)

**15.** Why does microwave oven heats up a food item containing water molecules most efficiently?

 [Watch Video Solution](#)

**16.** The charge on a parallel plate capacitor varies as  $= q_0 \cos 2\pi ft$ . The plates are very large and close together (area= $a$ , separation= $d$ ). Neglecting the edge effects, find the displacement current through the capacitor.



[Watch Video Solution](#)

**17.** A variable frequency AC source is connected to a capacitor. How will the displacement current change with decrease in frequency?



[Watch Video Solution](#)



18. The magnetic field of a beam emerging from a filter facing a floodlight is given by

$$B = 12 \times 10^{-8} \sin(1.20 \times 10^7 z - 3.60 \times 10^{14} t) T.$$

What is the average intensity of the beam?

 [Watch Video Solution](#)

19. Poynting vectors  $\vec{S}$  is defined as a vector whose magnitude is equal to the wave intensity and whose direction is along the direction of wave propagation.

Mathematically, it is given by  $\vec{S} = \frac{1}{\mu_0} \vec{E} \times \vec{B}$ . Show

the nature of  $S$  vs  $t$  graph


 [Watch Video Solution](#)

20. Professor C.V Raman surprised his students by suspending freely a tiny light ball in a transparent vacuum chamber by shining a laser beam on it. Which property of EM waves was he exhibiting? Give one more example of this property.



[Watch Video Solution](#)

21. Show that the magnetic field  $B$  at a point in between the plates of a parallel plate capacitor during charging is  $\frac{\mu_0 \epsilon_0 r}{2} \frac{dE}{dt}$  (symbols having usual meaning).



[Watch Video Solution](#)

22. Electromagnetic waves with wavelength

(i)  $\lambda_1$  is used in satellite communication.

(ii)  $\lambda_2$  used to kill germs in water purifier.

$\lambda$  used to detect leakage of oil in underground pipelines.

$\lambda_4$  used to improve visibility in runways during fog and

mist conditions.

(a) Identify and name the part of e.m. spectrum to which these radiations belong.

(b) Arrange these wavelengths in ascending order of their magnitude.

(c) Write one more application of each.



[Watch Video Solution](#)

**23.** Show that average value of radiant flux density  $S$

over a single period 'T' is given by  $S = \frac{1}{2c\mu_0} E_0^2$ .



[Watch Video Solution](#)

**24.** You are given a  $2\mu F$  parallel plate capacitor. How would you establish an instantaneous displacement current fo 1mA in the space between its plates?

 [Watch Video Solution](#)

**25.** Show that the radiation pressure exerted by an EM wave of intensity  $I$  on a surface kept in vacuum is  $I/c$ .

 [Watch Video Solution](#)

**26.** What happens to the intensity of light from a bulb if the distance from the bulb is doubled ? As a laser

beam travels across the length of room, its intensity essentially remain constant.

What geometrical characteristic of LASER beam is responsible for the constant intensity which is missing in the case of light from the bulb?



[Watch Video Solution](#)


27. Even though an electric field  $\vec{E}$  exerts a force  $q\vec{E}$  on a charged particle yet the electric field of an EM wave does not contribute to the radiation pressure (but transfer energy). Explain.



[Watch Video Solution](#)

**28.** An infinitely long thin wire carrying a uniform linear static charge density  $\lambda$  is placed along the z-axis Fig. The wire is set into motion along its length with a uniform velocity  $\vec{v} = v\hat{k}$ . Calculate the Poynting

$$\text{vector } \vec{S} = \frac{1}{\mu_0} (\vec{E} \times \vec{B}).$$



 Watch Video Solution



**29.** Sea water at frequency  $\nu = 4 \times 10^8$  Hz has permittivity  $\epsilon \approx 80\epsilon_0$  permeability  $\mu = \mu_0$  and resistivity  $\rho = 0.25\text{M}$ . Imagine a parallel plate capacitor immersed in sea water and driven by an alternating voltage source  $V(t) = V_0 \sin(2\pi\nu t)$ . What fraction of the conduction current density is the displacement current density?



**Watch Video Solution**

**30.** A long straight cable of length  $l$  is placed symmetrically along z-axis and has radius  $a$  ( $a \ll l$ ). The cable consists of a thin wire and a co-axial

conducting tube. An alternating current

$I(t) = I_0 \sin(2\pi vt)$ . Flows down the central thin

wire and returns along the co-axial conducting tube.

the induced electric at a distance  $s$  from the wire

inside the cable is

$$E(s, t) = \mu_0 I_0 v \cos(2\pi vt) \ln\left(\frac{s}{a}\right) \hat{k}.$$

(i) Calculate the displacement current density inside the cable.

(ii) Integrate the displacement current density across the cross-section of the cable to find the total displacement current  $I^d$ .

(iii) compare the conduction current  $I_0$  with the displacement current  $I_0^d$ .

 [Watch Video Solution](#)

31. A plane EM wave travelling in vacuum along z-direction is given by

$$E = E_0 \sin(kz - \omega t) \hat{i} \quad \text{and} \quad B = B_0 \sin(kz - \omega t) \hat{j}$$


(i) Evaluate  $\int E \cdot dl$  over the rectangular loop 1234 shown in figure.

(ii) Evaluate  $\int B \cdot ds$  over the surface bounded by loop 1234.

(iii) Use equation  $\int E \cdot dl = \frac{-d\phi_B}{dt}$  to prove  $\frac{E_0}{B_0} = c$ .

(iv) By using similar proces and the equation

$$\int \mathbf{B} \cdot d\mathbf{l} = \mu_0 I + \varepsilon_0 \frac{d\phi_E}{dt}, \text{ prove that } c = \frac{1}{\sqrt{\mu_0 \varepsilon_0}}$$



[Watch Video Solution](#)

32. A plane EM wave travelling along z direction is described by

$$\mathbf{E} = E_0 \sin(kz - \omega t) \hat{i} \text{ and } \mathbf{B} = B_0 \sin(kz - \omega t) \hat{j}.$$

show that

(i) The average energy density of the wave is given by

$$u_{av} = \frac{1}{4} \epsilon_0 E_0^2 + \frac{1}{4} \frac{B_0^2}{\mu_0}.$$

(ii) The time averaged intensity of the wave is given by

$$I_{av} = \frac{1}{2} c \epsilon_0 E_0^2.$$



**Watch Video Solution**