



MATHS

BOOKS - NAGEEN MATHS (HINGLISH)

APPLICATIONS OF DERIVATIVES

Solved Examples

1. Find the rate of change of area of a circle with respect to its radius 'r' when $r=7\text{cm}$.



[Watch Video Solution](#)

2. The side of a cube is increasing at a rate of 4 cm/sec . Find the rate of change of the volume of the cube when its side is 5 cm .

A. $350\text{ cm}^3/\text{sec}$

B. $450 \text{ cm}^3/\text{sec}$

C. $200 \text{ cm}^3/\text{sec}$

D. $300 \text{ cm}^3/\text{sec}$

Answer: D



Watch Video Solution

3. A stone is dropped in a quiet lake. If the produced circular waves increase at a rate of $4 \text{ cm}/\text{sec}$, then find the rate of increase in its area when the radius of circular wave is 7 cm .

A. $172 \text{ cm}^2/\text{sec}$

B. $174 \text{ cm}^2/\text{sec}$

C. $178 \text{ cm}^2/\text{sec}$

D. $176 \text{ cm}^2/\text{sec}$

Answer: D

 [Watch Video Solution](#)

4. In a cylindrical tank, rice is increasing at a rate of $314m^3/hr$. Find the rate of increase of the height of the rice in cylindrical tank when the radius of the base of tank is 5 m.

A. $1m/hr$

B. $2m/hr$

C. $3m/hr$

D. $4m/hr$

Answer: D

 [Watch Video Solution](#)

5. Find a point on the curve $y^2 = 2x$ at which the abscissa and ordinates are increasing at the same rate.

 [Watch Video Solution](#)

6. A ladder 13 m long is leaning against a wall. The bottom of the ladder is pulled along the ground away from the wall at the rate of 4 m/sec. Find the rate of decreasing at which the top of the ladder moving downwards on wall when the foot of the ladder is 5 m away from the wall.

A. $-\frac{5}{3}m/\text{sec}.$

B. $\frac{5}{3}m/\text{sec}.$

C. $-\frac{10}{3}m/\text{sec}.$

D. $\frac{10}{3}m/\text{sec}.$

Answer: A



[Watch Video Solution](#)

7. A man of height 1.7 m walks at a uniform speed of 6.6 m/min from a lamp post which is 5m high. Find the rate at which the length of his shadow increases.



[Watch Video Solution](#)

8. Show that the function $f(x) = 2x+1$ is strictly increasing on \mathbb{R} .



[Watch Video Solution](#)

9. Prove that $f(x) = e^{3x}$ is strictly increasing function on \mathbb{R} .



[Watch Video Solution](#)

10. Show that the function $f(x) = x - \sin x$ is an increasing function $\forall x \in \mathbb{R}$.



[Watch Video Solution](#)

11. Show that the function $f(x) = 3 - 2x$ is strictly decreasing function on \mathbb{R} .



[Watch Video Solution](#)

12. Show that the function $f(x) = x^3 - 3x^2 + 3x - 1$ is an increasing function on \mathbb{R} .

 [Watch Video Solution](#)

13. Find the intervals in which the function $f(x) = 2x^3 - 15x^2 + 36x + 6$ is (i) increasing, (ii) decreasing.

 [Watch Video Solution](#)

14. Find the intervals in which $f(x) = \sin 3x$, $x \in \left[0, \frac{\pi}{2}\right]$ is (i) increasing, (ii) decreasing.

A. $f(x)$ is increasing in $\left[0, \frac{\pi}{2}\right]$

$f(x)$ is decreasing in $\left[\frac{\pi}{6}, \frac{\pi}{2}\right]$

B. $f(x)$ is increasing in $\left[0, \frac{\pi}{6}\right]$

$f(x)$ is decreasing in $\left[\frac{\pi}{4}, \frac{\pi}{2}\right]$

C. $f(x)$ is increasing in $\left[0, \frac{\pi}{6}\right]$

$f(x)$ is decreasing in $\left[\frac{\pi}{6}, \frac{\pi}{2}\right]$

D. $f(x)$ is increasing in $\left[0, \frac{\pi}{6}\right]$

$f(x)$ is decreasing in $\left[\frac{\pi}{6}, \frac{\pi}{4}\right]$

Answer: C

 [Watch Video Solution](#)

15. Find the intervals in which function $f(x) = \sin x - \cos x$, $0 < x < 2\pi$ is

(i) increasing, (ii) decreasing.

 [Watch Video Solution](#)

16. Using differentials, find the approximate value of $\sqrt[3]{127}$.

 [Watch Video Solution](#)

17. Using differentials, find the approximate value of $\sqrt{0.037}$.



Watch Video Solution

18. Using differentials, find the approximate value of $\log_{10} 10.1$ when

$$\log_{10} e = 0.4343.$$



Watch Video Solution

19. Find the approximate value of $f(3.01)$ when

$$y = f(x) = x^2 + 3x + 1.$$



Watch Video Solution

20. Find the approximate percentage increase in a cube of side x metre if

the percentage increase in its side is 1%.



Watch Video Solution

21. The time period 'T' of a simple pendulum of length 'l' is given by $T =$

$$2\pi\sqrt{\frac{l}{g}}.$$

Find the percentage error in the value of 'T' if the percentage error in the value of 'l' is 2%.



[Watch Video Solution](#)

22. In ΔABC , the side c and angle C are constant. If there are small changes in the remaining sides and angles, then show that

$$\frac{da}{\cos A} + \frac{db}{\cos B} = 0.$$



[Watch Video Solution](#)

23. Find the slope of tangent at point $(1, 1)$ of the curve $x^2 = y$.



[Watch Video Solution](#)

24. Find the inclination from X - axis of the tangent at point $(a, 2a)$ of the curve $y^2 = 4ax$.

 [Watch Video Solution](#)

25. Find the equation of tangent of the curve $6y = 9 - 3x^2$ at point $(1,1)$.

 [Watch Video Solution](#)

26. Find the equation of tangent of tangent of the curve $y = b \cdot e^{-x/a}$ at that point at which the curve meets the Y-axis.

 [Watch Video Solution](#)

27. Find the equation of tangent of the curve $x = at^2, y = 2at$ at point t .

A. $x - ty + at^3 = 0$

B. $x - y + at^2 = 0$

C. $x - ty + at^2 = 0$

D. $tx - y + at^2 = 0$

Answer: C



Watch Video Solution

28. Find the point on the curve $y = x^2 + 1$ at which the tangent drawn makes an angle of 45° from X-axis.

A. $\left(\frac{1}{2}, \frac{5}{4}\right)$.

B. $\left(\frac{1}{2}, \frac{3}{4}\right)$.

C. $\left(-\frac{1}{2}, \frac{5}{4}\right)$.

D. None of these

Answer: A



Watch Video Solution

29. Find the equation of tangent of the curve $yx^2 + x^2 - 5x + 6 = 0$ at that point at which curve crosses the X-axis.

 [Watch Video Solution](#)

30. Find a point on the curve $y = 3x^2 - 2x - 4$ at which the tangent is parallel to the line $10x - y + 7 = 0$.

A. (2, 4)

B. (1, 4)

C. (-2, 4)

D. (2, -4)

Answer: A

 [Watch Video Solution](#)

31. Find the angle of intersection of the curves $2y^2 = x^3$ and $32x$.

 [Watch Video Solution](#)

32. Prove that the curves $y = x^3$ and $xy = k$ cut each other orthogonally, if $3k = 1$.

 [Watch Video Solution](#)

33. Find the equation of normal at point (a, a) of the curve $xy = a^2$.

 [Watch Video Solution](#)

34. Find the equation of normal to the curve $x = \cos \theta, y = \sin \theta$ at point $\theta = \frac{\pi}{4}$.

 [Watch Video Solution](#)

35. If a normal of curve $x^{2/3} + y^{2/3} = a^{2/3}$ makes an angle ϕ from X -axis then show that its equation is $y \cos \phi - x \sin \phi = a \cos 2\phi$.



Watch Video Solution

36. Find the value of x for which the function $(x^3 - 3x + 4)$ is maximum or minimum.



Watch Video Solution

37. Find the maximum or minimum values of the function $y = x + \frac{1}{x}$ or $x > 0$.



Watch Video Solution

38. Find the maximum value of the function $f(x) = \sin x + \cos x$.



Watch Video Solution

39. Find the minimum value of the function x^x .



Watch Video Solution

40. If x lies in first quadrant, find the maximum or minimum values of the function $4 \sin x + 3 \cos x$.



Watch Video Solution

41. If the function $x^2 - 2kx + 3$ is minimum at $x=3$ then find the value of k .



Watch Video Solution

42. Find two number whose sum is 8 and the sum of their cubes is minimum.



Watch Video Solution

[Watch Video Solution](#)

43. The sum of two number is constant. Show that their product will be maximum if each number is half of their sum.

[Watch Video Solution](#)

44. The perimeter of a rectangle is 40 cm. Find the dimensions of the rectangle if its area is maximum.

[Watch Video Solution](#)

45. If the sum of a side and hypotenuse of a right angled triangle is given then show that area of triangle will be maximum when the angle between side and the hypotenuse is $\pi/3$.

[Watch Video Solution](#)

46. Prove that the height of a cylinder of maximum volume, inscribed in a sphere of radius 'a' is $2a / \sqrt{3}$.

 [Watch Video Solution](#)

47. Prove that the cylinder inscribed in a right circular cone has maximum curved surface area if the radius of cylinder is half the radius of cone.

 [Watch Video Solution](#)

Exercise 6 A

1. Find the rate of change of area of the circle with respect to its radius 'r' when $r = 3.5$ cm.

 [Watch Video Solution](#)

2. (i) The radius of a circle is increasing at the rate of 5 cm/sec. Find the rate of increasing of its perimeter.

(ii) If the area of a circle increases at a constant rate, then show that the rate of increase of its circumference is inversely proportional to its circumference is iversely proportional to its radius.



[Watch Video Solution](#)

3. The side of a square is increasing at a rate of 3 cm/sec. Find the rate of increasing of its perimeter in cm/sec when the side of square is 5cm.



[Watch Video Solution](#)

4. The side of a square is increasing at a rate of 4cm/sec. Find the rate of increase of its area when the side of square is 10 cm.



[Watch Video Solution](#)

5. The rate of increase of the radius of an air bubble is 0.5 cm/sec . Find the rate of increase of its volume when the radius of air bubble is 2 cm .

 [Watch Video Solution](#)

6. A balloon which always remains spherical, is being inflated by pumping in 900 cubic centimetres of gas per second. Find the rate at which the radius of the balloon is increasing when the radius is 15 cm .

 [Watch Video Solution](#)

7. The volume of cube is increasing at a rate of $9 \text{ cm}^3 / \text{sec}$. Find the rate of increase of its surface area when the side of the cube is 10 cm .

 [Watch Video Solution](#)

8. The volume of a spherical balloon is increasing at a rate of $25 \text{ cm}^3 / \text{sec}$. Find the rate of increase of its curved surface when the radius of balloon

is 5 cm.

 [Watch Video Solution](#)

9. The surface of a spherical balloon is increase of its volume when its radius is 6 cm.

 [Watch Video Solution](#)

10. The length of a rectangle is decreasing at a rate of 3 cm/sec and breadth is increasing at a rate of 4 cm/sec. Find the rate of change of its (a) perimeter (b) area, when the length and breadth of rectangle are 7 cm and 8 cm respectively.

 [Watch Video Solution](#)

11. Find the point on the curve $y^2 = 8x$ for which the abscissa and ordinate change at the same rate.



[Watch Video Solution](#)

12. A particle moves along the curve $6y = x^3 + 2$. Find the points on the curve at which y-co-ordinate is changing 8 times as fast as the x-co-ordinate.



[Watch Video Solution](#)

13. The base of a cubical tank is $25m \times 40m$. The volume of water in the tank is increasing at the rate of $500m^3 / \text{min}$. Find the rate at which the height of water is increasing.



[Watch Video Solution](#)

14. The oil is leaking from a drum at a rate of $16cm^3 / \text{sec}$. If the radius and height of drum are 7 cm and 60 cm respectively, find the rate of change of the height of oil when height of oil in drum is 18 cm.



[Watch Video Solution](#)

15. The water is leaking from a conical funnel at a rate of $5\text{cm}^3 / \text{min}$. If the radius and height of the funnel are 5cm.and 10 cm respectively, find the rate of change of the surface of water in the funnel when height of water surface from the vertex is 7.5cm .

[View Text Solution](#)

16. A man of height 160 cm walks at a rate of 1.1 m/sec from a lamp post which 6m high. Find the rate at which the length of his shadow increases when he is 1m away from the lamp post.

[Watch Video Solution](#)

17. The total cost $C(x)$ in Rupees, associated with the production of x units of an item is given by $C(x) = 0.005x^3 - 0.02x^2 + 30x + 5000$. Find

the marginal cost when 3 units are produced, where by marginal cost we mean the instantaneous rate of cha

 [Watch Video Solution](#)

18. The total revenue of selling of x units of a product is represented by $R(x) = 2x^2 + x + 5$. Find its marginal revenue for 5 units.

 [Watch Video Solution](#)

19. A ladder is inclined to a wall making an angle of 30° with it. A man is ascending the ladder at the rate of 3 m/sec. How fast is he approaching the wall ?

 [Watch Video Solution](#)

20. The one end of a 20 m long ladder is on the floor and the other end is in the contact of vertical wall. Its lower end slides on the floor. Show that

the rate of sliding of the upper end is $\frac{4}{3}$ times the rate of sliding the lower end when the lower end is 16 m away from the wall.

 [Watch Video Solution](#)

Exercise 6 B

1. Show that the function $f(x) = 3x + 2$ is strictly increasing function on \mathbb{R} .

 [Watch Video Solution](#)

2. Show that the function $f(x) = x^2$ is strictly increasing function in the interval $]0, \infty[$.

 [Watch Video Solution](#)

3. Show that the function $f(x) = a^x$, $a > 1$ is strictly increasing on \mathbb{R} .

 [Watch Video Solution](#)

4. Show that the function $f(x) = -5x + 2$ is strictly decreasing function on \mathbb{R} .

 [Watch Video Solution](#)

5. Prove that the function given by $f(x) = \cos x$ is (a) strictly decreasing in $(0, \pi)$ (b) strictly increasing in $(\pi, 2\pi)$, and (c) neither increasing nor decreasing in $(0, 2\pi)$

 [Watch Video Solution](#)

6. Find the intervals in which the following functions are:

(i) increasing

(ii) decreasing.

(a) $f(x) = 10 - 6x + x^2$

(b) $f(x) = 2x^2 - 6x$

(c) $f(x) = 2x^3 - 3x^2 - 36x + 1$

$$(d) f(x) = x^3 + 2x^2 + x - 1$$

$$(e) f(x) = 4x + \frac{1}{x}, x \neq 0$$

$$(f) f(x) = 2x^3 - 9x^2 + 12x + 1$$

$$(g) f(x) = 5 + 36x + 3x^2 - 2x^3$$

$$(h) f(x) = (x + 2)^3(x - 3)^3$$



[View Text Solution](#)

7. Show that $y = \log(1 + x) - \frac{2x}{2 + x}$, $x > 1$ is an increasing function of x throughout its domain.



[Watch Video Solution](#)

8. (a) Find the intervals in which the function

$$f(x) = \log(1 + x) - \frac{x}{1 + x} \text{ is (i) increasing,}$$

(ii) decreasing function.

(b) Find the intervals in which the function $f(x) = \frac{x}{\log_e x}$, $x > 0$ is increasing or decreasing.



[View Text Solution](#)

 [View Text Solution](#)

9. Let I be an interval disjoint from $[-1, 1]$. Prove that the function $f(x) = x + \frac{1}{x}$ is increasing on I .

 [Watch Video Solution](#)

10. Show that the function $f(x) = \sin^4 x + \cos^4 x$

(i) is decreasing in the interval $\left[0, \frac{\pi}{4}\right]$.

(ii) is increasing in the interval $\left[\frac{\pi}{4}, \frac{\pi}{2}\right]$.

 [Watch Video Solution](#)

11. Find the intervals in which the function

$f(x) = \sin x + \cos x, x \in [0, 2\pi]$ is

(i) strictly increasing, (ii) strictly decreasing.

 [Watch Video Solution](#)

Exercise 6 C

1. Using differentials, find the approximate values of the following:

$$\sqrt{26}$$

 [Watch Video Solution](#)

2. Using differentials, find the approximate values of the following:

$$(0.007)^{1/3}$$

 [View Text Solution](#)

3. Use differentials and find approximate value of $(29)^{1/3}$

 [Watch Video Solution](#)

4. Using differentials, find the approximate values of the following:

$$\sqrt{0.24}$$



[Watch Video Solution](#)

5. Using differentials, find the approximate value of $\sqrt{0.48}$



[Watch Video Solution](#)

6. Using differentials, find the approximate values of the following:

$$(i) \sqrt[4]{15}(ii) (82)^{1/4}$$



[View Text Solution](#)

7. Using differentials, find the approximate values of the following:

$$\frac{1}{(2.002)^2}$$



[View Text Solution](#)

8. Use differentials to approximate $\sqrt{25.2}$



[Watch Video Solution](#)

9. Using differentials, find the approximate value of $(0.009)^{1/3}$



[Watch Video Solution](#)

10. Use differentials to find the approximate value of $(\log)_e(4.01)$, having given that $(\log)_e 4 = 1.3863$.



[Watch Video Solution](#)

11. Using differentials, find the approximate value of $(\log)_e 4.04$, it being given that $(\log)_{10} 4 = 0.6021$ and $(\log)_{10} e = 0.4343$.



[Watch Video Solution](#)

12. If $f(x) = 2x^2 + 5x + 2$, then find the approximate value of $f(2.01)$.



[Watch Video Solution](#)

13. If $f(x) = 3x^2 + 4x - 1$, then find the approximate value of $f(3.1)$.



[Watch Video Solution](#)

14. The radius of a circular plate increases by 2% on heating. If its radius is 10 cm before heating, find the approximate increase in its area.



[Watch Video Solution](#)

15. The radius of a sphere decreases from 10 cm to 9.9 cm. Find

(i) approximate decrease in its volume.

(ii) approximate decrease in its surface.



[View Text Solution](#)

16. The time t of a complete oscillation of a simple pendulum of length l is

given by the equation $T = 2\pi\sqrt{\frac{l}{g}}$ where g is constant. What is the

percentage error in T when l is increased by 1%?



[Watch Video Solution](#)

17. There is an error of 0.2% in measurement of the radius of a sphere.

Find the percentage error in its

(i) volume, (ii) surface.



[Watch Video Solution](#)

18. The radius of a sphere is 8 cm and 0.02 cm is the error in its

measurement. Find the approximate error in its volume.



[Watch Video Solution](#)

19. The semi-vertical angle of a cone remains constant. If its height increases by 2%, then find the approximate increase in its volume.



[Watch Video Solution](#)

Exercise 6 D

1. Find the slope of tangents drawn of the following curves at the given points:

(i) Curve $y = x^3 + 1$ at point $(0, 1)$

(ii) Curve $x^2 - y^2 = 20$ at point $(6, 4)$

(iii) Curve $y^2 = 4x$ at point $(1, 2)$

(iv) Curve $y^2 = 4ax$ at point $\left(\frac{a}{m^2}, \frac{2a}{m}\right)$



[Watch Video Solution](#)

2. Find the inclination from X-axis of the tangent drawn of the following curves at the given points:

(i) Curve $x^2 - 2y^2 = 8$ at point (4, 2)

(ii) Curve $y = (x - 1)(x - 2)$ at point (2, 0)

(iii) Curve $y^2 = 2x^3$ at point (2, 4)



[View Text Solution](#)

3. Find the equation of tangents of the following curves at the given points:

(i) Curve $x^2 = 25$ at point (3, 4)

(ii) Curve $y = 2x^3 + 2x^2 - 8x + 7$ at point (1, 3)

(iii) Curve $y = x + \frac{2}{x}$ at point (2, 3)

(iv) Curve $y^2(x - 1) = 4x^2$ at point (5, 5)



[View Text Solution](#)

4. Find the equation of tangent of the curve $x^2 + y^2 = 5$ at point (1, 2).



Watch Video Solution

5. Find the equation of tangent of the curve $y^2 = 4x + 5$ which is parallel to the line $2x - y = 5$.



Watch Video Solution

6. Find the equation of tangent of the curve $9x^2 + 16y^2 = 144$ at those points at which tangents are parallel to (i) X-axis, (ii) Y-axis.

A. (i) $x = \pm 6$

(ii) $y = \pm 3$

B. (i) $x = \pm 2$

(ii) $y = \pm 3$

C. (i) $x = \pm 4$

(ii) $y = \pm 9$

D. (i) $x = \pm 4$

(ii) $y = \pm 3$

Answer: D

 [Watch Video Solution](#)

7. Find the co-ordinates of that point on the curve $x^3 + y^3 = a^3$ at which the tangent drawn is parallel to X-axis.

 [Watch Video Solution](#)

8. Find the co-ordinates of that point on the curve $y^2 = x^2(1 - x)$ at which the tangent drawn is perpendicular to X-axis.

 [Watch Video Solution](#)

9. Find the co-ordinates of that point on the curve $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at which the tangent drawn is parallel to Y-axis.

 [Watch Video Solution](#)

10. Prove that the equation of tangent of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at point (x_1, y_1) is $\frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1$.

 [Watch Video Solution](#)

11. Find the value of $n \in \mathbb{N}$ such that the curve $\left(\frac{x}{a}\right)^n + \left(\frac{y}{b}\right)^n = 2$ touches the straight line $\frac{x}{a} + \frac{y}{b} = 2$ at the point (a, b) .

 [Watch Video Solution](#)

12. Show that the line $\frac{x}{a} + \frac{y}{b} = 1$ touches the curve $y = be^{-\frac{x}{a}}$ at the point where it crosses the y-axis.

 [Watch Video Solution](#)

13. Find the point on the curve $y^2 = x$ at which the tangent drawn makes an angle of 45° from X-axis.

 [Watch Video Solution](#)

14. Find the coordinates of the points on the curve $y = x^2 + 3x + 4$, the tangents at which pass through the origin.

 [Watch Video Solution](#)

15. The tangent drawn at any point of the curve $\sqrt{x} + \sqrt{y} = \sqrt{a}$ meets the OX and OY axes at points P and Q respectively, prove that $OP + OQ = a$.

 [Watch Video Solution](#)

16. If p and q are the intercept on the axis cut by the tangent of

$$\sqrt{\left(\frac{x}{a}\right)} + \sqrt{\left(\frac{y}{b}\right)} = 1, \text{ prove that } \frac{p}{a} + \frac{q}{b} = 1$$

 [Watch Video Solution](#)

17. Find the angle of intersection of the curves

$$xy = a^2 \text{ and } x^2 + y^2 = 2a^2$$

 [Watch Video Solution](#)

18. Prove that the curves $x^2 - y^2 = 16$ and $xy = 15$ intersect each other at 90° angle.

 [Watch Video Solution](#)

19. If two curves $ax^2 + by^2 = 1$ and $a'x^2 + b'y^2 = 1$ intersect orthogonally, then show that $\frac{1}{a} - \frac{1}{b} = \frac{1}{a'} - \frac{1}{b'}$

 [Watch Video Solution](#)

20. Prove that the curves $x = y^2$ and $xy = k$ intersect at right angles if $8k^2 = 1$.

 [Watch Video Solution](#)

21. Find the equation of the tangent and the normal at the point 't', on the curve $x = a \sin^3 t, y = b \cos^3 t$.

 [Watch Video Solution](#)

22. Prove that points of the curve $y^2 = 4a \left\{ x + a \sin \left(\frac{x}{a} \right) \right\}$ at which tangents are parallel to x-axis lie on the parabola.

 [Watch Video Solution](#)

23. Prove that the tangents drawn on the parabola $y^2 = 4ax$ at points $x = a$ intersect at right angle.

 [Watch Video Solution](#)

24. Prove that the curve $y^2 = 4x$ and $x^2 + y^2 - 6x + 1 = 0$ touches each other at the point $(1, 2)$, find the equation of the common tangents.

 [Watch Video Solution](#)

Exercise 6 E

1. Find the equation of normals of the following curves at the given points:

(i) Curve $y^2 = 4ax$ at point $(at^2, 2at)$.

(ii) Curve $y = e^x$ at point $(0, 1)$

(iii) Curve $y = x^3$ at point $(1, 1)$.

(iv) Curve $2y = 3 - x^2$ at point $(1, 1)$.

(v) Curve $16x^2 - 9y^2 = 432$ at point $(6, 4)$.



[View Text Solution](#)

2. Find the equation of the normal to the curve $y = 5x + x^2$ which makes an angle 45° with x axis.

A. $x - y = 3$

B. $2x - y = 3$

C. $x - 2y = 3$

D. $x - y = 2$

Answer: A



[Watch Video Solution](#)

3. Find the equation of normal to the curve $y(x - 2)(x - 3) - x + 7 = 0$ at that point at which the curve meets X-axis.

 [Watch Video Solution](#)

4. Find the equation of normal to the curve $x = at^2, y = 2at$ at point 't'.

 [Watch Video Solution](#)

5. Find the equation of the normal to the curve $x = a \cos \theta$ and $y = b \sin \theta$ at θ

 [Watch Video Solution](#)

6. Find the equation of normal to the curve $x = a \cos^3 \theta, y = b \sin^3 \theta$ at point ' θ '.



[Watch Video Solution](#)

7. Find the equations of tangent and normal to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at (x_1, y_1)

[Watch Video Solution](#)

8. (i) Find the co-ordinates of the points on the curve $xy = 16$ at which the normal drawn meet at origin.

(ii) Find the points on the curve $4x^2 + 9y^2 = 36$ at which the normal drawn is parallel to X-axis.

[Watch Video Solution](#)

9. Find the equation of normal at point $(4, 3)$ for the hyperbola $3x^2 - 4y^2 = 14$.

[Watch Video Solution](#)

10. Find the equation of normal to the curves $x = t^2$, $y = 2t + 1$ at point

A. $tx + y = t^3 + 3t + 1$

B. $tx + y = t^3 + 2t + 1$

C. $tx + 4y = t^3 + t + 1$

D. $tx + y = t^3 + 2t + 2$

Answer: B



[Watch Video Solution](#)

11. Find the point on the curve $9y^2 = x^3$, where the normal to the curve makes equal intercepts on the axes.



[Watch Video Solution](#)

12. If the normal at point 't' of the curve $xy = c^2$ meets the curve again at point ' t_1 ', then prove that

$$t^3 \cdot t_1 = -1.$$

 [Watch Video Solution](#)

13. Find the area area angle which made by the tangent to the curve $y(2a - x) = x^2$ at point (a, a) its normal and x-axis.

 [Watch Video Solution](#)

14. Find the co-ordinates of the point on the curve $y = x^3 + 3x^2 - 4x - 12$ at which the normal's inclination is $-\frac{1}{7}$. Also find the equation of the normal.

 [View Text Solution](#)

15. If the normal drawn on the curve $y = x^2 + 4$ at point $(1, 5)$ makes an angle θ from X-axis then show that:

$$\tan \theta = -\frac{1}{2}$$

 [Watch Video Solution](#)

16. Find the equation of normal of the curve $2y = 7x - 5x^2$ at those points at which the curve intersects the line $x = y$.

 [Watch Video Solution](#)

Exercise 6 F

1. Find the values of x for which the following functions are maximum or minimum:

(i) $x^3 - 3x^2 - 9x$

(ii) $4x^3 - 15x^2 + 12x + 1$

(iii) $1 - x - x^2$

(iv) $x^4 - 8x^3 + 22x^2 - 24x + 7$

(v) $\frac{x^2 - x + 1}{x^2 + x + 1}$

 [View Text Solution](#)

2. Find the maximum or minimum values of the following functions:

(i) $x^3 - 2x^2 + x + 3$

(ii) $x^3 - 6x^2 + 9x + 4$

(iii) $(x + 1)(x - 2)^2 + 1$

(iv) $2x^3 - 9x^2 + 12x + 1$



[View Text Solution](#)

3. For what values of x , the function $f(x) = x^5 - 5x^4 + 5x(3) - 1$ is maximum or minimum? Prove that at $x = 0$, the function is neither maximum nor minimum.



[Watch Video Solution](#)

4. Find the maximum and minimum values of the function $f(x) = \sin x + \cos 2x$.



[Watch Video Solution](#)

5. Find the maximum and minimum values of the function

$$f(x) = x + \sin 2x, (0 < x < \pi).$$

 [Watch Video Solution](#)

6. Find the maximum and minimum values of the function

$$f(x) = \frac{\sin x}{1 + \tan x}, (0 < x < 2\pi).$$

 [Watch Video Solution](#)

7. Show that $s \in^p \theta \cos^q \theta$ attains a maximum, when $\theta = \tan^{-1} \sqrt{\frac{p}{q}}$.

 [Watch Video Solution](#)

8. Find the maximum value of the function $\frac{\log x}{x}$ when $x > 0$.

 [Watch Video Solution](#)

9. Find the maximum value of the function $x^{1/x}$.

 [Watch Video Solution](#)

10. Show that the maximum value of $\left(\frac{1}{x}\right)^x$ is $e^{\frac{1}{e}}$.

 [Watch Video Solution](#)

11. Show that the function $x^3 - 3x^2 + 3x + 1$ has neither a maxima nor a minima.

 [Watch Video Solution](#)

12. Show that $f(x) = \sin x(1 + \cos x)$ is maximum at $x = \frac{\pi}{3}$ in the interval $[0, \pi]$.

 [Watch Video Solution](#)

13. If the function $f(x) = x^3 - 24x^2 + 6kx - 8$ is maximum at $x = 2$.

Then, find value of k .

 [Watch Video Solution](#)

14. Prove that value of the function $xy(y - x) = 2a^3$ is minimum at $x = a$.

 [Watch Video Solution](#)

15. Show that the function $(-x^2 \log x)$ is minimum at $x = \frac{1}{\sqrt{e}}$.

 [Watch Video Solution](#)

16. Find the maximum value of the function $x \cdot e^{-x}$.

 [Watch Video Solution](#)

17. Show that the maximum value of the function $\sqrt{2}(\sin x + \cos x)$ is 2.

 [Watch Video Solution](#)

18. Show that the maximum and minimum values of the function $\frac{(x + 1)^2}{(x + 3)^3}$ are respectively given by $\frac{2}{27}$ and 0.

 [Watch Video Solution](#)

19. If the function $y = ae^x + bx^2 + 3x$ is maximum at $x = 0$ and minimum at $x = -3$, then find the values of a and b .

 [View Text Solution](#)

Exercise 6 G

1. Divide 16 into two parts such that the sum of their cubes is minimum.

 [Watch Video Solution](#)

2. If $x + y = 1$ then find the maximum value of the function xy^2

 [Watch Video Solution](#)

3. Find two number whose sum is 100 and the sum of twice of first part and square of the second is minimum.

 [Watch Video Solution](#)

4. Find two numbers whose sum is 12 and the product of the square of one part and the 4th power of other is maximum.

 [Watch Video Solution](#)

5. Divide 15 into two parts such that product of square of one part and cube of other is maximum

 [Watch Video Solution](#)

6. (i) The two sides of a rectangle are x units and $(10 - x)$ units. For what value of x , the area of rectangle will be maximum?

(ii) Prove that a rectangle, whose area is constant has minimum perimeter if it is a square.

 [Watch Video Solution](#)

7. Show that the triangle of maximum area that can be inscribed in a given circle is an equilateral triangle.

 [Watch Video Solution](#)

8. If the surface area of an open cylinder is 100cm^2 , prove that its maximum volume will be $\frac{1000}{3\sqrt{3\pi}}\text{cm}^3$.

 [Watch Video Solution](#)

9. (i) The base of an open rectangular box is square and its volume is 256cm^3 . Find the dimensions of this box if we want to use least material for construction:

(ii) A window is in the form of a rectangle surmounted by a semi-circle. Its perimeter is 40m. Find the dimensions of this window from which maximum light can admit.

 [View Text Solution](#)

10. An open tank of square base is to be constructed which has a given quantity of water. Prove that for least material used for the construction of tank, the depth will be half of the width of tank.

 [Watch Video Solution](#)

[Watch Video Solution](#)

11. The base of a cuboid is square and its volume is given. Show that its total surface will be minimum, if it is a cube.

 [Watch Video Solution](#)

12. Find the maximum area of the isosceles triangle inscribed in the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, \text{ with its vertex at one end of major axis.}$$

 [Watch Video Solution](#)

13. The volume of a closed square based rectangular box is 1000 cubic metre. The cost of constructing the base is 15 paise per square metre and the cost of constructing the top is 25 paise per square metre. The cost of constructing its sides is 20 paise per square metre and the cost of constructing the box is Rs. 3. Find the dimensions of box for minimum cost of construction.



[Watch Video Solution](#)

14. Show that height of the cylinder of greatest volume which can be inscribed in a right circular cone of height h and semi vertical angle is one-third that of the cone and the greatest volume of cylinder is

$$\frac{4}{27} \pi h^3 \tan^2 \alpha.$$



[Watch Video Solution](#)

15. The sum of perimeter of a square and circumference of a circle is given. Prove that the sum of their areas will be minimum when the side of square is equal to the diameter of the circle.



[Watch Video Solution](#)

16. A square-based tank of capacity 250 cu m has to be dug out. The cost of land is Rs 50 per sq m. The cost of digging increases with the depth

and for the whole tank the cost is Rs $400 \times (\text{depth})^2$. Find the dimensions of the tank for the least total cost.

 [Watch Video Solution](#)

17. The stiffness of a beam of rectangular cross-section varies as the product of the breadth and square of the depth. What must be the breadth and depth of the stiffest beam that can be cut from a leg of diameter 'd'?

 [Watch Video Solution](#)

18. The expenditure on fuels in running a train varies as the square of its speed measured in km/hr and it is Rs. 48 per hour at the speed of 16 km/hr. If the constant expenditure is Rs. 300 per hour find the best speed of the train economically.

 [View Text Solution](#)

19. The combined resistance of two resistors R_1 and R_2 is given by $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2}$, where $R_1 + R_2 = C$ (constant), show that R is maximum when $R_1 = R_2$.

 [Watch Video Solution](#)

20. Prove that the area of right-angled triangle of given hypotenuse is maximum when the triangle is isosceles.

 [Watch Video Solution](#)

21. A wire of length 28 m is to be cut into two pieces. One of the pieces is to be made into a square and the other into a circle. What should be the length of the two pieces so that the combined area of the square and the circle is minimum?

 [Watch Video Solution](#)

Exercise 6 H Multiple Choice Questions

1. Using differentials, find the approximate value of $(82)^{\frac{1}{4}}$ upto 3 places of decimal .

A. 3.008

B. 3.009

C. 3.010

D. None of these

Answer: B



[Watch Video Solution](#)

2. Find the equation of the tangent to the curve

$$x = \theta + \sin \theta, y = 1 + \cos \theta \text{ at } \theta = \frac{\pi}{4}$$

A. $y = (1 + \sqrt{2})x + \frac{(\sqrt{2} - 1)\pi}{4} + 2$

B. $y = (1 - \sqrt{2})x + \frac{(\sqrt{2} - 1)\pi}{4} + 2$

C. $y = (1 + \sqrt{2})x + (\sqrt{2} - 1)\pi$

D. None of the above

Answer: B

 [Watch Video Solution](#)

3. The equation of normal to the curve $\left(\frac{x}{a}\right)^n + \left(\frac{y}{b}\right)^n = 2(n \in \mathbb{N})$ at the point with abscissa equal to 'a' can be

A. $\frac{x}{a} + \frac{y}{b} + 1 = 0$

B. $\frac{x}{a} - \frac{y}{b} = 1$

C. $\frac{x}{a} + \frac{y}{b} = 1$

D. None of these

Answer: C

 [Watch Video Solution](#)

4. Find the point on the curve $9y^2 = x^3$, where the normal to the curve makes equal intercepts on the axes.

A. $\left(3, \pm \frac{3}{8}\right)$

B. $\left(4, \pm \frac{8}{3}\right)$

C. $\left(3, \pm \frac{8}{3}\right)$

D. $\left(4, \pm \frac{3}{8}\right)$

Answer: D



Watch Video Solution

5. A balloon, which always remains spherical, has a variable diameter $\frac{3}{2}(2x + 1)$. Find the rate of change of its volume with respect to x .

A. $3\pi(2x + 1)^2$

B. $\frac{8}{9}\pi(2x + 1)^2$

C. $\frac{27}{8}\pi(2x + 1)^2$

D. $\frac{8}{3}\pi(2x + 1)^2$

Answer: C



Watch Video Solution

6. In the interval $(-1, 1)$, the function $f(x) = x^2 - x + 4$ is :

A. increasing

B. decreasing

C. neither increasing nor decreasing

D. None of the above

Answer: C



Watch Video Solution

7. If an error of $k\%$ is made in measuring the radius of a sphere, then percentage error in its volume. $k\%$ (b) $3k\%$ (c) $3k\%$ (d) $\frac{k}{3}\%$

A. 2%

B. 4%

C. 6%

D. 8%

Answer: C



[Watch Video Solution](#)

8. Minimum value of the function $f(x) = \left(\frac{1}{x}\right)^{1/x}$ is:

A. e

B. $\frac{1}{e}$

C. e^e

D. $e^{-1/e}$

Answer: D



[Watch Video Solution](#)

9. If the sum of two +ve numbers is 18 then the maximum value of their product is

A. 8,10

B. 9,9

C. 6,12

D. None of these

Answer: B



[Watch Video Solution](#)

10. For minimum curved surface area and given volume, the ratio of the height and radius of base of a cone is :

A. $1 : \sqrt{2}$

B. $\sqrt{2} : 1$

C. $1 : 2$

D. None of these

Answer: B



[View Text Solution](#)

Exercise 6 | Multiple Choice Questions

1.26. The points of contact of the tangents drawn from the origin to the curve $y = \sin x$, lie on the curve

A. $x^2 - y^2 = x^2 y^2$

B. $x^2 + y^2 = x^2y^2$

C. $y^2 - x^2 = x^2y^2$

D. None of these

Answer: A



[Watch Video Solution](#)

2. Equation of tangent of the curve $y = 1 - e^{x/2}$ at that point at which the curve crosses the y-axis, is :

A. $x + y = 1$

B. $2x + y = 1$

C. $x = -2y$

D. None of these

Answer: C



[View Text Solution](#)

3. Curve $b^2x^2 + a^2y^2 = a^2b^2$ and $m^2x^2 - y^2l^2 = l^2m^2$ intersect each other at right-angle if:

A. $a^2 + b^2 = l^2 + m^2$

B. $a^2 - b^2 = l^2 - m^2$

C. $a^2 - b^2 = l^2 + m^2$

D. $a^2 + b^2 = l^2 - m^2$

Answer: C

 [Watch Video Solution](#)

4. The function $x^{100} + \sin x - 1$ is decreasing in the interval:

A. $\left(0, \frac{\pi}{2}\right)$

B. $(0, 1)$

C. $\left(\frac{\pi}{2}, \pi\right)$

D. None of these

Answer: D



[View Text Solution](#)

5. The function $\log(1 + x) - \frac{2x}{x + 2}$ is increasing in the interval:

A. $(-\infty, 0]$

B. $(-1, \infty)$

C. $(-\infty, 1]$

D. None of these

Answer: B



[Watch Video Solution](#)

6. If the function $f(x) = a \sin x + \frac{1}{3} \sin 3x$ is maximum at $x = \frac{\pi}{3}$ a=?

A. 1

B. 2

C. 3

D. 4

Answer: B



[View Text Solution](#)

7. If $x + y = 8$, then the maximum value of xy is (a) 8 (b) 16 (c) 20 (d) 24

A. 8

B. 12

C. 16

D. 7

Answer: C



[Watch Video Solution](#)

8. An isosceles triangle of vertical angle 2θ is inscribed in a circle of radius a . Show that the area of the triangle is maximum when $\theta = \frac{\pi}{6}$.

A. π^3

B. $\frac{\pi}{4}$

C. $\frac{\pi}{6}$

D. None of these.

Answer: C



Watch Video Solution

9. Find minimum value of $px + qy$ where $p > 0, q > 0, x > 0, y > 0$ when $xy = r,^2$ without using derivatives.

A. $pq\sqrt{r}$

B. $2pq\sqrt{r}$

C. $2r\sqrt{pq}$

D. None of these

Answer: C



Watch Video Solution

10. The maximum area of rectangle, inscribed in a circle of radius 'r', is :

A. πr^2

B. r^2

C. $2r^2$

D. $\frac{1}{4}\pi r^2$

Answer: C



Watch Video Solution

1. Find the rate of change of the area of a circle with respect to its radius r when (a) $r = 3\text{ cm}$ (b) $r = 4\text{ cm}$

 [Watch Video Solution](#)

2. The volume of a cube is increasing at the rate of $8\text{ cm}^3 / \text{s}$. How fast is the surface area increasing when the length of an edge is 12 cm ?

 [Watch Video Solution](#)

3. The radius of a circle is increasing uniformly at the rate of 3 cm/s . Find the rate at which the area of the circle is increasing when the radius is 10 cm .

 [Watch Video Solution](#)

4. An edge of a variable cube is increasing at the rate of 3 cm per second. How fast is the volume of the cube increasing when the edge is 10 cm long?

 [Watch Video Solution](#)

5. A stone is dropped into a quiet lake and waves move in circles at the speed of 5 cm/s. At the instant when the radius of the circular wave is 8 cm, how fast is the enclosed area increasing?

 [Watch Video Solution](#)

6. The radius of a circle is increasing at the rate of 0.7 cm/sec. What is the rate of increase of its circumference?

 [Watch Video Solution](#)

7. The length x of a rectangle is decreasing at the rate of 5 cm/minute and the width y is increasing at the rate of 4 cm/minute. When $x = 8$ cm and $y = 6$ cm, find the rates of change of (a) the perimeter, and (b) the area of the rectangle

 [Watch Video Solution](#)

8. A balloon, which always remains spherical on inflation, is being inflated by pumping in 900 cubic centimetres of gas per second. Find the rate at which the radius of the balloon increases when the radius is 15 cm.

 [Watch Video Solution](#)

9. A balloon, which always remains spherical, has a variable radius. Find the rate at which its volume is increasing with the radius when the later is 10 cm

 [Watch Video Solution](#)

10. A ladder 5 m long is leaning against a wall. The bottom of the ladder is pulled along the ground, away from the wall, at the rate of 2 cm/s. How fast is its height on the wall decreasing when the foot of the ladder is 4 m away from the wall?

 [Watch Video Solution](#)

11. A particle moves along the curve $6y = x^3 + 2$. Find the points on the curve at which the y-coordinate is changing 8 times as fast as the x-coordinate.

 [Watch Video Solution](#)

12. The radius of an air bubble is increasing at the rate of $\frac{1}{2}$ cm/s. At what rate is the volume of the bubble increasing when the radius is 1 cm?

 [Watch Video Solution](#)

13. A balloon, which always remains spherical, has a variable diameter $\frac{3}{2}(2x + 1)$. Find the rate of change of its volume with respect to x .



[Watch Video Solution](#)

14. Sand is pouring from a pipe at the rate of $12 \text{ cm}^3 / \text{s}$. The falling sand forms a cone on the ground in such a way that the height of the cone is always one-sixth of the radius of the base. How fast is the height of the sand cone increasing when t



[Watch Video Solution](#)

15. The total cost $C(x)$ in Rupees associated with the production of x units of an item is given by $C(x) = 0.007x^3 - 0.003x^2 + 15x + 4000$. Find the marginal cost when 17 units are produced



[Watch Video Solution](#)

16. The total revenue in Rupees received from the sale of x units of a product is given by $R(x) = 13x^2 + 26x + 15$. Find the marginal revenue when $x = 7$.



[Watch Video Solution](#)

17. The rate of change of the area of a circle with respect to its radius r at $r = 6$ cm is equal to $A\pi$. then find value of A



[Watch Video Solution](#)

18. The total revenue in Rupees received from the sale of x units of a product is given by $R(x) = 3x^2 + 36x + 5$. The marginal revenue, when $x = 15$ is

A. 116

B. 96

C. 90

Answer: D



[Watch Video Solution](#)

Exercise 6 2

1. Show that the function given by $f(x) = 3x + 17$ is strictly increasing on \mathbb{R} .



[Watch Video Solution](#)

2. Show that the function given by $f(x) = e^{2x}$ is strictly increasing on \mathbb{R} .



[Watch Video Solution](#)

3. Show that $f(x) = \sin x$ is increasing on $(0, \pi/2)$ and decreasing on $(\pi/2, \pi)$ and neither increasing nor decreasing in $(0, \pi)$.

 [Watch Video Solution](#)

4. Find the intervals in which the function f given by $f(x) = 2x^2 - 3x$ is (a) strictly increasing (b) strictly decreasing

 [Watch Video Solution](#)

5. Find the intervals in which the function f given by $f(x) = 2x^3 - 3x^2 - 36x + 7$ is (a) strictly increasing (b) strictly decreasing

 [Watch Video Solution](#)

6. Find the intervals in which the following functions are strictly increasing or decreasing: (a) $x^2 + 2x - 5$ (b) $10 - 6x - 2x^2$ (c) $-2x^3 - 9x^2 - 12x + 1$ (d) $6 - 9x - x^2$ (e) $(x + 1)^3(x - 3)^3$

 [Watch Video Solution](#)

7. Show that $y = \log(1 + x) - \frac{2x}{2 + x}$, $x > 1$ is an increasing function of x throughout its domain.

 [Watch Video Solution](#)

8. Find the values of x for which $y = [x(x - 2)]^2$ is an increasing function

 [Watch Video Solution](#)

9. Prove that $y = \frac{4 \sin \theta}{2 + \cos \theta} - \theta$ is an increasing function of $\theta \in \left[0, \frac{\pi}{2}\right]$

 [Watch Video Solution](#)

10. Prove that the logarithmic function is strictly increasing on $(0, \infty)$.

 [Watch Video Solution](#)

11. Prove that the function $f(x) = x^2 - x + 1$ is neither increasing nor decreasing on $(-1, 1)$.

 [Watch Video Solution](#)

12. Which of the following functions are strictly decreasing on $\left[0, \frac{\pi}{2}\right]$ (A) $\cos x$ (B) $\cos 2x$ (C) $\cos 3x$ (D) $\tan x$

 [Watch Video Solution](#)

13. On which of the following intervals is the function $x^{100} + \sin x - 1$ decreasing? (a) $\left(0, \frac{\pi}{2}\right)$ (b) $(0, 1)$ (c) $\left(\frac{\pi}{2}, \pi\right)$ (d) none of these

 [Watch Video Solution](#)



Watch Video Solution

14. Find the least value of a such that the function f given by $f(x) = x^2 + ax + 1$ is strictly increasing on $(1, 2)$.



Watch Video Solution

15. Let I be any interval disjoint from $(1, 1)$. Prove that the function f given by $f(x) = x + \frac{1}{x}$ is strictly increasing on I .



Watch Video Solution

16. Prove that the function f given by $f(x) = \log \sin x$ is increasing on $\left(0, \frac{\pi}{2}\right)$ and decreasing on $\left(\frac{\pi}{2}, \pi\right)$.



View Text Solution

17. Prove that the function f given by $f(x) = \log|\cos x|$ is decreasing on $\left(0, \frac{\pi}{2}\right)$ and increasing on $\left(\frac{3\pi}{2}, 2\pi\right)$.

 [View Text Solution](#)

18. Prove that the function given by $f(x) = x^3 - 3x^2 + 3x - 100$ is increasing in \mathbb{R} .

 [Watch Video Solution](#)

19. The interval in which $y = x^2e^{-x}$ is increasing is

A. $(-\infty, \infty)$

B. $(-2, 0)$

C. $(2, \infty)$

D. $(0, 2)$

Answer: D

 [Watch Video Solution](#)

Exercise 6 3

1. Find the slope of the tangent to the curve $y = 3x^4 - 4x$ at $x = 4$.

 [Watch Video Solution](#)

2. Find the slope of the tangent to the curve $y = \frac{x - 1}{x - 2}, x \neq 2$ at $x = 10$.

 [Watch Video Solution](#)

3. Find the slope of the tangent to curve $y = x^3 - x + 1$ at the point whose x-coordinate is 2.

 [Watch Video Solution](#)

 Watch Video Solution

4. Find the slope of the tangent to the curve $y = x^3 - 3x + 2$ at the point whose x-coordinate is 3.

 Watch Video Solution

5. Find the slope of the normal to the curve $x = a \cos^3 \theta, y = \sin^3 \theta$ at $\theta = \frac{\pi}{4}$.

 Watch Video Solution

6. Find the slope of the normal to the curve $x = 1 - a \sin \theta, y = b \cos^2 \theta$ at $\theta = \frac{\pi}{2}$.

 Watch Video Solution

7. Find points at which the tangent to the curve $y = x^3 - 3x^2 - 9x + 7$ is parallel to the x-axis.

 [Watch Video Solution](#)

8. Find a point on the curve $y = (x - 2)^2$ at which the tangent is parallel to the chord joining the points (2, 0) and (4, 4).

 [Watch Video Solution](#)

9. Find the point on the curve $y = x^3 - 11x + 5$ at which the tangent is $y = x - 11$.

 [Watch Video Solution](#)

10. Find the equation of all lines having slope 1 that are tangents to the curve $y = \frac{1}{x - 1}$, $x \neq 1$.





Watch Video Solution

11. Find the equation of all lines having slope 2 which are tangents to the

curve $y = \frac{1}{x - 3}, x \neq 3$



Watch Video Solution

12. Find the equations of all lines having slope 0 which are tangent to the

curve $y = \frac{1}{x^2 - 2x + 3}$.



Watch Video Solution

13. Find points on the curve $\frac{x^2}{9} + \frac{y^2}{16} = 1$ at which the tangents are (i) parallel to x-axis (ii) parallel to y-axis.



Watch Video Solution

14. Find the equations of the tangent and normal to the given curves at the indicated points: (i) $y = x^4 - 6x^3 + 13x^2 - 10x + 5$ at $(0, 5)$ (ii) $y = x^4 - 6x^3 + 13x^2 - 10x + 5$ at $(1, 3)$ (iii) $y = x^3$ at $(1, 1)$ (iv) $y = x^2$ at $(0, 0)$ (v) $y = x^2$ at $(0, 0)$

 [Watch Video Solution](#)

15. Find the equation of the tangent line to the curve $y = x^2 - 2x + 7$ which is (a) parallel to the line $2x - y + 9 = 0$ (b) perpendicular to the line $5y - 15x = 13$.

 [Watch Video Solution](#)

16. Show that the tangents to the curve $y = 7x^3 + 11$ at the points where $x = 2$ and $x = -2$ are parallel.

 [Watch Video Solution](#)

17. Find the points on the curve $y = x^3$ at which the slope of the tangent is equal to the y-coordinate of the point.

 [Watch Video Solution](#)

18. For the curve $y = 4x^3 - 2x^5$, find all the points at which the tangent passes through the origin.

 [Watch Video Solution](#)

19. Find the points on the curve $x^2 + y^2 - 2x - 3 = 0$ at which the tangents are parallel to the x-axis.

 [Watch Video Solution](#)

20. Find the equation of the normal at the point (am^2, am^3) for the curve $ay^2 = x^3$.

 [Watch Video Solution](#)

21. Find the equation of the normals to the curve $y = x^3 + 2x + 6$ which are parallel to the line $x + 14y + 4 = 0$.

 [Watch Video Solution](#)

22. Find the equations of the tangent and normal to the parabola $y^2 = 4ax$ at the point $(at^2, 2at)$.

 [Watch Video Solution](#)

23. Show that the curves $x = y^2$ and $xy = k$ cut at right angles; if $8k^2 = 1$

 [Watch Video Solution](#)

24. Find the equations of the tangent and normal to the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1. \text{ at the point } (x_0, y_0)$$



Watch Video Solution

25. Find the equation of the tangent to the curve $y = \sqrt{3x - 2}$ which is parallel to the line $4x - 2y + 5 = 0$.



Watch Video Solution

26. The slope of the normal to the curve $y = 2x^2 + 3\sin x$ at $x = 0$ is

A. 3

B. $\frac{1}{3}$

C. -3

D. $-\frac{1}{3}$

Answer: D



Watch Video Solution

27. The line $y = x + 1$ is a tangent to the curve $y^2 = 4x$ at the point(A)

(1, 2) (B)(2, 1) (C) (1, 2) (D) (1, 2)

A. (1, 2)

B. (2, 1)

C. (1, - 2)

D. (- 1, 2)

Answer: A



Watch Video Solution

1. Using differentials, find the approximate value of each of the following up to 3 places of decimal.

$(i) \sqrt{25.3}$

$(ii) \sqrt{49.5}$

$(iii) \sqrt{0.6}$

$(iv) (0.009)^{\frac{1}{3}}$

$(v) (0.999)^{\frac{1}{10}}$

$(vi) (15)^{\frac{1}{4}}$

$(vii) (26)^{\frac{1}{3}}$

$(viii) (255)^{\frac{1}{4}}$

$(ix) (82)^{\frac{1}{4}}$

$(x) (401)^{\frac{1}{2}}$

$(xi) (0.0037)^{\frac{1}{2}}$

$(xii) (26.57)^{\frac{1}{3}}$

$(xiii) (81.5)^{\frac{1}{4}}$

$(xiv) (3.968)^{\frac{3}{2}}$

$(xv) (32.15)^{\frac{1}{5}}$



[View Text Solution](#)

2. Find the approximate value of $f(2.01)$, where $f(x) = 4x^2 + 5x + 2$.



[Watch Video Solution](#)

3. Find the approximate value of $f(5.001)$, where $f(x) = x^3 - 7x^2 + 15$.



[Watch Video Solution](#)

4. Find the approximate change in the volume V of a cube of side x metres caused by increasing the side by 1%.



[Watch Video Solution](#)

5. Find the approximate change in the surface area of a cube of side x metres caused by decreasing the side by 1%.



[Watch Video Solution](#)

6. If the radius of a sphere is measured as 7 m with an error of 0.02 m, find the approximate error in calculating its volume.



[Watch Video Solution](#)

7. If the radius of a sphere is measured as 9 cm with an error of 0.03 m, find the approximate error in calculating its surface area.

 [Watch Video Solution](#)

8. If $f(x) = 3x^2 + 15x + 5$, then the approximate value of $f(3.02)$ is :

A. 47.66

B. 57.66

C. 67.66

D. 77.66

Answer: D

 [Watch Video Solution](#)

9. The approximate change in the volume of a cube of side x metres caused by increasing the side by 3% is (A) $0.06 x^3 m^3$ (B) $0.6 x^3 m^3$ (C) 0.09

$$x^3m^3 \text{ (D) } 0.9x^3m^3$$

A. $0.06x^3m^3$

B. $0.6x^3m^3$

C. $0.09x^3m^3$

D. $0.9x^3m^3$

Answer: C



Watch Video Solution

Exercise 6 5

1. Find the maximum and minimum values, if any, of the following functions given by (i) $f(x) = (2x - 1)^2 + 3$ (ii) $f(x) = 9x^2 + 12x + 2$ (iii) $f(x) = -(x - 1)^2 + 10$ (iv) $g(x) = x^3 + 1$



Watch Video Solution

2. Find the maximum and minimum values, if any, of the following functions given by
- (i) $f(x) = |x + 2| - 1$ (ii)
- $g(x) = |x + 1| + 3$ (iii)
- $h(x) = s \in (2x) + 5$ (iv) 'f' '(x)' '=' '|''s in' '4x' '+'



Watch Video Solution

3. Find the local maxima and local minima, if any, of the following functions. Find also the local maximum and the local minimum values, as the case may be:

(i) $f(x) = x^2$

(ii) $g(x) = x^3 - 3x$

(iii) $h(x) = \sin x + \cos x, 0 < x < \frac{\pi}{2}$

(iv) $f(x) = \sin x - \cos x, 0 < x < 2\pi$

(v) $f(x) = x^3 - 6x^2 + 9x + 15$

(vi) $g(x) = \frac{x}{2} + \frac{2}{x}, x > 0$

(vii) $g(x) = \frac{1}{x^2 + 2}$

(viii) $f(x) = x\sqrt{1-x}, 0 < x < 1$



Watch Video Solution

[View Text Solution](#)

4. Prove that the following functions do not have maxima or minima:(i)

$$f(x) = ex \text{ (ii) } g(x) = \log x \text{ (iii) } h(x) = x^3 + x^2 + x + 1$$

 [Watch Video Solution](#)

5. Find the absolute maximum value and the absolute minimum value of the following functions in the given intervals:(i) $f(x) = x^2, x \in [-2, 2]$

(ii) $f(x) = \sin x + \cos x, x \in [0, \pi]$ (iii)

$f(x) = 4x - \frac{1}{2}x^2, x \in \left[-2, \frac{9}{2}\right]$ (iv) $f(x) = (x-1)$

 [Watch Video Solution](#)

6. Find the maximum profit that a company can make, if the profit function is given by $p(x) = 41 - 24x - 18x^2$

 [Watch Video Solution](#)

7. Find the maximum value and the minimum value and the minimum value of $3x^4 - 8x^3 + 12x^2 - 48x + 25$ on the interval $[0, 3]$.

 [Watch Video Solution](#)

8. At what points in the interval $[0, 2\pi]$, does the function $\sin 2x$ attain its maximum value?

 [Watch Video Solution](#)

9. What is the maximum value of the function $\sin x + \cos x$?

 [Watch Video Solution](#)

10. Find the maximum value of $2x^3 - 24x + 107$ in the interval $[1, 3]$. Find the maximum value of the same function in $[3, 1]$.

 [Watch Video Solution](#)

11. It is given that at $x = 1$, the function $x^4 - 62x^2 + ax + 9$ attains its maximum value, on the interval $[0, 2]$. Find the value of a .



[Watch Video Solution](#)

12. Find the maximum and minimum values of $x + \sin 2x$ on $[0, 2\pi]$.



[Watch Video Solution](#)

13. Find two numbers whose sum is 24 and whose product is as large as possible.



[Watch Video Solution](#)

14. Find two positive numbers x and y such that $x + y = 60$ and xy^3 is maximum.

A. $x = 15, y = 45$

B. $x = 15, y = 40$

C. $x = 10, y = 50$

D. $x = 20, y = 40$

Answer: A



Watch Video Solution

15. Find two positive numbers x and y such that their sum is 35 and the product x^2y^5 is a maximum.

A. $x = 15, y = 25$

B.

C.

D.

Answer:

 [Watch Video Solution](#)

16. Find two positive numbers whose sum is 16 and the sum of whose cubes is minimum.

 [Watch Video Solution](#)

17. A square piece of tin of side 18 cm is to be made into a box without top, by cutting a square from each corner and folding up the flaps to form the box. What should be the side of the square to be cut off so that the volume of the box is the maxi


 [Watch Video Solution](#)

18. A rectangular sheet of tin 45 cm by 24 cm is to be made into a box without top, by cutting off square from each corner and folding up the flaps. What should be the side of the square to be cut off so that the volume of the box is maximum ?



[Watch Video Solution](#)

19. Show that of all the rectangles in a given fixed circle, the square has the maximum area.

 width="80%">



[View Text Solution](#)

20. Show that the right circular cylinder of given surface and maximum volume is such that its height is equal to the diameter of the base.



[Watch Video Solution](#)

21. Of all the closed cylindrical cans (right circular), of a given volume of 100 cubic centimetres, find the dimensions of the can which has the minimum surface area?



[Watch Video Solution](#)

22. A wire of length 28 m is to be cut into two pieces. One of the pieces is to be made into a square and the other into a circle. What should be the length of the two pieces so that the combined area of the square and the circle is minimum?

 [Watch Video Solution](#)

23. Prove that the volume of the largest cone, that can be inscribed in a sphere of radius R , is $\frac{8}{27}$ of the volume of the sphere.

 [Watch Video Solution](#)

24. Show that the right circular cone of least curved surface and given volume has an altitude equal to $\sqrt{2}$ time the radius of the base.

 [Watch Video Solution](#)

25. Show that the semi-vertical angle of the cone of the maximum volume and of given slant height is $\tan^{-1} \sqrt{2}$.

 [Watch Video Solution](#)

26. Show that semi-vertical angle of right circular cone of given surface area and maximum volume is $\sin^{-1} \left(\frac{1}{3} \right)$.

 [Watch Video Solution](#)

27. The point on the curve $x^2 = 2y$ which is nearest to the point (0, 5) is (A) $(2\sqrt{2}, 4)$ (B) $(2\sqrt{2}, 0)$ (C) (0, 0) (D) (2, 2)

A. $(2\sqrt{2}, 4)$

B. $(2\sqrt{2}, 0)$

C. (0, 0)

D. (2, 2)

Answer: A



Watch Video Solution

28. For all real values of x , the minimum value of $\frac{1 - x + x^2}{1 + x + x^2}$ is

A. 0

B. 1

C. 3

D. $\frac{1}{3}$

Answer: D



Watch Video Solution

29. The maximum value of $[x(x - 1) + 1]^{\frac{1}{3}}$, $0 \leq x \leq 1$ is:

A. $\left(\frac{1}{3}\right)^{\frac{1}{3}}$

B. $\frac{1}{2}$

C. 1

D. 0

Answer: C

 [View Text Solution](#)

Miscellaneous Exercise

1. Using differentials, find the approximate value of each of the following:

(a) $\left(\frac{17}{81}\right)^{\frac{1}{4}}$ (b) $(33)^{-\frac{1}{5}}$

 [View Text Solution](#)

2. Show that the function given by $f(x) = \frac{\log x}{x}$ has maximum at $x = e$.

 [Watch Video Solution](#)

3. The two equal sides of an isosceles triangle with fixed base b are decreasing at the rate of 3 cm per second. How fast is the area decreasing when the two equal sides are equal to the base ?

 [Watch Video Solution](#)

4. Find the equation of the normal to the curve $x^2 = 4y$ which passes through the point $(1, 2)$.

 [Watch Video Solution](#)

5. Show that the normal at any point θ to the curve $x = a \cos \theta + a\theta \sin \theta$, $y = a \sin \theta - a \theta \cos \theta$ is at a constant distance from the origin.

 [Watch Video Solution](#)

6. Find the intervals in which the function f given by

$$f(x) = \frac{4 \sin x - 2x - x \cos x}{2 + \cos x} \text{ is (i) increasing (ii) decreasing.}$$

 [Watch Video Solution](#)

7. Find the intervals in which the function f given by

$$f(x) = x^3 + \frac{1}{x^3}, \quad x \neq 0 \text{ is (i) increasing (ii) decreasing.}$$

 [Watch Video Solution](#)

8. Find the maximum area of an isosceles triangle inscribed in the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ with its vertex at one end of the major axis.}$$

 [Watch Video Solution](#)

9. A tank with rectangular base and rectangular sides, open at the top is to be constructed so that its depth is 2 m and volume is 8 m³. If building

of tank costs Rs 70 per square metre for the base and Rs 45 per square metre for sides, what is the cost of least expensive tank?

 [Watch Video Solution](#)

10. The sum of the perimeter of a circle and square is k , where k is some constant. Prove that the sum of their areas is least when the side of square is double the radius of the circle.

 [Watch Video Solution](#)

11. A window is in the form of a rectangle surmounted by a semicircular opening. The total perimeter of the window is 10 m. Find the dimensions of the window to admit maximum light through the whole opening.

 [Watch Video Solution](#)

12. A point on the hypotenuse of a triangle is at distance a and b from the sides of the triangle. Show that the maximum length of the hypotenuse is

$$\left(a^{\frac{2}{3}} + b^{\frac{2}{3}}\right)^{\frac{3}{2}}.$$



[Watch Video Solution](#)

13. Find the points at which the function f given by $f(x) = (x - 2)^4(x + 1)^3$ has local maxima local minima point of inflexion



[Watch Video Solution](#)

14. Find the absolute maximum and minimum values of the function f given by $f(x) = \cos^2 x + \sin x$, $x \in [0, \pi]$.



[Watch Video Solution](#)

15. Show that the altitude of the right circular cone of maximum volume that can be inscribed in a sphere of radius r is $\frac{4r}{3}$.

 [Watch Video Solution](#)

16. Let f be a function defined on $[a, b]$ such that $f'(x) > 0$, for all $x \in (a, b)$. Then prove that f is an increasing function on (a, b) .

 [Watch Video Solution](#)

17. Show that the height of the cylinder of maximum volume that can be inscribed in a sphere of radius R is $\frac{2R}{\sqrt{3}}$. Also find the maximum volume.

 [Watch Video Solution](#)

18. Show that height of the cylinder of greatest volume which can be inscribed in a right circular cone of height h and semi vertical angle is

one-third that of the cone and the greatest volume of cylinder is

$$\frac{4}{27}\pi h^3 \tan^2 \alpha.$$

 [Watch Video Solution](#)

19. A cylindrical tank of radius 10 m is being filled with wheat at the rate of 314 cubic metre per hour. Then the depth of the wheat is increasing at the rate of

- A. 1 m/h
- B. $0.1m / h$
- C. $1.1m / h$
- D. $0.5m / h$

Answer: A

 [Watch Video Solution](#)

20. The slope of the tangent to the curve $x = t^2 + 3t - 8$,
 $y = 2t^2 - 2t - 5$ at the point $(2, 1)$

A. $\frac{22}{7}$

B. $\frac{6}{7}$

C. $\frac{7}{6}$

D. $\frac{-6}{7}$

Answer: B



Watch Video Solution

21. The line $y = mx + 1$ is a tangent to the curve $y^2 = 4x$ if the value of m is

A. 1

B. 2

C. 3

D. $\frac{1}{2}$

Answer: A



Watch Video Solution

22. The normal at the point $(1, 1)$ on the curve $2y + x^2 = 3$ is

A. $x+y=0$

B. $x-y= 0$

C. $x+y+1=0$

D. $x-y=1$

Answer: B



Watch Video Solution

23. The normal to the curve $x^2 = 4y$ passing $(1, 2)$ is

A. $x+y = 3$

B. $x - y = 3$

C. $x+ y = 1$

D. $x- y = 1$

Answer: A



Watch Video Solution

24. Find the point on the curve $9y^2 = x^3$, where the normal to the curve makes equal intercepts on the axes.

A. $\left(4, \pm \frac{8}{3}\right)$

B. $4, \frac{-8}{3}$

C. $\left(4, \pm \frac{3}{8}\right)$

D. $\left(\pm 4, \frac{3}{8}\right)$

Answer: A



Watch Video Solution