



MATHS

BOOKS - NAGEEN MATHS (HINGLISH)

PROBABILITY

Solved Example Type

1. For two events A and B $P(A) = \frac{7}{13}$, $P(B) = \frac{9}{13}$ and $(A \cap B) = \frac{4}{13}$.

Find the value of $P(A/B)$.

A. $\frac{4}{11}$

B. $\frac{2}{9}$

C. $\frac{4}{9}$

D.

Answer: C



[Watch Video Solution](#)

2. If A and B are two events such that $P(A) = 0.4$, $P(B) = 0.8$ and $P(B/a) = 0.6$, then find:

(i) $P(A \cap B)$ (ii) $P(A \cup B)$ (iii) $P(A/B)$



[Watch Video Solution](#)

3. In one throw of a dice, the result is an even number. Find the probability that it is prime.



[Watch Video Solution](#)

4. There are two children in a family. Find the probability that both boys if it is known that:

(i) one of the children is a boy

(ii) Older child is a by



[Watch Video Solution](#)



Watch Video Solution

5. A dice is thrown twice and the sum of numbers appear to be 6. Find the probability of getting 4 on at least one throw.



Watch Video Solution

6. 20 cards whose number are from 1 to 20 are mixed after placing in a box. A card is drawn at random and it is found that the number on the card is greater than 11. Find the probability that the number on the card is an even number.

A. $\frac{5}{9}$

B. $\frac{5}{20}$

C. $\frac{9}{20}$

D. $\frac{9}{5}$

Answer: A



 [Watch Video Solution](#)

7. Two integers are selected at random from the integers 1 to 11. If their sum is even find the probability that both integers are odd.

 [Watch Video Solution](#)

8. If $P(A) = \frac{2}{5}$ and $P(B) = \frac{1}{3}$ and A and B are independent events, then find $P(A \cap B)$.

 [Watch Video Solution](#)

9. If E and F are two independent events and $P(E) = \frac{1}{3}$, $P(F) = \frac{1}{4}$, then find $P(E \cup F)$.

 [Watch Video Solution](#)

10. If A and B are two independent events and $P(A) = 0.2$, $P(B) = 0.3$, then find the values of the following:

- (i) $P(A \text{ and } B)$ (ii) $P(A \text{ and not } B)$
(iii) $P(A \text{ or } B)$ (iv) $P(\text{none of } A \text{ and } B)$

 [Watch Video Solution](#)

11. There are 5 red and 7 white balls in a bag. Two balls are drawn one by one without replacement. Find the probability that first ball is red and second is white.

 [Watch Video Solution](#)

12. A dice is thrown once. If E is the event of getting a multiple of 3 and F is the event of getting an even number, then find whether E and F are independent or not?

 [Watch Video Solution](#)

13. There are 5 red, 4 black and 3 white in a bag. Three balls are drawn one by one without replacement find the probability that all three balls are red.



[Watch Video Solution](#)

14. Two cards are drawn one by one without replacement. Find the probability that both are hearts.



[Watch Video Solution](#)

15. A man A truth in 70% of the cases while another man B speaks truth in 60% of the cases. Find the probability of an event in which they agree with on another.



[Watch Video Solution](#)

16. There are 3 white, 6 black balls in one bag and 6 white, 3 black balls in second bag. A bag is selected at random and a ball is drawn from it. Find the probability that the ball drawn is black.

A. $\frac{3}{7}$

B. $\frac{3}{10}$

C. $\frac{1}{4}$

D. $\frac{1}{2}$

Answer: D



[Watch Video Solution](#)

17. A man takes the contract of construction work. The probability is 0.30, if there will be no strike. The probability that the construct work will be completed on time are 0.8 and 0.3 in cases of no strike and in case strike respectively. Find the probability that the construction work will be completed in time.

A. 0.34

B. 0.56

C. 0.64

D. 0.45

Answer: D



[Watch Video Solution](#)

18. Two thirds of the students in a class are boys and the rest girls. It is known that the probability of a girl getting a first class is 0.25 and that of a boy getting a first class is 0.28. Find the probability that a student chosen at random will get first class marks in the subject.



[Watch Video Solution](#)

19. A company has two plants to manufacture bicycles. The first plant manufactures 60% of the bicycles and the second plant 40%. Out of that

80% of the bicycles are rated of standard quality at the first plant and 90% of standard quality at the second plant. A bicycle is picked up at random and found to be standard quality. Find the probability that it comes from the second plant.

 [Watch Video Solution](#)

20. One bag a contains 4 red and 5 black balls. The other bag B contains 6 red and 3 black balls. A ball is taken rom bag A and transferred to bag B. Now a ball is from bag B. Find the probability that the ball drawn is red.

 [Watch Video Solution](#)

21. Find the mean and variance of the number of heads in the two tosses of a coin.

 [Watch Video Solution](#)

22. Two cards are drawn simultaneously (or successively without replacement) from a well shuffled pack of 52 cards. Find the mean, variance and standard deviation of the number of kings.

 [Watch Video Solution](#)

23. There are 4 white and 5 red balls in a bag. Three balls are drawn with replacement. Find the probability distribution for number of red balls.

 [Watch Video Solution](#)

24. A coin is tossed 5 times. If X is the number of heads observed, find the probability distribution of X .

 [Watch Video Solution](#)

25. An unbiased coin is tossed 8 times. Find , by using binomial distribution, the probability of getting at least 6 heads.

 [Watch Video Solution](#)

26. There are 5 questions in a multiple choice examination in which each question has 3 possible answers. Find the probability that a student gives 4 correct answers by guess only.

 [Watch Video Solution](#)

27. The probability of a shooter hitting a target is $\frac{3}{4}$. How many minimum number of times must he/she fire so that the probability of hitting the target at least once is more than 0.99?

 [Watch Video Solution](#)

1. If A and B are two events such that $P(A) = 0.5$, $P(B) = 0.6$ and $P(A \cup B) = 0.8$, find $P(A/B)$ and $P(B/A)$.

 [Watch Video Solution](#)

2. For two events A and B $P(A) = \frac{6}{13}$, $P(B) = \frac{5}{13}$ and $P(A \cup B) = \frac{7}{13}$. Find the values of following:

(i) $P(A \cap B)$ (ii) $P(A/B)$

(iii) $P(B/A)$

 [Watch Video Solution](#)

3. For two events A and B $P(A) = 0.7$, $P(B) = 0.5$ and $P(A \cap B) = 0.3$.

Find the values of following:

(i) $P(A \cup B)$ (ii) $P(A/B)$

(iii) $P(\bar{A}/\bar{B})$ (iv) $P(\bar{B}/\bar{A})$

 [Watch Video Solution](#)

4. In a throw of a dice a number divisible by 3 appears. Find the probability that it is an even number.

 [Watch Video Solution](#)

5. If $P(A) = 0.3$, $P(B) = 0.6$ and $P(A/B) = 0.4$ then find:

(i) $P(A \cap B)$ (ii) $P(B/A)$

 [Watch Video Solution](#)

6. A card is drawn from a well shuffled pack of 52 cards and it is found to be red. Find the probability that it is a king.

 [View Text Solution](#)

7. In one toss of three coins, at least one head appears. Find the probability of getting three heads.



[View Text Solution](#)

8. In three throws of a dice, the events A and B are defined as follows:

A : 4 appears on 3rd throw

B : 3 appears on 1st throw and 5 appears on 2nd throw.

Find the probability of the occurrence of event A if it is known that event B is already occur.



[View Text Solution](#)

9. There are two children in a family. It is known that at least one child is a boy, then find the probability that both are boys.



[View Text Solution](#)

10. 12 cards, numbered 1 to 12 are given. A card is drawn at random. It is known that the number on this card is greater than 6. Find the probability that it is odd.



[Watch Video Solution](#)

11. The sum of numbers on two throws of a dice is 8. Find the probability of getting 3 in at least one throw.



[Watch Video Solution](#)

12. It is known that the numbers getting in one throw of two dice are different. Find the probability of getting a sum of 6.



[Watch Video Solution](#)

13. In a school 25% students are failed in Mathematics, 15% in Physics and 5% students fail in both Mathematics and Physics. A student is selected

at random.

(i) If he is fail in Physics, find the probability that he is fail in Mathematics.

(ii) If he is fail in Mathematics, find the probability that he in fail in Physics.

(iii) Find the probability that he is fail either in Mathematics or in Physics.

 [Watch Video Solution](#)

14. Two integers are selected at random from the integers 1 to 9. If their sum is even, then find the probability that both integers are odd.

 [Watch Video Solution](#)

15. In a throw of two dice, the numbers obtained are different. Find the probability that their sum is 4.

 [Watch Video Solution](#)

1. If A and B are independent events such that $P(A) = 0.4$ and $P(B) = 0.5$, then find $P(A \cap B)$.

 [Watch Video Solution](#)

2. If A and B are two events such that $P(A) = \frac{1}{2}$, $P(A \cup B) = \frac{3}{5}$ and $P(B) = p$, then find the value of p when:

- (i) A and B are mutually exclusive
- (ii) A and B are independent events.

 [Watch Video Solution](#)

3. If A and B are independent events such that $P(A) = 0.3$ and $P(B) = 0.5$, then find the values of:

- (i) $P(A \cap B)$ (ii) $P(A \cup B)$
- (iii) $P(A/B)$ (iv) $P(B/A)$

 [Watch Video Solution](#)

4. A and B are two events such that

$$P(A) = \frac{1}{4}, P(B) = \frac{1}{2} \text{ and } P(A \cap B) = \frac{1}{8}$$

Find the value of $P(\text{not A and not B})$.



[Watch Video Solution](#)

5. If A and B are two independent events, then prove that the probability that at least one event will happen will be $1 - P(A') \cdot P(B')$.



[Watch Video Solution](#)

6. A and b are two events such that $P(A) = \frac{3}{5}$, $P(B) = \frac{3}{10}$ and $P(A \cup B) = \frac{1}{2}$

Find whether A and B are independent or not.



[Watch Video Solution](#)

7. There are 10 black and 8 white balls in a bag. Two balls are drawn without replacement. Find the probability that:

- (i) both balls are black
- (ii) first ball is black and second is white
- (iii) one ball is black and other is white.



[Watch Video Solution](#)

8. A card is drawn from a well shuffled pack of 52 cards. In which of the following cases, the events E and F are independent events?

(i) E = card drawn is heart.

F = card drawn is a king.

(ii) E = card drawn is red.

F = card drawn is a king.



[Watch Video Solution](#)

9. The probabilities that A and b can solve a problem independently are $\frac{1}{3}$ and $\frac{1}{6}$ respectively. If both try to solve the problem independently, find the probability that:

- (i) problem will be solved
- (ii) both can solve the problem.



[Watch Video Solution](#)

10. Three cards are drawn one by one without replacement from a well shuffled pack of 52 cards. Find the probability that the first two cards drawn are queen and third card drawn is a king.



[Watch Video Solution](#)

11. In three throws of a dice, find the probability of getting odd number at least in one throw.



[Watch Video Solution](#)

12. In a hostel 60% students read Hindi newspaper, 40% read English newspaper and 20% read both newspaper. A student is selected at random. Find the probability that student neither read Hindi newspaper nor read English newspaper.

 [Watch Video Solution](#)

13. A person A speaks truth in 75% cases while other person B in 80% cases. In what percent of cases they contradict each other?

 [Watch Video Solution](#)

14. The odds in favour of A whose age is 45 years will live upto 60 years are 7:10. The odds against of B whose age is 50 years, will live upto 65 years are 3:2. Find the probability that at least one will live upto next 15 years.

 [Watch Video Solution](#)

15. In a company there are two vacancies. A man and his wife come for interview. The probability of selection of mans is $\frac{1}{10}$ and his wife is $\frac{1}{8}$.

Find the probabilty that:

- (i) both will be selected
- (ii) only one will be selected
- (iii) none of them will be selected.

 [Watch Video Solution](#)

16. A bag contains 50 tickets marked with numbers 1,2,.....50. Five tickets are selected at random from the bag and arranged in ascending order $x_1 < x_2 < x_3 < x_4 < x_5$. Find the probability that $x_3 = 30$.

 [Watch Video Solution](#)

17. In a company there are two vacancies. A man and his wife come for interview. The probability of selection of mans is $\frac{1}{10}$ and his wife is $\frac{1}{8}$.

Find the probabilty that:

- (i) both will be selected
- (ii) only one will be selected
- (iii) none of them will be selected.

 [Watch Video Solution](#)

Exercise 13 C

1. There are 3 red and 2 black, 2 red and 3 black, 4 red and 1 black balls in three bags respectively. The probability of selecting each bag is same. A ball is drawn at random from one bag. Find the probability that it is black.

 [Watch Video Solution](#)

2. There are two pockets in a purse. First pocket contains 3,50 paise coin and 2, one rupee coins while second pocket contains 2,50 paise coins and 3, one rupee coins. Find the probability of getting a one rupee coin.

 [Watch Video Solution](#)

3. A bag contains 4 white and 5 red balls. Second bag contains 6 white and 3 red balls. A ball is transferred from first bag to second and then a ball is drawn from the second bag. Find the probability that this ball drawn is red.



[Watch Video Solution](#)

4. One bag contains 5 white balls and 3 black balls and a second bag contains 2 white balls and 4 black balls. One ball drawn from the first bag and placed unseen in the second bag. What is the probability that a ball now drawn from second bag is black?



[Watch Video Solution](#)

5. A bag contains 3 black and 4 white balls. Second bag contains 5 black and 6 white balls. A ball is drawn at random from one of the bags and it is

found to be black. What is the probability that this ball is drawn from second bag?

 [Watch Video Solution](#)

6. In a bolt factory, machines A, B and C manufacture 25%, 35% and 40% respectively of the total output from which 5%, 4% and 2% respectively are defective. A bolt is drawn and is found to be defective. What is the probability that it was manufactured by the machine C?

 [Watch Video Solution](#)

7. In a factory, two machines A and B manufacture 60% and 40% respectively of the total output from which 2% and 1% respectively are defective. If randomly selected one item is defective, find the probability that it was manufactured by machine B.

 [Watch Video Solution](#)

8. A doctor is to visit a patient. From the past experience, it is known that the probabilities that he will come by train, bus, scooter or by other means of transportation are respectively $\frac{3}{10}$, $\frac{1}{5}$, $\frac{1}{10}$ or $\frac{2}{5}$. The probabilities that he will be late are $\frac{1}{4}$, $\frac{1}{3}$ and $\frac{1}{12}$ if he comes by train, bus and scooter respectively but if he comes by other means of transport, then he will not be late. If he arrives late, find the probability that he comes by train.

 [Watch Video Solution](#)

9. Probability that A speaks truth is $\frac{4}{5}$. A coin is tossed and A reports that a tail appears. Find the probability that actually there was tail.

 [Watch Video Solution](#)

10. A man is known to speak the truth 3 out of 4 times. He throws a dice and reports that it is a six. Find the probability that it is actually a six.

 [Watch Video Solution](#)

11. There are three coins . One con has tails on both side. Another is a biased coin that shows tails in 75% cases. The third coins is unbiased.. One of the three coins is chosen at random and tossed it showed tail. What is the probability that it is first coin.

 [Watch Video Solution](#)

12. Given that 5% of men and 0.25% of women have white hair. A white haired person is selected at random. What is the probability that this person is a man. Assume that there are equal number of men and women.

 [Watch Video Solution](#)

13. In answering a question on a multiple choice test a student either knows the answer or guesses. Let the probability that he knows the answer is $\frac{3}{4}$ and probability that he guesses is $\frac{1}{4}$. Assuming that a student who guesses the answer and given correct answer is $\frac{1}{4}$. What is

the probability that student knows the answer given that he answered it correctly?

 [Watch Video Solution](#)

14. Suppose a girl throws a dice. If she gets 5 or 6 she tosses a coin three times and note the number of tails. If she gets 1,2,3 or 4, she tossed a coin once and notes whether a head or tails is obtained. If she is obtained exactly one tail, what is the probability that she threw 1,2,3 or 4 with the dice?

 [Watch Video Solution](#)

15. Two groups are competing for the position of the Board of Directors of a Corporation. The probabilities that the first and the second groups will win are 0.6 and 0.4 respectively. Further, if the first group wins the probability of introducing a new product is 0.7 and the corresponding probability is 0.3, if the second group wins. Find the probability that the new product introduced was by the second group.



[Watch Video Solution](#)

Exercise 13 D

1. Find the probability distribution for number of heads obtained in two tosses of a coin.



[Watch Video Solution](#)

2. Find the probability distribution for number of heads obtained in three tosses of a coin.



[Watch Video Solution](#)

3. Two cards are drawn successively with replacement from a well shuffled pack of 52 cards. Find the probability distribution of the number of aces.



[Watch Video Solution](#)

4. Which of the following probability distribution does not exist for random variable? Give reason?

(i)	X	0	1	2	
	$P(X)$	0.3	0.2	0.5	
(ii)	X	0	1	2	3
	$P(X)$	0.1	0.3	-0.2	0.8
(iii)	X	-1	0	1	2
	$P(X)$	0.4	0.2	0.3	0.15

 [Watch Video Solution](#)

5. The probability distribution for a random variable X is as follows:

X	0	1	2	3	4	5	6	7
$P(X)$	0	k	$2k$	$2k$	$3k$	k^2	$2k^2$	$7k^2 + k$

Find:

(i) k (ii) $P(X \geq 3)$

(iii) $P(X \geq 6)$ (iv) $P(0 < X < 3)$

 [Watch Video Solution](#)

6. Let X denote the number of hours you study during a randomly selected school day. The probability that X take the values of x has the following form, where k is some unknown constant.

$$P(X = x) = \begin{cases} 0.1 & \text{if } x = 0, \\ kx & \text{if } x = 1 \text{ or } 2, \\ k(5 - x) & \text{if } x = 3 \text{ or } 4, \\ 0 & \text{otherwise} \end{cases}$$

Find:

- (i) k
- (ii) the probability that you study exactly two hours,
- (iii) the probability that you study at most two hours.



[View Text Solution](#)

7. If X represents the difference of number of heads and number of tails when a coin is tossed 6 times. What are the possible values of X ?



[Watch Video Solution](#)

8. Find the probability distribution of number of doublets in three throws of a pair of dice.



[Watch Video Solution](#)

9. There are 5 red and 2 black balls in a bag. Two balls are drawn at random. If X denotes the number of black balls, what are the possible values of X ?



[Watch Video Solution](#)

10. Find the probability of getting 5 at least once in there throws of a dice.



[Watch Video Solution](#)

11. 30 बल्बों के एक ढेर से, जिसमें 6 बल्ब खराब है 4 बल्बों का एक नमूना (प्रतिदर्श) यादृच्छ्या बिना प्रतिस्थापना के निकाला जाता है। खराब बल्बों की संख्या का प्रायिकता बंटन ज्ञात कीजिए।



[Watch Video Solution](#)

12. मान लीजिए दो पासों को फेंकने पर प्राप्त संख्याओं के योग को X से व्यक्त किया गया है। X का प्रसारण और मानक विचलन ज्ञात कीजिए।

 [Watch Video Solution](#)

13. Find the mean and variance of the number of heads in the two tosses of a coin.

 [Watch Video Solution](#)

14. Two cards are drawn at random from a well shuffled pack of 52 cards without replacement, find the mean and variance of number of aces.

 [View Text Solution](#)

15. There are 3 defective bulbs, in a group of 10 bulbs. If X is the number of defective bulbs in a random draw of 3 bulbs, then find its mean and

variance.

 [View Text Solution](#)

16. There are 5 white and 4 red balls in a bag. Three balls are drawn one by one without replacement. Find the probability distribution for the number of white balls drawn.

 [Watch Video Solution](#)

17. In a game 3 coins are tossed. A person is paid Rs. 5, if he gets all head or all tail and be in supposed pay Rs. 3 if he gets are head or 2 heads. What can be expert to win on an arrange per game.

 [Watch Video Solution](#)

1. A pair of dice is thrown 4 times. If getting a doublet is considered a success, find the probability of two successes.

 [Watch Video Solution](#)

2. Suppose X has a binomial distribution $B\left(6, \frac{1}{2}\right)$. Show that $X = 3$ is the most likely outcome. (Hint: $P(x = 3)$ is the maximum among all $P(x_i)$, $x_i = 0, 1, 2, 3, 4, 5, 6$)

 [Watch Video Solution](#)

3. Find the probability of getting 5 exactly twice in 7 throws of a die.

 [Watch Video Solution](#)

4. Find the probability of throwing at most 2 sixes in 6 throws of a single die.

 [Watch Video Solution](#)

5. A fair coin is tossed four times. Let X denote the number of heads occurring. Find the probability distribution, mean and variance of X .

 [Watch Video Solution](#)

6. Find the probability distribution of number of doublets in three throws of a pair of dice.

 [Watch Video Solution](#)

7. If a fair coin is tossed 10 times, find the probability of (i) exactly six heads (ii) at least six heads (iii) at most six heads

 [Watch Video Solution](#)

8. A bag consists of 10 balls each marked with one of the digits 0 to 9. If four balls are drawn successively with replacement from the bag, what is the probability that none is marked with the digit 0?



[Watch Video Solution](#)

9. The probability that a student is not a swimmer is $\frac{1}{5}$. Then the probability that out of five students, four are swimmers is (A)

${}^5C_4 \left(\frac{4}{5}\right)^4$ (B) $\left(\frac{4}{5}\right)^4$ (C) ${}^5C_1 \frac{1}{5} \left(\frac{4}{5}\right)^4$ (D) None of these



[Watch Video Solution](#)

10. There are 10 defective bulbs in a group of 100 bulbs. If a sample of 8 bulbs are selected at random then find the probability that:

(i) in this sample 3 bulbs are defective and 5 are non-defective.

(ii) in this sample at least one bulb is defective.



[Watch Video Solution](#)

11. Ten eggs are drawn successively with replacement from a lot containing 10% defective eggs. Find the probability that there is at least one defective egg.



[Watch Video Solution](#)

12. A and B throw a dice alternately til any one gets a 6 on the and win the game. If A begins the game find the probabilities of winning the game.



[View Text Solution](#)

Exercise 13 F Multiple Choice Questions

1. If $P(A) = 0.4$, $P(B) = 0.8$, $P\left(\frac{B}{A}\right) = 0.6$. Find $P\left(\frac{A}{B}\right)$

A. 0.2

B. 0.3

C. 0.4

D. None of these

Answer: B



[Watch Video Solution](#)

2. A dice is thrown twice and the sum of numbers appear to be 6. Find the probability of getting 4 on at least one throw.

A. $\frac{2}{5}$

B. $\frac{3}{5}$

C. $\frac{1}{5}$

D. $\frac{4}{5}$

Answer: A



[Watch Video Solution](#)

3. Events E and F are independent. Find

$P(F)$, if $P(E) = 0.35$ and $P(E \cup F) = 0.6$.

A. $3/13$

B. $4/13$

C. $6/13$

D. $5/13$

Answer: D



Watch Video Solution

4. If A and B are two independent events, then the probability that at least one event will happen will be

A. $P(A')P(B')$

B. $1 - P(A')P(B')$

C. $1 + P(A')P(B')$

D. None of these

Answer: B



[Watch Video Solution](#)

5. A speaks truth in 60% cases and B speaks truth in 70% cases. The probability that they will say the same thing while describing a single event is`

A. 0.42

B. 0.58

C. 0.54

D. 0.46

Answer: C



[Watch Video Solution](#)

6. In a class two third students are boys and remaining are girls. The probability of getting first class by a boy is 0.30 and the probability of getting first class by a girl is 0.24. The probability of getting first class by a randomly selected student is:

A. 0.28

B. 0.26

C. 0.24

D. None of these

Answer: A



[Watch Video Solution](#)

7. One bag contains 5 white balls and 3 black balls and a second bag contains 2 white balls and 4 black balls. One ball drawn from the first bag and placed unseen in the second bag. What is the probability that a ball now drawn from second bag is black?

A. $\frac{29}{63}$

B. $\frac{5}{8}$

C. $\frac{7}{13}$

D. None of these

Answer: B



[Watch Video Solution](#)

8. The mean of getting tail two tosses of a coin is:

A. $\frac{1}{4}$

B. $\frac{1}{3}$

C. $\frac{1}{2}$

D. 1

Answer: D



[Watch Video Solution](#)

9. A coin is tossed 5 times. What is the probability of getting at least 3 heads.

A. $11/32$

B. $21/32$

C. $5/32$

D. None of these

Answer: A



[Watch Video Solution](#)

10. The probability of any ship return safely to the port is $1/5$. Find the probability that the returning out of 5 ships, at least 3 ships returns safely. It is given that returning ships are independently.

A. $256/625$

B. $\frac{369}{625}$

C. $\frac{184}{3125}$

D. $\frac{181}{3125}$

Answer: D



Watch Video Solution

Exercise 13 G Questions For Competitive Examinations

1. A lottery consists 3 prize tickets and 6 blank tickets. A choose three tickets from this lottery. In another lottery there are 1 prize ticket and 2 blank tickets. B choose one ticket from this lottery. Find the ratio in favour of A and B.

A. 7: 16

B. 1: 28

C. 16: 9

Answer: B



Watch Video Solution

2. Ten eggs are drawn successively with replacement from a lot containing 10% defective eggs. Find the probability that there is at least one defective egg.

A. $\left(\frac{9}{10}\right)^{10}$

B. $1 - \left(\frac{9}{10}\right)^{10}$

C. $\left(\frac{1}{10}\right)^{10}$

D. None of these

Answer: B



Watch Video Solution

3. Two integers are selected at random from integers 1 through 11. If the sum is even, find the probability that both the numbers are odd.

A. $\frac{1}{5}$

B. $\frac{2}{5}$

C. $\frac{3}{5}$

D. $\frac{4}{5}$

Answer: C



[Watch Video Solution](#)

4. A bag contains 50 tickets numbered 1, 2, 3, ..., 50 of which five are drawn at random and arranged in ascending order of magnitude x_1

A. $\frac{653}{15134}$

B. $\frac{949}{15134}$

C. $\frac{551}{15134}$

D. None of these

Answer: C



[Watch Video Solution](#)

5. A card from a pack of 52 cards is lost. From the remaining cards of the pack, two cards are drawn and are found to be both diamonds. Find the probability of the lost card being a diamond.

A. $\frac{11}{50}$

B. $\frac{17}{50}$

C. $\frac{33}{50}$

D. $\frac{39}{50}$

Answer: A



[Watch Video Solution](#)

6. A man is known to speak truth 3 out of 4 times. He throws a die and report that it is a 6. Find the probability that it is actually 6.

A. $\frac{1}{8}$

B. $\frac{7}{8}$

C. $\frac{5}{8}$

D. $\frac{3}{8}$

Answer: D



[Watch Video Solution](#)

7. Two cards are drawn from 52 playing cards without replacement. The variance of the number of ace is

A. $\frac{400}{2873}$

B. $\frac{600}{2873}$

C. $\frac{200}{2873}$

D. None of these

Answer: A



[Watch Video Solution](#)

8. Let A and B be two events such that $p(\bar{A} \cup B) = \frac{1}{6}$, $p(A \cap B) = \frac{1}{4}$ and $p(\bar{A}) = \frac{1}{4}$, where \bar{A} stands for the complement of the event A.

Then the events A and B are (1) mutually exclusive and independent (2) equally likely but not independent (3) independent but not equally likely (4) independent and equally likely

A. $1/4$

B. $1/3$

C. $1/2$

D. $1/6$

Answer: B



[Watch Video Solution](#)

9. 13. If A and B are two events such that $P(A) = \frac{1}{2}$ and $P(B) = \frac{2}{3}$, then

A. $1/6$

B. $1/3$

C. $2/3$

D. $1/2$

Answer: B



[Watch Video Solution](#)

10. A pair of unbiased die is thrown 3 times independently. The probability of getting the sum of number 9 two times is:

A. $5/243$

B. $7/243$

C. 8/243

D. 10/243

Answer: C



[Watch Video Solution](#)

Exercise 13 1

1. Given that E and F are events such that $P(E) = 0.6$, $P(F) = 0.6$, $P(F) = 0.3$ and $P(E \cap F) = 0.2$, find $P(E | F)$ and $P(F | E)$.



[Watch Video Solution](#)

2. Compute $P(A | B)$. if $P(B) = 0.5$ and $P(A \cap B) = 0.32$



[Watch Video Solution](#)

3. If $P(A) = 0.8$, $P(B) = 0.5$ and $P(B | A) = 0.4$, find (i) $P(A \cap B)$
(ii) $P(A | B)$ (iii) $P(A \cup B)$

 [Watch Video Solution](#)

4. Evaluate $P(A \cup B)$, if $2P(A) = P(B) = \frac{5}{13}$ and $P(A | B) = \frac{2}{5}$

 [Watch Video Solution](#)

5. If $P(A) = \frac{6}{11}$, $P(B) = \frac{5}{11}$ and $P(A \cup B) = \frac{7}{11}$, find (i) $P(A \cap B)$
(ii) $P(A | B)$ (iii) $P(B | A)$

 [Watch Video Solution](#)

6. Determine $P(E | F)$ in : A coin is tossed three times, where (i) E : Head on third toss, F : heads on first two tosses (ii) E : at least two heads, F : at most two heads (iii) E : at most two tails, F : at least one tail





[Watch Video Solution](#)

7. Determine $P(E | F)$ in : Two coins are tossed once, where (i) E: tail appears on one coin, F : one coin shows head (ii) E : no tail appears, F : no head appears



[Watch Video Solution](#)

8. Determine $P(E|F)$ in : A die is thrown three times, E : 4 appears on the third toss, F: 6 and 5 appears respectively on first two tosses



[Watch Video Solution](#)

9. Determine $P(E|F)$ in : Mother father and son line up at random for a family picture E: son on one end. F: father in middle



[Watch Video Solution](#)

10. A black and a red dice are rolled. (a) Find the conditional probability of obtaining a sum greater than 9. Given that the black die resulted in a 5. (b) Find the conditional probability of obtaining the sum 8? given that the red die resulted in a



[Watch Video Solution](#)

11. A fair die is rolled. Consider events $E = \{1, 3, 5\}$, $F = \{2, 3\}$ and $G = \{2, 3, 4, 5\}$ Find (i) $P(E | F)$ and $P(F | E)$ (ii) $P(E | G)$ and $P(G | E)$ (iii) $P((E \cup F) | G)$ and $P((E \cap F) | G)$



[Watch Video Solution](#)

12. Assume that each child born is equally likely to be a boy or a girl. If a family has two children, what is the conditional probability that both are girls given that i. the youngest is a girl, ii. at least one is a girl?



[Watch Video Solution](#)

13. An instructor has a question bank consisting of 300 easy True / False questions. 200 difficult True / False questions. 500 easy multiple choice questions and 400 difficult multiple choice questions. If a question is selected at random from the

 [Watch Video Solution](#)

14. Given that the two number appearing on throwing two dice are different. Find the probability of the event the sum of numbers on the dice is 4.

 [Watch Video Solution](#)

15. Consider the experiment of throwing a die. if a multiple of 3 comes tip. throw the die again and if any other number comes, toss a coin Find the conditional probability of the event the coin shows a tail, given that at least one die shows a 3.

 [Watch Video Solution](#)

16. In each of the Exercises choose the correct answer: If $P(A) = \frac{1}{2}$, $P(B) = 0$, then $P(A | B)$ is

A. 0

B. $\frac{1}{2}$

C. not defined

D. 1

Answer: C



[Watch Video Solution](#)

17. If A and B are two events such that $A \cap B \neq \phi$, $P\left(\frac{A}{B}\right) = P\left(\frac{B}{A}\right)$.

Then,

A. $A \subset B$ but $A \neq B$

B. $A = B$

C. $A \cap B = \phi$

D. $P(A) = P(B)$

Answer: A



[Watch Video Solution](#)

Exercise 13 2

1. If $P(A) = \frac{3}{5}$ and $P(B) = \frac{1}{5}$, find $P(A \cap B)$ if A and B are independent events.



[Watch Video Solution](#)

2. Two cards are drawn at random and without replacement from a pack of 52 playing cards. Find the probability that both the cards are black.



[Watch Video Solution](#)

3. A box of oranges is inspected by examining three randomly selected oranges drawn without replacement. If all the three oranges are good, the box is approved for sale, otherwise, it is rejected. Find the probability that a box containing 15 oranges

 [Watch Video Solution](#)

4. A fair coin and an unbiased die are tossed. Let A be the event head appears on the coin and B be the event 3 on the die. Check whether A and B are independent events or not.

 [Watch Video Solution](#)

5. A die marked 1, 2, 3 in red and 4, 5, 6 in green is tossed. Let A be the event, the number is even, and B be the event, the number is red. Are A and B independent?

 [Watch Video Solution](#)

6. Let E and F be events with $P(E) = \frac{3}{5}$, $P(F) = \frac{3}{10}$ and $P(E \cap F) = \frac{1}{5}$. Are E and F independent?

 [Watch Video Solution](#)

7. Given that the events A and B are such that $P(A) = \frac{1}{2}$, $P(A \cup B) = \frac{3}{5}$ and $P(B) = p$. Find p if they are i. mutually exclusive, ii. independent.

 [Watch Video Solution](#)

8. Let A and B be independent events with $P(A) = 0.3$ and $P(B) = 0.4$. Find (i) $P(A \cap B)$ (ii) $P(A \cup B)$ (iii) $P(A | B)$ (iv) $P(B | A)$

 [Watch Video Solution](#)

9. If A and B are two events such that $P(A) = \frac{1}{4}$, $P(B) = \frac{1}{2}$ and $P(A \cap B) = \frac{1}{8}$, find P (not A and not B).

 [Watch Video Solution](#)

10. If A and B are two events such that $P(A) = \frac{1}{2}$, $P(B) = \frac{7}{12}$ and $P(\text{not A or not B}) = \frac{1}{4}$. State whether A and B are independent?

 [Watch Video Solution](#)

11. Given two independent events A and B such that $P(A) = 0.3$, $P(B) = 0.6$. Find (i) $P(A \text{ and } B)$ (ii) $P(A \text{ and not } B)$ (iii) $P(A \text{ or } B)$ (iv) $P(\text{neither } A \text{ nor } B)$

 [Watch Video Solution](#)

12. A die is tossed thrice. Find the probability of getting an odd number at least once.

 [Watch Video Solution](#)

13. Two balls are drawn at random with replacement from a box containing 10 black and 8 red balls. Find the probability that i. both balls are red, ii. first ball is black and second is red, iii. One of them is black and other is red.

 [Watch Video Solution](#)

14. Probability of solving specific problem independently by A and B are $\frac{1}{2}$ and $\frac{1}{3}$ respectively. If both try to solve the problem independently find the probability that (i) the problem is solved (ii) exactly one of them solves the problem.

 [Watch Video Solution](#)

15. One card is drawn at random from a well shuffled deck of 52 cards. In which of the following cases are the events E and F independent? (i) E : the card drawn is a spade F : the card drawn is an ace (ii) E : the card drawn is black F : the card dr



[Watch Video Solution](#)

16. In a hostel 60% of the students read Hindi news paper, 40% read English news paper and 20% read both Hindi and English news papers. A student is selected at random. (a) Find the probability that she reads neither Hindi nor English news papers. (b



[Watch Video Solution](#)

17. The probability of obtaining an even prime number on each die, when a pair of dice is rolled is

A. 0

B. $\frac{1}{3}$

C. $\frac{1}{12}$

D. $\frac{1}{36}$

Answer: A



Watch Video Solution

18. Two events A and B will be independent, if (A) A and B are mutually exclusive (B) $P(A' B') = [1 - P(A)][1 - P(B)]$ (C) $P(A) = P(B)$ (D) $P(A) + P(B) = 1$

A. A and B are mutually exclusive

B. $P(A' B') = [1 - P(A)][1 - P(B)]$

C. $P(A) = P(B)$

D. $P(A) + P(B) = 1$

Answer: A



[Watch Video Solution](#)

Exercise 13 3

1. An urn contains 5 red and 5 black balls. A ball is drawn at random, its colour is noted and is returned to the urn. Moreover, 2 additional balls of the colour drawn are put in the urn and then a ball is drawn at random.

What is the probability that



[Watch Video Solution](#)

2. A bag contains 4 red and 4 black balls, another bag contains 2 red and 6 black balls. One of the two bags is selected at random and a ball is drawn from the bag which is found to be red. Find the probability that the ball is drawn from the first



[Watch Video Solution](#)

3. Of the students in a college, it is known that 60% reside in hostel and 40% are day scholars (not residing in hostel). Previous year results report that 30% of all students who reside in hostel attain A grade and 20% of day scholars attain A grad



[Watch Video Solution](#)

4. In answering a question on a multiple choice test, a student either knows the answer or guesses. Let $\frac{3}{4}$ be the probability that he knows the answer and $\frac{1}{4}$ be the probability that he guesses. Assuming that a student who guesses at the ans



[Watch Video Solution](#)

5. A laboratory blood test is 99% effective in detecting a certain disease when it is in fact, present. However, the test also yields a false positive result for 0.5% of the healthy person tested (i.e. if a healthy person is tested, then, with proba



[Watch Video Solution](#)

6. There are three coins. One is a two headed coin (having head on both faces), another is a biased coin that comes up heads 75% of the time and third is an unbiased coin. One of the three coins is chosen at random and tossed, it shows heads, what i



[Watch Video Solution](#)

7. An insurance company insured 2000 scooter drivers, 4000 car drivers and 6000 truck drivers. The probabilities of an accident for them are 0.01, 0.03 and 0.15 respectively. One of the insured persons meets with an accident. What is the probability that he is a scooter driver or a car driver?



[Watch Video Solution](#)

8. A factory has two machines A and B. Past record shows that machine A produced 60% of the items of output and machine B produced 40% of the items. Further, 2% of the items produced by machine A and 1% produced by machine B were defective. All the

 [Watch Video Solution](#)

9. Two groups are competing for the position on the Board of directors of a corporation. The probabilities that the first and the second groups will win are 0.6 and 0.4 respectively. Further, if the first group wins, the probability of introducing a

 [Watch Video Solution](#)

10. Suppose a girl throws a die. If she gets a 5 or 6, she tosses a coin three times and notes the number of heads. If she gets 1, 2, 3 or 4, she tosses a coin once and notes whether a head or tail is obtained. If she obtained exactly one head, what



Watch Video Solution

11. A manufacturer has three machine operators A , B and C . The first operator A produces 1% defective items, where as the other two operators B and C produce 5% and 7% defective items respectively. A is on the job for 50% of the tune, B is on the job for 30% of the time and C is on the job for 20% of the time. A defective item is produced, what is the probability that it was produced by A ?

A. $\frac{5}{36}$

B. $\frac{5}{38}$

C. $\frac{7}{34}$

D. $\frac{5}{34}$

Answer: D



Watch Video Solution

12. A card from a pack of 52 cards is lost. From the remaining cards of the pack, two cards are drawn and are found to be both diamonds. Find the probability of the lost card being a diamond.

A. $\frac{11}{50}$

B. $\frac{13}{50}$

C. $\frac{17}{50}$

D. $\frac{21}{50}$

Answer: A



[Watch Video Solution](#)

13. Probability that A speaks truth is $\frac{4}{5}$. A coin is tossed. A reports that a appears. The probability that actually there was head is

A. $\frac{4}{5}$

B. $\frac{1}{2}$

C. $\frac{1}{5}$

D. $\frac{2}{5}$

Answer: A



Watch Video Solution

14. If A and B are two events such that $A \subset B$ and $P(B) \neq 0$ when of the following is correct?

A. $P(A/B) = \frac{P(B)}{P(A)}$

B. $P(A/B) < P(A)$

C. $P(A/B) \geq P(A)$

D. None of these

Answer: C



Watch Video Solution

Exercise 13 4

1. State which of the following are not the probability distributions of a random variable. Give reasons for your answer.



[Watch Video Solution](#)

2. An urn contains 5 red and 2 black balls. Two balls are randomly drawn. Let X represent the number of black balls. What are the possible values of X ? Is X a random variable?



[Watch Video Solution](#)

3. Let X represent the difference between the number of heads and the number of tails obtained when a coin is tossed 6 times. What are possible values of X ?



[Watch Video Solution](#)

4. Find the probability distribution of (i) number of heads in two tosses of a coin. (ii) number of tails in the simultaneous tosses of three coins. (iii) number of heads in four tosses of a coin.



[Watch Video Solution](#)

5. Find the probability distribution of the number of successes in two tosses of a die, where a success is defined as (i) number greater than 4 (ii) six appears on at least one die



[Watch Video Solution](#)

6. From a lot of 30 bulbs which include 6 defectives, a sample of 4 bulbs is drawn at random with replacement. Find the probability distribution of the number of defective bulbs.



[Watch Video Solution](#)

7. A coin is biased so that the head is 3 times as likely to occur as tail. If the coin is tossed twice, find the probability distribution of number of tails.



[Watch Video Solution](#)

8. A random variable X has the following probability distribution:

X	0	1	2	3	4	5	6	7
$P(X)$	0	k	$2k$	$2k$	$3k$	k^2	$2k^2$	$7k^2 + k$

Determine :

- (i) k
- (ii) $P(X < 3)$
- (iii) $P(X > 6)$
- (iv) $P(0 < X < 3)$



[Watch Video Solution](#)

9. The random variable X has a probability distribution $P(X)$ of the following form, where k is some number : $P(X) = \{k, \text{ if } x = 02k, \text{ if } x = 13k, \text{ if } x = 20, \text{ otherwise}$ (a) Determine the value of k . (b) Find

 [Watch Video Solution](#)

10. Find the mean number of heads in three tosses of a fair coin.

 [Watch Video Solution](#)

11. Two dice are thrown simultaneously. If X denotes the number of sixes, find the expectation of X .

 [Watch Video Solution](#)

12. Two numbers are selected at random (without replacement) from the first six positive integers. Let X denote the larger of the two numbers obtained. Find $E(X)$.



[Watch Video Solution](#)

13. Let X denote the sum of the numbers obtained when two fair dice are rolled. Find the variance and standard deviation of X .



[Watch Video Solution](#)

14. A class has 15 students whose ages are 14, 17, 15, 14, 21, 17, 19, 20, 16, 18, 20, 17, 16, 19 and 20 years. One student is selected in such a manner that each has the same chance of being chosen and the age X of the selected student is recorded.



[Watch Video Solution](#)

15. In a meeting, 70% of the members favour and 30% oppose a certain proposal. A member is selected at random and we take $X = 0$ if he opposed, and $X = 1$ if he is in favour. Find $E(X)$ and $\text{Var}(X)$.

 [Watch Video Solution](#)

16. The mean of the numbers obtained on throwing a die having written 1 on three faces, 2 on two faces and 5 on one face is

A. 1

B. 2

C. 5

D. $\frac{8}{5}$

Answer: B

 [Watch Video Solution](#)

17. Suppose that two cards are drawn at random from a deck of cards. Let X be the number of aces obtained. Then the value of $E(X)$ is

A. $\frac{37}{221}$

B. $\frac{5}{13}$

C. $\frac{1}{13}$

D. $\frac{2}{13}$

Answer: D



[Watch Video Solution](#)

Exercise 13 5

1. A die is thrown 6 times. If "getting an odd number" is a success, what is the probability of (i) 5 successes? (ii) at least 5 successes? (iii) at most 5 successes?



[Watch Video Solution](#)

2. A pair of dice is thrown 4 times. If getting a doublet is considered a success, find the probability of two successes.



[Watch Video Solution](#)

3. There are 5% defective items in a large bulk of items. What is the probability that a sample of 10 items will include not more than one defective item?



[Watch Video Solution](#)

4. Five cards are drawn successively with replacement from a well-shuffled deck of 52 cards. What is the probability that (i) all the five cards are spades? (ii) only 3 cards are spades? (iii) none is a spade?



[Watch Video Solution](#)

5. The probability that a bulb produced by a factory will fuse after 150 days of use is 0.05. Find the probability that out of 5 such bulbs (i) none (ii) not more than one (iii) more than one (iv) at least one will fuse after 150 days of use.

 [Watch Video Solution](#)

6. A bag consists of 10 balls each marked with one of the digits 0 to 9. If four balls are drawn successively with replacement from the bag, what is the probability that none is marked with the digit 0?

 [Watch Video Solution](#)

7. In an examination, 20 questions of true-false type are asked. Suppose a student tosses a fair coin to determine his answer to each question. If the coin falls heads, he answers "true"; if it falls tails, he answers "false". Find the probability that

 [Watch Video Solution](#)

8. Suppose X has a binomial distribution $B\left(6, \frac{1}{2}\right)$. Show that $X = 3$ is the most likely outcome. (Hint: $P(x = 3)$ is the maximum among all $P(x_i)$, $x_i = 0, 1, 2, 3, 4, 5, 6$)

 [Watch Video Solution](#)

9. On a multiple choice examination with three possible answers (out of which only one is correct) for each of the five questions, what is the probability that a candidate would get four or more correct answers just by guessing?

 [Watch Video Solution](#)

10. A person buys a lottery ticket in 50 lotteries, in each of which his chance of winning a prize is $\frac{1}{100}$. What is the probability that he will win a prize (a) at least once (b) exactly once (c) at least twice?

 [Watch Video Solution](#)



[Watch Video Solution](#)

11. Find the probability of getting 5 exactly twice in 7 throws of a die.



[Watch Video Solution](#)

12. Find the probability of throwing at most 2 sixes in 6 throws of a single die.



[Watch Video Solution](#)

13. It is known that 10% of certain articles manufactured are defective. What is the probability that in a random sample of 12 such articles, 9 are defective?



[Watch Video Solution](#)

14. In a box containing 100 bulbs, 10 are defective. The probability that out of a sample of 5 bulbs, none is defective is

A. 10^{-1}

B. $\left(\frac{1}{2}\right)^5$

C. $\left(\frac{9}{10}\right)^5$

D. $\frac{9}{10}$

Answer: C



[Watch Video Solution](#)

15. The probability that a student is not a swimmer is $\frac{1}{5}$. Then the probability that out of five students, four are swimmers is

A. ${}^5C_4 \left(\frac{4}{5}\right)^4 \frac{1}{5}$

B. $\left(\frac{4}{5}\right)^4 \frac{1}{5}$

C. ${}^5C_3 \frac{1}{5} \left(\frac{4}{5}\right)^4$

D. None of these

Answer: A

 [Watch Video Solution](#)

Miscellaneous Exercise

1. A and B are two events such that $P(A) \neq 0$. Find $P(B | A)$, if (i) A is a subset of B (ii) $A \cap B = \varnothing$

 [Watch Video Solution](#)

2. A couple has two children. Find the probability that both the children are (i) males, if it is known that at least one of the children is male. (ii) females, if it is known that the elder child is a female.

 [Watch Video Solution](#)

3. Suppose that 5% of men and 0.25% of women have grey hair. A grey haired person is selected at random. What is the probability of these person being male? Assume that there are equal number of males and females.

 [Watch Video Solution](#)

4. Suppose that 90% of people are right-handed. What is the probability that at most 6 of a random sample of 10 people are right-handed?

 [Watch Video Solution](#)

5. An urn contains 25 balls of which 10 bear a mark 'X' and the remaining 15 bear a mark 'Y'. A ball is drawn at random from the urn, its mark noted down and it is replaced. If 6 balls are drawn in this way, find the probability that (i) all will bear mark 'X', (ii) not more than two will bear 'Y' mark, (iii) at least one ball will bear 'X' k and 'Y' mark will be equal

 [Watch Video Solution](#)

6. In a hurdle race, a player has to cross 10 hurdles. The probability that he will clear each hurdle is $\frac{5}{6}$. What is the probability that he will knock down fewer than 2 hurdles?

 [Watch Video Solution](#)

7. A die is thrown again and again until three sixes are obtained. Find the probability of obtaining the third six in the sixth throw of the die.

 [Watch Video Solution](#)

8. If a leap year is selected at random, what is the chance that it will contain 53 Tuesdays?

 [Watch Video Solution](#)

9. An experiment succeeds twice as often as it fails. Find the probability that in the next six trials, there will be atleast 4 successes.

 [Watch Video Solution](#)

10. How many times must a man toss a fair coin so that the probability of having at least one head is more than 90%?

 [Watch Video Solution](#)

11. In a game, a man wins a rupee for a six and loses a rupee for any other number when a fair die is thrown. The man decided to throw a die thrice but to quit as and when he gets a six. Find the expected value of the amount he wins / loses.

 [Watch Video Solution](#)

12. Suppose we have four boxes A,B,C and D containing coloured marbles as given below : One of the boxes has been selected at random and a single marble is drawn from it. If the marble is red, what is the probability that it was drawn from box A? bo



[Watch Video Solution](#)

13. Assume that the chances of a patient having a heart attack is 40%. It is also assumed that a meditation and yoga course reduce the risk of heart attack by 30% and prescription of certain drug reduces its chances by 25%. At a time a patient can ch



[Watch Video Solution](#)

14. If each element of a second order determinant is either zero or one, what is the probability that the value of the determinant is positive? (Assume that the individual entries of the determinant are chosen independently, each value being assumed



[Watch Video Solution](#)

15. An electronic assembly consists of two subsystems, say, A and B. From previous testing procedures, the following probabilities are assumed to be known: $P(A \text{ fails}) = 0.2$ $P(B \text{ fails alone}) = 0.15$ $P(A \text{ and } B \text{ fail}) = 0.15$
Evaluate the following probability



[Watch Video Solution](#)

16. Bag I contains 3 red and 4 black balls and Bag II contains 4 red and 5 black balls. One ball is transferred from Bag I to Bag II and then a ball is drawn from Bag II. The ball so drawn is found to be red in colour. Find the probability that the t



[Watch Video Solution](#)

17. If A and B are two events such that $P(A) \neq 0$ and $P(B | A) = 1$; then (A) $A \subset B$ (B) $B \subset A$ (C) $B = \varnothing$ (D) $A = \varnothing$

A. $A \subset B$ but $A \neq B$

B. $B \subset A$

C. $B = \phi$

D. $A = \phi$

Answer: A



Watch Video Solution

18. If $P(A|B) > P(A)$, then which of the following is correct: (A)

$P(B | A) < P(B)$ (B) $P(A \cap B) > P(A)P(B)$ (C) $P(B | A) = P(B)$ (D) $P(A \cap B) = P(A)P(B)$

A. $P(B/A) < P(B)$

B. $P(A \cap B) < P(A) \cdot P(B)$

C. $P(B/A) > P(B)$

D. $P(B/A) = P(B)$

Answer: A

 [Watch Video Solution](#)

19. If A and B are any two events such that $P(A) + P(B) - P(A \text{ and } B) = P(A)$, then:

A. $P(B/A) = 1$

B. $P(A/B) = 1$

C. $P(B/A) = 0$

D. $P(A/B) = 0$

Answer: A

 [Watch Video Solution](#)