



MATHS

BOOKS - DHANPAT RAI & CO MATHS (HINGLISH)

PAIR OF STRAIGHT LINES

Illustration

1. If the equation $2x^2 + \lambda xy + 2y^2 = 0$ represents a pair of real and distinct lines, then

A. $\lambda \in (-4, 4)$

B. $\lambda \in R$

C. $\lambda \in (-\infty, -4) \cup (4, \infty)$

D. $\lambda = 4, -4$

Answer: C



Watch Video Solution

2. If the pair of lines represented by $ax^2 + 2hxy + by^2 = 0, b \neq 0$, are such that the sum of the slopes of the lines is three times the product of their slopes, then

A. $3b + 2h = 0$

B. $3a + 2h = 0$

C. $2a + 3h = 0$

D. none of these

Answer: B



Watch Video Solution

3. If two lines represented by $x^2(\tan^2 \theta + \cos^2 \theta) - 2xy \tan \theta + y^2 \sin^2 \theta = 0$ make angles α, β with x-axis then

A. 4

B. 3

C. 2

D. none of these

Answer: C



Watch Video Solution

4. If the sum of the slopes of the lines given by $4x^2 + 2kxy - 7y^2 = 0$ is equal to the product of the slopes, then $k =$

A. -4

B. 4

C. -2

D. 2

Answer: C



[Watch Video Solution](#)

5. If the sum of the slopes of the lines given by $x^2 + 2cxy - y^2 = 0$ is four times their product, then c has the

value

A. -2

B. -1

C. 2

D. 1

Answer: A



Watch Video Solution

6. If the slopes of the lines given by $ax^2 + 2hxy + by^2 = 0$ are in the ratio $3:1$, then $h^2 =$

A. $\frac{ab}{3}$

B. $\frac{4ab}{3}$

C. $\frac{4a}{3b}$

D. none of these

Answer: B



Watch Video Solution

7. If the slope of one line in the pair $ax^2 + 4xy + y^2 = 0$ is three times the other, then a =

A. 3

B. 1

C. -3

D. -1

Answer: A



Watch Video Solution

8. The combined equation of the pair of lines through the origin and perpendicular to the pair of lines given by $ax^2 + 2hxy + by^2 = 0$, is

A. $ax^2 - 2hxy + by^2 = 0$

B. $bx^2 + 2hxy + ay^2 = 0$

C. $bx^2 - 2hxy + ay^2 = 0$

D. $bx^2 + 2hxy - ay^2 = 0$

Answer: C



Watch Video Solution

9. Equation of pair of straight lines drawn through (1, 1) and perpendicular to the pair of lines $3x^2 - 7xy + 2y^2 = 0$, is

A. $2x^2 + 7xy - 11x + 6 = 0$

B. $2(x - 1)^2 + 7(x - 1)(y - 1) - 3y^2 = 0$

C. $2(x - 1)^2 + 7(x - 1)(y - 1) - 3(y - 1)^2 = 0$

D. none of these

Answer: D



Watch Video Solution

10. The equation to the pair of lines perpendicular to the pair of lines $3x^2 - 4xy + y^2 = 0$, is

A. $x^2 + 4xy + 3y^2 = 0$

B. $x^2 - 4xy - 3y^2 = 0$

C. $x^2 + 4xy + y^2 = 0$

D. none of these

Answer: A



Watch Video Solution

11. If the slope of one of the lines given by $ax^2 + 2hxy + by^2 = 0$ is 5 times the other, then

A. $5h^2 = ab$

B. $5h^2 = 9ab$

C. $9h^2 = 5ab$

D. $h^2 = ab$

Answer: B



Watch Video Solution

12. If one of the lines represented by the equation $ax^2 + 2hxy + by^2 = 0$ is coincident with one of the lines represented by $a'x^2 + 2h'xy + b'y^2 = 0$, then

- A. $(h'b - hb')(ha' - h'a)$
- B. $4(h'b - hb')(ha' - h'a)$
- C. $2(h'b - hb')(ha' - h'a)$
- D. $4(h'b + hb')(ha' + h'a)$

Answer: B



Watch Video Solution

13. If one of the lines represented by the equation $ax^2 + 2hxy + by^2 = 0$ is coincident with one of the lines represented by $a'x^2 + 2h'xy + b'y^2 = 0$, then

- A. $4(ha' + h'a)(hb' + h'b)$
- B. $-4(ha' + h'a)(hb' + h'b)$
- C. $-4(ha' + h'a)(hb' + h'b)$
- D. $-4(ha' - h'a)(hb - h'b)$

Answer: B



Watch Video Solution

14. If the slope of one of the lines represented by $ax^2 + 2hxy + by^2 = 0$ be the square of the other, then

$$\frac{a+b}{h} + \frac{8h^2}{ab} =$$

A. 2

B. 4

C. 6

D. 1

Answer: C



Watch Video Solution

15. The image of the pair of lines represented by

$ax^2 + 2hxy + by^2 = 0$ by the line mirror $y = 0$ is: (A)

$ax^2 - 2hxy + by^2 = 0$ (b) $bx^2 - 2hxy + ay^2 = 0$ (c)

$bx^2 + 2hxy + ay^2 = 0$ (d) $ax^2 - 2hxy - by^2 = 0$

A. $ax^2 - 2hxy + by^2 = 0$

B. $bx^2 + 2hxy + ay^2 = 0$

C. $bx^2 - 2hxy + ay^2 = 0$

D. none of these

Answer: A



Watch Video Solution

16. Difference of slopes of the lines represented by the equation $x^2(\sec^2 \theta - \sin^2 \theta) - 2xy \tan \theta + y^2 \sin^2 \theta = 0$ is

(A) 4

(B) 3

(C) 2

(D) None of these

A. 4

B. 2

C. -4

D. 6

Answer: B



Watch Video Solution

17. The product of perpendiculars let fall from the point (x_1, y_1) upon the lines represented by $ax^2 + 2hxy + by^2$, is

A. $\frac{|ax_1^2 + 2hx_1y_1 + by_1^2|}{\sqrt{(a-b)^2 + 4h^2}}$

B. $\frac{|ax_1^2 + 2hx_1y_1 + by_1^2|}{\sqrt{(a-b)^2 + h^2}}$

$$C. \frac{|ax_1^2 + 2hx_1y_1 + by_1^2|}{\sqrt{(a+b)^2 + 4h^2}}$$

$$D. \frac{|ax_1^2 - 2hx_1y_1 + by_1^2|}{\sqrt{(a-b)^2 + 4h^2}}$$

Answer: A



Watch Video Solution

18. The angle between the pair of straight lines $y^2 \sin^2 \theta - xy \sin \theta + x^2 (\cos^2 \theta - 1) = 0$ is

A. $\frac{\pi}{3}$

B. $\frac{\pi}{4}$

C. $\frac{2\pi}{3}$

D. none of these

Answer: D



Watch Video Solution

19. The two lines represented by $3ax^2 + 5xy + (a^2 - 2)y^2 = 0$ are perpendicular to each other for two values of a (b) a for one value of a (d) for no values of a

- A. two values of a
- B. for one value of a
- C. for one value of a
- D. for no value of a

Answer: A



Watch Video Solution

20. The angle between the lines $2x^2 - 7xy + 3y^2 = 0$ is

A. 60°

B. 45°

C. $\tan^{-1}\left(\frac{7}{6}\right)$

D. 30°

Answer: B



Watch Video Solution

21. Find the angle between the lines represented by

$$x^2 + 2xy \sec \theta + y^2 = 0$$

A. 4θ

B. 2θ

C. θ

D. none of these

Answer: C



Watch Video Solution

22. If the angle θ is acute, then the acute angle between

$$x^2(\cos \theta - \sin \theta) + 2xy \cos \theta + y^2(\cos \theta + \sin \theta) = 0, \text{ is}$$

A. 2θ

B. $\theta/3$

C. θ

D. $\theta/2$

Answer: C



Watch Video Solution

23. If θ is the acute angle between the lines given by

$$x^2 - 2pxy + y^2 = 0, \text{ then}$$

A. $\cos \theta = p$

B. $\tan \theta = p$

C. $\sec \theta = p$

D. $\cot \theta = p$

Answer: C



Watch Video Solution

24. If the coordinate axes are the bisectors of the angles between the pair of lines $ax^2 + 2hxy + by^2 = 0$, then

A. $a + b = 0$

B. $h = 0$

C. $h \neq 0, a + b = 0$

D. $a + b \neq 0$

Answer: B



Watch Video Solution

25. If $x^2 - 2pxy - y^2 = 0$ and $x^2 - 2qxy - y^2 = 0$ bisect angles between each other, then find the condition.

A. $p + q = 1$

B. $pq = 1$

C. $pq + 1$

D. $p^2 + pq + q^2 = 0$

Answer: C



Watch Video Solution

26. If the equation

$$2x^2 + 7xy + 3y^2 - 9x - 7y + k = 0$$

represents a pair of lines, then k is equal to

A. 4

B. 2

C. 1

D. -4

Answer: A



[Watch Video Solution](#)

27. The equation $x^3 - yx^2 + x - y = 0$ represents

A. a hyperboal

B. an ellipse

C. a pair of straight lines

D. a rectangular hyperbola

Answer: C



[Watch Video Solution](#)

28. Distance between the pair of lines represented by the equation $x^2 - 6xy + 9y^2 + 3x - 9y - 4 = 0$, is

A. $\frac{15}{\sqrt{10}}$

B. $\frac{1}{2}$

C. $\sqrt{\frac{5}{2}}$

D. $\frac{1}{\sqrt{10}}$

Answer: C



Watch Video Solution

29. The equation $x^2 - 3xy + \lambda y^2 + 3x - 5y + 2 = 0$ where λ is a real number, represents a pair of straight lines. If θ is the

angle between the lines, then $\cos ec^2\theta =$

A. 2

B. 0

C. 3

D. 1

Answer: B



Watch Video Solution

30. The equation $ax^2 + 2\sqrt{ab}xy + by^2 + 2gx + 2fy + c = 0$ represents a pair of parallel straight lines, if

A. $ag = bf^2$

B. $a^2g = b^2f$

$$C. bg^2 = af^2$$

$$D. b^2g = a^2f$$

Answer: C



Watch Video Solution

31. If the equation

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$$

represents two straight lines, then the product of the perpendicular from the origin on these straight lines, is

A. $\frac{c}{\sqrt{(a-b)^2 - 4h^2}}$

B. $\frac{c}{\sqrt{(a-b)^2 + 4h^2}}$

C. $\frac{c}{\sqrt{(a+b)^2 - 4h^2}}$

D. none of these

Answer: B



Watch Video Solution

32. If $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents two parallel straight lines, then

A. $h^2 = ab$

B. $bg^2 = af^2$

C. $hg = af$

D. all of these

Answer: D



Watch Video Solution

33. $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$

A. $2\sqrt{\frac{g^2 - ac}{a(a + b)}}$

B. $2\sqrt{\frac{g^2 - ac}{h^2 - a^2}}$

C. $2\sqrt{\frac{f^2 - bc}{h^2 + b^2}}$

D. all of these

Answer: D



Watch Video Solution

34. The equation of pair of lines joining origin to the points of intersection of $x^2 + y^2 = 9$ and $x + y = 3$

A. $x^2 + (3 - x)^2 = 9$

B. $xy = 0$

C. $(3 + y)^2 + y^2 = 9$

D. $(x - y)^2 = 9$

Answer: B



Watch Video Solution

35. Prove that the angle between the lines joining the origin to the points of intersection of straight line $y = 3x + 2$ with the curve $x^2 + 2xy + 3y^2 + 4x + 8y - 11 = 0$ is

A. $\frac{\sqrt{2}}{3}$

B. $\frac{2\sqrt{2}}{3}$

C. $\frac{2}{3}$

D. $2\sqrt{2}$

Answer: B



Watch Video Solution

36. The acute angle between lines joining origin and intersection points of line $\sqrt{3}x + y - 2 = 0$ and the circle $x^2 + y^2 = 4$ is

A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{2}$

Answer: C



Watch Video Solution

37. Prove that the straight lines joining the origin to the point of intersection of the straight line $hx + ky = 2hk$ and the curve $(x - k)^2 + (y - h)^2 = c^2$ are perpendicular to each other if $h^2 + k^2 = c^2$.

A. $h^2 + k^2 = c^2$

B. $h^2 + k^2 = 2c^2$

C. $h^2 - k^2 = c^2$

D. none of these

Answer: A



Watch Video Solution

38. If the lines joining the origin to the points of intersection of the line $y = mx + 2$ and the curve $x^2 + y^2 = 1$ are at right-angles, then

A. $m^2 = 1$

B. $m^2 = 2$

C. $m^2 = 7$

D. $2m^2 = 1$

Answer: C



[Watch Video Solution](#)

1. The equation $ax^2 + by^2 + cx + cy = 0$ represents a pair of straight lines if

A. $a + b = 0$

B. $a + c = 0$

C. $b + c = 0$

D. none of these

Answer: A



[Watch Video Solution](#)

2. The equation $3x^2 + 2hxy + 3y^2 = 0$

represents a pair of straight lines passing through the origin.

The two lines are

A. real and distinct if $h^2 > 3$

B. real and distinct if $h^2 > 9$

C. real and coincident if $h^2 = 3$

D. real and coincident if $h^2 > 3$

Answer: B



Watch Video Solution

3. If the pairs of straight lines $ax^2 + 2hxy - ay^2 = 0$ and $bx^2 + 2gxy - by^2 = 0$ be such that each bisects the angles between the other, then

A. $hg + ab = 0$

B. $ah + bg = 0$

C. $h^2 - ab = 0$

D. $ag + bh = 0$

Answer: A



Watch Video Solution

4. If $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents parallel straight lines, then

A. $hf = bg$

B. $h^2 = bc$

C. $a^2f = b^2g$

D. none of these

Answer: A



Watch Video Solution

5. If the equation $12x^2 + 7xy - py^2 - 18x + qy + 6 = 0$ represents a pair of perpendicular straight lines, then

A. $p = 12, q = -1$

B. $p = -12, q = 1$

C. $p = 12, q = 1$

D. $p = 1, q = 12$

Answer: C



Watch Video Solution

6. The equation $8x^2 + 8xy + 2y^2 + 26x + 13y + 15 = 0$ represents a pair of straight lines. The distance between them is

A. $\frac{7}{\sqrt{5}}$

B. $\frac{7}{2\sqrt{5}}$

C. $\sqrt{\frac{7}{5}}$

D. none of these

Answer: B



[Watch Video Solution](#)

7. 1. Let PQR be a right angled isosceles triangle, right angled at P (2,1). If the equation of the line QR is $2x + y - 3 = 0$, then the

equation representing the pair of (1999, 2M) lines P and PR is

A. $3x^2 - 3y^2 + 8xy + 20x + 10y + 25 = 0$

B. $3x^2 - 3y^2 + 8xy - 20x - 10y + 25 = 0$

C. $3x^2 - 3y^2 + 8xy + 10x + 15y + 20 = 0$

D. $3x^2 - 3y^2 - 8xy - 10x - 15y - 20 = 0$

Answer: B



Watch Video Solution

8. If the gradient of one of the lines given by

$x^2 + hxy + 2y^2 = 0$ is twice that of the other, then $h =$

A. ± 2

B. ± 3

C. ± 1

D. $\pm 3/2$

Answer: B



Watch Video Solution

9. The set of value of h for which the equation $4x^2 + hxy - 3y^2 = 0$ represents a pair of real and distinct lines is

A. \mathbb{R}

B. $(3, 4)$

C. $(-3, 4)$

D. $(4, \infty)$

Answer: A



Watch Video Solution

10. If one of the lines of $my^2 + (1 - m^2)xy - mx^2 = 0$ is a bisector of the angle between the lines $xy = 0$, then m is 1 (b) 2 (c) $-\frac{1}{2}$ (d) -1

A. 1

B. 2

C. $-1/2$

D. -2

Answer: A



Watch Video Solution

11. If one of the lines of the pair $ax^2 + 2hxy + by^2 = 0$ bisects the angle between positive direction of the axes, then a , b and h satisfy the relation.

A. $a + b = 2|h|$

B. $a + b = -2h$

C. $a - b = 2|h|$

D. $(a - b)^2 = 4h^2$

Answer: B



Watch Video Solution

12. The equation $kx^2 + 4xy + 5y^2 = 0$ represents two lines inclined at an angle π if k is

A. $5/4$

B. $4/5$

C. -45

D. none of these

Answer: B



Watch Video Solution

13. If the equation $x^2 + (\lambda + \mu)xy + \lambda\mu y^2 + x + \mu y = 0$ represents two parallel straight lines, then prove that $\lambda = \mu$.

A. $\lambda \mid \mu = 0$

B. $\lambda = 4\mu$

C. $\lambda - \mu$

D. none of these

Answer: C



Watch Video Solution

14. The gradient of one of the lines given by $ax^2 + 2hxy + by^2 = 0$ is twice that of the other, then

A. $h^2 = ab$

B. $h = a + b$

C. $8h^2 = 9ab$

D. $9h^2 = 8ab$

Answer: C



Watch Video Solution

15. The equation $x^3 + y^3 = 0$ represents

- A. three real straight lines
- B. three points
- C. the combined equation of a straight line and a circle
- D. none of these

Answer: D



[Watch Video Solution](#)

16. One bisector of the angle between the lines given by $a(x - 1)^2 + 2h(x - 1)y + by^2 = 0$ is $2x + y - 2 = 0$. The equation of the other bisector is

A. $x - 2y + 1 = 0$

B. $x - 2y - 2 = 0$

C. $x - 2y - 1 = 0$

D. none of these

Answer: C



Watch Video Solution

17. The equation $2x^2 - 3xy - py^2 + x + qy - 1 = 0$ represent two mutually perpendicular lines if

A. $p = 3, q = 2$

B. $p = 2, q = 3$

C. $p = -2, q = 3$

D. $p = 2, q = -9/2$.

Answer: B::D



Watch Video Solution

18. The product of the perpendiculars drawn from the point (1,2) to the pair of lines $x^2 + 4xy + y^2 = 0$ is

A. $9/4$

B. $3/4$

C. $9/16$

D. none of these

Answer: D



Watch Video Solution

19. The three lines whose combined equation is $y^3 - 4x^2y = 0$ form a triangle which is

- A. isosceles
- B. equilateral
- C. right angled
- D. none of these

Answer: D



[Watch Video Solution](#)

20. The angle between the pair of lines whose equation is $4x^2 + 10xy + my^2 + 5x + 10y = 0$, is

A. $\tan^{-1} \cdot \frac{3}{8}$

B. $\tan^{-1} \cdot \frac{3}{4}$

C. $\tan^{-1} \cdot \left(\frac{2\sqrt{25 - 4m}}{m + 4} \right), m \in R$

D. none of these

Answer: B



Watch Video Solution

21. If two of the three lines represented by $ax^3 + bx^2y + cxy^2 + dy^3 = 0$ may be at right angles then

A. $a^2 + ac + bd - d^2 = 0$

B. $a^2 + ac - bd + d^2 = 0$

C. $a^2 - ac + bd + d^2 = 0$

$$D. a^2 + ac + bd + d^2 = 0$$

Answer: D



Watch Video Solution

22. The orthocentre of the triangle formed by the pair of lines $2x^2 - xy - y^2 + x + 2y - 1 = 0$ and the line $x + y + 1 = 0$, is

A. $(-1, 0)$

B. $(0, 1)$

C. $(-1, 1)$

D. none of these

Answer: A



Watch Video Solution

23. If the distance of a point (x_1, y_1) from each of the two straight lines, which pass through the origin of coordinates, is δ , then the two lines are given by

A. $(x_1y - xy_1)^2 = \delta^2(x^2 + y^2)$

B. $(x_1y + xy_1)^2 = \delta^2(x^2 + y^2)$

C. $(x_1y - xy_1)^2 = \delta^2(x^2 - y^2)$

D. none of these

Answer: A



Watch Video Solution

24. The equation of two straight lines through the point (x_1, y_1) and perpendicular to the lines given by $ax^2 + 2hxy + by^2 = 0$, is

A. $b(x - x_1)^2 - 2h(x - x_1)(y - y_1) + a(y - y_1)^2 = 0$

B. $b(x - x_1)^2 + 2h(x - x_1)(y - y_1) + a(y - y_1)^2 = 0$

C. $a(x - x_1)^2 - 2h(x - x_1)(y - y_1) + b(y - y_1)^2 = 0$

D. none of these

Answer: A



[Watch Video Solution](#)

25. The equation of the straight lines through the point (x_1, y_1) and parallel to the lines given by

$$ax^2 + 2xy + ay^2 = 0, \text{ is}$$

A. $a(y - y_1)^2 + 2h(x - x_1)(y - y_1) + b(x - x_1)^2 = 0$

B. $a(y - y_1)^2 - 2h(x - x_1)(y - y_1) + b(x - x_1)^2 = 0$

C. $b(y - y_1)^2 + 2h(x - x_1)(y - y_1) + a(x - x_1)^2 = 0$

D. none of these

Answer: C



Watch Video Solution

26. The triangle formed by the lines whose combined equation

is $(y^2 - 4xy - x^2)(x + y - 1) = 0$ is

A. equilateral

B. right angled

C. isosceles

D. obtuse angled

Answer: B



Watch Video Solution

27. The combined equation of the pair of lines through the point $(1, 0)$ and perpendicular to the lines represented by $2x^2 - xy - y^2 = 0$, is

A. $2x^2 - xy - y^2 - x + y - 1 = 0$

B. $2y^2 + xy - x^2 + 2x - y - 1 = 0$

C. $2y^2 + xy - x^2 - x - xy + 2 = 0$

D. none of these

Answer: B



Watch Video Solution

28. The equation $x^3 + ax^2y + bxy^2 + y^3 = 0$ represents three straight lines, two of which are perpendicular, then the equation of the third line, is

A. $y = ax$

B. $y = bx$

C. $y = x$

D. $y = -x$

Answer: C



Watch Video Solution

29. The combined equation of the lines L_1 and L_2 is $2x^2 + 6xy + y^2 = 0$ and that lines L_3 and L_4 is $4x^2 + 18xy + y^2 = 0$. If the angle between L_1 and L_4 be α , then the angle between L_2 and L_3 will be

A. $\frac{\pi}{2} - \alpha$

B. 2α

C. $\frac{\pi}{4} + \alpha$

D. α

Answer: D



Watch Video Solution

30. The lines represented by $x^2 + 2\lambda xy + 2y^2 = 0$ and the lines represented by $1 + \lambda_x^2 - 8xy + y^2 = 0$ are equally inclined, then

A. λ is any real number

B. $\lambda > 2$

C. $\lambda = \pm 2$

D. $\lambda < -2$

Answer: C



Watch Video Solution

31. The equation $\lambda(x^3 - 3xy^2) + y^3 - 3x^2y = 0$ represent three straight lines passing through the origin such that

A. they are equally inclined to one another

B. two of which are at right angles

C. two of which are coincident

D. none of these

Answer: A



Watch Video Solution

32. If the equation $x^4 + bx^3y + cx^2y^2 + dxy^3 + ey^4 = 0$ represent two pairs of perpendicular lines, then

A. $b + d = 1$ and $e = -1$

B. $b + d = 0$ and $e = -1$

C. $b + d = 0$ and $e = 1$

D. none of these

Answer: B



Watch Video Solution

33. The equation $x^3 + x^2y - xy^2 - y^3 = 0$ represents three straight lines passing through the origin such that

A. two of them are coincident and two of them are perpendicular

B. two of them are coincident but not two are perpendicular

C. two of them are perpendicular but no two are coincident

D. none of these

Answer: A



Watch Video Solution

34. If one of the lines represented by the equation $\lambda y^2 + (1 - \lambda^2)xy - \lambda x^2 = 0$ is bisector of the angle between the lines $xy = 0$, then, $\lambda =$

A. 1, -1

B. 2, $-\frac{1}{2}$

C. 1, 2

D. -1, $-\frac{1}{2}$

Answer: A



Watch Video Solution

35. The equation $x^2 - 3xy + \lambda y^2 + 3x - 5y + 2 = 0$ where λ is a real number, represents a pair of straight lines. If θ is the angle between the lines, then $\cos ec^2\theta =$

A. 9

B. 10

C. 15

D. 26

Answer: B



[Watch Video Solution](#)

36. If the line $y = mx$ bisects the angle between the lines $ax^2 + 2hxy + by^2 = 0$ then m is a root of the quadratic

equation

A. $h(m^2 - 1) + m(b - 1) = 0$

B. $h(m^2 - 1) + m(a - b)$

C. $h(m^2 + 1) + m(a - b) = 0$

D. none of these

Answer: B



Watch Video Solution

37. If two pairs of straight lines having equations

$y^2 + xy - 12x^2 = 0$ and $ax^2 + 2hxy + by^2 = 0$ have one line

common, then a =

A. $-3(2h + 3b)$

B. $8(h - 2b)$

C. $2(b + h)$

D. $-3(b + h)$

Answer: A::B



Watch Video Solution

38. The point of intersection of the pair of straight lines given by $6x^2 + 5xy - 4y^2 + 7x + 13y - 2 = 0$, is

A. $(1, 1)$

B. $(1, -1)$

C. $(-1, 1)$

D. $(-1, -1)$

Answer: C



Watch Video Solution

39. The straight lines represented by

$x^2 + mxy - 2y^2 + 3y - 1 = 0$ meet at $\left(-\frac{1}{3}, \frac{2}{3}\right)$ (b)

$\left(-\frac{1}{3}, -\frac{2}{3}\right)$ $\left(\frac{1}{3}, \frac{2}{3}\right)$ (d) none of these

A. $-\frac{1}{3}, \frac{2}{3}$

B. $(-\frac{1}{3}, -\frac{2}{3})$

C. $(\frac{1}{3}, \frac{2}{3})$

D. none of these

Answer: A:C



Watch Video Solution

40. The square of the distance between the origin and the point of intersection of the lines given by

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0, \text{ is}$$

A. $\frac{c(a + b) + f^2 + g^2}{ab - h^2}$

B. $\frac{c(a + b) - f^2 - g^2}{h^2 - ab}$

C. $\frac{c(a + b) - f^2 - g^2}{ab - h^2}$

D. none of these

Answer: C



Watch Video Solution

41. The centroid of the triangle whose three sides are given by the combined equation $(x^2 + 7xy + 2y^2)(y - 1) = 0$, is

A. $(\frac{2}{3}, 0)$

B. $(\frac{7}{3}, \frac{2}{3})$

C. $(-\frac{7}{3}, \frac{2}{3})$

D. none of these

Answer: C



Watch Video Solution

42. If first degree terms and constant term are to be removed from the equation $12x^2 + 7xy - 12y^2 - 17x - 31y - 7 = 0$, then the origin must be shifted at shifted at the point .

A. $(1, -1)$

B. $(-1, 1)$

C. $(-1, -1)$

D. none of these

Answer: A



Watch Video Solution

43. The combined equation of three sides of a triangle is $(x^2 - y^2)(2x + 3y - 6) = 0$. If $(-2, a)$ is an interior point and $(b, 1)$ is an exterior point of the triangle, then

A. $a \in (2, 10/3)$

B. $a \in (2, 10/3)$

C. $b \in (-1, 9/2)$

D. $b \in (-1/1)$

Answer: A::D



Watch Video Solution

44. The angle between the straight lines joining the origin to the point of intersection of $3x^2 + 5xy - 3y^2 + 2x + 3y = 0$ and $3x - 2y = 1$, is

A. $\frac{\pi}{3}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{6}$

D. $\frac{\pi}{2}$

Answer: D



Watch Video Solution

45. All chords of the curve $3x^2 - y^2 - 2x + 4y = 0$ which subtend a right angle at the origin, pass through the fixed point

A. $(1, 2)$

B. $(-1, 2)$

C. $(1, -2)$

D. $(-1, -2)$

Answer: C



Watch Video Solution

46. If the pair of lines $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ intersect on the y-axis then

A. $2fgh = bg^2 + ch^2$

B. $bg^2 + ch^2 = fgh$

C. $abc = 2fgh$

D. none of these

Answer: A



Watch Video Solution

47. If the $y = mx + 1$, of the circle $x^2 + y^2 = 1$ subtends an angle of measure 45° of the major segment of the circle then value of m is -

A. ± 2

B. 1, 2

C. ± 1

D. none of these

Answer: C



Watch Video Solution

48. The straight lines represented by $(y - mx)^2 = a^2(1 + m^2)$ and $(y - nx)^2 = a^2(1 + n^2)$ form a rectangle (b) rhombus trapezium (d) none of these

A. rectangle

B. trapezium

C. rhombus

D. none of these

Answer: C



Watch Video Solution

49. The equation of the image of the pair of rays $y = |x|$ in the line mirror $x = 1$, is

A. $|y| = x + 2$

B. $|y| + 2 = x$

C. $y = |x - 2|$

D. none of these

Answer: C



Watch Video Solution

50. Two lines represented by the equation.

$x^2 - y^2 - 2x + 1 = 0$ are rotated about the point $(1, 0)$, the

line making the bigger angle with the position direction of the

x-axis being turned by 45° in the clockwise sense and the

other line being turned by 15° in the anti-clockwise sense. The

combined equation of the pair of lines in their new position is

A. $\sqrt{3}x^2 - xy + 2\sqrt{3}x - y + \sqrt{3} = 0$

B. $\sqrt{3}x^2 - xy - 2\sqrt{3}x + y + \sqrt{3} = 0$

C. $\sqrt{3}x^2 - xy - 2\sqrt{3}x + \sqrt{3} = 0$

D. none of these

Answer: B



Watch Video Solution

51. The value of λ for which the lines joining the point of intersection of curves C_1 and C_2 to the origin are equally inclined to the axis of x.

$$C_1: \lambda x^2 + 3y^2 - 2\lambda xy + 9x = 0, C_2: 3x^2 - 4y^2 + 8xy - 3x = 0$$

A. $\lambda = \frac{4}{3}$

B. $\lambda = 12$

C. $\lambda = 1$

D. none of these

Answer: B



Watch Video Solution

52. If one of the lines given by the equation $2x^2 + axy + 3y^2 = 0$ coincide with one of those given by $2x^2 + bxy - 3y^2 = 0$ and the other lines represented by them be perpendicular, then

A. $a = -5, b = 1$

B. $a = 5, b = -1$

C. $a = 5, b = 1$

D. none of these

Answer: C



Watch Video Solution

53. If the pair of lines $x^2 + 2xy + ay^2 = 0$ and $ax^2 + 2xy + y^2 = 0$ have exactly one line in common, then $a =$

A. 1

B. -3

C. -1

D. 3

Answer: B



[Watch Video Solution](#)

54. If one of the lines given by $6x^2 - xy + 4cy^2 = 0$ is $3x + 4y = 0$, then $c =$

A. -3

B. -1

C. 3

D. 1

Answer: A



Watch Video Solution

55. Area of the triangle formed by the line $x + y = 3$ and angle bisectors of the pair of straight lines $x^2 - y^2 + 2y = 1$ is *2squnits* b. *4squnits* c. *6squnits* d. *8squnits*

A. 2 sq. units

B. 4 sq. units

C. 6 sq. units

D. 8 sq. units

Answer: A



Watch Video Solution

56. If the pair of lines $ax^2 + 2hxy + by^2 = 0$ ($h^2 > ab$) forms an equilateral triangle with the line $lx + my + n = 0$ then $(a + 3b)(3a + b) =$

A. H^2

B. $-H^2$

C. $2H^2$

D. $4H^2$

Answer: D



Watch Video Solution

57. The area (in square units) of the quadrilateral formed by two pairs of lines $l^2x^2 - m^2y^2 - n(lx + my) = 0$ and $l^2x^2 - m^2y^2 + n(lx - my) = 0$, is

A. $\frac{n^2}{2|lm|}$

B. $\frac{n^2}{|m|}$

C. $\frac{n}{2|lm|}$

D. $\frac{n^2}{4|lm|}$

Answer: A



Watch Video Solution

58. The equation $4x^2 - 24xy + 11y^2 = 0$ represents

- A. two parallel lines
- B. two perpendicular lines
- C. two lines through the origin
- D. a circle

Answer: C



[Watch Video Solution](#)

59. If the pair of lines $ax^2 + 2(a + b)xy + by^2 = 0$ lie along diameters of a circle and divide the circle into four sectors such that the area of one of the sectors is thrice the area of another sector then

A. $3a^2 + 2ab + 3b^2 = 0$

B. $3a^2 + 10ab + 3b^2 = 0$

C. $3a^2 - 2ab + 3b^2 = 0$

D. $3a^2 - 10ab + 3b = 0$

Answer: A



Watch Video Solution

60. If the equation of the pair of straight lines passing through the point $(1, 1)$, one making an angle θ with the positive direction of the x-axis and the other making the same angle with the positive direction of the y-axis, is $x^2 - (a + 2)xy + y^2 + a(x + y - 1) = 0, a \neq 2$, then the value of $\sin 2\theta$ is $a - 2$ (b) $a + 2$ $2(a + 2)$ (d) $\frac{2}{a}$

A. $a - 2$

B. $a + 2$

C. $\frac{2}{a + 2}$

D. $\frac{2}{a}$

Answer: C

 [Watch Video Solution](#)

61. If θ_1 and θ_2 are the angles made by the lines $(x^2 + y^2)(\cos^2 \theta \sin^2 \theta + \sin^2 \theta) = (x \tan \alpha - y \sin \theta)^2$ with positive axis of x in the anti-clock wise direction, if $\theta = \frac{\pi}{6}$, $\tan \theta_1 + \tan \theta_2$ equals

A. $-\frac{8}{3} \sin^2 \alpha$

B. $-\frac{8}{2} \operatorname{cosec} 2\alpha$

C. $-8\sqrt{3}\operatorname{cosec} 2\alpha$

D. $-4\sec 2\alpha$

Answer: B



Watch Video Solution

62. If the product of the perpendiculars drawn from the point (1,1) on the lines $ax^2 + 2hxy + by^2 = 0$ is 1, then

A. $h(a - b) + ab = 0$

B. $h(a + b) - ab = 0$

C. $h(a + b) + ab = 0$

D. $h(a - b) - ab = 0$

Answer: C



Watch Video Solution

63. If the equation $2x^2 + 2hxy + 6y^2 - 4x + 5y - 6 = 0$ represents a pair of straight lines, then the length of intercept on the x-axis cut by the lines is equal to

A. 2

B. 4

C. $\sqrt{7}$

D. 0

Answer: B



Watch Video Solution

1. If $ax^3 + by^3 + cx^2y + dxy^2 = 0$, represents three distinct straight lines, such that each line bisects the angle between the other two, then which of the following is true:

A. $3b + c = 0$

B. $3a + d = 0$

C. $d + 3a = 0$

D. $b + 3c = 0$

Answer: A



[Watch Video Solution](#)

Section II Assertion Reason Type

1. Statement -1 : If $a > b > c$, then the lines represented by $(a - b)x^2 + (b - c)xy + (c - a)y^2 = 0$ are real and distinct.

Statement-2 : Pair of lines represented by $ax^2 + 2hxy + by^2 = 0$ are real and distinct if $h^2 > ab$.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1.

B. Statement-1 is True, Statement-2 is True, Statement-2 is not a correct explanation for Statement-1.

C. Statement-1 is True, Statement-2 is False.

D. Statement-1 is False, Statement-2 is True.

Answer: A



Watch Video Solution

2. Statement-1: All chords of the curve $3x^2 - y^2 - 2x + 4y = 0$ which subtend a right angle at the origin pass through a fixed point.

Statement-2: The equation $ax + by + c = 0$ represents a family of straight lines passing through a fixed point iff there is a linear relation between a, b and c.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1.
- B. Statement-1 is True, Statement-2 is True, Statement-2 is not a correct explanation for Statement-1.
- C. Statement-1 is True, Statement-2 is False.
- D. Statement-1 is False, Statement-2 is True.

Answer: A



Watch Video Solution

3. Statement-1: If a, b are non-zero real number such that $a + b = 2$, then $ax^2 + 2xy + by^2 + 2ax + 2by = 0$ represents a pair of straight lines. Statement-2: If a, b are of opposite signs, then $ax^2 + 2xt + by^2 = 0$ represents a pair of distinct lines passing through the origin.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1.
- B. Statement-1 is True, Statement-2 is True, Statement-2 is not a correct explanation for Statement-1.
- C. Statement-1 is True, Statement-2 is False.
- D. Statement-1 is False, Statement-2 is True.

Answer: B



Watch Video Solution

Exercise

1. For what value of k is $4x^2 + 8xy + ky^2 = 9$ the equation of a pair of straight lines?

A. 0

B. 4

C. 9

D. -9

Answer: B



Watch Video Solution

2. The angle between the lines represented by $x^2 - y^2 = 0$ is

A. 0°

B. 45°

C. 90°

D. 180°

Answer: C



[Watch Video Solution](#)

3. If the angle between the two lines represented by $2x^2 + 5xy + 3y^2 + 6x + 7y + 4 = 0$ is $\tan^{-1}(m)$, then m is equal to

A. $|1/5|$

B. -1

C. $-\frac{2}{3}$

D. none of these

Answer: A



Watch Video Solution

4. If one of the lines of $ax^2 + 2hxy + by^2 = 0$ bisects the angle between the axes, in the first quadrant, then

A. $h^2 - ab = 0$

B. $h^2 + ab = 0$

C. $(a + b)^2 = h^2$

D. $(a + b)^2 = 4h^2$

Answer: D



Watch Video Solution

5. The point of intersection of the two lines given by

$$2x^2 - 5xy + 2y^2 - 3x + 3y + 1 = 0 \text{ is}$$

A. $(1/2, 1/3)$

B. $(-1/7, -1/7)$

C. $(-1/3, 1/3)$

D. none of these

Answer: D



Watch Video Solution

6. The value of 'a' for which the lines represented by

$ax^2 + 5xy + 2y^2 = 0$ are mutually perpendicular is

A. 2

B. -2

C. $\frac{25}{8}$

D. none of these

Answer: B



[Watch Video Solution](#)

7. Joint equation of the diagonals of the square formed by the

pairs of lines $xy + 4x - 3y - 12 = 0$ and

$xy - 3x + 4y - 12 = 0$, is

A. $x^2 - y^2 + x - y = 0$

B. $x^2 - y^2 + x + y = 0$

C. $x^2 + 2xy + y^2 + x + y = 0$

D. $x^2 - 2xy + y^2 + x - y = 0$

Answer: A



Watch Video Solution

8. Which of the following pair of straight lines intersect at right angle?

A. $2x^2 = y(x + 2y)$

B. $(x + y)^2 = x(y + 3x)$

C. $2y(x + y) = xy$

D. $y = \pm 2x$

Answer: A



Watch Video Solution

9. The equation of the lines parallel to the line common to the pair of lines given by $6x^2 - xy - 12y^2 = 0$ and $15x^2 + 14xy - 8y^2 = 0$ and the sum of whose intercepts on the axes is 7, is

A. $2x - 3y = 42$

B. $3x + 4y = 12$

C. $5x - 2y = 10$

D. none of these

Answer: B



Watch Video Solution

10. The equation $x^2 + k_1y^2 + k_2xy = 0$ represents a pair of perpendicular lines if

A. $k_1 = -1$

B. $k_1 = 2k_2$

C. $2k_1 = k_2$

D. none of these

Answer: A



Watch Video Solution

11. If the equation $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents a pair of parallel lines, then

A. $\frac{a}{f} = \frac{b}{h} = \frac{f}{g}$

B. $\frac{a}{h} = \frac{h}{b} = \frac{f}{g}$

C. $\frac{a}{h} = \frac{h}{b} = \frac{g}{f}$

D. none of these

Answer: C



Watch Video Solution

12. If the slope of one of the lines given by $ax^2 - 6xy + y^2 = 0$ is twice the other, then a =

A. 1

B. 2

C. 4

D. 8

Answer: D



Watch Video Solution

13. If the equation $12x^2 + 7xy - py^2 - 18x + qy + 6 = 0$ represents a pair of perpendicular straight lines, then

A. $p = 12, q = 1$

B. $p = 1, q = 12$

C. $p = -1, q = 12$

D. $p = 1, q = -12$

Answer: A



Watch Video Solution

14. If θ is the angle between the lines given by the equation $6x^2 + 5xy - 4y^2 + 7x + 13y - 3 = 0$, then find the equation of the line passing through the point of intersection of these lines and making an angle θ with the positive x-axis.

A. $2x + 11y + 13 = 0$

B. $11x - 2y + 13 = 0$

C. $2x - 11y + 2 = 0$

D. $11x + 2y - 11 = 0$

Answer: B



Watch Video Solution

15. If the pair of straight lines $ax^2 + 2hxy + by^2 = 0$ is rotated about the origin through 90° , then their equations in the new position are given by

A. $ax^2 - 2hxy + by^2 = 0$

B. $ax^2 - 2hxy - by^2 = 0$

C. $bx^2 - 2hxy + ay^2 = 0$

D. $bx^2 + 2hxy + ay^2 = 0$

Answer: C



Watch Video Solution

16. If the lines represented by $x^2 - 2pxy - y^2 = 0$ are rotated about the origin through an



[View Text Solution](#)

17. The difference of the tangents of the angles which the lines $x^2(\sec^2 - \sin^2 \theta) - 2xy \tan \theta + y^2 \sin^2 \theta = 0$ make with X-axis, is

A. $2 \tan \theta$

B. 2

C. $2 \cot \theta$

D. $\sin 2\theta$

Answer: B



Watch Video Solution

18. If the two pairs of line $x^2 - 2mxy - y^2 = 0$ and $x^2 - 2nxy - y^2 = 0$ are such that one of them represent the bisector of the angles between the other, then: (A) $mn + 1 = 0$ (B) $mn - 1 = 0$ (C) $1/m + 1/n = 0$ (D) $1/m - 1/n = 0$

A. $mn + 1 = 0$

B. $mn - 1 = 0$

C. $\frac{1}{m} + \frac{1}{n} = 0$

D. $\frac{1}{m} - \frac{1}{n} = 0$

Answer: A



Watch Video Solution

19. If $\lambda x^2 - 10xy + 12y^2 + 5x - 16y - 3 = 0$, represents a pair of straight lines, then the value of λ is

A. 4

B. 3

C. 2

D. 1

Answer: C



[Watch Video Solution](#)

20. The equation $y^2 - x^2 + 2x - 1 = 0$, represents

A. a pair of st. lines

B. a circle

C. a parabola

D. an ellipse

Answer: A



Watch Video Solution

21. The angle between the straight lines $x^2 - y^2 - 2x - 1 = 0$

, is

A. 90°

B. 60°

C. 75°

D. 36°

Answer: A



Watch Video Solution

22. If the angle between the two lines represented by $2x^2 + 5xy + 3y^2 + 6x + 7y + 4 = 0$ is $\tan^{-1}(m)$, then m is equal to

A. $1/5$

B. 1

C. $7/5$

D. 7

Answer: A



Watch Video Solution

23. The diagonal of the rectangle formed by the lines $x^2 - 7x + 6 = 0$ and $y^2 - 14y + 40 = 0$ is

A. $5x + 6y = 0$

B. $5x - 6y = 0$

C. $6x - 5y + 14 = 0$

D. $6x - 5y - 14 = 0$

Answer: C



Watch Video Solution

24. The angle between the pair of straight lines $2x^2 + 5xy + 2y^2 + 3x + 3y + 1 = 0$ is

(A) $\cos^{-1}\left(\frac{4}{5}\right)$

(A) $\tan^{-1}\left(\frac{4}{5}\right)$

(A) $\frac{\pi}{2}$

(D) 0

A. $\cos^{-1}(4/5)$

B. $\tan^{-1}(4/5)$

C. 0

D. $\pi/2$

Answer: A



Watch Video Solution

25. The circumcentre of the triangle formed by the lines,

$xy + 2x + 2y + 4 = 0$ and $x + y + 2 = 0$ is-

A. $(0, 0)$

B. $(-2, -2)$

C. $(-1, -1)$

D. $(-1, -2)$

Answer: C



Watch Video Solution

26. Distance between the lines represented by

$9x^2 - 6xy + y^2 + 18x - 6y + 8 = 0$, is $\frac{1}{\sqrt{10}}$ 2. $\frac{2}{\sqrt{10}}$ 3. $\frac{4}{\sqrt{10}}$

4. $\sqrt{10}$ 5. $\frac{2}{\sqrt{5}}$

A. $\frac{1}{\sqrt{10}}$

B. $\frac{2}{\sqrt{10}}$

C. $\frac{4}{\sqrt{10}}$

D. $\sqrt{10}$

Answer: B



Watch Video Solution

27. The joint equation of the straight lines $x + y = 1$ and $x - y = 4$, is

A. $x^2 - y^2 = -4$

B. $x^2 - y^2 = 4$

C. $(x + y - 1)(x - y - 4) = 0$

D. $(x + y + 1)(x - y + 4) = 0$

Answer: C



Watch Video Solution

28. If the slope of one of the lines given by $ax^2 - 6xy + y^2 = 0$ is square of the other, then a =

A. 8, - 27

B. - 8, 27

C. 1, 8

D. - 8, - 27

Answer: A



Watch Video Solution

29. The value of k such that

$$3x^2 - 11xy + 10y^2 - 7x + 13y + k = 0$$

may represent a pair of straight lines, is

A. 3

B. 4

C. 6

D. 8

Answer: B



[Watch Video Solution](#)

30. If $x^2 - kxy + y^2 + 2y + 2 = 0$ denotes a pair of straight lines then $k =$

A. 2

B. $1/\sqrt{2}$

C. $2\sqrt{2}$

D. $\sqrt{2}$

Answer: D



Watch Video Solution

31. The equation of the lines which are parallel to the line common to the pair of the lines given by $6x^2 - xy - 12y^2 = 0$ and $15x^2 + 14xy - 8y^2 = 0$ and at a distance of 7 units from it are

A. $3x + 4y = \pm 35$

B. $5x - 2y = \pm 7$

C. $2x - 3y = \pm 7$

D. none of these

Answer: A



Watch Video Solution

32. If the slope of one of the lines given by $36x^2 + 2hxy + 72y^2 = 0$ is four times the other, then $h^2 =$

A. 5040

B. 4050

C. 8100

D. none of these

Answer: B



Watch Video Solution

33. The combined equation of the pair of the straight lines through the point $(1, 0)$ and parallel to the lines represented by

$$2x^2 - xy - y^2 = 0 \text{ is}$$

A. $2x^2 - xy - 2y^2 + 4x - y = 6$

B. $2x^2 - xy - y^2 - 4x + y + 2 = 0$

C. $2x^2 - xy - y^2 - 4x - y + 2 = 0$

D. none of these

Answer: C



Watch Video Solution

34. The equation $x^2 - 6x^2y + 11xy^2 - 6y^3 = 0$ represents three straight lines passing through the origin, the slopes of which form an

A. A.P.

B. G.P.

C. H.P.

D. none of these

Answer: C



[Watch Video Solution](#)

Chapter Test

1. If the lines given by $ax^2 + 2hxy + by^2 = 0$ are equally inclined to the lines given by $ax^2 + 2hxy + by^2 + \lambda(x^2 + y^2) = 0$, then

- A. λ is any real number
- B. $\lambda = 2$
- C. $\lambda = 1$
- D. none of these

Answer: A



[Watch Video Solution](#)

2. The equation to the straight lines passing through the origin and making an angle α with straight line $y + x = 0$ are

given by

A. $x^2 + 2xy \sec 2\alpha + y^2 = 0$

B. $x^2 - 2xy \sec 2\alpha + y^2 = 0$

C. $x^2 + 2xy \cos 2\alpha + y^2 = 0$

D. none of these

Answer: A



Watch Video Solution

3. The lines joining the origin to the point of intersection of $x^2 + y^2 + 2gx + c = 0$ and $x^2 + y^2 + 2fy - c = 0$ are at right angles if

A. $g^2 + f^2 = c$

B. $g^2 - f^2 = c$

C. $g^2 - f^2 = 2c$

D. $g^2 + f^2 = c^2$

Answer: C



Watch Video Solution

4. If the area of the triangle formed by the pair of lines

$8x^2 - 6xy + y^2 = 0$ and the line $2x + 3y = a$ is 7 then $a =$

A. 14

B. $14\sqrt{2}$

C. 28

D. none of these

Answer: C



Watch Video Solution

5. The lines bisecting the angle between the bisectors of the angles between the lines $ax^2 + 2hxy + by^2 = 0$ are given by

A. $(a - b)(x^2 - y^2) - 4hxy = 0$

B. $(a - b)(x^2 + y^2) + 4hxy = 0$

C. $(a - b)(x^2 - y^2) + 4hxy = 0$

D. none of these

Answer: C



Watch Video Solution

6. The pair of lines $\sqrt{3}x^2 - 4xy + \sqrt{3}y^2 = 0$ are rotated about the origin by $\pi/6$ in anticlockwise sense. The equation of the pair in the new position, is

A. $\sqrt{3}x^2 - xy = 0$

B. $x^2 - \sqrt{3}xy = 0$

C. $xy - \sqrt{3}y^2 = 0$

D. none of these

Answer: A



Watch Video Solution

7. If two of the lines given by the equation

$$ax^3 + bx^2y + cxy^2 + dy^3 = 0 (a \neq 0)$$

make complementary angles with x-axis in anticlockwise sense,

then

A. $(a - c) = d(b - d)$

B. $d(a - c) = a(d - b)$

C. $a(a - c) = d(d - b)$

D. none of these

Answer: C



Watch Video Solution

8. Two of the lines represented by the equation

$$ay^4 + bxy^3 + cx^2y^2 + dx^3y + ex^4 = 0$$
 will be perpendicular,

then

A. $(c + a + e)(e - a)^2 + (b + d)(ad + be) = 0$

B. $(c + a + e)(e - a)^2 - (b + d)(ad + be) = 0$

C. $(c + a + e)(e + a)^2 + (b + d)(ab + be) = 0$

D. none of these

Answer: A



Watch Video Solution

9. The pairs of straight lines $ax^2 - 2hxy - ay^2 = 0$ and $hx^2 - 2axy - gy^2 = 0$ are such that

A. one pair bisects the angle between the other pair

B. the lines of one pair are equally inclined to the lines of the other pair

C. the lines of each pair are perpendicular to other pair

D. all of these

Answer: D



Watch Video Solution

10. The equation $a^2x^2 + 2h(a + b)xy + b^2y^2 = 0$ and $ax^2 + 2hxy + by^2 = 0$ represent two pairs of perpendicular straight lines two pairs of parallel straight lines two pairs of straight lines which are equally inclined to each other none of these

A. two pairs of perpendicular straight lines

B. two pairs of parallel straight lines

- C. two pairs of straight lines which are equally inclined to each other
- D. none of these

Answer: C



Watch Video Solution

11. if $\frac{x^2}{a} + \frac{y^2}{b} + \frac{2xy}{h} = 0$ represent pair of straight lines and slope of one line is twice the other line then $ab:h^2$.

- A. 9: 8
- B. 8: 9
- C. 1: 2
- D. 2: 1

Answer: A



Watch Video Solution

12. If the lines represented by the equation $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ are equidistant from the origin, then

A. $f^4 - g^4 = c(bf^2 - ag^2)$

B. $f^4 - g^4 = c(af^2 - bg^2)$

C. $f^4 - g^4 = c(ag^2 - bf^2)$

D. none of these

Answer: A



Watch Video Solution

13. The distance between the two lines represented by the sides of an equilateral triangle a right-angled triangle an isosceles triangle none of these

A. $\frac{8}{5}$

B. $\frac{6}{5}$

C. $\frac{11}{5}$

D. none of these

Answer: A



[Watch Video Solution](#)

14. The equation of the image of the lines $y = |x|$ in the line mirror $x = 2$ is

A. $y = |x - 4|$

B. $|y| = x + 4$

C. $|y| + 4 = x$

D. none of these

Answer: A



Watch Video Solution

15. If the equation $3x^2 + xy - y^2 - 3x + 6y + k = 0$ represents a pair of straight lines, then the value of k , is

A. 9

B. 1

C. -9

D. 0

Answer: C



Watch Video Solution

16. The equation of second degree

$x^2 + 2\sqrt{2}x + 2y^2 + 4x + 4\sqrt{2}y + 1 = 0$ represents a pair of straight lines. The distance between them is a. 4 b. $\frac{4}{\sqrt{3}}$ c. 2 d.

$2\sqrt{3}$

A. 4

B. $2\sqrt{3}$

C. $4\sqrt{3}$

D. 2

Answer: D



Watch Video Solution

17. The value of λ for which the equation $x^2 - y^2 - x - \lambda y - 2 = 0$ represent a pair of straight line, are

A. $-3, 1$

B. $3, -3$

C. $-1, 1$

D. $3, 1$

Answer: B



Watch Video Solution

18. Distance between the pair of lines represented by the equation $x^2 - 6xy + 9y^2 + 3x - 9y - 4 = 0$, is

A. $\frac{15}{\sqrt{10}}$

B. $\frac{1}{2}$

C. $\sqrt{\frac{5}{2}}$

D. $\frac{1}{\sqrt{10}}$

Answer: C



Watch Video Solution

19. The equation $x^2 - 3xy + \lambda y^2 + 3x - 5y + 2 = 0$ where λ is a real number, represents a pair of straight lines. If θ is the angle between the lines, then $\cos ec^2\theta =$

A. 2

B. 0

C. 3

D. 1

Answer: B



Watch Video Solution