



MATHS

BOOKS - OBJECTIVE RD SHARMA MATHS VOL I (HINGLISH)

COMPLEX NUMBERS

Illustration

1. If $n \in N$, then find the value of $i^n + i^{n+1} + i^{n+2} + i^{n+3}$.

A. 1

B. i

C. i^n

D. 0

Answer: D





Watch Video Solution

2. If $i = \sqrt{-1}$, then $\{i^n + i^{-n}, n \in \mathbb{Z}\}$ is equal to

- A. $\{0, 2\}$
- B. $\{0, -2\}$
- C. $\{0, -2, 2\}$
- D. $\{0, -2i\}$

Answer: C



Watch Video Solution

3. The value of $\sum_{n=1}^{13} (i^n + i^{n+1})$, where $i = \sqrt{-1}$ equals

- A. i
- B. $i - 1$
- C. $-i$

D. 0

Answer: B



[Watch Video Solution](#)

4. If n is an odd integer, then $(1 + i)^{6n} + (1 - i)^{6n}$ is equal to

A. 0

B. 2

C. -2

D. none of these

Answer: A



[Watch Video Solution](#)

5. If m, n, p, q are consecutive integers then the value of $i^m + i^n + i^p + i^q$ is

A. 1

B. 4

C. 0

D. none of these

Answer: C



Watch Video Solution

6. The value of $i^2 + i^4 + i^6 + i^8 \dots$ upto $(2n+1)$ terms , where $i^2 = -1$, is equal to:

A. -1

B. 1

C. $-i$

D. i

Answer: A

 [Watch Video Solution](#)

7. If $a, b \in R$ such that $ab > 0$, then $\sqrt{a}\sqrt{b}$ is equal to

A. $\sqrt{|a||b|}$

B. $-\sqrt{|a||b|}$

C. \sqrt{ab}

D. none of these

Answer: D

 [Watch Video Solution](#)

8. If $ab < 0$, then $\sqrt{a} \cdot \sqrt{b}$ is equal to :

A. $i\sqrt{|ab|}$

B. $i\sqrt{|a||b|}$

C. $i\sqrt{|a||b|}$

D. $-\sqrt{|a||b|}$

Answer: C



Watch Video Solution

9. $\sin^{-1} \left\{ \frac{1}{i}(z - 1) \right\}$, where z is non real and $i = \sqrt{-1}$, can be the angle of a triangle if:

A. $\text{Re}(z)=1, \text{Im}(z)=2$

B. $\text{Re}(z)=1, -1 \leq \text{Im}(z) \leq 1$

C. $\text{Re}(z)+\text{Im}(z)=0$

D. None of these

Answer: B



Watch Video Solution

10. If $\sqrt{3} + i = (a + ib)/(c + id)$, then find the value of $\tan^{-1}(b/a)\tan^{-1}(d/c)$.

A. $\frac{\pi}{3}$

B. $\frac{\pi}{6}$

C. $-\frac{\pi}{6}$

D. $\frac{5\pi}{6}$

Answer: B



Watch Video Solution

11. The conjugate of a complex number is $\frac{1}{i-1}$. Then the complex number is

A. $-\frac{1}{i+1}$

B. $\frac{1}{i-1}$

C. $-\frac{1}{i-1}$

D. $\frac{1}{i+1}$

Answer: A

 [Watch Video Solution](#)

12. If $Im\left(\frac{z-1}{2z+1}\right) = -4$, then locus of z is

A. an ellipse

B. a parabola

C. a straight line

D. a circle

Answer: D

 [Watch Video Solution](#)

13. Let z be a complex number such that the imaginary part of z is nonzero and $a = z^2 + z + 1$ is real. Then a cannot take the value (A) -1 (B) 1 (C) $1/2$ (D) $3/4$

A. -1

B. $\frac{1}{3}$

C. $\frac{1}{2}$

D. $\frac{3}{4}$

Answer: D



[Watch Video Solution](#)

14. The number of solutions of $z^2 + \bar{z} = 0$ is

A. 1

B. 2

C. 3

D. 4

Answer: D



[Watch Video Solution](#)

15. If z_1, z_2 and z_3 be unimodular complex numbers, then the maximum value of $|z_1 - z_2|^2 + |z_2 - z_3|^2 + |z_3 - z_1|^2$, is

A. 6

B. 9

C. 12

D. 3

Answer: B



[Watch Video Solution](#)

16. If $|z_1| = 2$, $|z_2| = 3$, $|z_3| = 4$ and $|2z_1 + 3z_2 + 4z_3| = 4$ then the expression $|8z_2z_3 + 27z_3z_1 + 64z_1z_2|$ equals

A. 24

B. 48

C. 72

D. 96

Answer: D

 [Watch Video Solution](#)

17. Let $|z_i| = i$, $i = 1, 2, 3, 4$ and $|16z_1z_2z_3 + 9z_1z_2z_4 + 4z_1z_3z_4 + z_2z_3z_4| = 48$

, then the value of $\left| \frac{1}{\bar{z}_1} + \frac{4}{\bar{z}_2} + \frac{9}{\bar{z}_3} + \frac{16}{\bar{z}_4} \right|$

A. 1

B. 2

C. 4

D. 8

Answer: B



[Watch Video Solution](#)

18. If z_1, z_2, z_3 are complex numbers such that

$$|z_1| = |z_2| = |z_3| = \left| \frac{1}{z_1} + \frac{1}{z_2} + \frac{1}{z_3} \right| = 1 \text{ then } |z_1 + z_2 + z_3| \text{ is equal to}$$

A. equal to 1

B. less than 1

C. greater than 1

D. equal to 3

Answer: A



[Watch Video Solution](#)

19. The number of solutions of the equation $z^3 + \bar{z} = 0$, is

A. 2

B. 3

C. 4

D. 5

Answer: D



Watch Video Solution

20. If $|z_1| = |z_2| = |z_3| = 1$ and $z_1 + z_2 + z_3 = \sqrt{2} + i$, then the complex number $z_2\bar{z}_3 + z_3\bar{z}_1 + z_1\bar{z}_2$, is

A. purely real

B. purely imaginary

C. a positive real number

D. none of these

Answer: B



[Watch Video Solution](#)

21. If z is a complex number satisfying the equation $|z - (1 + i)|^2 = 2$ and $\omega = \frac{2}{z}$, then the locus traced by ' ω ' in the complex plane is

A. $(x - y + 1) = 0$

B. $x - y - 1 = 0$

C. $x + y - 1 = 0$

D. $x + y + 1 = 0$

Answer: B



[Watch Video Solution](#)

22. If $\left| \frac{z+i}{z-i} \right| = \sqrt{3}$, then z lies on a circle whose radius, is

A. $\frac{2}{\sqrt{21}}$

B. $\frac{1}{\sqrt{21}}$

C. $\sqrt{3}$

D. $\sqrt{21}$

Answer: C



[Watch Video Solution](#)

23. The smallest positive integral value of n for which $\left(\frac{1-i}{1+i} \right)^n$ is purely imaginary with positive imaginary part is

A. 1

B. 3

C. 5

D. none of these

Answer: B



[Watch Video Solution](#)

24. The least positive integer n for which $\left(\frac{1+i}{1-i}\right)^n$ is real, is

A. 2

B. 4

C. 8

D. none of these

Answer: A



[Watch Video Solution](#)

25. Find the smallest positive integer value of n for which $\frac{(1+i)^n}{(1-i)^{n-2}}$ is a real number.

A. 2

B. 1

C. 3

D. 4

Answer: B



[Watch Video Solution](#)

26. If $\left(\frac{1+i}{1-i}\right)^x = 1$, then

A. $x = 2n + 1$, where n is any positive integer.

B. $x=4n$, where n is any positive integer

C. $x=2n$, where n is any positive integer

D. $x=4n+1$, where n is any positive integer.

Answer: B



[Watch Video Solution](#)

27. If $z = x - iy$ and $z^{\frac{1}{3}} = p + iq$, then $\frac{1}{p^2 + q^2} \left(\frac{x}{p} + \frac{y}{q} \right)$ is equal to

A. -2

B. -1

C. 2

D. 1

Answer: A



[Watch Video Solution](#)

28. If $z = x + iy$, $z^{\frac{1}{3}} = a - ib$ and $\frac{x}{a} - \frac{y}{b} = \lambda(a^2 - b^2)$, then λ is equal to

A. 2

B. 4

C. 6

D. 1

Answer: B



[Watch Video Solution](#)

29. Let $z = x + iy$ be a complex number where x and y are integers. Then, the area of the rectangle whose vertices are the roots of the equation $zz^3 + \bar{z}\bar{z}^3 = 350$ is 48 (b) 32 (c) 40 (d) 80

A. 48

B.

C. 32

D. 40



Watch Video Solution

30. Taking the value of the square root with positive real part only, the value of $\sqrt{7 + 24i} + \sqrt{-7 - 24i}$, is

A. $1 + 7i$

B. $-1 - 7i$

C. $7 - i$

D. $-7 + i$

Answer: C



Watch Video Solution

31. If $(x + iy)^2 = 7 + 24i$, then the value of $(7 + \sqrt{-576})^{1/2} - (7 - \sqrt{-576})^{1/2}$, is

A. $-6i$

B. $-3i$

C. $2i$

D. 6

Answer: A



Watch Video Solution

32.
$$\frac{\sqrt{5+12i} + \sqrt{5-12i}}{\sqrt{5+12i} - \sqrt{5-12i}}$$

A. $\frac{3}{2}i$

B. $-\frac{3}{2}i$

C. $-3 + \frac{2}{5}i$

D. None of these

Answer: B



Watch Video Solution

33. Principal argument of complex number $z = \frac{\sqrt{3} + i}{\sqrt{3} - i}$ equal

A. $-\frac{\pi}{3}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{6}$

D. None of these

Answer: B



[Watch Video Solution](#)

34. Let z be a purely imaginary number such that $\text{Im}(z) > 0$. Then, $\arg(z)$ is equal to

A. π

B. $\pi/2$

C. 0

D. $-\pi/2$

Answer: B



[Watch Video Solution](#)

35. Let z be a purely imaginary number such that $\text{Im}(z) > 0$. Then, $\arg(z)$ is equal to

A. π

B. $\pi/2$

C. 0

D. $-\pi/2$

Answer: D



[Watch Video Solution](#)

36. If z is a purely real complex number such that $\operatorname{Re}(z) < 0$, then, $\arg(z)$ is equal to

A. π

B. $\pi/2$

C. 0

D. $-\pi/2$

Answer: A



[Watch Video Solution](#)

37. Let z be any non-zero complex number. Then $\operatorname{pr. arg}(z) + \operatorname{pr. arg}(\bar{z})$ is equal to

A. π

B. $-\pi$

C. 0

D. $\pi/2$

Answer: C



[Watch Video Solution](#)

38. If $z = x + iy$ such that $|z + 1| = |z - 1|$ and $\arg\left(\frac{z - 1}{z + 1}\right) = \frac{\pi}{4}$ then

A. $x^2 - y^2 - 2x - 1 = 0$

B. $x^2 + y^2 - 2x - 1 = 0$

C. $x^2 + y^2 - 2y - 1 = 0$

D. $x^2 + y^2 + 2x - 1 = 0$

Answer: C



[Watch Video Solution](#)

39. If z is a complex number of unit modulus and argument q , then

$$\arg\left(\frac{1+z}{1+\bar{z}}\right) \text{ equal}$$

A. $-\theta$

B. $\frac{\pi}{2} - \theta$

C. θ

D. $\pi - \theta$

Answer: C



Watch Video Solution

40. Find the amplitude of $\sin\left(\frac{\pi}{5}\right) + i\left(1 - \cos\left(\frac{\pi}{5}\right)\right)$

A. $\frac{2\pi}{5}$

B. $\frac{\pi}{15}$

C. $\frac{\pi}{10}$

D. $\frac{\pi}{5}$

Answer: C



Watch Video Solution

41. The value of $\sum_{n=1}^{10} \left\{ \frac{\sin(2n\pi)}{11} - i \frac{\cos(2n\pi)}{11} \right\}$, is

A. -1

B. 0

C. $-i$

D. i

Answer: D



Watch Video Solution

42. The value of

$$1 + \sum_{k=0}^{14} \left\{ \frac{\cos((2k+1)\pi)}{15} + i \frac{\sin((2k+1)\pi)}{15} \right\}, \text{ is}$$

A. 0

B. -1

C. 1

D. i

Answer: C



Watch Video Solution

43. For any integer k , let $\alpha_k = \frac{\cos(k\pi)}{7} + i \frac{\sin(k\pi)}{7}$, where $i = \sqrt{-1}$. Value of

the expression $\frac{\sum_{k=1}^{12} |\alpha_{k+1} - \alpha_k|}{\sum_{k=1}^{13} |\alpha_{4k-1} - \alpha_{4k-2}|}$ is

A. 8

B. 6

C. 4

D. 2

Answer: C



Watch Video Solution

44. If z is a complex number of unit modulus and argument θ , then the

real part of $\frac{z(1 - \bar{z})}{\bar{z}(1 + z)}$, is

A. $2\cos^2\left(\frac{\theta}{2}\right)$

B. $1 - \cos\left(\frac{\theta}{2}\right)$

C. $1 + \sin\left(\frac{\pi}{2}\right)$

D. $-2\sin^2\left(\frac{\theta}{2}\right)$

Answer: D



Watch Video Solution

45. For any two complex numbers z_1, z_2 the values of $|z_1 + z_2|^2 + |z_1 - z_2|^2$, is

A. $|z_1|^2 + |z_2|^2$

B. $2(|z_1|^2 + |z_2|^2)$

C. $(|z_1| + |z_2|)^2$

D. none of these

Answer: B



Watch Video Solution

46. For any two complex numbers, z_1, z_2

$\left| \frac{1}{2}(z_1 + z_2) + \sqrt{z_1 z_2} \right| + \left| \frac{1}{2}(z_1 + z_2) - \sqrt{z_1 z_2} \right|$ is equal to

A. $|z_1 + z_2|$

B. $|z_1 - z_2|$

C. $|z_1| + |z_2|$

D. $|z_1| - |z_2|$

Answer: C



Watch Video Solution

47. Let z_1, z_2 be two complex numbers such that $|z_1 + z_2| = |z_1| + |z_2|$.

Then,

A. $\arg(z_1) = \arg(z_2)$

B. $\arg(z_1) + \arg(z_2) = \frac{\pi}{2}$

C. $|z_1| = |z_2|$

D. $z_1 z_2 = 1$

Answer: A



Watch Video Solution

48. For any two complex numbers z_1 and z_2 , we have

$$|z_1 + z_2|^2 = |z_1|^2 + |z_2|^2, \text{ then}$$

A. $\operatorname{Re}\left(\frac{z_1}{z_2}\right) = 0$

B. $\operatorname{Im}\left(\frac{z_1}{z_2}\right) = 0$

C. $\operatorname{Re}(z_1 z_2) = 0$

D. $\operatorname{Im}(z_1 z_2) = 0$

Answer: A



Watch Video Solution

49. If z_1 and z_2 , are two non-zero complex numbers such that

$$|z_1 + z_2| = |z_1| + |z_2| \text{ then } \arg(z_1) - \arg(z_2) \text{ is equal to}$$

A. $-\pi$

B. $\pi/2$

C. 0

D. $\pi/2$

Answer: C



[Watch Video Solution](#)

50. If z_1 and z_2 are two complex numbers such that two

$$|z_1| = |z_2| + |z_1 - z_2|, \text{ then } \arg(z_1) - \arg(z_2)$$

A. 0

B. $\pi/2$

C. $-\pi/2$

D. none of these

Answer: A



[Watch Video Solution](#)

51. If $|z + 4| \leq 3$ then the maximum value of $|z + 1|$ is

A. 6

B. 0

C. 4

D. 10

Answer: A



[Watch Video Solution](#)

52. If $|z| < \sqrt{2} - 1$, then $|z^2 + 2z\cos\alpha|$ is less than

A. 1

B. $\sqrt{2} + 1$

C. $\sqrt{2} - 1$

D. $\sqrt{2}$

Answer: A



[Watch Video Solution](#)

53. Let z_1, z_2 and z_3 be three points on $|z| = 1$. If θ_1, θ_2 and θ_3 be the arguments of z_1, z_2, z_3 respectively, then

$$\cos(\theta_1 - \theta_2) + \cos(\theta_2 - \theta_3) + \cos(\theta_3 - \theta_1)$$

A. $\geq -\frac{3}{2}$

B. $\leq -\frac{3}{2}$

C. $\geq \frac{3}{2}$

D. none of these

Answer: A



[Watch Video Solution](#)

54. If z and ω are two non-zero complex numbers such that $|z\omega| = 1$ and $\arg(z) - \arg(\omega) = \frac{\pi}{2}$, then $\bar{z}\omega$ is equal to

A. $-i$

B. 1

C. -1

D. i

Answer: A



[Watch Video Solution](#)

55. If $A(z_1)$ and $B(z_2)$ are two fixed points in the Argand plane the locus of point $P(z)$ satisfying $|z - z_1| + |z - z_2| = |z_1 - z_2|$, is

A. line passing through A and B

B. line segment joining A and B

C. an ellipse

D. a circle

Answer: B

 [Watch Video Solution](#)

56. If $A(z_1)$ and $A(z_2)$ are two fixed points in the Argand plane and a point $P(z)$ moves in the Argand plane in such a way that $|z - z_1| = |z - z_2|$, then the locus of P , is

- A. the line passing through A and B
- B. the perpendicular bisector of the line segment joining A and B
- C. a line passing through the mid-point of AB
- D. a circle

Answer: B

 [Watch Video Solution](#)

57. The inequality $|z - 2| < |z - 4|$ represent the half plane

A. $\operatorname{Re}(z) \geq 3$

B. $\operatorname{Re}(z) = 3$

C. $\operatorname{Re}(z) \leq 3$

D. None of these

Answer: D



[Watch Video Solution](#)

58. If $\log_{\frac{1}{3}}|z + 1| > \log_{\frac{1}{3}}|z - 1|$ then prove that $\operatorname{Re}(z) < 0$.

A. $\operatorname{Re}(z) \geq 0$

B. $\operatorname{Re}(z) < 0$

C. $\operatorname{Im}(z) > 0$

D. None of these

Answer: B



Watch Video Solution

59. The complex numbers $z = x + iy$ which satisfy the equation

$$\left| \frac{z - 5i}{z + 5i} \right| = 1 \text{ lie on}$$

- A. the axis of x
- B. the straight line $x=5$
- C. the circle passing through the origin.
- D. none of these

Answer: A



Watch Video Solution

60. If $\omega = \frac{z}{z - \left(\frac{1}{3}\right)i}$ and $|\omega| = 1$, then find the locus of z .

A. a parabola

B. a straight line

C. a circle

D. an ellipse

Answer: B



Watch Video Solution

61. The region of the complex plane for which $\left| \frac{z - a}{z + \bar{a}} \right| = 1, (Re(a) \neq 0)$ is

A. x-axis

B. y-axis

C. the straight line $x = a$

D. none of these

Answer: B



Watch Video Solution

62. $A(z_1)$ and $B(z_2)$ are two fixed points in the Argand plane and a point $P(z)$ moves in the plane such that $|z - z_1| + |z - z_2| = \text{Constant}$ ($\neq |z_1 - z_2|$), then the locus of P , is

- A. a circle
- B. a parabola
- C. an ellipse
- D. a hyperbola

Answer: C



Watch Video Solution

63. The region of argand diagram defined by $|z - 1| + |z + 1| \leq 4$ is

- A. interior of an ellipse

B. exterior of a circle

C. interior and boundary of an ellipse

D. none of these

Answer: C



Watch Video Solution

64. $A(z_1)$ and $B(z_2)$ are two fixed points in the Argand plane and a point $P(z)$ moves in the plane such that $|z - z_1| + |z - z_2| = \text{Constant}$ ($\neq |z_1 - z_2|$), then the locus of P, is

A. a circle

B. a parabola

C. an ellipse

D. a hyperbola

Answer: D



Watch Video Solution

65. The point z in the complex plane satisfying $|z + 2| - |z - 2| = +3$ lies on

- A. a circle
- B. a parabola
- C. an ellipse
- D. a hyperbola



Watch Video Solution

66. $A(z_1)$ and $B(z_2)$ are two given points in the complex plane. The locus of a point $P(z)$ in the complex plane satisfying $|z - z_1| - |z - z_2| = |z_1 - z_2|$, is

- A. a circle
- B. an ellipse

C. a hyperbola

D. none of these

Answer: D



[Watch Video Solution](#)

67. $A(z_1)$ and $B(z_2)$ are two fixed points in the Argand plane and $P(z)$ is variable point satisfying $|z - z_1| = k|z - z_2|$, where $k > 0$ and $k \neq 1$. The locus of is

A. a circle

B. a parabola

C. an ellipse

D. a hyperbola

Answer: D



[Watch Video Solution](#)

68. If $z = x + iy$, then the equation $|(2z - i)/(z + 1)| = m$ represents a circle, then m can be 1/2 b. 1 c. 2 d. 3

A. 1/2

B. 1

C. 3

D. 2

Answer: C



[Watch Video Solution](#)

69. Points z in the complex plane satisfying $\operatorname{Re}(z + 1)^2 = |z|^2 + 1$ lie on

A. a circle

B. a parabola

C. an ellipse

D. a hyperbola

Answer: B

 [Watch Video Solution](#)

70. If z_1, z_2, z_3 be the affixes of the vertices A, B and C of a triangle having centroid at G such that $z = 0$ is the mid point of AG then

$$4z_1 + z_2 + z_3 =$$

A. $4z_1 + z_2 + z_3 = 0$

B. $z_1 + 4z_1 + z_3 = 0$

C. $z_1 + z_2 + 4z_3 = 0$

D. $z_1 + z_2 + z_3 = 0$

Answer: A

 [Watch Video Solution](#)

71. Find the relation if z_1, z_2, z_3, z_4 are the affixes of the vertices of a parallelogram taken in order.

A. $z_1 + z_3 = z_2 + z_4$

B. $z_1 + z_2 = z_3 + z_4$

C. $z_1 - z_3 = z_2 - z_4$

D. none of these

Answer: A



[Watch Video Solution](#)

72. If z_1, z_2, z_3 are the affixes of the vertices of a triangle having its circumcenter at the origin. If z is the affix of its orthocenter, then

A. $z_1 + z_2 + z_3 + z = 0$

B. $z_1 + z_2 + z_3 - z = 0$

C. $z_1 - z_2 + z_3 + z = 0$

$$D. z_1 + z_2 - z_3 + z = 0$$

Answer: B



Watch Video Solution

73. The equation $z\bar{z} + a\bar{z} + \bar{a}z + b = 0$, $b \in R$ represents circle, if

A. $|a|^2 = b$

B. $|a|^2 > b$

C. $|a|^2 < b$

D. none of these

Answer: B



Watch Video Solution

74. The center of the circle .

$z\bar{z} + (1 + i)z + (1 + i)\bar{z} - 7 = 0$ are respectively.

A. $1 + i$

B. $-1 + i$

C. $-1 - i$

D. 1

Answer: C



[Watch Video Solution](#)

75. The radius of the circle $\left| \frac{z - i}{z + i} \right| = 3$, is

A. $\frac{5}{4}$

B. $\frac{3}{4}$

C. $\frac{1}{4}$

D. none of these

Answer: B



[Watch Video Solution](#)

76. The set of values of k for which the equation

$$z\bar{z} + (-3 + 4i)\bar{z} - (3 + 4i)z + k = 0$$

represents a circle, is

A. $(-\infty, 25]$

B. $[25, \infty)$

C. $[5, \infty)$

D. $(-\infty, 5)$

Answer: A



[Watch Video Solution](#)

77. if the complex no z_1, z_2 and z_3 represents the vertices of an equilateral triangle such that $|z_1| = |z_2| = |z_3|$ then relation among z_1, z_2 and z_3

A. $z_1 + z_2 = z_3$

B. $z_2 + z_3 = z_1$

C. $z_1 + z_3 = z_2$

D. $z_1 + z_2 + z_3 = 0$

Answer: D



[Watch Video Solution](#)

78. if $|z| = 3$ then the points representing the complex numbers $-1 + 4z$ lie on a

A. line

B. circle

C. parabola

D. none of these

Answer: B



Watch Video Solution

79. If z is a complex number having least absolute value and $|z - 2 + 2i| = \sqrt{2}$, then $z =$

A. $\left(2 - \frac{1}{\sqrt{2}}\right)(1 - i)$

B. $\left(2 - \frac{1}{\sqrt{2}}\right)(1 + i)$

C. $\left(2 + \frac{1}{\sqrt{2}}\right)(1 - i)$

D. $\left(2 + \frac{1}{\sqrt{2}}\right)(1 + i)$

Answer: A



Watch Video Solution

80. The least value of p for which the two curves $\arg z = \frac{\pi}{6}$ and $|z - 2\sqrt{3}i| = p$ intersect is

A. $\sqrt{3}$

B. 3

C. $1/\sqrt{3}$

D. $1/3$

Answer: B



[Watch Video Solution](#)

81. Let a be a complex number such that $|a| < 1$ and z_1, z_2, \dots be vertices of a polygon such that $z_k = 1 + a + a^3 + a^{k-1}$.

Then, the vertices of the polygon lie within a circle.

A. $|z - a| = a$

$$\text{B. } \left| z - \frac{1}{1-a} \right| = |1-a|$$

$$\text{C. } \left| z - \frac{1}{1-a} \right| = \frac{1}{|1-a|}$$

$$\text{D. } |z - (1-a)| = |1-a|$$

Answer: C



Watch Video Solution

82. The complex number having least positive argument and satisfying

$$|z - 5i| \leq 3, \text{ is}$$

A. $12 + 16i$

B. $\frac{12}{5} + \frac{16i}{5}$

C. $\frac{16}{5} + \frac{12i}{5}$

D. $-\frac{12}{5} + \frac{16i}{5}$

Answer: B



Watch Video Solution

83. If $|z - 3 + 2i| \leq 4$, (where $i = \sqrt{-1}$) then the difference of greatest and least values of $|z|$ is

A. $2\sqrt{11}$

B. $3\sqrt{11}$

C. $2\sqrt{13}$

D. $3\sqrt{13}$

Answer: C



[Watch Video Solution](#)

84. The least distance between the circles $|z| = 12$ and $|z - 3 - 4i| = 5$, is

A. 0

B. 2

C. 7

Answer: B [Watch Video Solution](#)

85. z_1, z_2, z_3 are the vertices of an equilateral triangle taken in counter clockwise direction. If its circumference is at the origin and $z_1 = 1 + i$, then

A. $z_2 = z_1 e^{i2\pi/3}, z_3 = e^{\pi/3}$

B. $z_2 = z_1 e^{i2\pi/3}, z_3 = z_1 e^{i4\pi/3}$

C. $z_2 = z_1 e^{i4\pi/3}, z_3 = z_1 e^{i2\pi/3}$

D. $z_2 = z_1 e^{i\pi/3}, z_3 = z_1 e^{i2\pi/3}$

Answer: B [Watch Video Solution](#)

86. z_1, z_2, z_3 are the vertices of an equilateral triangle taken in counter clockwise direction. If its circumcenter is at $(1 - 2i)$ and $(z_1 = 2 + i)$, then

$z_2 =$

A. $\frac{1 - 3\sqrt{3}}{2} + \frac{\sqrt{3} - 7}{2}i$

B. $\frac{1 + 3\sqrt{3}}{2} - \frac{7 + \sqrt{3}}{2}i$

C. $\frac{1 + 3\sqrt{3}}{2}, \frac{\sqrt{3} - 7}{2}i$

D. $\frac{1 + 3\sqrt{3}}{2} + \frac{7 + \sqrt{3}}{2}i$

Answer: A



Watch Video Solution

87. The complex number z_1, z_2 and z_3 satisfying $\frac{z_1 - z_3}{z_2 - z_3} = \frac{1 - i\sqrt{3}}{2}$ are the vertices of a triangle which is :

A. of area zero

B. right angled isosceles

C. equilateral

D. obtuse-angled isosceles

Answer: C



[Watch Video Solution](#)

88. The area of the triangle on the Argand plane formed by the complex numbers z , iz and $z+iz$ is?

A. $|z|^2$

B. $\frac{1}{2}|z|^2$

C. $\frac{1}{4}|z|^2$

D. $\frac{\sqrt{3}}{4}|z|^2$

Answer: B



[Watch Video Solution](#)

89. If z is any complex number, then the area of the triangle formed by the complex number z , wz and $z+wz$ as its sides, is

A. $\frac{1}{2}|z|^2$

B. $\frac{3}{2}|z|^2$

C. $\frac{\sqrt{3}}{4}|z|^2$

D. $\frac{1}{2}|z|^2$

Answer: C



Watch Video Solution

90. The area of the triangle whose vertices are represented by 0 , z , $ze^{i\alpha}$

A. $\frac{1}{2}|z|^2\cos\alpha$

B. $\frac{1}{|z|^2}\sin\alpha$

C. $\frac{1}{2}|z|^2\sin\alpha\cos\alpha$

D. $\frac{1}{2}|z|^2$

Answer: B



Watch Video Solution

91. If z_1, z_2 are vertices of an equilateral triangle with z_0 its centroid, then

$$z_1^2 + z_2^2 + z_3^2 =$$

A. $z_1^2 + z_2^2 + z_3^2 = z_1z_2 + z_2z_3 + z_3z_1$

B. $z_1^2 + z_2^2 + z_3^2 = 2(z_1z_2 + z_2z_3 + z_3z_1)$

C. $z_1^2 + z_2^2 + z_3^2 + z_1z_2 + z_2z_3 + z_3z_1 = 0$

D. None of these

Answer: A



Watch Video Solution

92. The vertices of a square are z_1, z_2, z_3 and z_4 taken in the anticlockwise order, then $z_3 =$

A. $-iz_1 + (1 + i)z_2$

B. $iz_1 + (1 - i)z_2$

C. $z_1 + (1 + i)z_2$

D. $(1 + i)z_1 + z_2$

Answer: A



[Watch Video Solution](#)

93. ABCD is a rhombus in the Argand plane. If the affixes of the vertices are z_1, z_2, z_3 and z_4 respectively, and $\angle CBA = \pi/3$, then

A. $z_1 + \omega z_2 + \omega^2 z_3 = 0$

B. $z_1 - \omega z_2 - \omega^2 z_3 = 0$

C. $\omega z_1 + z_2 + \omega^2 z_3 = 0$

$$D. \omega^2 z_1 + \omega z_2 + z_3 = 0$$

Answer: A



[Watch Video Solution](#)

94. If two triangles whose vertices are respectively the complex numbers z_1, z_2, z_3 and a_1, a_2, a_3 are similar, then the determinant.

$$\begin{vmatrix} z_1 & a_1 & 1 \\ z_2 & a_2 & 1 \\ z_3 & a_3 & 1 \end{vmatrix} \text{ is equal to}$$

A. $z_1 z_2 z_3$

B. $a_1 a_2 a_3$

C. 1

D. 0

Answer: D



[Watch Video Solution](#)

95. The point representing the complex number z for which \arg

$$(z - 2)(z + 2) = \frac{\pi}{3} \text{ lies on}$$

- A. a circle
- B. a straight line
- C. a parabola
- D. an ellipse

Answer: A



[Watch Video Solution](#)

96. If z be any complex number ($z \neq 0$) then $\arg\left(\frac{z - i}{z + i}\right) = \frac{\pi}{2}$ represents the curve

- A. $|z| = 1$

B. $|z| = 1, \operatorname{Re}(z) > 0$

C. $|z| = 1, \operatorname{Re}(z) < 0$

D. none of these

Answer: C



[Watch Video Solution](#)

97. If $\arg \frac{z - a}{z + a} = \pm \frac{\pi}{2}$, where a is a fixed real number, then the locus of z is

A. a straight line

B. a circle with center at the origin and radius a

C. a circle with center on y -axis

D. none of these

Answer: B



[Watch Video Solution](#)

98. The length of perpendicular from $P(2 - 3i)$ on the line $(3 + 4i)Z + (3 - 4i)\bar{Z} + 9 = 0$ is equal to

A. 9

B. $9/4$

C. $9/2$

D. none of these

Answer: C



Watch Video Solution

99. Find $\left\{ \frac{1 + \cos\pi/8 + i\sin\pi/8}{1 + \cos\pi/8 - i\sin\pi/8} \right\}^8 =$

A. $1 + i$

B. $1 - i$

C. 1

D. -1

Answer: D



[Watch Video Solution](#)

100. $\frac{(\sin\pi/8 + i\cos\pi/8)^8}{(\sin\pi/8 - i\cos\pi/8)^8} =$

A. -1

B. 0

C. 1

D. 2i

Answer: C



[Watch Video Solution](#)

101. The principal amplitude of

$$\left(\sin 40^\circ + i\cos 40^\circ\right)^5, \text{ is}$$

A. 70°

B. -110°

C. 110°

D. -70°

Answer: B



[Watch Video Solution](#)

102. If $\cos\alpha + \cos\beta + \cos\gamma = \sin\alpha + \sin\beta + \sin\gamma = 0$, then the value of $\cos 3\alpha + \cos 3\beta + \cos 3\gamma$ is

A. 0

B. $\cos(\alpha + \beta + \gamma)$

C. $3\cos(\alpha + \beta + \gamma)$

D. $3\sin(\alpha + \beta + \gamma)$

Answer: C

 [Watch Video Solution](#)

103. If $x_n = \cos\left(\frac{\pi}{2^n}\right) + i\sin\left(\frac{\pi}{2^n}\right)$, $n \in N$ then $x_1, x_2, x_3, \dots, x_\infty$.

Is equal to

A. 1

B. -1

C. 0

D. none of these

Answer: B

 [Watch Video Solution](#)

104. If $(\cos\theta + i\sin\theta)(\cos2\theta + i\sin2\theta)\dots(\cos n\theta + i\sin n\theta) = 1$, then the value of θ , is

A. $4m\pi$

B. $\frac{2m\pi}{n(n+1)}$

C. $\frac{4m\pi}{n(n+1)}$

D. $\frac{m\pi}{n(n+1)}$

Answer: C



[Watch Video Solution](#)

105. If $x + \frac{1}{x} = 2\cos\theta$, then $x^n + \frac{1}{x^n}$ is equal to

A. $2\cos n\theta$

B. $2\sin n\theta$

C. $\cos n\theta$

D. $\sin n\theta$

Answer: A



Watch Video Solution

106. Let $z = \cos\theta + i\sin\theta$. Then the value of $\sum_{m \rightarrow 1-15} \text{Im}g(z^{2m-1})$ at $\theta = 2^\circ$ is:

A. $\frac{1}{\sin 2^\circ}$

B. $\frac{1}{3\sin 2^\circ}$

C. $\frac{1}{2\sin 2^\circ}$

D. $\frac{1}{4\sin 2^\circ}$

Answer: D



Watch Video Solution

107. The number of roots of the equation $z^6 = -64$ whose real parts are non-negative,

- A. 2
- B. 3
- C. 4
- D. 5

Answer: C



[Watch Video Solution](#)

108. If z_1 and z_2 are two n^{th} roots of unity, then $\arg\left(\frac{z_1}{z_2}\right)$ is a multiple of

- A. $n\pi$
- B. $\frac{3\pi}{n}$
- C. $\frac{2\pi}{n}$

D. none of these

Answer: C



Watch Video Solution

109. If $1, \alpha_1, \alpha_2, \dots, \alpha_{n-1}$ are nk^{th} roots of unity, then the value of

$(1 - \alpha_1)(1 - \alpha_2)(1 - \alpha_3) \dots (1 - \alpha_{n-1})$ is equal to

A. $\sqrt{3}$

B. $1/2$

C. n

D. 0

Answer: C



Watch Video Solution

110. If $1, \alpha_1, \alpha_2, \dots, \alpha_{n-1}$ are n^{th} roots of unity and n is an even natural number, then

$$(1 + \alpha_1)(1 + \alpha_2)(1 + \alpha_3) \dots (1 + \alpha_{n-1}) \text{ equals}$$

- A. 1
- B. 0
- C. -1
- D. none of these

Answer: B

 [Watch Video Solution](#)

111. If α is an n^{th} roots of unity, then $1 + 2\alpha + 3\alpha^2 + \dots + n\alpha^{n-1}$ equals

- A. $\frac{n}{1 - \alpha}$
- B. $-\frac{n}{1 - \alpha}$
- C. $-\frac{n}{(1 - \alpha)^2}$

D. none of these

Answer: B



[Watch Video Solution](#)

112. if $1, \omega, \omega^2$ root of the unity then The roots of the equation

$(x - 1)^3 + 8 = 0$ are

A. $-1, 1 + 2\omega, 1 + 2\omega^2$

B. $-1, 1 - 2\omega, 1 - 2\omega^2$

C. $2, 2\omega, 2\omega^2$

D. $2, 1 + 2\omega, 1 + 2\omega^2$

Answer: B



[Watch Video Solution](#)

113. The argument of $\frac{1 - i\sqrt{3}}{1 + i\sqrt{3}}$, is

A. $\frac{\pi}{3}$

B. $\frac{2\pi}{3}$

C. $\frac{7\pi}{6}$

D. $\frac{4\pi}{3}$

Answer: D



Watch Video Solution

114. If ω is an imaginary cube root of unity, then $(1 + \omega - \omega^2)^7$ equals

A. 128ω

B. -128ω

C. $128\omega^2$

D. $-128\omega^2$

Answer: D



Watch Video Solution

115. If $\omega (\neq 1)$ be a cube root of unity and $(1 + \omega^2)^n = (1 + \omega^4)^n$, then the least positive value of n , is

A. 2

B. 3

C. 5

D. 6

Answer: B



Watch Video Solution

116. If $i = \sqrt{-1}$, then $4 + 5\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^{334} + 3\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^{365}$ is equal to

A. $1 - i\sqrt{3}$

B. $-1 + i\sqrt{3}$

C. $i\sqrt{3}$

D. $-i\sqrt{3}$

Answer: C

 [Watch Video Solution](#)

117. If $\left(\frac{3}{2} + \frac{i\sqrt{3}}{2}\right)^{50} = 3^{25}(x + iy)$, where x and y are reals, then the ordered pair (x,y) is given by

A. $(0,3)$

B. $\left(1/2, \sqrt{3}/2\right)$

C. $(-3, 0)$

D. $(0, -3)$

Answer: B



Watch Video Solution

118. $x + iy = (1 - i\sqrt{3})^{100}$, then $(x, y) =$

A. $(2^{99}, 2^{99}\sqrt{3})$

B. $(2^{99}, -2^{99}\sqrt{3})$

C. $(-2^{99}, 2^{99}\sqrt{3})$

D. none of these

Answer: C



Watch Video Solution

119. If $z(2 - 2\sqrt{3}i)^2 = i(\sqrt{3} + i)^4$, then $\arg(z) =$

A. $\frac{5\pi}{6}$

B. $-\frac{\pi}{6}$

C. $\frac{\pi}{6}$

D. $\frac{7\pi}{6}$

Answer: B



Watch Video Solution

120. If ω is a complex cube root of unity, then $\arg(i\omega) + \arg(i\omega^2) =$

A. 0

B. $\pi/2$

C. π

D. $\pi/4$

Answer: C



Watch Video Solution

121. The value of the expression

$$1. (2 - \omega) \cdot (2 - \omega^2) + 2. (3 - \omega)(3 - \omega^2) + \dots + (n - 1)(n - \omega)(n - \omega^2), \quad \text{where}$$

ω is an imaginary cube root of unity, is.....

A. $\left\{ \frac{n(n+1)}{2} \right\}^2$

B. $\left\{ \frac{n(n+1)}{2} \right\}^2 - n$

C. $\left\{ \frac{n(n+1)}{2} \right\}^2 + n$

D. none of these

Answer: B



Watch Video Solution

122. If $z^2 + z + 1 = 0$ where z is a complex number, then the value of

$$\left(z + \frac{1}{z} \right)^2 + \left(z^2 + \frac{1}{z^2} \right)^2 + \dots + \left(z^6 + \frac{1}{z^6} \right)^2 \text{ is}$$

A. 54

B. 6

C. 12

D. 18

Answer: C



[Watch Video Solution](#)

123. If $\omega (\neq 1)$ is a cube root of unity, and $(1 + \omega)^7 = A + B\omega$. Then (A, B) equals

A. (0,1)

B. (1,1)

C. (1,0)

D. (-1,1)

Answer: B



[Watch Video Solution](#)

124. If α and β are the roots of the equation $x^2 - x + 1 = 0$, then $\alpha^{2009} + \beta^{2009} =$

- A. 1
- B. 2
- C. -2
- D. -1

Answer: A



[Watch Video Solution](#)

125. Let $\omega \neq 1$ be a complex cube root of unity. If

$$(3 - 3\omega + 2\omega^2)^{4n+3} + (2 + 3\omega - 3\omega^2)^{4n+3} + (-3 + 2\omega + 2\omega^2)^{4n+3} = 0,$$

then the set of possible value(s) of n is are

- A. N

B. $\{3k: k \in \mathbb{N}\}$

C. $\mathbb{N} - \{3k: k \in \mathbb{N}\}$

D. $\{6k: k \in \mathbb{N}\}$

Answer: C



Watch Video Solution

126. If z_1, z_2 are vertices of an equilateral triangle with z_0 its centroid, then

$$z_1^2 + z_2^2 + z_3^2 =$$

A. z_0^2

B. $3z_0^2$

C. $2z_0^2$

D. 0

Answer: B



Watch Video Solution

127. The origin and the roots of the equation $z^2 + pz + q = 0$ form an equilateral triangle If -

A. $p^2 = q$

B. $p^2 = 3q$

C. $q^2 = 3p$

D. $q^2 = p$

Answer: B



[Watch Video Solution](#)

128. If $A(z_1)$ and $B(z_2)$ are two points in the Argand plane such that $z_1^2 + z_2^2 + z_1z_2 = 0$, then $\triangle OAB$, is

A. equilateral

B. isosceles with $\angle AOB = \frac{\pi}{2}$

C. isosceles with $\angle AOB = \frac{2\pi}{3}$

D. isosceles with $\angle AOB = \frac{\pi}{4}$

Answer: C



[Watch Video Solution](#)

129. If $A(z_1)$, $B(z_2)$ and $C(z_3)$ are three points in the Argand plane such that $z_1 + \omega z_2 + \omega^2 z_3 = 0$, then

- A. A, B, C are collinear triangle
- B. $\triangle ABC$ is a right triangle
- C. $\triangle ABC$ is an equilateral triangle
- D. $\triangle ABC$ is right angled isosceles triangle.

Answer: C



[Watch Video Solution](#)

130. The value of i^i , is

A. $-\frac{\pi}{2}$

B. $e^{-\frac{\pi}{2}}$

C. $e^{\frac{\pi}{2}}$

D. none of these

Answer: B



[Watch Video Solution](#)

Section I - Solved Mcqs

1. The smallest positive integral value of n for which $(1 + \sqrt{3}i)^{\frac{n}{2}}$ is real is

A. 3

B. 6

C. 12

D. 0

Answer: B



[Watch Video Solution](#)

2. The least positive integral value of n for which $(\sqrt{3} + i)^n = (\sqrt{3} - i)^n$,
is

A. 3

B. 4

C. 6

D. none of these

Answer: C



[Watch Video Solution](#)

3. If $(\sqrt{3} - i)^n = 2^n$, $n \in I$, the set of integers, then n is a multiple of

- A. 6
- B. 10
- C. 9
- D. 12

Answer: D



[Watch Video Solution](#)

4. If $(1 + i)z = (1 - i)\bar{z}$, then z is equal to

- A. $t(1 - i)$, $t \in R$
- B. $t(1 + i)$, $t \in R$
- C. $\frac{t}{1 + i}$, $t \in R^+$
- D. none of these

Answer: A



Watch Video Solution

5. Let $z = \frac{\cos\theta + i\sin\theta}{\cos\theta - i\sin\theta}$, $\frac{\pi}{4} < \theta < \frac{\pi}{2}$. Then $\arg(z) =$

A. 2θ

B. $2\theta - \pi$

C. $\pi + 2\theta$

D. none of these

Answer: A



Watch Video Solution

6. If $\arg(z) < 0$, then $\arg(-z) - \arg(z) =$

A. π

B. $-\pi$

C. $\frac{\pi}{2}$

D. $\frac{\pi}{2}$

Answer: A



Watch Video Solution

7. The value of $\{\sin(\log i^i)\}^3 + \{\cos(\log i^i)\}^3$, is

A. 1

B. -1

C. 2

D. 2i

Answer: B



Watch Video Solution

8. If $z = a + ib$ satisfies $\arg(-1) = \arg(z + 3i)$, then $(a - 1) : b =$

A. 2 : 1

B. 1 : 3

C. -1 : 3

D. none of these

Answer: B



[Watch Video Solution](#)

9. If the area of the triangle on the complex plane formed by the points z , iz and $z+iz$ is 50 square units, then $|z|$ is

A. 5

B. 10

C. 15

D. none of these

Answer: B



[Watch Video Solution](#)

10. If the area of the triangle on the complex plane formed by complex numbers $z, \omega z$ is $4\sqrt{3}$ square units, then $|z|$ is

A. 4

B. 2

C. 6

D. 3

Answer: A



[Watch Video Solution](#)

11. If $x^2 + x + 1 = 0$ then the value of

$$\left(x + \frac{1}{x}\right)^2 + \left(x^2 + \frac{1}{x^2}\right)^2 + \dots + \left(x^{27} + \frac{1}{x^{27}}\right)^2 \text{ is}$$

A. 27

B. 72

C. 45

D. 54

Answer: D



Watch Video Solution

12. If $x^2 - x + 1 = 0$ then the value of $\sum_{n=1}^5 \left[x^n + \frac{1}{x^n}\right]^2$ is:

A. 8

B. 10

C. 12

D. none of these

Answer: A



[Watch Video Solution](#)

13. The value of $\alpha^{-n} + \alpha^{-2n}$, $n \in \mathbb{N}$ and α is a non-real cube root of unity, is

A. 3, if n is a multiple of 3

B. -1, if n is a multiple of 3

C. 2, if n is a multiple of 3

D. none of these

Answer: C



[Watch Video Solution](#)

14. If α is a non-real fourth root of unity, then the value of $\alpha^{4n-1} + \alpha^{4n-2} + \alpha^{4n-3}, n \in N$ is

A. 0

B. -1

C. 3

D. none of these

Answer: B



[Watch Video Solution](#)

15. If $1, \alpha, \alpha^2, \dots, \alpha^{n-1}$ are n^{th} root of unity, the value of $(3 - \alpha)(3 - \alpha^2)(3 - \alpha^3) \dots (3 - \alpha^{n-1})$, is

A. n

B. 0

C. $\frac{3n - 1}{2}$

D. $\frac{3n + 1}{2}$

Answer: C



Watch Video Solution

16. If ω is an imaginary cube root of unity, then find the value of

$$(1 + \omega)(1 + \omega^2)(1 + \omega^3)(1 + \omega^4)(1 + \omega^5) \dots \dots \cdot (1 + \omega^{3n}) =$$

A. 2^{3n}

B. 2^{2n}

C. 2^n

D. none of these

Answer: C



Watch Video Solution

17. If α is a non-real fifth root of unity, then the value of $3 \left| 1 + \alpha + \alpha^2, \alpha^{-2} - \alpha^{-1} \right|$, is

A. 9

B. 1

C. 11/3

D. none of these

Answer: A



[Watch Video Solution](#)

18. If $Z_r = \cos\left(\frac{2r\pi}{5}\right) + i\sin\left(\frac{2r\pi}{5}\right)$, $r = 0, 1, 2, 3, 4, \dots$ then $z_1 z_2 z_3 z_4 z_5$ is equal to

A. -1

B. 0

C. 1

D. none of these

Answer: C



[Watch Video Solution](#)

19. z is a complex number satisfying $z^4 + z^3 + 2z^2 + z + 1 = 0$, then $|z|$ is equal to

A. $\frac{1}{2}$

B. $\frac{3}{4}$

C. 1

D. none of these

Answer: C



[Watch Video Solution](#)

20. if $\frac{5z_2}{7z_1}$ is purely imaginary number then $\left| \frac{2z_1 + 3z_2}{2z_1 - 3z_2} \right|$ is equal to

A. $5/7$

B. $7/9$

C. $\frac{25}{49}$

D. none of these

Answer: D



Watch Video Solution

21. The locus of point z satisfying $Re\left(\frac{1}{z}\right) = k$, where k is a nonzero real number, is a. a straight line b. a circle c. an ellipse d. a hyperbola

A. a straight line

B. a circle

C. an ellipse

D. a hyperbola

Answer: B



[Watch Video Solution](#)

22. If z lies on the circle $|z| = 1$, then $2/z$ lies on

A. a circle

B. an ellipse

C. a straight line

D. a parabola

Answer: A



[Watch Video Solution](#)

23. The maximum value of $|z|$ where z satisfies the condition $\left|z + \left(\frac{2}{z}\right)\right| = 2$ is

A. $\sqrt{3} - 1$

B. $\sqrt{3}$

C. $\sqrt{3} + 1$

D. $\sqrt{2} + \sqrt{3}$

Answer: C



Watch Video Solution

24. If $\left|z - \frac{4}{z}\right| = 2$, then the maximum value of $|z|$ is equal to (1) $\sqrt{3} + 1$ (2)

$\sqrt{5} + 1$ (3) 2 (4) $2 + \sqrt{2}$

A. $\sqrt{5}$

B. $\sqrt{5} + 1$

C. $\sqrt{5} - 1$

D. none of these

Answer: B



[Watch Video Solution](#)

25. if $|z^2 - 1| = |z|^2 + 1$ then z lies on

A. a circle

B. a parabola

C. an ellipse

D. none of these

Answer: D



[Watch Video Solution](#)

26. If the number $\frac{z - 1}{z + 1}$ is purely imaginary, then

A. $|z| = 1$

B. $|z| > 1$

C. $|z| < 1$

D. $|z| > 2$

Answer: A



Watch Video Solution

27. If $|z| = k$ and $\omega = \frac{z - k}{z + k}$, then $\text{Re}(\omega) =$

A. 0

B. k

C. $\frac{1}{k}$

D. $-\frac{1}{k}$

Answer: A



Watch Video Solution

28. If $k > 0$, $|z| = |w| = k$, and $\alpha = \frac{z - \bar{w}}{k^2 + z\bar{w}}$, $Re(\alpha)$

A. 0

B. $k/2$

C. k

D. none of these

Answer: A



Watch Video Solution

29. The region in the Argand diagram defined by $|z - 2i| + |z + 2i| < 5$ is the ellipse with major axis along

A. the real axis

B. the imaginary axis

C. $y = x$

D. $y = -x$

Answer: B



Watch Video Solution

30. Prove that $|Z - Z_1|^2 + |Z - Z_2|^2 = a$ will represent a real circle [with center $(\frac{|Z_1 + Z_2|}{2} +)$] on the Argand plane if $2a \geq |Z_1 - Z_2|^2$

A. $k < |z_1 - z_2|^2$

B. $k = |z_1 - z_2|^2$

C. $k \geq \frac{1}{2}|z_1 - z_2|^2$

D. $k < \frac{1}{2}|z_1 - z_2|^2$

Answer: C



[Watch Video Solution](#)

31. The equation $|z - 1|^2 + |z + 1|^2 = 2$, represent

- A. a circle of radius one unit
- B. a straight line
- C. the ordered pair (0,0)
- D. none of these

Answer: C



[Watch Video Solution](#)

32. The points representing the complex numbers z for which

$|z + 4|^2 - |z - 4|^2 = 8$ lie on

- A. a straight line parallel to x-axis
- B. a straight line parallel to y-axis

C. a circle with center as origin

D. a circle with center other than the origin.

Answer: B



[Watch Video Solution](#)

33. If $|z + \bar{z}| = |z - \bar{z}|$, then value of locus of z is

A. a pair of straight line

B. a rectangular hyperbola

C. a line

D. a set of four lines

Answer: A



[Watch Video Solution](#)

34. If $|z + \bar{z}| + |z - \bar{z}| = 2$, then z lies on

- A. a straight line
- B. a square
- C. a circle
- D. none of these

Answer: A



[Watch Video Solution](#)

35. The closest distance of the origin from a curve given as

$A\bar{z} + \bar{A}z + A\bar{A} = 0$ is: (A is a complex number).

A. 1 unit

B. $\frac{\operatorname{Re}(A)}{|A|}$

C. $\frac{\operatorname{Im}(A)}{|A|}$

D. $\frac{1}{2}|A|$

Answer: D



Watch Video Solution

36. If $z_1 = 1 + 2i$, $z_2 = 2 + 3i$, $z_3 = 3 + 4i$, then z_1, z_2 and z_3 represent the vertices of a/an.

- A. equilateral triangle
- B. right angled triangle
- C. isosceles triangle
- D. none of these

Answer: D



Watch Video Solution

37. If z_1 and z_2 are two of the 8^{th} roots of unity such that $\arg\left(\frac{z_1}{z_2}\right)$ is least positive, then $\frac{z_1}{z_2}$ is

A. $1 + i$

B. $1 - i$

C. $\frac{1 + i}{\sqrt{2}}$

D. $\frac{1 - i}{\sqrt{2}}$

Answer: C

 [View Text Solution](#)

38. The number of roots of the equation $z^{15} = 1$ satisfying $|\arg(z)| \leq \pi/2$, is

A. 6

B. 7

C. 8

D. none of these

Answer: B



[View Text Solution](#)

39. If z_1, z_2, \dots, z_n lie on the circle $|z| = R$, then

$$\left| z_1 + z_2 + \dots + z_n - R^2 \left(\frac{1}{z_1} + \frac{1}{z_2} + \dots + \frac{1}{z_n} - (n) \right) \right| \text{ is equal to}$$

A. nR

B. $-nR$

C. 0

D. n

Answer: C



[Watch Video Solution](#)

40. Q. Let z_1 and z_2 be n th roots of unity which subtend a right angle at the origin, then n must be the form $4k$.

A. $4k + 1$

B. $4k + 2$

C. $4k + 3$

D. $4k$

Answer: D



Watch Video Solution

41. The complex number z_1, z_2 and z_3 satisfying $\frac{z_1 - z_3}{z_2 - z_3} = \frac{1 - i\sqrt{3}}{2}$ are the vertices of a triangle which is :

A. of area zero

B. right-angled isosceles

C. equilateral

D. obtuse-angled isosceles

Answer: C



Watch Video Solution

42. Let $\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$, then the value of the determinant

$$\begin{vmatrix} 1 & 1 & 1 \\ 1 & -1 - \omega^2 & \omega^2 \\ 1 & \omega^2 & \omega^4 \end{vmatrix}, \text{ is}$$

A. 3ω

B. $3\omega(\omega - 1)$

C. $3\omega^2$

D. $3\omega(1 - \omega)$

Answer: B



Watch Video Solution

43. For all complex numbers z_1, z_2 satisfying $|z_1| = 12$ and $|z_2 - 3 - 4i| = 5$, find the minimum value of $|z_1 - z_2|$

A. 0

B. 2

C. 7

D. 17

Answer: B

 Watch Video Solution

44. Let z_1, z_2 be two complex numbers represented by points on the circle $|z_1| = 4$ and $|z_2| = 2$ are then

A. $\max |2z_1 + z_2| = 4$

B. $\min |z_1 - z_2| = 1$

C. $\left| z_2 + \frac{1}{z_1} \right| \leq 3$

D. all of the above.

Answer: D

 [Watch Video Solution](#)

45. If z lies on unit circle with center at the origin, then $\frac{1+z}{1+\bar{z}}$ is equal to

A. z

B. \bar{z}

C. $z + \bar{z}$

D. none of these

Answer: A

 [Watch Video Solution](#)

46. If $|z_1 - 1| < 1$, $|z_2 - 2| < 2$, $|z_3 - 3| < 3$ then $|z_1 + z_2 + z_3|$

A. is less than 6

B. is more than 3

C. is less than 12

D. lies between 6 and 12

Answer: C



[Watch Video Solution](#)

47. Complex numbers z_1 and z_2 lie on the rays $\arg(z_1) = \theta$ and $\arg(z_1) = -\theta$ such that $|z_1| = |z_2|$. Further, image of z_1 in y-axis is z_3 . Then, the value of $\arg(z_1 z_3)$ is equal to

A. $\frac{\pi}{2}$

B. $-\frac{\pi}{2}$

C. π

D. none of these

Answer: C



[Watch Video Solution](#)

48. If z is a complex number satisfying $|z|^2 - |z| - 2 < 0$, then the value of $|z^2 + z\sin\theta|$, for all values of θ , is

A. equal to 4

B. equal to 6

C. more than 6

D. less than 6

Answer: D



[Watch Video Solution](#)

49. $|z - i| \leq 2$ and $z_0 = 5 + 3i$ then max. value of $|iz + z_0|$ is :

A. $2 + \sqrt{31}$

B. 7

C. $\sqrt{31} - 2$

D. none of these

Answer: B



[Watch Video Solution](#)

50. If $|z| = \max\{|z - 2|, |z + 2|\}$, then

A. $|z + \bar{z}| = 2$

B. $z + \bar{z} = 4$

C. $|z + \bar{z}| = 1$

D. none of these

Answer: A



Watch Video Solution

51. if $\left| \frac{z-6}{z+8} \right| = 1$, then the value of $x \in R$, where

$$z = x + i \begin{vmatrix} -3 & 2i & 2+i \\ -2i & 2 & 4-3i \\ 2-i & 4+3i & 7 \end{vmatrix}, \text{ is}$$

A. 5

B. 7

C. 9

D. 0

Answer: B



Watch Video Solution

52. If $|z - 1| + |z + 3| \leq 8$, then the range of values of $|z - 4|$ is

A. (0,8)

B. [0,9]

C. [1,9]

D. [5,9]

Answer: C



[Watch Video Solution](#)

53. The equation $|z - i| + |z + i| = k, k > 0$ can represent an ellipse, if $k =$

A. 1

B. 2

C. 4

D. none of these

Answer: C



Watch Video Solution

54. Find the range of K for which the equation $|z + i| - |z - i| = K$ represents a hyperbola.

A. $k \in (-2, 2)$

B. $k \in [2, 2]$

C. $k \in (0, 2)$

D. $k \in (-2, 0)$

Answer: A



Watch Video Solution

55. If $|z + 3i| + |z - i| = 8$, then the locus of z , in the Argand plane, is

- A. an ellipse of eccentricity $\frac{1}{2}$ and major axis along x-axis.
- B. an ellipse of eccentricity $\frac{1}{2}$ and major axis of along y-axis.
- C. an ellipse of eccentricity $\frac{1}{\sqrt{2}}$ and major axis along y-axis
- D. none of these

Answer: A

 [Watch Video Solution](#)

56. In Fig. 42, a point 'z' is equidistant from three distinct points z_1, z_2 and z_3 in the Argand plane. If z, z_1 and z_2 are collinear, then $\arg\left(\frac{z_3 - z_1}{z_3 - z_2}\right)$. Will be (z_1, z_2, z_3) are in anticlockwise sense).

- A. $\frac{\pi}{2}$
- B. $-\frac{\pi}{2}$
- C. $\frac{\pi}{6}$
- D. $\frac{2\pi}{3}$

Answer: B



Watch Video Solution

57. Let $P(e^{i\theta_1})$, $Q(e^{i\theta_2})$ and $R(e^{i\theta_3})$ be the vertices of a triangle PQR in the Argand Plane. The orthocenter of the triangle PQR is

A. $e^{i(\theta_1 + \theta_2 + \theta_3)}$

B. $\frac{2}{3}e^{i(\theta_1 + \theta_2 + \theta_3)}$

C. $e^{i\theta_1} + e^{i\theta_2} + e^{i\theta_3}$

D. none of these

Answer: C



Watch Video Solution

58. If $A(z_1)$, $B(z_2)$, $C(z_3)$ are the vertices of an equilateral triangle ABC ,

then $\arg \frac{2z_1 - z_2 - z_3}{z_3 - z_2} =$

A. $\frac{\pi}{4}$

B. $\frac{\pi}{2}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{6}$

Answer: B

 [Watch Video Solution](#)

59. If $A(z_1)$, $B(z_2)$ and $C(z_3)$ are three points in the argand plane where

$$|z_1 + z_2| = |z_1 - z_2| \text{ and } |(1 - i)z_1 + iz_3| = |z_1| + |z_3| - |z_1|, \text{ where } i = \sqrt{-1}$$

then

A. A, B and C lie on a circle with center $\frac{z_2 + z_3}{2}$

B. A, B and C are collinear points.

C. A, B, C form an equilateral triangle.

D. A, B, C form an obtuse angle triangle.

Answer: A



Watch Video Solution

60. If a_1, a_2, \dots, a_n are n th roots of unity then

$\frac{1}{1-a_1} + \frac{1}{1-a_2} + \frac{1}{1-a_3} \dots + \frac{1}{1-a_n}$ is equal to

A. $\frac{n-1}{2}$

B. $\frac{n}{2}$

C. $\frac{2^n - 1}{2}$

D. none of these

Answer: A



Watch Video Solution

61. Let $A(z_1)$ and $B(z_2)$ be such that $\angle AOB = \theta$ (O' being the origin). If

we define $z_1 \times z_2 = |z_1| |z_2| \sin \theta$, then $z_1 \times z_2$ is also equal to

A. $\operatorname{Re}(z_1 \bar{z}_2) = 0$

B. $\operatorname{Re}(\bar{z}_1 z_2) = 0$

C. $\operatorname{Im}(\bar{z}_1 z_2) = 0$

D. none of these

Answer: C



Watch Video Solution

62. If one root of $z^2 + (a + i)z + b + ic = 0$ is real, where $a, b, c \in R$, then

$c^2 + b - ac =$

A. 0

B. -1

C. 1

D. none of these

Answer: A

 [Watch Video Solution](#)

63. If A and B represent the complex numbers z_1 and z_2 such that $|z_1 + z_2| = |z_1 - z_2|$, then the circumcenter of $\triangle OAB$, where O is the origin, is

A. $\frac{z_1 + z_2}{3}$

B. $\frac{z_1 + z_2}{2}$

C. $\frac{z_1 - z_2}{2}$

D. none of these

Answer: B

 [Watch Video Solution](#)

64. If z_1 , lies in $|z - 3| \leq 4$, z_2 on $|z - 1| + |z + 1| = 3$ and $A = |z_1 - z_2|$, then :

A. $0 \leq A \leq \frac{15}{2}$

B. $0 < A < \frac{15}{2}$

C. $0 \leq A \leq \frac{17}{2}$

D. $0 \leq A < \frac{17}{2}$

Answer: D



Watch Video Solution

65. Let O, A, B be three collinear points such that $OA \cdot OB = 1$. If O and B represent the complex numbers O and z , then A represents

A. $\frac{1}{z}$

B. \bar{z}

C. $\frac{1}{\bar{z}}$

D. none of these

Answer: C



Watch Video Solution

66. If z_0, z_1 represent points P and Q on the circle $|z - 1| = 1$ taken in anticlockwise sense such that the line segment PQ subtends a right angle at the center of the circle, then $z_1 =$

A. $1 + i(z_0 - 1)$

B. iz_0

C. $1 - i(z_0 - 1)$

D. $i(z_0 - 1)$

Answer: A



Watch Video Solution

67. The center of a square ABCD is at the origin and point A is represented by z_1 . The centroid of $\triangle BCD$ is represented by

A. $\frac{z_1}{3}$

B. $-\frac{z_1}{3}$

C. $\frac{iz_1}{3}$

D. $-\frac{iz_1}{3}$

Answer: B



Watch Video Solution

68. The value of k for which the inequality $|Re(z)| + |Im(z)| \leq \lambda|z|$ is true for all $z \in C$, is

A. 2

B. $\sqrt{2}$

C. 1

D. none of these

Answer: B



Watch Video Solution

69. The value of λ for which the inequality $\left| \frac{z_1}{|z_1|} + \frac{z_2}{|z_2|} \right| \leq \lambda$ is true for all

$z_1, z_2 \in C$, is

A. 1

B. 2

C. 3

D. none of these

Answer: B



Watch Video Solution

70. If z_1 and z_2 both satisfy $z + z = 2|z - 1|$ and $\arg(z_1 - z_2) = \frac{\pi}{4}$, then find

$\text{Im}(z_1 + z_2)$.

A. 0

B. 1

C. 2

D. none of these

Answer: C



[Watch Video Solution](#)

71. If z satisfies $|z + 1| < |z - 2|$, then $w = 3z + 2 + i$

A. $|\omega + 1| < |\omega - 8|$

B. $|\omega + 1| < |\omega - 7|$

C. $\omega + \bar{\omega} > 7$

D. $|\omega + 5| < |\omega - 4|$

Answer: A



[Watch Video Solution](#)

72. If z complex number satisfying $|z - 1| = 1$, then which of the following is correct

A. $\arg(z - 1) = 2\arg(z)$

B. $2\arg(z) = \frac{2}{3}\arg(z^2 - z)$

C. $\arg(z - 1) = 2\arg(z + 1)$

D. $\arg z = 2\arg(z + 1)$

Answer: A



[Watch Video Solution](#)

73. If z_1, z_2, z_3 are the vertices of an isoscles triangle right angled at z_2 , then

A. $z_1^2 + 2z_2^2 + z_3^2 = 2z_2(z_1 + z_3)$

B. $z_1^2 + z_2^2 + z_3^2 = 2z_2(z_1 + z_3)$

C. $z_1^2 + z_2^2 + 2z_3^2 = 2z_2(z_1 + z_3)$

D. $2z_1^2 + z_2^2 + z_3^2 = 2z_2(z_1 + z_3)$

Answer: A



Watch Video Solution

74. All the roots of the equation $a_1z^3 + a_2z^2 + a_3z + a_4 = 3$ where $|a_i| < 1, i = 1, 2, 3, 4$, lie outside the circle with center at the origin and radius equal to

A. 1

B. $1/3$

C. $2/3$

D. none of these

Answer: C



Watch Video Solution

75. If z is a point on the Argand plane such that $|z - 1| = 1$, then $\frac{z - 2}{z}$ is equal to

- A. $\tan(\arg z)$
- B. $\cot(\arg z)$
- C. $i \tan(\arg z)$
- D. none of these

Answer: C



[View Text Solution](#)

76. If z is a non-real complex number lying on the circle $|z| = 1$, then $z =$

A.
$$\frac{1 - i \tan\left(\frac{\arg z}{2}\right)}{1 + i \tan\left(\frac{\arg z}{2}\right)}$$

$$1 + i \tan\left(\frac{\arg z}{2}\right)$$

B. $\frac{1 + i \tan\left(\frac{\arg z}{2}\right)}{1 - i \tan\left(\frac{\arg z}{2}\right)}$

$$1 - i \tan\left(\frac{\arg z}{2}\right)$$

C. $\frac{1 - i \tan(\arg z)}{1 + i \tan\left(\frac{\arg z}{2}\right)}$

$$1 + i \tan\left(\frac{\arg z}{2}\right)$$

D. none of these

Answer: B



Watch Video Solution

77. If $|z| = 2$ and the locus of $5z-1$ is the circle having radius 'a' and

$$z_1^2 + z_2^2 - 2z_1z_2\cos\theta = 0 \text{ then } |z_1| : |z_2| =$$

A. a:1

B. 2a:1

C. a:10

D. none of these

Answer: C



[Watch Video Solution](#)

78. If $|z + \bar{z}| + |z - \bar{z}| = 8$, then z lies on

- A. a circle
- B. a straight line
- C. a square
- D. an ellipse

Answer: C



[Watch Video Solution](#)

79. If a point z_1 is the reflection of a point z_2 through the line

$b\bar{z} + \bar{b}z = c, b \in \mathbb{C}, c \in \mathbb{R}$, in the Argand plane, then $b\bar{z}_2 + \bar{b}z_1 =$

A. $4c$

B. $2c$

C. c

D. none of these

Answer: C



Watch Video Solution

80. If z is a complex number satisfying $\left|z^2 + 1\right| = 4|z|$, then the minimum value of $|z|$ is

A. $2\sqrt{5} + 4$

B. $2\sqrt{5} - 4$

C. $\sqrt{5} - 2$

D. none of these

Answer: C



Watch Video Solution

81. If z_1 and z_2 are two complex numbers satisfying the equation.

$$\left| \frac{iz_1 + z_2}{iz_1 - z_2} \right| = 1, \text{ then } \frac{z_1}{z_2} \text{ is}$$

- A. 0
- B. purely real
- C. negative real
- D. purely imaginary

Answer: D



Watch Video Solution

82. If α is an imaginary fifth root of unity, then $\log_2 \left| 1 + \alpha + \alpha^2 + \alpha^3 - \frac{1}{\alpha} \right| =$

- A. 1

B. 0

C. 2

D. -1

Answer: A



[Watch Video Solution](#)

83. The roots of the equation $(1 + i\sqrt{3})^x - 2^x = 0$ form

A. an A.P.

B. a G.P.

C. an H.P.

D. none of these

Answer: A



[Watch Video Solution](#)

84. If $|z| = 1$ and $\omega = \frac{z-1}{z+1}$ (where $z \neq -1$), then $\operatorname{Re}(\omega)$ is

A. 0

B. $-\frac{1}{|z+1|^2}$

C. $\left| \frac{z}{z+1} \right| \frac{1}{|z+1|^2}$

D. $\frac{\sqrt{2}}{|z+1|^2}$

Answer: A



Watch Video Solution

85. Let z, w be complex numbers such that $\bar{z} + i\bar{w} = 0$ and $\arg(zw) = \pi$

.Then $\arg(z)$ equals

A. $\frac{5\pi}{4}$

B. $\frac{\pi}{2}$

C. $\frac{3\pi}{4}$

D. $\frac{\pi}{4}$

Answer: C



Watch Video Solution

86. Let $OP \cdot OQ = 1$ and let O, P and Q be three collinear points. If O and Q represent the complex numbers of origin and z respectively, then P represents

A. $\frac{1}{z}$

B. \bar{z}

C. $\frac{1}{\bar{z}}$

D. $-z$

Answer: C



Watch Video Solution

87. If $|z| = 1$ and $z \neq \pm 1$, then all the values of $\frac{z}{1-z^2}$ lie on a line not passing through the origin $|z| = \sqrt{2}$ the x-axis (d) the y-axis

A. a line not passing through the origin

B. $|z| = \sqrt{2}$

C. the x-axis

D. the y-axis

Answer: D



Watch Video Solution

88.

Let

$A = \{z: \text{Im}(z) \geq 1\}$, $B = \{z: |z - 2 - i| = 3\}$, $C = \{z: \text{Re}\{(1 - i)z\} = \sqrt{2}\}$ be

three sides of complex numbers. Then, the number of elements in the set

$A \cap B \cap C$, is

A. 0

B. 1

C. 2

D. ∞

Answer: B



Watch Video Solution

89. Let $S = S_1 \cap S_2 \cap S_3$ where

$$S_1 = \left\{ z \in \mathbb{C} : |z| < 4 \right\}, S_2 = \left\{ z \in \mathbb{C} : \operatorname{Im} \left[\frac{z - 1 + \sqrt{3}i}{1 - \sqrt{3}i} \right] > 0 \right\}$$

and

$$S_3 = \{ z \in \mathbb{C} : \operatorname{Re} z < 0 \}$$

Let z be any point in $A \cap B \cap C$

The $|z + 1 - i|^2 + |z - 5 - i|^2$ lies between

A. 25 and 29

B. 30 and 34

C. 35 and 39

D. 40 and 44

Answer: C



[Watch Video Solution](#)

90. In Q.no. 88, if z be any point in $A \ B \ C$ and ω be any point satisfying

$|\omega - 2 - i| < 3$. Then, $|z| - |\omega| + 3$ lies between

A. -6 and 3

B. -3 and 6

C. -6 and 6

D. -3 and 9

Answer: D



[Watch Video Solution](#)

91. A particle P starts from the point $z_0 = 1 + 2i$, where $i = \sqrt{-1}$. It moves first horizontally away from origin by 5 units and then vertically away

from origin by 3 units in the direction of the vector $\hat{i} + \hat{j}$ and then it moves through an angle $\frac{\pi}{2}$ in anticlockwise direction on a circle with center at origin, to reach a point z_2 . The point z_2 is given by

A. $6 + 7i$

B. $-7 + 6i$

C. $7 + 6i$

D. $-6 + 7i$

Answer: D



Watch Video Solution

92. If $w = \alpha + i\beta$ where $\beta \neq 0$ and $z \neq 1$ satisfies the condition that $\left(\frac{w - \bar{w}z}{1 - z} \right)$

is purely real then the set of values of z is

A. $\{z : |z| = 1\}$

B. $\{z : z = \bar{z}\}$

C. $\{z: z \neq 1\}$

D. $\{z: |z| = 1, z \neq 1\}$

Answer: D



[Watch Video Solution](#)

93. If z_1 and \bar{z}_1 represent adjacent vertices of a regular polygon of n sides where centre is origin and if $\frac{\text{Im}(z)}{\text{Re}(z)} = \sqrt{2} - 1$, then n is equal to:

A. 8

B. 16

C. 24

D. 32

Answer: A



[Watch Video Solution](#)

94. If $|z| = \max \{|z - 1|, |z + 1|\}$, then

A. $|z + \bar{z}| = \frac{1}{2}$

B. $z + \bar{z} = 1$

C. $|z + \bar{z}| = 1$

D. $z - \bar{z} = 5$

Answer: C



Watch Video Solution

95. The minimum value of $|a + b\omega + c\omega^2|$, where a, b, c are all not equal integers and $\omega (\neq 1)$ is a cube root of unity, is

A. $\sqrt{3}$

B. $1/2$

C. 1

D. 0

Answer: C



Watch Video Solution

96. The shaded region, where $P = (-1, 0)$, $Q = (-1 + \sqrt{2}, \sqrt{2})$, $R = (-1 + \sqrt{2}, -\sqrt{2})$, $S = (1, 0)$ is represented by Figure $|z+1| > 2, |\arg(z+1)| > \pi/4$

A. $|z + 1| > 2, |\arg(z + 1)| < \frac{\pi}{4}$

B. $|z + 1| < 2, |\arg(z + 1)| < \frac{\pi}{4}$

C. $|z - 1| > 2, |\arg(z + 1)| > \frac{\pi}{4}$

D. $|z - 1| < 2, |\arg(z + 1)| > \frac{\pi}{2}$

Answer: A



Watch Video Solution

97. If a, b, c are distinct integers and $\omega (\neq 1)$ is a cube root of unity, then the minimum value of $\left| a + b\omega + c\omega^2 \right| + \left| a + b\omega^2 + c\omega \right|$ is

A. $2\sqrt{3}$

B. 3

C. $4\sqrt{2}$

D. 2

Answer: A



Watch Video Solution

98. Let a and b be two positive real numbers and z_1 and z_2 be two non-zero complex numbers such that $a|z_1| = b|z_2|$. If $z = \frac{az_1}{bz_2} + \frac{bz_2}{az_1}$, then

A. $\text{Re}(z)=0$

B. $\text{Im}(z)=0$

C. $|z| = \frac{a}{b}$

D. $|z| > 2$

Answer: B



Watch Video Solution

99. If points having affixes z , $-iz$ and 1 are collinear, then z lies on

A. a straight line

B. a circle

C. an ellipse

D. a pair of straight lines.

Answer: B



Watch Video Solution

100. If $0 \leq \arg(z) \leq \frac{\pi}{4}$, then the least value of $|z - i|$, is

A. 1

B. $\frac{1}{\sqrt{2}}$

C. $\sqrt{2}$

D. none of these

Answer: B



Watch Video Solution

101. If $|z_1| + |z_2| = 1$ and $z_1 + z_2 + z_3 = 0$ then the area of the triangle whose vertices are z_1, z_2, z_3 is

A. $\frac{3\sqrt{3}}{4}$

B. $\frac{\sqrt{3}}{4}$

C. 1

Answer: A


102. Let Z_1 and Z_2 , be two distinct complex numbers and let $w = (1 - t)z_1 + tz_2$ for some number "t" with $0 < t < 1$

A. $|z - z_2| + |z - z_1| = |z_1 - z_2|$

B. $\arg(z - z_1) = \arg(z - z_2)$

C. $\begin{vmatrix} z - z_1 & \bar{z} - \bar{z}_1 \\ z_2 - z_1 & \bar{z}_2 - \bar{z}_1 \end{vmatrix} = 0$

D. $\arg(z - z_1) = \arg(z_2 - z_1)$

Answer: B


103. Let ω be the complex number $\cos\left(\frac{2\pi}{3}\right) + i\sin\left(\frac{2\pi}{3}\right)$. Then the number of distinct complex numbers z satisfying

$$\Delta = \begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 0$$
 is

- A. 1
- B. 0
- C. 2
- D. 3

Answer: A



[Watch Video Solution](#)

104. The set of points z in the complex plane satisfying $|z - i|z|| = |z + i|z||$ is contained or equal to the set of points z satisfying

A. $\text{Im}(z) = 0$

B. $\text{Im}(z) \leq 1$

C. $|\text{Re}(z)| \leq 2$

D. $|z| \leq 3$

Answer: A

 [Watch Video Solution](#)

105. The set of points z satisfying $|z + 4| + |z - 4| = 10$ is contained or equal to

A. an ellipse with eccentricity $= \frac{4}{5}$

B. the set of points z satisfying $|z| \leq 3$

C. the set of points z satisfying $|\text{Re}(z)| \leq 2$

D. the set of points z satisfying $|\text{Im}(z)| < 1$

Answer: A



Watch Video Solution

106. If $|\omega| = 2$, then the set of points $z = \omega - \frac{1}{\omega}$ is contained in or equal to the set of points z satisfying

A. $\text{Im}(z) = 0$

B. $|\text{Im}(z)| \leq 1$

C. $|\text{Re}(z)| \leq 2$

D. $|z| \leq 3$

Answer: D



Watch Video Solution

107. If $|\omega| = 1$, then the set of points $z = \omega + \frac{1}{\omega}$ is contained in or equal to the set of points z satisfying.

A. $\text{Re}(z) \leq 2$ and $\text{Im}(z) = 0$

B. $\operatorname{Re}(z) \leq 1$ and $\operatorname{Im}(z) = 0$

C. $\operatorname{Re}(z) \leq 2$ and $\operatorname{Im}(z) = 0$

D. $\operatorname{Re}(z) \leq 1$ and $\operatorname{Im}(z) = 0$

Answer: C

 [Watch Video Solution](#)

108. The number of complex numbers z such that $|z - 1| = |z + 1| = |z - i|$ is

A. 2

B. ∞

C. 0

D. 1

Answer: D

 [Watch Video Solution](#)

109. Let α and β be real numbers and z be a complex number. If $z^2 + \alpha z + \beta = 0$ has two distinct non-real roots with $\operatorname{Re}(z)=1$, then it is necessary that

- A. $\beta \in (0, 1)$
- B. $\beta \in (-1, 0)$
- C. $|\beta| < 1$
- D. $\beta \in (1, \infty)$

Answer: D



[Watch Video Solution](#)

110. If $\omega \neq 1$ is the complex cube root of unity and matrix $H = \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix}$, then H^{70} is equal to

- A. $-H$
- B. H^2

C. H

D. O

Answer: C



[Watch Video Solution](#)

111. The maximum value of $\left| \arg\left(\frac{1}{1-z}\right) \right|$ or $|z| = 1, z \neq 1$ is given by.

A. $\frac{\pi}{6}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{2}$

D. π

Answer: C



[Watch Video Solution](#)

112. If z is any complex number satisfying $|z - 3 - 2i| \leq 2$ then the maximum value of $|2z - 6 + 5i|$ is

A. 3

B. 4

C. 5

D. $5/2$

Answer: C



[Watch Video Solution](#)

113. Let ω be the solution of $x^3 - 1 = 0$ with $\text{Im}(\omega) > 0$. If $a=2$ with b and c

satisfying $[abc] \begin{bmatrix} 1 & 9 & 7 \\ 2 & 8 & 7 \\ 7 & 3 & 7 \end{bmatrix} = [0, 0, 0]$, then the value of $\frac{3}{\omega^a} + \frac{1}{\omega^b} + \frac{1}{\omega^c}$ is

equal to

A. -2

B. 2

C. 3

D. -3

Answer: A



Watch Video Solution

114. The set $\left\{ \operatorname{Re} \left(\frac{2iz}{1-z^2} \right) : z \text{ is a complex number, } |z| = 1, z \neq \pm 1 \right\}$ is _____.

A. $(-\infty, -1) \cup (1, \infty)$

B. $(-\infty, 0) \cup (1, \infty)$

C. $[2, \infty)$

D. $(-\infty, -1) \cup [1, \infty)$

Answer: D



Watch Video Solution

115. Let $\omega = e^{\frac{i\pi}{3}}$ and a, b, c, x, y, z be non-zero complex numbers such that $a + b + c = x, a + b\omega + c\omega^2 = y, a + b\omega^2 + c\omega = z$. Then, the value of $\frac{|x|^2 + |y|^2 + |z|^2}{|a|^2 + |b|^2 + |c|^2}$

A. 3

B. 6

C. 9

D. 1

Answer: A



Watch Video Solution

116. The minimum value of $|z_1 - z_2|$ as z_1 and z_2 vary over the curves $|\sqrt{3}(1 - 2z) + 2i| = 2\sqrt{7}$ and $|\sqrt{3}(-1 - z) - 2i| = |\sqrt{3}(9 - z) + 18i|$ respectively is

- $7\sqrt{7}$
- A. $\frac{7\sqrt{7}}{2\sqrt{3}}$
- $5\sqrt{7}$
- B. $\frac{5\sqrt{7}}{2\sqrt{3}}$
- $14\sqrt{7}$
- C. $\frac{14\sqrt{7}}{\sqrt{3}}$
- $7\sqrt{7}$
- D. $\frac{7\sqrt{7}}{5\sqrt{3}}$

Answer: B



[View Text Solution](#)

117. Let complex numbers α and $\frac{1}{\alpha}$ lies on circle

$(x - x_0)^2 + (y - y_0)^2 = r^2$ and $(x - x_0)^2 + (y - y_0)^2 = 4r^2$ respectively. If

$z_0 = x_0 + iy_0$ satisfies the equation $2|z_0|^2 = r^2 + 2$ then $|\alpha|$ is equal to (a)

- (a) $\frac{1}{\sqrt{2}}$ (b) $\frac{1}{2}$ (c) $\frac{1}{\sqrt{7}}$ (d) $\frac{1}{3}$

A. $\frac{1}{\sqrt{2}}$

B. $\frac{1}{2}$

C. $\frac{1}{\sqrt{7}}$

D. $\frac{1}{3}$

Answer: C



Watch Video Solution

118. Let $w = \left(\sqrt{3} + \frac{i}{2}\right)$ and $P = \{w^n : n = 1, 2, 3, \dots\}$, Further

$H_1 = \left\{z \in \mathbb{C} : \operatorname{Re}(z) > \frac{1}{2}\right\}$ and $H_2 = \left\{z \in \mathbb{C} : \operatorname{Re}(z) < -\frac{1}{2}\right\}$ Where \mathbb{C} is

set of all complex numbers. If $z_1 \in P \cap H_1, z_2 \in P \cap H_2$ and O represent the origin, then $\angle Z_1 O Z_2 =$

A. $\frac{\pi}{2}, \frac{5\pi}{6}$

B. $\pi, \frac{2\pi}{3}$

C. $\frac{2\pi}{3}, \frac{5\pi}{3}$

D. $\frac{5\pi}{3}, \frac{7\pi}{3}$

Answer: B



Watch Video Solution

119. Let $S = S_1 \cap S_2 \cap S_3$, where

$$S_1 = \{z \in \mathbb{C} : |z| < 4\}, S_2 = \left\{ z \in \mathbb{C} : \ln \left[\frac{z - 1 + \sqrt{3}i}{1 - \sqrt{31}} \right] > 0 \right\} \text{ and } S_3 = \{z \in \mathbb{C} : \operatorname{Re} z > 0\}$$

A. $\frac{10\pi}{3}$

B. $\frac{20\pi}{3}$

C. $\frac{16\pi}{3}$

D. $\frac{32\pi}{3}$

Answer: B



Watch Video Solution

120. Let $S = S_1 \cap S_2 \cap S_3$, where

$$S_1 = \{z \in \mathbb{C} : |z| < 4\}, S_2 = \left\{ z \in \mathbb{C} : \ln \left[\frac{z - 1 + \sqrt{3}i}{1 - \sqrt{31}} \right] > 0 \right\} \text{ and } S_3 = \{z \in \mathbb{C} : \operatorname{Re} z > 0\}$$

$$A. \frac{2 - \sqrt{3}}{2}$$

$$B. \frac{2 + \sqrt{3}}{2}$$

$$C. \frac{3 - \sqrt{3}}{2}$$

$$D. \frac{3 + \sqrt{3}}{2}$$

Answer: C



Watch Video Solution

121. Let $z_k = \frac{\cos(2k\pi)}{10} + i \frac{\sin(2k\pi)}{10}$, $k = 1, 2, \dots, 9$. Then,

$\frac{1}{10} \{ |1 - z_1| |1 - z_2| \dots |1 - z_9| \}$ equals

A. 0

B. 1

C. 2

D. 3

Answer: B



[Watch Video Solution](#)

122. In Q. No. 121, $1 - \sum_{k=1}^9 \frac{\cos(2k\pi)}{10}$ equals

A. 0

B. 1

C. 2

D. 10

Answer: C



[Watch Video Solution](#)

123. If z is a complex number such that $|z| \geq 2$ then the minimum value of

$$\left| z + \frac{1}{2} \right|$$
 is

A. is strictly greater than $\frac{5}{2}$

B. is strictly greater than $\frac{3}{2}$ but less than $\frac{5}{2}$

C. is equal to $\frac{5}{2}$

D. lies in the interval (1,2)

Answer: D

 [Watch Video Solution](#)

124. A complex number z is said to be uni-modular if $|z| = 1$. Suppose z_1 and z_2 are complex numbers such that $\frac{z_1 - 2z_2}{2 - z_1\bar{z}_2}$ is uni-modular and z_2 is not uni-modular. Then the point z_1 lies on a:

A. circle of radius 2

B. circle of radius $\sqrt{2}$

C. straight line parallel to x-axis.

D. straight line parallel to y-axis.

Answer: A



Watch Video Solution

125. If $|z - 2 - i| = |z| \sin\left(\frac{\pi}{4} - \arg z\right)$, where $i = \sqrt{-1}$, then locus of z , is

A. pair of straight lines

B. circle of radius $\sqrt{2}$

C. parabola

D. ellipse

Answer: C



Watch Video Solution

126. $f(n) = \cot^2\left(\frac{\pi}{n}\right) + \cot^2 \frac{2\pi}{n} + \dots + \cot^2 \frac{(n-1)\pi}{n}$, ($n > 1, n \in N$)

then $\lim_{n \rightarrow \infty} \frac{f(n)}{n^2}$ is equal to (A) $\frac{1}{2}$ (B) $\frac{1}{3}$ (C) $\frac{2}{3}$ (D) 1

A. $\frac{1}{2}$

B. $\frac{1}{3}$

C. $\frac{2}{3}$

D. 1

Answer: B

 [Watch Video Solution](#)

127. If z_1 and z_2 are lying on $|z - 3| \leq 4$ and $|z - 1| = |z + 1| = 3$ respectively.

Then $d = |z_1 - z_2|$ satisfies.

A. $0 \leq d < \frac{15}{2}$

B. $0 < d \leq \frac{15}{2}$

C. $0 \leq d \leq \frac{17}{2}$

D. $0 < d < \frac{17}{2}$

Answer: C



Watch Video Solution

128. If $|z - 1| = 1$ and $\arg(z) = \theta$, where $z \neq 0$ and θ is acute, then $\left(1 - \frac{2}{z}\right)$ is equal to

A. $\tan\theta$

B. $I\tan\theta$

C. $\frac{\tan\theta}{2}$

D. $I\frac{\tan\theta}{2}$

Answer: B



Watch Video Solution

129. If z is a complex number lying in the first quadrant such that $\operatorname{Re}(z) + \operatorname{Im}(z) = 3$, then the maximum value of $\{\operatorname{Re}(z)\}^2 \operatorname{Im}(z)$, is

A. 1

B. 2

C. 3

D. 4

Answer: D



Watch Video Solution

130. The maximum area of the triangle formed by the complex coordinates z, z_1, z_2 which satisfy the relations $|z - z_1| = |z - z_2|$ and

$$\left| z - \frac{z_1 + z_2}{2} \right| \leq r, \text{ where } r > |z_1 - z_2| \text{ is}$$

A. $\frac{1}{2} |z_1 - z_2|^2$

B. $\frac{1}{2} |z_1 - z_2| r$

C. $\frac{1}{2} |z_1 - z_2|^2 r^2$

D. $\frac{1}{2} |z_1 - z_2| r^2$

Answer: B



[Watch Video Solution](#)

131. If z is a complex number satisfying $|z|^2 + 2(z + 2) + 3i(z - \bar{z}) + 4 = 0$, then complex number $z + 3 + 2i$ lies on

- A. circle with center $1-5i$ and radius 4
- B. circle with center $1+5i$ and radius 4
- C. circle with center $1+5i$ and radius 3
- D. circle with center $1-5i$ and radius 3

Answer: B



[View Text Solution](#)

132. Locus of z if $\arg[z - (1 + i)] = \begin{cases} \frac{3\pi}{4} & \text{when } |z| \leq |z - 2| \\ \frac{-\pi}{4} & \text{when } |z| > |z - 4| \end{cases}$ is

A. a straight line passing through (2,0)

B. a straight line passing through (2,0) and (1,1)

C. a line segment

D. a set of two rays

 [Watch Video Solution](#)

133. Let $z \in \mathbb{C}$ and if $A = \left\{ z : \arg(z) = \frac{\pi}{4} \right\}$ and $B = \left\{ z : \arg(z - 3 - 3i) = \frac{2\pi}{3} \right\}$

. Then $n(A \cap B) =$

A. 1

B. 2

C. 3

D. 0

Answer: D

[Watch Video Solution](#)

134. Let $S = \{z \in \mathbb{C} : z \left(iz_1 + 1, |z_1| < 1 \right)\}$. Then, for all $z \in S$, which one of the following is always true?

A. $\operatorname{Re}(z) - \operatorname{Im}(z) < 0$

B. $\operatorname{Re}(z) + \operatorname{Im}(z) < 0$

C. $\operatorname{Re}(z) < 0$

D. $\operatorname{Re}(z) - \operatorname{Im}(z) > 0$

Answer: A

[Watch Video Solution](#)

135. Let $z = 1 + ai$ be a complex number, $a > 0$, such that z^3 is a real number. Then the sum $1 + z + z^2 + \dots + z^{11}$ is equal to:

A. $-1250\sqrt{3}i$

B. $1250\sqrt{3}i$

C. $-1365\sqrt{3}i$

D. $1365\sqrt{3}i$

Answer: C



Watch Video Solution

136. Let $0 \neq a, 0 \neq b \in R$. Suppose

$$S = \left\{ z \in C, z = \frac{1}{a + ibt} t \in R, t \neq 0 \right\}, \text{ where } i = \sqrt{-1}. \text{ If } z = x + iy \text{ and}$$

$z \in S$, then (x, y) lies on

A. on the circle with radius $\frac{1}{2a}$ and center $\left(-\frac{1}{2a}, 0 \right)$

B. on the circle with radius $\frac{1}{2a}$ and center $\left(\frac{1}{2a}, 0 \right)$

C. on the x -axis

D. on the y -axis.

Answer: B



Watch Video Solution

137. Let $a, b \in \mathbb{R}$ and $a^2 + b^2 \neq 0$. Suppose

$S = \left\{ z \in \mathbb{C} : z = \frac{1}{a + ibt}, t \in \mathbb{R}, t \neq 0 \right\}$, where $i = \sqrt{-1}$. If $z = x + iy$ and z in

S , then (x, y) lies on

A. the x -axis for $a \neq 0, b = 0$

B. the y -axis for $a \neq 0, b = 0$

C. the y -axis for $a \neq 0, b \neq 0$

D. the x - axis for $a=0, b \neq 0$



Watch Video Solution

138. Let $a, b \in \mathbb{R}$ and $a^2 + b^2 \neq 0$. Suppose

$S = \left\{ z \in \mathbb{C} : z = \frac{1}{a + ibt}, t \in \mathbb{R}, t \neq 0 \right\}$, where $i = \sqrt{-1}$. If $z = x + iy$ and z in

S , then (x, y) lies on

A. $a = 0, b \neq 0$

B. $a \neq 0, b = 0$

C. $a \neq 0, b \neq 0$

D. all $a, b \in \mathbb{R}$



Watch Video Solution

139. The point represented by $2 + i$ in the Argand plane moves 1 unit eastwards, then 2-units northwards and finally from there $2\sqrt{2}$ units in the south-westwards direction. Then its new position in the Argand plane is at the point represented by

A. $2 + 2i$

B. $-2 - 2i$

C. $1 + i$

D. $-1 - i$

Answer: C



Watch Video Solution

140. Let ω be a complex number such that $2\omega + 1 = \sqrt{3}i$. If

$$\begin{vmatrix} 1 & 1 & 1 \\ 1 & -\omega^2 - 1 & \omega^2 \\ 1 & \omega^2 & \omega^7 \end{vmatrix} = 3k, \text{ then } k \text{ is equal to}$$

A. -1

B. 1

C. $-i\sqrt{3}$

D. $i\sqrt{3}$

Answer: C



Watch Video Solution

141. Let a, b, x and y be real numbers such that $a - b = 1$ and $y \neq 0$. If the

complex number $z = x + iy$ satisfies $\operatorname{Im}\left(\frac{az + b}{z + 1}\right) = y$, then which of the

following is (are) possible value(s) of x ? (a) $-1 - \sqrt{1 - y^2}$ (b) $1 + \sqrt{1 + y^2}$

(c) $-1 + \sqrt{1 - y^2}$ (d) $-1 - \sqrt{1 + y^2}$

A. $-1 - \sqrt{1 - y^2}$

B. $1 + \sqrt{1 + y^2}$

C. $1 - \sqrt{1 + y^2}$

D. $-1 + \sqrt{1 - y^2}$

Answer: A::D



Watch Video Solution

1. Statement-1, For any two complex numbers z_1 and z_2

$$\left| z_1 + \sqrt{z_1^2 - z_2^2} \right| + \left| z_1 - \sqrt{z_1^2 - z_2^2} \right| = \left| z_1 + z_2 \right| + \left| z_1 - z_2 \right|$$

Statement-2: For any two complex numbers z_1 and z_2

$$\left| z_1 + z_2 \right|^2 + \left| z_1 - z_2 \right|^2 = 2 \left(\left| z_1 \right|^2 + \left| z_2 \right|^2 \right)$$

- A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.
- B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.
- C. Statement-1 is True, statement-2 is false,
- D. statement-1 is False, Statement-2 is true.

Answer: a



View Text Solution

2. Statement-1: for any two complex numbers z_1 and z_2

$$|z_1 + z_2|^2 \leq \left(1 + \frac{1}{\lambda}\right) |z_2|^2, \text{ where } \lambda \text{ is a positive real number.}$$

Statement-2 $AM \geq GM$.

- A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.
- B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.
- C. Statement-1 is True, statement-2 is false,
- D. statement-1 is False, Statement-2 is true.

Answer: a



[Watch Video Solution](#)

3. Statement-1, If $z_1, z_2, z_3, \dots, z_n$ are uni-modular complex numbers, then

$$\left| z_1 + z + (2) + \dots + z_n \right| = \left| \frac{1}{z_1} + \frac{1}{z_2} + \dots + \frac{1}{z_n} \right|$$

Statement-2: For any complex number z , $z\bar{z} = |z|^2$

- A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.
- B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.
- C. Statement-1 is True, statement-2 is false,
- D. statement-1 is False, Statement-2 is true.

Answer: b

 [Watch Video Solution](#)

4. Statement-1, if z_1 and z_2 are two complex numbers such that

$$\left| z_1 \right| \leq 1, \left| z_2 \right| \leq 1, \text{ then}$$

$$|z_1 - z_2|^2 \leq (|z_1| - |z_2|)^2 + \arg(z_2)^2$$

Statement-2 $\sin\theta > \theta$ for all $\theta > 0$.

- A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.
- B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.
- C. Statement-1 is True, statement-2 is false,
- D. statement-1 is False, Statement-2 is true.

Answer: c

 [Watch Video Solution](#)

5. Statement -1: for any complex number z , $|\operatorname{Re}(z)| + |\operatorname{Im}(z)| \leq |z|$

Statement-2: $|\sin\theta| \leq 1$, for all θ

A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. Statement-1 is True, statement-2 is false,

D. statement-1 is False, Statement-2 is true.

Answer: d



Watch Video Solution

6. Statement-1: for any non-zero complex number z , $\left| \frac{z}{|z|} - 1 \right| \leq \arg(z)$

Statement-2 : $\sin\theta \leq \theta$ for $\theta \geq 0$

A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. Statement-1 is True, statement-2 is false,

D. statement-1 is False, Statement-2 is true.

Answer: a

 [Watch Video Solution](#)

7. Statement-1: for any non-zero complex number $|z - 1| \leq ||z| - 1| + |z| \arg(z)$

Statement-2 : For any non-zero complex number z

$$\left| \frac{z}{|z|} - 1 \right| \leq \arg(z)$$

A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. Statement-1 is True, statement-2 is false,

D. statement-1 is False, Statement-2 is true.

Answer: a

 [Watch Video Solution](#)

8. Statement-1: If z_1, z_2 are affixes of two fixed points A and B in the Argand plane and P(z) is a variable point such that " $\arg \frac{z - z_1}{z - z_2} = \frac{\pi}{2}$ ", then the locus of z is a circle having z_1 and z_2 as the end-points of a diameter.

Statement-2 : $\arg \frac{z_2 - z_1}{z_1 - z} = \angle APB$

A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. Statement-1 is True, statement-2 is false,

D. statement-1 is False, Statement-2 is true.

Answer: d

 [Watch Video Solution](#)

9. Statement-1: If z is a complex number satisfying $(z - 1)^n = z^n$, $n \in \mathbb{N}$, then the locus of z is a straight line parallel to imaginary axis.

Statement-2: The locus of a point equidistant from two given points is the perpendicular bisector of the line segment joining them.

A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. Statement-1 is True, statement-2 is false,

D. statement-1 is False, Statement-2 is true.

Answer: a



Watch Video Solution

10. Let z_0 be the circumcenter of an equilateral triangle whose affixes are z_1, z_2, z_3 .

Statement-1 : $z_1^2 + z_2^2 + z_3^2 = 3z_0^2$

Statement-2: $z_1^2 + z_2^2 + z_3^2 = 2(z_1z_2 + z_2z_3 + z_3z_1)$

A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. Statement-1 is True, statement-2 is false,

D. statement-1 is False, Statement-2 is true.

Answer: c



Watch Video Solution

11. Let z_1 and z_2 be the roots of the equation $z^2 + pz + q = 0$. Suppose z_1 and z_2 are represented by points A and B in the Argand plane such that $\angle AOB = \alpha$, where O is the origin.

Statement-1: If $OA=OB$, then $p^2 = 4q \frac{\cos^2 \alpha}{2}$

Statement-2: If affix of a point P in the Argand plane is z , then ze^{ia} is represented by a point Q such that $\angle POQ = \alpha$ and $OP = OQ$.

A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. Statement-1 is True, statement-2 is false,

D. statement-1 is False, Statement-2 is true.

Answer: a



Watch Video Solution

12. Statement-1: The locus of point z satisfying $\left| \frac{3z + i}{2z + 3 + 4i} \right| = \frac{3}{2}$ is a straight line.

Statement-2 : The locus of a point equidistant from two fixed points is a straight line representing the perpendicular bisector of the segment joining the given points.

- A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.
- B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.
- C. Statement-1 is True, statement-2 is false,
- D. statement-1 is False, Statement-2 is true.

Answer: a



Watch Video Solution

13. Statement-1: If a, b, c are distinct real number and $\omega (\neq 1)$ is a cube root

of unity, then $\left| \frac{a + b\omega + c\omega^2}{a\omega^2 + b + c\omega} \right| = 1$ **Statement-2:** For any non-zero complex

number $z, |z / \bar{z}| = 1$

A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. Statement-1 is True, statement-2 is false,

D. statement-1 is False, Statement-2 is true.

Answer: b



Watch Video Solution

14. Let z be a unimodular complex number.

Statement-1: $\arg(z^2 + \bar{z}) = \arg(z)$

Statement-2: $\bar{z} = \cos(\arg z) - i\sin(\arg z)$

A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. Statement-1 is True, statement-2 is false,

D. statement-1 is False, Statement-2 is true.

Answer: d



Watch Video Solution

15. Let z and ω be complex numbers such that $|z| = |\omega|$ and $\arg(z)$ denote the principal of z .

Statement-1: If $\arg z + \arg \omega = \pi$, then $z = -\bar{\omega}$

Statement -2: $|z| = |\omega|$ implies $\arg z - \arg \bar{\omega} = \pi$, then $z = -\bar{\omega}$

A. Statement-1 is True, Statement-2 is True: Statement-2 is a correct explanation for statement-1.

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. Statement-1 is True, statement-2 is false,

D. statement-1 is False, Statement-2 is true.

Answer: c



[Watch Video Solution](#)

Exercise

1. Which of the following is correct?

A. $1 + i > 2 - i$

B. $2 + i > 1 + i$

C. $2 - i > 1 + i$

D. none of these

Answer: D



[Watch Video Solution](#)

2. If $a = \sqrt{2}i$, then which of the following is correct?

A. $a = 1 + i$

B. $a = 1 - i$

C. $a = -2(\sqrt{2})i$

D. none of these

Answer: A



[Watch Video Solution](#)

3. Let z_1, z_2 be two complex numbers such that $z_1 + z_2$ and $z_1 z_2$ both are real, then

A. $z_1 = -z_2$

B. $z_1 = \bar{z}_2$

C. $z_1 = -\bar{z}_2$

D. $z_1 = z_2$

Answer: B



[Watch Video Solution](#)

4. If the complex numbers z_1, z_2, z_3 are in AP, then they lie on

A. a circle

B. a parabola

C. a line

D. an ellipse

Answer: C



Watch Video Solution

5. The locus of complex number z for which $\left(\frac{z-1}{z+1}\right) = k$, where k is non-zero real, is

A. a circle with center on y -axis

B. a circle with center on x -axis

C. a straight line parallel to y -axis

D. a straight line making $\pi/3$ angle with the x -axis.

Answer: c



Watch Video Solution

6. The locus of the points z satisfying the condition $\arg\left(\frac{z-1}{z+1}\right) = \frac{\pi}{3}$ is, a

A. parabola

B. Statement-1 is true, statement -2 is true, Statement-2 is not a correct explanation for statement-1.

C. circle

D. pair of straight line

Answer: a



Watch Video Solution

7. If $\sqrt{x+iy} = \pm(a+ib)$, then find $\sqrt{-x-iy}$.

A. $\pm(b+ia)$

B. $\pm(a - ib)$

C. $\pm(b - ia)$

D. $\pm(a + ib)$

Answer: C



Watch Video Solution

8. The locus of the points z satisfying the condition $\arg\left(\frac{z-1}{z+1}\right) = \frac{\pi}{3}$ is, a

A. parabola

B. circle

C. pair of straight lines

D. none of these

Answer: d



Watch Video Solution

9. If $(\sqrt{3} + i)^{10} = a + ib$, then a and b are respectively

A. 128 & $128\sqrt{3}$

B. 64 and $64\sqrt{3}$

C. 512 and $512\sqrt{3}$

D. none of these

Answer: C



Watch Video Solution

10. If $\operatorname{Re}\left(\frac{z - 8i}{z + 6}\right) = 0$, then lies on the curve

A. $x^2 + y^2 + 6x - 8y = 0$

B. $4x - 3y + 24 = 0$

C. $x^2 + y^2 - 8 = 0$

D. none of these

Answer: A



Watch Video Solution

11. If $z = \left(\frac{\sqrt{3}}{2} + \frac{i}{2}\right)^5 + \left(\frac{\sqrt{3}}{2} - \frac{i}{2}\right)^5$, then

A. $\text{Re}(z)=0$

B. $\text{Im}(z)=0$

C. $\text{Re}(z) > 0, \text{Im}(z) > 0$

D. $\text{Re}(z) > 0, \text{Im}(z) < 0$

Answer: B



Watch Video Solution

12. If $z = x + iy$ and $\omega = \frac{1 - iz}{z - i}$, then $|\omega| = 1$ implies that in the complex plane

A. z lies on imaginary axis

B. z lies on real axis

C. z lies on unit circle

D. none of these

Answer: b



[Watch Video Solution](#)

13. Let $3 - i$ and $2 + i$ be affixes of two points A and B in the Argand plane and P represents the complex number $z = x + iy$. Then, the locus of the P if $|z - 3 + i| = |z - 2 - i|$, is

A. circle on AB as diameter

B. the line AB

C. the perpendicular bisector of AB

D. none of these

Answer: c



Watch Video Solution

14. POQ is a straight line through the origin O,P and Q represent the complex numbers $a+ib$ and $c+id$ respectively and $OP=OQ$. Then, which one of the following is true?

A. $|a + ib| = |c + id|$

B. $a + b = c + d$

C. $\arg(a + ib) = \arg(c + id)$

D. none of these

Answer: a



Watch Video Solution

15. If $z_1 = a + ib$ and $z_2 = c + id$ are complex numbers such that $|z_1| = |z_2| = 1$ and $\operatorname{Re}(z_1 \bar{z}_2) = 0$, then the pair of complex numbers $\omega = a + ic$ and $\omega_2 = b + id$ satisfies

A. $|\omega_1| = 1$

B. $|\omega_2| = 1$

C. $\operatorname{Re}(\omega_1 \omega_2^{-2}) = 0$

D. all of these

Answer: d



Watch Video Solution

16. Let z_1 and z_2 be complex numbers such that $z_1 \neq z_2$ and $|z_1| = |z_2|$. If z_1 has positive real part and z_2 has negative imaginary part, then $\frac{z_1 + z_2}{z_1 - z_2}$ may be zero (b) real and positive real and negative (d) purely imaginary

A. cannot be zero

B. is real and positive

C. is real and negative

D. is purely imaginary

Answer: d



Watch Video Solution

$$17. \sum_{k=1}^6 \left(\frac{\sin(2\pi k)}{7} - i \frac{\cos(2\pi k)}{7} \right) = ?$$

A. -1

B. 0

C. $-i$

D. i

Answer: D



Watch Video Solution

18. The equation $\bar{b}z + b\bar{z} = c$, where b is a non-zero complex constant and c is a real number, represents

- A. a circle
- B. a straight line
- C. a pair of straight line
- D. none of these

Answer: b



[Watch Video Solution](#)

19. If $|a_i| < 1$, $\lambda_i \geq 0$ for $i = 1, 2, \dots, n$ and $\lambda_1 + \lambda_2 + \dots + \lambda_n = 1$, then the value of $|\lambda_1 a_1 + \lambda_2 a_2 + \dots + \lambda_n a_n|$ is

- A. equal to 1
- B. less than 1
- C. greater than 1

D. none of these

Answer: b

 [Watch Video Solution](#)

20. For any two complex numbers, z_1, z_2 and any two real numbers a and

$$b, \left| az_1 - bz_2 - (2) \right|^2 + \left| bz_1 + az_2 \right|^2 =$$

A. $(a + b) \left(|z_1|^2 + |z_2|^2 \right)$

B. $(a^2 + b^2) \left(|z_1|^2 + |z_2|^2 \right)$

C. $(a^2 + b^2) \left(|z_1| + |z_2| \right)$

D. none of these

Answer: B

 [Watch Video Solution](#)

21. Common roots of the equation $z^3 + 2z^2 + 2z + 1 = 0$ and $z^{2020} + z^{2018} + 1 = 0$, are

- A. ω, ω^2
- B. $1, \omega, \omega^2$
- C. $-1, \omega, \omega^2$
- D. $-\omega, -\omega^2$

Answer: a



[Watch Video Solution](#)

22. If z_1 and z_2 are two complex numbers such that $\left| \frac{z_1 - z_2}{1 - \bar{z}_1 z_2} \right| = 1$, then

which one of the following is true?

- A. $|z_1| = 1, |z_2| = 1$
- B. $z_1 = e^{i\theta}, \theta \in R$

C. $z_2 = e^{i\theta}, \theta \in R$

D. all of these

Answer: b



Watch Video Solution

23. The points representing cube roots of unity

A. are collinear

B. lie on a circle of radius $\sqrt{3}$

C. form an equilateral triangle

D. none of these

Answer: c



Watch Video Solution

24. If z_1 and z_2 are two complex numbers such that $\left| \frac{z_1 - z_2}{z_1 + z_2} \right| = 1$, then

A. $z_1 = kz_2, k \in R$

B. $z_1 = ikz_2, k \in R$

C. $z_1 = z_2$

D. none of these

Answer: B



Watch Video Solution

25. If z_1, z_2 are two complex numbers such that $\left| \frac{z_1 - z_2}{z_1 + z_2} \right| = 1$ and

$iz_1 = Kz_2$, where $K \in R$, then the angle between $z_1 - z_2$ and $z_1 + z_2$ is

A. $\frac{\tan^{-1}(2k)}{k^2 + 1}$

B. $\frac{\tan^{-1}(2k)}{1 - k^2}$

C. $-2\tan^{-1}k$

D. none of these

Answer: c



[Watch Video Solution](#)

26. If n is a positive integer greater than unity z is a complex number satisfying the equation $z^n = (z + 1)^n$, then

A. $\operatorname{Re}(z) < 0$

B. $\operatorname{Re}(z) > 0$

C. $\operatorname{Re}(z) = 0$

D. none of these

Answer: A



[Watch Video Solution](#)

27. If n is a positive integer greater than unity z is a complex number satisfying the equation $z^n = (z + 1)^n$, then

A. $\text{Im}(z) < 0$

B. $\text{Im}(z) > 0$

C. $\text{Im}(z) = 0$

D. none of these

Answer: d



Watch Video Solution

28. If at least one value of the complex number $z = x + iy$ satisfies the condition $|z + \sqrt{2}| = \sqrt{a^2 - 3a + 2}$ and the inequality $|z + i\sqrt{2}| < a$, then

A. $a > 2$

B. $a = 2$

C. $a < 2$

D. $a > 1$

Answer: a



[Watch Video Solution](#)

29. Given z is a complex number with modulus 1. Then the equation $[(1 + ia)/(1 - ia)]^4 = z$ has all roots real and distinct two real and two imaginary three roots two imaginary one root real and three imaginary

- A. all roots, real and distinct
- B. two real and two imaginary
- C. three roots real and one imaginary
- D. one root real and three imaginary

Answer: a



[Watch Video Solution](#)

30. The center of a regular polygon of n sides is located at the point $z=0$, and one of its vertex z_1 is known. If z_2 be the vertex adjacent to z_1 , then z_2 is equal to

A. $z_1 \left(\cos 2\frac{\pi}{n} \pm i \sin 2\frac{\pi}{n} \right)$

B. $z_1 \left(\frac{\cos \pi}{n} \pm i \frac{\sin \pi}{n} \right)$

C. $z_1 \left(\frac{\cos \pi}{2n} \pm \frac{\sin \pi}{2n} \right)$

D. none of these

Answer: a



Watch Video Solution

31. If the points z_1, z_2, z_3 are the vertices of an equilateral triangle in the Argand plane, then which one of the following is not correct?

A. $\frac{1}{z_1 - z_2} + \frac{1}{z_2 - z_3} + \frac{1}{z_3 - z_1} = 0$

B. $z_1^2 + z_2^2 + z_3^2 = z_1z_2 + z_2z_3 + z_3z_1$

$$C. (z_1 - z_2)^2 + (z_2 - z_3)^2 + (z_3 - z_1)^2 = 0$$

$$D. z_1^3 + z_2^3 + z_3^3 + 3z_1z_2z_3 = 0$$

Answer: d



Watch Video Solution

32. For any complex number z , the minimum value of $|z| + |z - 1|$

A. $\text{Re}(z) < 0$

B. $\text{Re}(z) > 0$

C. $\text{Re}(z) > 2$

D. $\text{Re}(z) > 3$

Answer: a



Watch Video Solution

33. The inequality $|z - 4| < |z - 2|$ represents

A. $\operatorname{Re}(z) \geq 0$

B. $\operatorname{Re}(z) < 0$

C. $\operatorname{Re}(z) > 0$

D. None of these

Answer: d



Watch Video Solution

34. Number of non-zero integral solution of the equation $|1 - i|^n = 2^n$, is

A. 1

B. 2

C. infinite

D. none of these

Answer: D



Watch Video Solution

35. If $\text{Im} \frac{2z + 1}{iz + 1} = -2$, then locus of z , is

- A. a circle
- B. a parabola
- C. a straight line
- D. none of these

Answer: A



Watch Video Solution

36. If $z (\neq -1)$ is a complex number such that $\frac{z - 1}{z + 1}$ is purely imaginary, then $|z|$ is equal to

A. 1

B. 2

C. 3

D. 4

Answer: a



Watch Video Solution

37. If $x = -5 + 2\sqrt{-4}$, find the value of $x^4 + 9x^3 + 35x^2 - x + 4$.

A. 0

B. -160

C. 160

D. -164

Answer: b



Watch Video Solution

38. If z_1, z_2, z_3 are vertices of an equilateral triangle with z_0 its centroid, then $z_1^2 + z_2^2 + z_3^2 =$

A. z_0^2

B. $9z_0^2$

C. $3z_0^2$

D. $2z_0^2$

Answer: c



Watch Video Solution

39. If z_1, z_2 are two complex numbers such that $Im(z_1 + z_2) = 0, Im(z_1 z_2) = 0$, then:

A. $z_1 = -z_2$

B. $z_1 = z_2$

C. $z_1 = \bar{z}_2$

D. $z_1 = -\bar{z}_2$

Answer: c



Watch Video Solution

40. If $z^2 + z|z| + |z^2| = 0$, then the locus z is a. a circle b. a straight line c. a pair of straight line d. none of these

A. a circle

B. a straight line

C. a pair of straight lines

D. none of these

Answer: c



Watch Video Solution

41. If $\log \sqrt{3} \left(\frac{|z|^2 - |z| + 1}{2 + |z|} \right) > 2$, then the locus of z is

A. $|z| = 5$

B. $|z| < 5$

C. $|z| > 5$

D. none of these

Answer: c



Watch Video Solution

42. Let $g(x)$ and $h(x)$ are two polynomials such that the polynomial $P(x) = g(x^3) + xh(x^3)$ is divisible by $x^2 + x + 1$, then which one of the following is not true?

A. $g(1) = h(1) = 0$

B. $g(1) = h(1) \neq 0$

C. $g(1) = -h(1)$

D. $g(1) + h(1) = 0$

Answer: a



Watch Video Solution

43. If $g(x)$ and $h(x)$ are two polynomials such that the polynomials $P(x) = g(x^3) + xh(x^3)$ is divisible by $x^2 + x + 1$, then which one of the following is not true?

A. $g(1) = h(1) = 0$

B. $g(1) = h(1) \neq 0$

C. $g(1) = -h(1)$

D. $g(1) + h(1) = 0$

Answer: b



Watch Video Solution

44. If $|z_1| = |z_2| = |z_3|$ and $z_1 + z_2 + z_3 = 0$, then z_1, z_2, z_3 are vertices of

A. a right angled triangle

B. an equilateral triangle

C. isosceles triangle

D. scalene triangle

Answer: b

 [Watch Video Solution](#)

45. If $x_n = \cos\left(\frac{\pi}{3^n}\right) + i\sin\left(\frac{\pi}{3^n}\right)$, then $x_1, x_2, x_3, \dots, x_\infty$ is

equal to

A. 1

B. -1

C. i

D. $-i$

Answer: C



Watch Video Solution

46. If $(a_1 + ib_1)(a_2 + ib_2) \dots (a_n + ib_n) = A + iB$, then

$(a_1^2 + b_1^2)(a_2^2 + b_2^2) \dots (a_n^2 + b_n^2)$ is equal to (A) 1 (B) $(A^2 + B^2)$ (C) $(A + B)$

(D) $\left(\frac{1}{A^2} + \frac{1}{B^2}\right)$

A. 1

B. $A^2 + B^2$

C. $A + B$

D. $\frac{1}{A^2} + \frac{1}{B^2}$

Answer: b



Watch Video Solution

47. If $(a_1 + ib_1)(a_2 + ib_2) \dots (a_n + ib_n) = A + iB$, then

$\sum_{i=1}^n \tan^{-1} \left(\frac{b_i}{a_i} \right)$ is equal to

A. $\frac{B}{A}$

B. $\tan \left(\frac{B}{A} \right)$

C. $\tan^{-1} \left(\frac{B}{A} \right)$

D. $\tan^{-1} \left(\frac{A}{B} \right)$

Answer: c



Watch Video Solution

48. If $\cos\alpha + 2\cos\beta + 3\cos\gamma = \sin\alpha + 2\sin\beta + 3\sin\gamma = 0$ then the value of $\sin 3\alpha + 8\sin 3\beta + 27\sin 3\gamma$, is

A. $\sin(\alpha + \beta + \gamma)$

B. $3\sin(\alpha + \beta + \gamma)$

C. $18\sin(\alpha + \beta + \gamma)$

D. $\sin(\alpha + 2\beta + 3\gamma)$

Answer: c

 [Watch Video Solution](#)

49. If α, β, γ are the cube roots of p , then for any x, y, z $\frac{x\alpha + y\beta + z\gamma}{x\beta + y\gamma + z\alpha} =$

A. ω, ω^2

B. $-\omega, -\omega^2$

C. $1, -1$

D. none of these

Answer: a

 [Watch Video Solution](#)

50. Prove that $\tan\left(i(\log)_e\left(\frac{a-ib}{a+ib}\right)\right) = \frac{2ab}{a^2-b^2}$ (where $a, b \in \mathbb{R}^+$)

A. $\frac{ab}{a^2+b^2}$

B. $\frac{2ab}{a^2-b^2}$

C. $\frac{ab}{a^2-b^2}$

D. $\frac{2ab}{a^2+b^2}$

Answer: b



Watch Video Solution

51. Find the relation if z_1, z_2, z_3, z_4 are the affixes of the vertices of a parallelogram taken in order.

A. $z_1 + z_4 = z_2 + z_3$

B. $z_1 + z_3 = z_2 + z_4$

C. $z_1 + z_2 = z_3 + z_4$

D. none of these

Answer: b



Watch Video Solution

52. The locus of the points representing the complex numbers z for which

$$|z| - 2 = |z - i| - |z + 5i| = 0, \text{ is}$$

- A. a circle with center at the origin
- B. a straight line passing through the origin
- C. the angle point $(0, -2)$
- D. none of these

Answer: c



Watch Video Solution

53. For $n = 6k, k \in \mathbb{Z}$, $\left(\frac{1 - i\sqrt{3}}{2}\right)^n + \left(\frac{-1 - i\sqrt{3}}{2}\right)^n$ has the value

A. -1

B. 0

C. 1

D. 2

Answer: d



Watch Video Solution

54. The product of all values of $(\cos\alpha + i\sin\alpha)^{3/5}$ is

A. 1

B. $\cos\alpha + i\sin\alpha$

C. $\cos 3\alpha + i\sin 3\alpha$

D. $\cos 5\alpha + i\sin 5\alpha$

Answer: C



Watch Video Solution

55. If $C^2 + S^2 = 1$, then $\frac{1 + C + iS}{1 + C - iS}$ is equal to

A. $C + iS$

B. $C - iS$

C. $S + iC$

D. $S - iC$

Answer: a



Watch Video Solution

56. The center of a square ABCD is at $z=0$. The affix of the vertex A is z_1 .

Then, the affix of the centroid of the triangle ABC is

A. $z_1(\cos\pi \pm i\sin\pi)$

B. $\frac{z_1}{3}(\cos\pi \pm i\sin\pi)$

C. $z_1\left(\cos\left(\frac{\pi}{2}\right) \pm i\sin\left(\frac{\pi}{2}\right)\right)$

D. $\frac{z_1}{3}\left(\cos\left(\frac{\pi}{2}\right) \pm i\sin\left(\frac{\pi}{2}\right)\right)$

Answer: d



Watch Video Solution

57. The number of solutions of the system of equations $\operatorname{Re}(z^2) = 0, |z| = 2$, is

A. 4

B. 3

C. 2

D. 1

Answer: a



Watch Video Solution

58. The vector $z = -4 + 5i$ is turned counter clockwise through an angle of 180° and stretched 1.5 times. The complex number corresponding to the newly obtained vector is

A. $6 - \frac{15}{2}i$

B. $-6 + \frac{15}{2}i$

C. $6 + \frac{15}{2}i$

D. $6 + \frac{15}{2}i$

Answer: a



Watch Video Solution

59. The value of $\left[\sqrt{2} \left(\cos(56^\circ 15') + i \sin(56^\circ 15') \right) \right]^8$, is

A. $4i$

B. $8i$

C. $16i$

D. $-16i$

Answer: c



Watch Video Solution

60. Find the complex number z satisfying the equations

$$\left| \frac{z - 12}{z - 8i} \right| = \frac{5}{3}, \quad \left| \frac{z - 4}{z - 8} \right| = 1$$

A. 6

B. $6 \pm 8i$

C. $6 + 8i, 6 + 17i$

D. $8 \pm 6i$

Answer: c



[Watch Video Solution](#)

61. The vertices B and D of a parallelogram are $1 - 2i$ and $4 - 2i$. If the diagonals are at right angles and $AC = 2BD$, the complex number representing A is

A. $\frac{5}{2}$

B. $3i - \frac{3}{2}$

C. $3i - 4$

D. $3i + 4$

Answer: b



[Watch Video Solution](#)

62. If for complex numbers z_1 and z_2 , $\arg z_1 - \arg(z_2) = 0$ then $|z_1 - z_2|$ is equal to

A. $|z_1| + |z_2|$

B. $|z_1| - |z_2|$

C. $||z_1| - |z_2||$

D. 0

Answer: c



Watch Video Solution

63. The join of $z_1 = a + ib$ and $z_2 = \frac{1}{-a + ib}$ passes through

A. $z=0$

B. $z = 1 + i0$

C. $z = 0 + i$

D. $z = 1 + i$

Answer: a



Watch Video Solution

64. If z_1, z_2, z_3, z_4 are the affixes of four point in the Argand plane, z is the affix of a point such that $|z - z_1| = |z - z_2| = |z - z_3| = |z - z_4|$, then prove that z_1, z_2, z_3, z_4 are concyclic.

- A. concylic
- B. vertices of a triangle
- C. vertices of a rhombus
- D. in a straight line

Answer: a



Watch Video Solution

65. The value of $\sum_{r=1}^8 \left(\frac{\sin(2r\pi)}{9} + i \frac{\cos(2r\pi)}{9} \right)$, is

- A. -1

B. 1

C. i

D. $-i$

Answer: d



Watch Video Solution

66. If $z_1, z_2, z_3, \dots, z_n$ are n n th roots of unity, then for $k = 1, 2, \dots, n$

A. $|z_k| = k|z_n + 1|$

B. $|z_{k+1}| = k|z_k|$

C. $|z_{K+1}| = |z_k|z_{k+1}|$

D. $|z_k| = |z_{k+1}|$

Answer: d



Watch Video Solution

67. If z_1, z_2 and z_3, z_4 are two pairs of conjugate complex numbers then

$$\arg\left(\frac{z_1}{z_4}\right) + \arg\left(\frac{z_2}{z_3}\right) =$$

A. 0

B. $\pi/2$

C. $3\pi/2$

D. π

Answer: A



Watch Video Solution

68. If $|z_1| = |z_2|$ and $\arg(z_1) + \arg(z_2) = 0$, then

A. $z_1 = z_2$

B. $z_1 = \bar{z}_2$

C. $z_1 z_2 = 1$

D. $z_1 \bar{z}_2 = 1$

Answer: B



Watch Video Solution

69. If one vertex of a square whose diagonals intersect at the origin is $3(\cos\theta + i\sin\theta)$, then find the two adjacent vertices.

A. $\pm 3(\sin\theta - i\sin\theta)$

B. $\pm(\sin\theta + i\cos\theta)$

C. $\pm(\cos\theta - i\sin\theta)$

D. $z_1 \bar{z}_2 = 1$

Answer: a



Watch Video Solution

70. The value of z satisfying the equation

$$\log z + \log z^2 + \dots + \log z^n = 0, \text{ is}$$

- A. $\frac{\cos(4m\pi)}{n(n+1)} + i \frac{\sin(4m\pi)}{n(n+1)}, m = 1, 2, \dots$
- B. $\frac{\cos(4m\pi)}{n(n+1)} - i \frac{\sin(4m\pi)}{n(n+1)}, m = 1, 2, \dots$
- C. $\frac{\sin(4m\pi)}{n} + i \frac{\cos(4m\pi)}{n}, m = 1, 2, \dots$
- D. 0

Answer: a



Watch Video Solution

71. If $|z_1| = |z_2| = \dots = |z - (n)| = 1$, then the value of $|z_1 + z_2 + \dots + z_n|$, is

A. n

B. $\left| \frac{1}{z_1} + \frac{1}{z_2} + \dots + \frac{1}{z_n} \right|$

C. 0

D. none of these

Answer: b



[Watch Video Solution](#)

72. If $\omega (\neq 1)$ be a cube root of unity and $(1 + \omega)^7 = A + B\omega$, then A and B are respectively the numbers.

A. 0,1

B. 1,1

C. 1,0

D. -1, 1

Answer: b



[Watch Video Solution](#)

73. If ω is the complex cube root of unity then

$$\begin{vmatrix} 1 & 1+i+\omega^2 & \omega^2 \\ 1-i & -1 & \omega^2-1 \\ -i & -i+\omega-1 & -1 \end{vmatrix} =$$

A. 0

B. 1

C. i

D. ω

Answer: A



[Watch Video Solution](#)

74. Let z and ω be two non-zero complex numbers, such that $|z| = |\omega|$ and $\arg(z) + \arg(\omega) = \pi$. Then, z equals

A. ω

B. $-\omega$

C. $\bar{\omega}$

D. $-\bar{\omega}$

Answer: ad

 [Watch Video Solution](#)

75. If $z \neq 0$ be a complex number and $\arg(z) = \pi/4$, then

A. $\operatorname{Re}(z) = \operatorname{Im}(z)$ only

B. $\operatorname{Re}(z) = \operatorname{Im}(z) > 0$

C. $\operatorname{Re}(z^2) = \operatorname{Im}(z^2)$

D. none of these

Answer: b

 [Watch Video Solution](#)

76. $(1 + i)^8 + (1 - i)^8 = ?$

A. 2^8

B. 2^5

C. $2^4 \frac{\cos \pi}{4}$

D. $2^8 \frac{\cos \pi}{8}$

Answer: B



Watch Video Solution

77. What is the smallest positive integer n for which $(1 + i)^{2n} = (1 - i)^{2n}$?

A. 4

B. 8

C. 3

D. 12

Answer: C



Watch Video Solution

78. If α and β are different complex numbers with $|\beta| = 1$, then find $\left| \frac{\beta - \alpha}{1 - \bar{\alpha}\beta} \right|$.

A. 0

B. 8

C. 2

D. 2

Answer: c



Watch Video Solution

79. For any complex number z , the minimum value of $|z| + |z - 1|$

A. 1

B. 0

C. $1/2$

D. $3/2$

Answer: a



[Watch Video Solution](#)

80. If $\frac{3\pi}{2} > \alpha > 2\pi$, find the modulus and argument of $(1 - \cos 2\alpha) + i\sin 2\alpha$.

A. $-2\cos\alpha[\cos(\pi + \alpha) + i\sin(\pi + \alpha)]$

B. $2\cos\alpha[\cos\alpha + i\sin\alpha]$

C. $2\cos\alpha[\cos(\pi - \alpha) + i\sin(\pi - \alpha)]$

D. $-2\cos\alpha[\cos(\pi - \alpha) + i\sin(\pi - \alpha)]$

Answer: a



[Watch Video Solution](#)

81. If the roots of $(z - 1)^n = i(z + 1)^n$ are plotted in the Argand plane, then prove that they are collinear.

- A. lie on a parabola
- B. are concyclic
- C. are collinear
- D. the vertices of a triangle

Answer: b



[Watch Video Solution](#)

82. Area of the triangle formed by 3 complex numbers, $1 + i$, $i - 1$, $2i$, in the Argand plane, is

- A. $1/2$
- B. 1
- C. $\sqrt{2}$

D. 2

Answer: B



[Watch Video Solution](#)

83. If ω is a complex cube root of unity, then

$$(1 - \omega + \omega^2)^6 + (1 - \omega^2 + \omega)^6 =$$

A. 0

B. 6

C. 64

D. 128

Answer: D



[Watch Video Solution](#)

84. The locus represented by the equation $|z - 1| = |z - i|$ is

- A. a circle of radius 1
- B. an ellipse with foci at 1 and $-i$
- C. a line through the origin
- D. a circle on the line joining 1 and $-i$ as diameter.

Answer: C



[Watch Video Solution](#)

85. If $z = i \log(23)$, then $\cos z =$ -1 b. $-1/2$ c. 1 d. $1/2$

- A. i
- B. $2i$
- C. 1
- D. 2

Answer: d



Watch Video Solution

86.

If

$a = \cos\alpha + i\sin\alpha$, $b = \cos\beta + i\sin\beta$, $c = \cos\gamma + i\sin\gamma$ and $\frac{b}{c} + \frac{c}{a} + \frac{a}{b} = 1$,

then $\cos(\beta - \gamma) + \cos(\gamma - \alpha) + \cos(\alpha - \beta) =$

A. $3/2$

B. $-3/2$

C. 0

D. 1

Answer: D



Watch Video Solution

87. If z_1, z_2, z_3 are vertices of an equilateral triangle inscribed in the circle $|z| = 2$ and if $z_1 = 1 + i\sqrt{3}$, then

A. $z_2 = -2, z_3 = 1 - i\sqrt{3}$

B. $z_2 = 2, z_3 = 1 - i\sqrt{3}$

C. $z_2 = -2, z_3 = -1 - i\sqrt{3}$

D. $z_2 = 1 - i\sqrt{3}, z_3 = 1 - i\sqrt{3}$

Answer: a



Watch Video Solution

88. The general value of θ which satisfies the equation $(\cos\theta + i\sin\theta)(\cos3\theta + i\sin3\theta)(\cos5\theta + i\sin5\theta)\dots\dots\dots((\cos2n - 1)\theta + i\sin(2n - 1)\theta) = 1$ is

A. $\frac{r\pi}{n^2}$

B. $\frac{(r - 1)\pi}{n^2}$

C. $\frac{(2r + 1)\pi}{n^3}$

D. $\frac{2r\pi}{n^2}$

Answer: d



Watch Video Solution

89. If z is a complex numbers such that $z \neq 0$ and $\operatorname{Re}(z) = 0$, then

A. $\operatorname{Re}(z^2) = 0$

B. $\operatorname{Im}(z^2) = 0$

C. $\operatorname{Re}(z^2) = \operatorname{Im}(z^2)$

D. none of these

Answer: b



Watch Video Solution

90. If $z + z^{-1} = 1$, then find the value of $z^{100} + z^{-100}$.

A. i

B. $-i$

C. 1

D. -1

Answer: d



Watch Video Solution

91. Let A, B and C represent the complex number z_1, z_2, z_3 respectively on the complex plane. If the circumcentre of the triangle ABC lies on the origin, then the orthocentre is represented by the number

A. $z_1 + z_2 - z_3$

B. $z_2 + z_3 - z_1$

C. $z_3 + z_1 - z_2$

D. $z_1 + z_2 + z_3$

Answer: d



Watch Video Solution

92. Number of solutions of the equation $z^2 + |z|^2 = 0$, where $z \in \mathbb{C}$, is

A. 1

B. 2

C. 3

D. infinity many

Answer: D



Watch Video Solution

93. The number of solutions of the equation $z^2 + z = 0$ where z is a complex number, is

A. 2

B. 4

C. 6

D. none of these

Answer: b



[Watch Video Solution](#)

94. The centre of a square is at the origin and one of the vertex is $1 - i$ extremities of diagonal not passing through this vertex are

A. $1 - I, -1 + i$

B. $1 - I, -1 - i$

C. $-1 + I, -1 - i$

D. none of these

Answer: a

 [Watch Video Solution](#)

95. Let z and ω be two complex numbers such that $|z| \leq 1$, $|\omega| \leq 1$ and $|z - i\omega| = |z + i\omega| = 2$, then z equals 1 or i b. i or $-i$ c. 1 or -1 d. i or -1

A. 1 or i

B. i or $-i$

C. 1 or -1

D. i or -1

Answer: b

 [Watch Video Solution](#)

96. The system of equation $|z + 1 + i| = \sqrt{2}$ and $|z| = 3$ }, (where $i = \sqrt{-1}$)

has

- A. no solutions
- B. one solution
- C. two solution
- D. none of these

Answer: a



[Watch Video Solution](#)

97. The triangle with vertices at the point z_1z_2 , $(1 - i)z_1 + iz_2$ is

- A. right angled but not isoscles
- B. isosceles but not right angled
- C. right angled and isosceles
- D. equilateral

Answer: C



Watch Video Solution

98. Let α and β be two fixed non-zero complex numbers and 'z' a variable complex number. If the lines $\alpha\bar{z} + \bar{\alpha}z + 1 = 0$ and $\beta\bar{z} + \bar{\beta}z - 1 = 0$ are mutually perpendicular, then

A. $\alpha\beta + \bar{\alpha}\bar{\beta} = 0$

B. $\alpha\beta - \bar{\alpha}\bar{\beta} = 0$

C. $\bar{\alpha} - \alpha\bar{\beta} = 0$

D. $\alpha\bar{\beta} + \bar{\alpha}\beta = 0$

Answer: D



Watch Video Solution

99. The center of a square is at $z=0$. A is z_1 , then the centroid of the triangle ABC is

A. $z_1(\cos\pi \pm i\sin\pi)$

B. $\frac{1}{3}z_1(\cos\pi \pm i\sin\pi)$

C. $z_1\left(\cos\left(\frac{\pi}{2}\right) \pm i\sin\left(\frac{\pi}{2}\right)\right)$

D. $\frac{1}{3}z_1\left(\cos\left(\frac{\pi}{2}\right) \pm i\sin\left(\frac{\pi}{2}\right)\right)$

Answer: D



Watch Video Solution

100. If $z = x + iy$, then the equation $|(2z - i)/(z + 1)| = m$ represents a circle, then m can be 1/2 b. 1 c. 2 d. '3

A. 1/2

B. 1

C. 2

D. 3

Answer: c



[Watch Video Solution](#)

101. If $x^2 - 2x\cos\theta + 1 = 0$, then the value of $x^{2n} - 2x^n\cos n\theta + 1, n \in N$ is equal to

A. $\cos 2n\theta$

B. $\sin 2n\theta$

C. 0

D. $\cos n\theta$

Answer: C



[Watch Video Solution](#)

102. If $p^2 - p + 1 = 0$, then the value of p^{3n} can be

- A. 1
- B. -1
- C. 0
- D. $\cos n\theta$

Answer: d



Watch Video Solution

103. If $n \in \mathbb{Z}$, then $\frac{2^n}{(1+i)^{2n}} + \frac{(1+i)^{2n}}{2^n}$ is equal to

- A. 0
- B. 2
- C. $[1 + (-1)^n]i^n$
- D. 1

Answer: d



Watch Video Solution

104. If $\arg(z_1 z_2) = 0$ and $|z_1| = |z_2| = 1$, then

A. $z_1 + z_2 = 0$

B. $z_1 \bar{z}_2 = 1$

C. $z_1 = \bar{z}_2$

D. $z_1 + \bar{z}_2 = 0$

Answer: C



Watch Video Solution

105. If ω is a complex cube root of unity, then $\frac{(1+i)^{2n} - (1-i)^{2n}}{(1+\omega^4 - \omega^2)(1-\omega^4 + \omega^2)}$

is equal to

A. 0, if n is an even integer

B. 0 for all $n \in \mathbb{Z}$

C. $2^{n-1}i$ for all $n \in \mathbb{N}$

D. none of these

Answer: A



[Watch Video Solution](#)

106. If z is a complex number satisfying $z + z^{-1} = 1$ then $z^n + z^{-n}$, $n \in \mathbb{N}$, has the value

A. $2(-1)^n$, where n is a multiple of 3

B. $(-1)^n$, where n is not a multiple of 3

C. $(-1)^{n+1}$, where n is not a multiple of 3

D. none of these

Answer: a



Watch Video Solution

107. $x^{3m} + x^{3n-1} + x^{3r-2}$, where, $m, n, r \in N$ is divisible by

- A. m, n, k are rational
- B. m, n, k are integers
- C. m, n, k are positive integers
- D. none of these

Answer: b



Watch Video Solution

108. If z is nonreal root of $[-1]^{\frac{1}{7}}$ then, find the value of $z^{86} + z^{175} + z^{289}$

- A. 0
- B. -1
- C. 3

D. 1

Answer: B



[Watch Video Solution](#)

109. The locus of point z satisfying $\operatorname{Re}(z^2) = 0$, is

- A. a pair of straight lines
- B. a circle
- C. a rectangular hyperbola
- D. none of these

Answer: A



[Watch Video Solution](#)

110. The curve represented by $\text{Im}(z^2) = k$, where k is a non-zero real number, is

- A. a pair of straight line
- B. an ellipse
- C. a parabola
- D. a hyperbola

Answer: D



[Watch Video Solution](#)

111. If $\log_{\tan 30^\circ} \left[\frac{2|z|^2 + 2|z| - 3}{|z| + 1} \right] < -2$ then $|z| =$

- A. $|z| < 3/2$
- B. $|z| > 3/2$
- C. $|z| > 2$

D. $|z| < 2$



Watch Video Solution

112. The roots of the cubic equation $(z + \alpha\beta)^3 = \alpha^3$, α is not equal to 0, represent the vertices of a triangle of sides of length

A. $\frac{1}{\sqrt{3}}|\alpha\beta|$

B. $\sqrt{3}|\alpha|$

C. $\sqrt{3}|\beta|$

D. $\frac{1}{\sqrt{3}}|\alpha|$

Answer: cb



Watch Video Solution

113. The roots of the cubic equation $(z + \alpha\beta)^3 = \alpha^3$, α is not equal to 0, represent the vertices of a triangle of sides of length

- A. represent sides of an equilateral triangle
- B. represent the sides of an isosceles triangle
- C. represent the sides of a triangle whose one side is of length $\sqrt{3}\alpha$
- D. none of these

Answer: d



Watch Video Solution

114. If α, β, γ and δ are the equation $x^4 - 1 = 0$, then the value of

$$\frac{a\alpha + b\beta + c\gamma + d\delta}{a\gamma + b\delta + c\alpha + d\beta} + \frac{a\gamma + b\delta + c\alpha + d\beta}{a\alpha + b\beta + c\gamma + d\delta}, \text{ is}$$

- A. 3β
- B. 0
- C. 2γ

D. 3α

Answer: d



[View Text Solution](#)

115. If ω is a complex cube root of unity, then the equation

$|z - \omega|^2 + |z - \omega^2|^2 = \lambda$ will represent a circle, if

A. $\gamma \in (0, 3/2)$

B. $\gamma \in [3/2, \infty)$

C. $\gamma \in (0, 3)$

D. $\gamma \in [3, \infty)$

Answer: b



[Watch Video Solution](#)

116. If ω is a complex cube root of unity, then the equation

$$|z - \omega|^2 + |z - \omega^2|^2 = \lambda \text{ will represent a circle, if}$$

A. 4

B. 3

C. 2

D. $\sqrt{2}$

Answer: B



[Watch Video Solution](#)

117. The equation $z\bar{z} + (4 - 3i)z + (4 + 3i)\bar{z} + 5 = 0$ represents a circle of radius

A. 5

B. $2\sqrt{5}$

C. $5/2$

D. none of these

Answer: B



Watch Video Solution

118. z is such that $\arg\left(\frac{z - 3\sqrt{3}}{z + 3\sqrt{3}}\right) = \frac{\pi}{3}$ then locus z is

A. $|z - 3i| = 6$

B. $|z - 3i| = 6, \text{Im}(z) > 0$

C. $|z - 3i| = 6, \text{Im}(z) < 0$

D. none of these

Answer: B



Watch Video Solution

119. Let $z = 1 - t + i\sqrt{t^2 + t + 2}$, where t is a real parameter. the locus of the z in argand plane is

- A. a hyperbola
- B. an ellipse
- C. a straight line
- D. none of these

Answer: A



[Watch Video Solution](#)

120. If $|z - 4 + 3i| \leq 1$ and m and n be the least and greatest values of $|z|$ and K be the least value of $\frac{x^4 + x^2 + 4}{x}$ on the interval $(0, \infty)$, then $K =$

- A. m
- B. n
- C. $m + n$

D. mn

Answer: b

 [Watch Video Solution](#)

121. If $1, \alpha, \alpha^2, \dots, \alpha^{n-1}$ are the n, n^{th} roots of unity and z_1 and z_2 are any two complex numbers such that $\sum_{r=0}^{n-1} |z_1 + \alpha^r z_2|^2 = \lambda (|z_1|^2 + |z_2|^2)$, then $\lambda =$

A. n

B. $(n - 1)$

C. $(n + 1)$

D. 2n

Answer: a

 [Watch Video Solution](#)

122. If z_r ($r = 0, 1, 2, \dots, 6$) be the roots of the equation

$$(z + 1)^7 + z^7 = 0, \text{ then } \sum_{r=0}^6 \operatorname{Re}(z_r) =$$

- A. 0
- B. $3/2$
- C. $7/2$
- D. $-7/2$

Answer: D



[Watch Video Solution](#)

123. The least positive integer n for which $\left(\frac{1+i}{1-i}\right)^n = \frac{2}{\pi} \sin^{-1}\left(\frac{1+x^2}{2x}\right)$,

where $x > 0$ and $i = \sqrt{-1}$ is

- A. 2
- B. 4

C. 8

D. 12

Answer: B



[Watch Video Solution](#)

124. The area of the triangle formed by the points representing $-z$, iz and $z - iz$ in the Argand plane, is

A. $\frac{1}{2}|z|^2$

B. $|z|^2$

C. $\frac{3}{2}|z|^2$

D. $\frac{1}{4}|z|^2$

Answer: c



[Watch Video Solution](#)

125. If $z_0 = \frac{1-i}{2}$, then the value of the product

$(1+z_0)(1+z_0^2)(1+z_0^{2^2})(1+z_0^{2^3})\dots(1+z_0^{2^n})$ must be

A. $(1-i)\left(1 + \frac{1}{\frac{2}{2^{n-1}}}\right)$, if $n > 1$

B. $(1-i)\left(1 - \frac{1}{2^{2^n}}\right)$, if $n > 1$

C. $(1-i)\left(1 - \frac{1}{2^{n-1}}\right)$, if $n > 1$

D. $(1-i)\left(1 + \frac{1}{2^{2^n}}\right)$, if $n > 1$

Answer: b



Watch Video Solution

126. The greatest positive argument of complex number satisfying

$|z - 4| = \operatorname{Re}(z)$ is

A. $\frac{\pi}{3}$

B. $\frac{2\pi}{3}$

C. $\frac{\pi}{2}$

D. $\frac{\pi}{4}$

Answer: D



Watch Video Solution

127. If the points in the complex plane satisfy the equations $\log_5(|z| + 3) - \log_{\sqrt{5}}(|z - 1|) = 1$ and $\arg(z - 1) = \frac{\pi}{4}$ are of the form $A_1 + iB_1$, then the value of $A_1 + B_1$, is

A. $2\sqrt{2}$

B. $\sqrt{2}$

C. $4\sqrt{2}$

D. 0

Answer: a



[Watch Video Solution](#)

128. A complex number z with $(\text{Im})(z)=4$ and a positive integer n be such

that $\frac{z}{z+n} = 4i$, then the value of n , is

A. 4

B. 16

C. 17

D. 32

Answer: C



[Watch Video Solution](#)

129. If $\arg \left(\frac{z_1 - \frac{z}{|z|}}{\frac{z}{|z|}} \right) = \frac{\pi}{2}$ and $\left| \frac{z}{|z|} - z_1 \right| = 3$, then $|z_1|$ equals to

A. $\sqrt{26}$

B. $\sqrt{10}$

C. $\sqrt{3}$

D. $2\sqrt{2}$

Answer: B



Watch Video Solution

130. If z_1 and z_2 satisfy the equation $2|z + 3| = |\operatorname{Re}(z)|$ and $\arg \frac{z + 3}{1 + i} = \frac{\pi}{2}$,

then $\arg \frac{z_1 + 3}{z_2 + 3}$ is equal to

A. 0

B. $\pm \frac{\pi}{2}$

C. $\pm\pi$

D. $\pm\frac{\pi}{4}$

Answer: c



[View Text Solution](#)

131. If $A = \{ z \in \mathbb{C} : z = x + ix - 1 \text{ for all } x \in \mathbb{R} \}$ and $|z| \leq |\omega|$ for all $z, \omega \in A$, then z is equal to

A. $\frac{1}{2}(1 + i)$

B. $-\frac{1}{2}(1 - i)$

C. $-\frac{1}{2}(1 + i)$

D. $\frac{1}{3}(1 - 2i)$

Answer: b



[Watch Video Solution](#)

1. The locus of the center of a circle which touches the circles

$$|z - z_1| = a, |z - z_2| = b \text{ externally will be}$$

- A. an ellipse
- B. a hyperbola
- C. a circle
- D. none of these

Answer: b



[Watch Video Solution](#)

2. If n_1, n_2 are positive integers, then

$(1 + i)^{n_1} + (1 + i^3)^{n_1} + (1 + i^5)^{n_2} + (1 + i^7)^{n_2}$ is real if and only if :

- A. $n_1 = n_2 + 1$

B. $n_1 = n_2 - 1$

C. $n_1 = n_2$

D. $n_1 > 0, n_2 > 0$

Answer: d



Watch Video Solution

3. The modulus of $\sqrt{2}i - \sqrt{-2}i$ is

A. 2

B. $\sqrt{2}$

C. 0

D. $2\sqrt{2}$

Answer: a



Watch Video Solution

4. Prove that the triangle formed by the points 1 , $\frac{1+i}{\sqrt{2}}$, and i as vertices in the Argand diagram is isosceles.

- A. scalene
- B. equilateral
- C. isosceles
- D. right-angled

Answer: c

 [Watch Video Solution](#)

5. The value of $\frac{1+i\sqrt{3}}{(1-i\sqrt{3})^6} + \frac{1-i\sqrt{3}}{(1+i\sqrt{3})^6}$ is

- A. 2
- B. -2
- C. 1

D. 0

Answer: a



Watch Video Solution

6. If $\alpha + i\beta = \tan^{-1}(z)$, $z = x + iy$ and α is constant, the locus of 'z' is

A. $x^2 + y^2 + 2x\cot 2\alpha = 1$

B. $\cot 2\alpha (x^2 + y^2) = 1 + x$

C. $x^2 + y^2 + 2y\tan\alpha = 1$

D. $x^2 + y^2 + 2x\sin 2\alpha = 1$

Answer: a



Watch Video Solution

7. If $\cos A + \cos B + \cos C = 0$, $\sin A + \sin B + \sin C = 0$ and $A + B + C = 180^\circ$, then the value of $\cos 3A + \cos 3B + \cos 3C$ is

A. 3

B. -3

C. $\sqrt{3}$

D. 0

Answer: b



Watch Video Solution

8. The value of the expression

$$1. (2 - \omega). (2 - \omega^2) + 2. (3 - \omega)(3 - \omega^2) + \dots + (n - 1)(n - \omega)(n - \omega^2), \text{ where}$$

ω is an imaginary cube root of unity, is.....

A. $\left\{ \frac{n(n+1)}{2} \right\}^2$

B. $\left\{ \frac{n(n+1)}{2} \right\}^2 - n$

C. $\left\{ \frac{n(n+1)}{2} \right\}^2 + n$

D. none of these

Answer: c

 [Watch Video Solution](#)

9. The value of the expression

$$\left(1 + \frac{1}{\omega}\right)\left(1 + \frac{1}{\omega^2}\right) + \left(2 + \frac{1}{\omega}\right)\left(2 + \frac{1}{\omega^2}\right) + \left(3 + \frac{1}{\omega}\right)\left(3 + \frac{1}{\omega^2}\right) + \dots +$$

, where ω is an imaginary cube root of unity, is

A. $\frac{n(n^2 + 2)}{3}$

B. $\frac{n(n^2 - 2)}{3}$

C. $\frac{n(n^2 + 1)}{3}$

D. none of these

Answer: A



Watch Video Solution

10. The condition that $x^{n+1} - x^n + 1$ shall be divisible by $x^2 - x + 1$ is that

A. $n = 6k + 1$

B. $n = 6k - 1$

C. $n = 3k + 1$

D. none of these

Answer: a



Watch Video Solution

11. The expression $(1 + i)^{n_1} + (1 + i^3)^{n_2}$ is real iff

A. $n_1 = -n_2$

B. $n_1 = 4r + (-1)^r n_2$

C. $n_1 = 2r + (-1)^r n_2$

D. none of these

Answer: b



Watch Video Solution

12. $\begin{vmatrix} 6i & -3i & 1 \\ 4 & 3i & -1 \\ 20 & 3 & i \end{vmatrix} = x+iy$ then

A. $x = 3, y = 1$

B. $x = 1, y = 3$

C. $x = 0, y = 3$

D. none of these

Answer: D



Watch Video Solution

13. The centre of a square ABCD is at z_0 . If A is z_1 , then the centroid of the

ABC is $2z_0 - (z_1 - z_0)$ (b) $\left(z_0 + i\left(\frac{z_1 - z_0}{3}\right)\frac{z_0 + iz_1}{3}\right)$ (d) $\frac{2}{3}(z_1 - z_0)$

A. $z_1(\cos\pi \pm i\sin\pi)$

B. $\frac{z_1}{3}(\cos\pi \pm i\sin\pi)$

C. $z_1\left(\cos\alpha \pm i\frac{\sin\pi}{2}\right)$

D. $\frac{z_1}{3}\left(\frac{\cos\pi}{2} \pm i\frac{\sin\pi}{2}\right)$

Answer: d



Watch Video Solution

14. If $\cos\alpha + 2\cos\beta + 3\cos\gamma = \sin\alpha + 2\sin\beta + 3\sin\gamma = 0$ and $\alpha + \beta + \gamma = 0$,

then $\cos 3\alpha + 8\cos 3\beta + 27\sin 3\gamma =$

A. 0

B. 3

C. 18

D. -18

Answer: c



[Watch Video Solution](#)

15. If $\cos\alpha + 2\cos\beta + 3\cos\gamma = \sin\alpha + 2\sin\beta + 3\sin\gamma = 0$ and $\alpha + \beta + \gamma = 0$, then $\cos 3\alpha + 8\cos 3\beta + 27\sin 3\gamma =$

A. 0

B. 3

C. 8

D. -18

Answer: a



[Watch Video Solution](#)

16. Sum of the series $\sum_{r=0}^n (-1)^r {}^n C_r [i^{5r} + i^{6r} + i^{7r} + i^{8r}]$ is

A. 2^n

B. $2^{n/2+1}$

C. $n^n + 2^{n/2+1}$

D. $2^n + 2^{n/2+1} \frac{\cos(n\pi)}{4}$

Answer: d



Watch Video Solution

17. If $az_1 + bz_2 + cz_3 = 0$ for complex numbers z_1, z_2, z_3 and real numbers a, b, c then z_1, z_2, z_3 lie on a

A. straight line

B. circle

C. depends on the choice of a, b, c

D. none of these

Answer: c



Watch Video Solution

18. If $2z_1 - 3z_2 + z_3 = 0$, then z_1, z_2 and z_3 are represented by

A. three vertices of a triangle

B. three collinear points

C. three vertices of a rhombus

D. none of these

Answer: B



Watch Video Solution

19. $Re\left(\frac{z+4}{2z-1}\right) = \frac{1}{2}$, then z is represented by a point lying on

A. a circle

B. an ellipse

C. a straight line

D. none of these

Answer: C

 [Watch Video Solution](#)

20. The vertices of a square are z_1, z_2, z_3 and z_4 taken in the anticlockwise order, then $z_3 =$

A. $z_1 + z_2 + z_3 + z_4 = 0$

B. $z_1 + z_2 = z_3 + z_4$

C. $\text{amp} \left(\frac{z_2 - z_4}{z_1 - z_3} \right) = \frac{\pi}{2}$

D. $\text{amp} \frac{z_1 - z_2}{z_3 - z_4} = \frac{\pi}{2}$

Answer: c



Watch Video Solution

21. Let $\lambda \in \mathbb{R}$. If the origin and the non-real roots of $2z^2 + 2z + \lambda = 0$ form the three vertices of an equilateral triangle in the Argand lane, then λ is 1

b. $\frac{2}{3}$ c. 2 d. -1

A. 1

B. 2

C. -1

D. none of these

Answer: d



Watch Video Solution

22. if the complex no z_1, z_2 and z_3 represents the vertices of an equilateral triangle such that $|z_1| = |z_2| = |z_3|$ then relation among z_1, z_2 and z_3

A. $z_1 + z_2 + z_3 = 0$ and $z_1 z_2 z_3 = 1$

B. $z_1 + z_2 + z_3 = 1$ and $z_1 z_2 z_3 = 1$

C. $z_1 z_2 + z_2 z_3 + z_3 z_1 = 0$ and $z_1 + z_2 + z_3 = 0$

D. $z_1 z_2 + z_2 z_3 + z_3 z_1 = 0$ and $z_1 z_2 z_3 = 1$

Answer: a



Watch Video Solution

23. If P, P' represent the complex number z_1 and its additive inverse respectively, then the equation of the circle with PP' as a diameter is

A. $\frac{z}{z_1} = \frac{\bar{z}_1}{z}$

B. $z\bar{z} + z_1\bar{z}_1 = 0$

C. $z\bar{z}_1 + \bar{z}z_1 = 0$

D. none of these

Answer: a



Watch Video Solution

24. Let $A(z_1), B(z_2), C(z_3)$ be the vertices of an equilateral triangle ABC

in the Argand plane, then the number $\frac{z_2 - z_3}{2z_1 - z_2 - z_3}$, is

A. purely real

B. purely imaginary

C. a complex number with non-zero real and imaginary parts

D. none of these

Answer: b



Watch Video Solution

25. The area of the triangle (in square units) whose vertices are i , ω and ω^2 where $i = \sqrt{-1}$ and ω, ω^2 are complex cube roots of unity, is

A. $\frac{3\sqrt{3}}{2}$

B. $\frac{3\sqrt{3}}{4}$

C. 0

D. $\frac{\sqrt{3}}{4}$

Answer: d



Watch Video Solution

26. The complex number z satisfying $|z + 1| = |z - 1|$ and $\arg \frac{z - 1}{z + 1} = \frac{\pi}{4}$, is

A. $(\sqrt{2} + 1) + 0i$

B. $0 + (\sqrt{2} + 1)i$

C. $0 + (\sqrt{2} - 1)i$

D. $(-\sqrt{2} + 1) + 0i$

Answer: B



Watch Video Solution

27. If A,B,C are three points in the Argand plane representing the complex numbers, z_1, z_2, z_3 such that $z_1 = \frac{\lambda z_2 + z_3}{\lambda + 1}$, where $\lambda \in R$, then the distance of A from the line BC, is

A. λ

B. $\frac{\lambda}{\lambda + 1}$

C. 1

D. 0

Answer: d



Watch Video Solution

28. If $z \left(\overline{z + \alpha} \right) + \bar{z}(z + \alpha) = 0$, where α is a complex constant, then z is represented by a point on

- A. a circle
- B. a straight line
- C. a parabola
- D. none of these

Answer: A



[Watch Video Solution](#)

29. Let A,B,C be three collinear points which are such that $AB.AC=1$ and the points are represented in the Argand plane by the complex numbers, $0, z_1$ and z_2 respectively. Then,

- A. $z_1 z_2 = 1$
- B. $z_1 \bar{z}_2 = 1$

C. $|z_1| |z_2| = 1$

D. $z_1 = \bar{z}_2$

Answer: b



Watch Video Solution

30. If z_1, z_2, z_3, z_4 are the four complex numbers represented by the vertices of a quadrilateral taken in order such that $z_1 - z_4 = z_2 - z_3$ and

$\text{amp} \frac{z_4 - z_1}{z_2 - z_1} = \frac{\pi}{2}$ then the quadrilateral is a

A. a rhombus

B. a square

C. a rectangle

D. not a cyclic quadrilateral

Answer: c



Watch Video Solution

31. If z be a complex number, then

$|z - 3 - 4i|^2 + |z + 4 + 2i|^2 = k$ represents a circle, if k is equal to

A. 30

B. 40

C. 55

D. 35

Answer: c



Watch Video Solution

32. In Argand diagram, O, P, Q represent the origin, z and $z + iz$ respectively

then $\angle OPQ =$

A. $\frac{\pi}{4}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{2}$

D. $\frac{2\pi}{3}$

Answer: c



Watch Video Solution

33. If $\frac{2z_1}{3z_2}$ is purely imaginary number, then $\left| \frac{z_1 - z_2}{z_1 + z_2} \right|^4$ is equal to

A. $3/2$

B. 1

C. $2/3$

D. $4/9$

Answer: B



Watch Video Solution

34. If ω is a cube root of unity then find the value of

$$\sin\left(\left(\omega^{10} + \omega^{23}\right)\pi - \frac{\pi}{4}\right)$$

A. $\frac{1}{\sqrt{2}}$

B. $\frac{\sqrt{3}}{2}$

C. $-\frac{1}{\sqrt{3}}$

D. $-\frac{\sqrt{3}}{2}$

Answer: A



Watch Video Solution

35. If center of a regular hexagon is at the origin and one of the vertices on the Argand diagram is $1 + 2i$, then its perimeter is`

A. $2\sqrt{5}$

B. $6\sqrt{2}$

C. $4\sqrt{5}$

D. $6\sqrt{5}$

Answer: D



[Watch Video Solution](#)

36. If $z^2 + (p + iq)z + (r + is) = 0$, where p, q, r, s are non-zero, has real roots, then

A. $pqs = s^2 + q^2r$

B. $pqr = r^2 + p^2s$

C. $prs = q^2 + r^2p$

D. $qrs = p^2 + s^2q$

Answer: A



[Watch Video Solution](#)

37. Let z_1, z_2, z_3 be three vertices of an equilateral triangle circumscribing the circle $|z| = \frac{1}{2}$, if $z_1 = \frac{1}{2} + \sqrt{3}\frac{i}{2}$ and z_1, z_2, z_3 are in anticlockwise sense then z_2 is

A. $1 + i\sqrt{3}$

B. $1 - i\sqrt{3}$

C. 1

D. -1

Answer: D



Watch Video Solution

38. If ω is the complex cube root of unity, then the value of

$$\omega + \omega \frac{1}{2} + \frac{3}{8} + \frac{9}{32} + \frac{27}{128} + \dots \dots \dots,$$

A. -1

B. 1

C. $-i$

D. i

Answer: A



[Watch Video Solution](#)

39. the locus of $z = i + 2\exp\left(i\left(\theta + \frac{\pi}{4}\right)\right)$ is

A. a circle

B. an ellipse

C. a parabola

D. hyperbola

Answer: A



[Watch Video Solution](#)

40. If z lies on the circle $|z - 1| = 1$, then $\frac{z - 2}{z}$ is

- A. purely real
- B. Purely imaginary
- C. positive real
- D. hyperbola

Answer: B



[Watch Video Solution](#)

41. If $a > 0$ and the equation $|z - a^2| + |z - 2a| = 3$, represents an ellipse, then 'a' belongs to the interval

- A. (1,3)
- B. $(\sqrt{2}, \sqrt{3})$
- C. (0,3)

D. $(1, \sqrt{3})$

Answer: C

 [Watch Video Solution](#)

42. For any complex number z , find the minimum value of $|z| + |z - 2i|$

A. 0

B. 1

C. 2

D. 4

Answer: C

 [Watch Video Solution](#)

43. Find the greatest and the least value of $|z_1 + z_2|$ if $z_1 = 24 + 7i$ and $|z_2| = 6$.

A. 31,19

B. 25,16

C. 31,25

D. 19,16

Answer: a



Watch Video Solution

44. For all complex numbers z_1, z_2 satisfying $|z_1| = 12$ and $|z_2 - 3 - 4i| = 5$, the minimum value of $|z_1 - z_2|$ is

A. 0

B. 2

C. 7

Answer: B



Watch Video Solution

45. If $k > 1$, $|z_1| < k$ and $\left| \frac{k - z_1 \bar{z}_2}{z_1 - kz_2} \right| = 1$, then

A. $|z_2| < k$

B. $|z_2| = k$

C. $z_2 = 0$

D. $|z_2| = 1$

Answer: d



Watch Video Solution

46. If $|z - i| = 1$ and $\arg(z) = \theta$ where $0 < \theta < \frac{\pi}{2}$, then $\cot\theta - \frac{2}{z}$ equals

A. $2i$

B. $-i$

C. i

D. $1 + i$

Answer: C



Watch Video Solution

47. If $\operatorname{Re}(z) < 0$ then the value of $(1 + z + z^2 + \dots + z^n)$ cannot exceed

A. $|z^n| - \frac{1}{|z|}$

B. $n|z|^n + 1$

C. $|z|^n - \frac{1}{|z|}$

D. $|z|^n + \frac{1}{|z|}$

Answer: d



Watch Video Solution

48. Let z_1, z_2, z_3 be three complex numbers satisfying $\frac{1}{z_1} + \frac{1}{z_2} + \frac{1}{z_3} = 0$.

Let $z_k = r_k(\cos\alpha_k + i\sin\alpha_k)$ and $\omega_k = \frac{\cos 2\alpha_k + i\sin 2\alpha_k}{z_k}$ for $k = 1, 2, 3$. If

ω_1, ω_2 and ω_3 are the affixes of points A_1, A_2 and A_3 respectively in the Argand plane, then $\Delta A_1 A_2 A_3$ has its

- A. incenter at the origin
- B. centroid at the origin
- C. circumcenter at the origin
- D. orthocenter at the origin

Answer: b



View Text Solution

49. a and b are real numbers between 0 and 1 such that the points $z_1 = a + i, z_2 = 1 + bi, z_3 = 0$ form an equilateral triangle, then a and b are equal to

A. $a = \sqrt{3} - 1, b = \frac{\sqrt{3}}{2}$

B. $a = 2 - \sqrt{3}, b = 2 - \sqrt{3}$

C. $a = 1/2, b = 3/4$

D. none of these

Answer: B



Watch Video Solution

50. If ω is a cube root of unity, then find the value of the following:

$$\frac{a + b\omega + c\omega^2}{b + c\omega + a\omega^2} + \frac{a + b\omega + c\omega^2}{b + c\omega + a\omega^2}$$

A. 1

B. 0

C. -1

D. 2

Answer: D



[Watch Video Solution](#)

51. If a, b, c and u, v, w are the complex numbers representing the vertices of two triangles such that $c = (1 - r)a + rb$ and $w = (1 - r)u + rv$, where r is a complex number, then the two triangles

A. have the same area

B. are similar

C. are congruent

D. none of these

Answer: B



[Watch Video Solution](#)

52. If $z = re^{i\theta}$ then $|e^{iz}|$ is equal to:

A. $e^{-r\sin\theta}$

B. $re^{-r\sin\theta}$

C. $e^{-r\cos\theta}$

D. $re^{-r\cos\theta}$

Answer: A



Watch Video Solution

53. If a complex number z lies in the interior or on the boundary of a circle of radius 3 and center at $(-4, 0)$, then the greatest and least values of $|z + 1|$ are

A. 5,0

B. 6,1

C. 6,0

D. none of these

Answer: C



[Watch Video Solution](#)

54. Let z_1 and z_2 be two non - zero complex numbers such that

$\frac{z_1}{z_2} + \frac{z_2}{z_1} = 1$ then the origin and points represented by z_1 and z_2

A. z_1, z_2 are collinear

B. z_1, z_2 are the origin from a right angled triangle

C. z_1, z_2 and the origin form an equilateral triangle

D. none of these

Answer: c



[Watch Video Solution](#)

55. If z_1, z_2, z_3 be vertices of an equilateral triangle occurring in the anticlockwise sense, then

A. $z_1^2 + z_2^2 + z_3^2 = 2(z_1z_2 + z_2z_3 + z_3z_1)$

B. $\frac{1}{z_1 + z_2} + \frac{1}{z_2 + z_3} + \frac{1}{z_3 + z_1} = 0$

C. $z_1 + \omega z_2 + \omega^2 z_3 = 0$

D. none of these

Answer: C



[Watch Video Solution](#)

56. Let z be a complex number satisfying $|z - 5i| \leq 1$ such that $\text{amp}(z)$ is minimum, then z is equal to

A. $\frac{2\sqrt{6}}{5} + \frac{24i}{5}$

B. $\frac{24}{5} + \frac{2\sqrt{6}i}{5}$

$$C. \frac{2\sqrt{6}}{5} - \frac{24i}{5}$$

D. none of these

Answer: A



Watch Video Solution

57. If $|z - 25i| \leq 15$. then $|\text{maximum } \arg(z) - \text{minimum } \arg(z)|$ equals

$$A. \cos^{-1}\left(\frac{3}{5}\right)$$

$$B. \pi - 2\cos^{-1}\left(-\frac{3}{5}\right)$$

$$C. \frac{\pi}{2} + \cos^{-1}\left(\frac{3}{5}\right)$$

D. none of these

Answer: B



Watch Video Solution

58. Let z be a complex number (not lying on x-axis) of maximum modulus

such that $\left|z + \frac{1}{z}\right| = 1$. Then,

A. $\text{Im}(z)=0$

B. $\text{Re}(z)=0$

C. $\text{amp}(z)=\pi$

D. $\text{Re}(z)=1$

Answer: b



Watch Video Solution

59. The maximum distance from the origin of coordinates to the point z

satisfying the equation $\left|z + \frac{1}{z}\right| = a$ is

A. $\frac{1}{2} \left(\sqrt{a^2 + 1} + a \right)$

B. $\frac{1}{2} \left(\sqrt{a^2 + 2} + a \right)$

$$\text{C. } \frac{1}{2} \left(\sqrt{a^2 + 4} + a \right)$$

$$\text{D. } \frac{1}{2} \left(\sqrt{a^2 + 1} - a \right)$$

Answer: c



Watch Video Solution