



MATHS

BOOKS - OBJECTIVE RD SHARMA MATHS VOL I (HINGLISH)

SCALAR AND VECTOR PRODUCTS OF TWO VECTORS

Illustration

1. if \vec{a} and \vec{b} are unit vectors such that $\vec{a} \cdot \vec{b} = \cos \theta$, then the value of $|\vec{a} + \vec{b}|$, is

A. $2 \sin \theta / 2$

B. $2 \sin \theta$

C. $2 \cos \theta / 2$

D. $2 \cos \theta$

Answer: C



Watch Video Solution

2. If \vec{a} and \vec{b} are unit vectors, then the greatest value of $|\vec{a} + \vec{b}| + |\vec{a} - \vec{b}|$ is

A. 2

B. 4

C. $2\sqrt{2}$

D. $\sqrt{2}$

Answer: C



Watch Video Solution

3. If the unit vectors \vec{a} and \vec{b} are inclined of an angle 2θ such that $|\vec{a} - \vec{b}| < 1$ and $0 \leq \theta \leq \pi$ then θ in the interval

A. $\left[0, \frac{\pi}{6}\right) \cup (5\pi/6, \pi]$

B. $[0, \pi]$

C. $[\pi/6, \pi/2]$

D. $[\pi/2, 5\pi/6]$

Answer: A



Watch Video Solution

4. Let \vec{a} and \vec{b} be two unit vectors and α be the angle between them, then $\vec{a} + \vec{b}$ is a unit vector, if $\alpha =$

A. $\pi/4$

B. $\pi/3$

C. $2\pi/3$

D. $\pi/2$

Answer: C



Watch Video Solution

5. If $|\vec{a} - \vec{b}| = |\vec{a}| = |\vec{b}| = 1$, then the angle between \vec{a} and \vec{b} , is

A. $\frac{\pi}{3}$

B. $\frac{3\pi}{4}$

C. $\frac{\pi}{2}$

D. 0

Answer: A



Watch Video Solution

6. Let \vec{a} and \vec{b} are two vectors inclined at an angle of 60° , If $|\vec{a}| = |\vec{b}| = 2$, the the angle between \vec{a} and $\vec{a} + \vec{b}$ is

A. 30°

B. 60°

C. 45°

D. none of these

Answer: A



[Watch Video Solution](#)

7. Consider a pyramid OPQRS located in the first octant ($x \geq 0, y \leq 0, z \leq 0$) with O as origin, and OP and OR along the x-axis and y-axis respectively. The base OPQR of the pyramid is a square with $OP=3$. The point S is directly above the mid-point T of the diagonal OQ such that $TS=3$, Then, the angle between OQ and OS, is

A. $\frac{\pi}{3}$

B. $\frac{\pi}{6}$

C. $\cos^{-1} \frac{1}{\sqrt{3}}$

D. $\cos^{-1} \frac{1}{3}$

Answer: C



[Watch Video Solution](#)

8. If \vec{a} , \vec{b} are unit vectors such that $\vec{a} - \vec{b}$ is also a unit vector, then the angle between \vec{a} and \vec{b} , is

A. $\frac{\pi}{6}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{4}$

D. $\frac{\pi}{2}$

Answer: B



[Watch Video Solution](#)

9. If \vec{a} , \vec{b} are unit vectors such that

$$|\vec{a} + \vec{b}| = -1 \text{ then } |2\vec{a} - 3\vec{b}| =$$

A. 19

B. $\sqrt{19}$

C. $\sqrt{13}$

D. 4

Answer: B



Watch Video Solution

10. If \vec{a}, \vec{b} are unit vectors such that

$$|\vec{a} + \vec{b}| = 1 \text{ and } |\vec{a} - \vec{b}| = \sqrt{3}, \text{ " then " } |3\vec{a} + 2\vec{b}| =$$

A. 7

B. 4

C. $\sqrt{7}$

D. $\sqrt{19}$

Answer: C



Watch Video Solution

11. If \vec{a} and \vec{b} are unit vectors inclined to x-axis at angle 30° and 120° then $|\vec{a} + \vec{b}|$ equals

A. $\sqrt{2/3}$

B. $\sqrt{2}$

C. $\sqrt{3}$

D. 2

Answer: B



Watch Video Solution

12. If \vec{a} , \vec{b} , \vec{c} are three vectors such that

$$\vec{a} + \vec{b} + \vec{c} = \vec{0}, |\vec{a}| = 1, |\vec{b}| = 2, |\vec{c}| = 3, \text{ then}$$

$\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$ is equal to

A. 1

B. 0

C. -7

D. 7

Answer: C



Watch Video Solution

13. Let \vec{a} , \vec{b} , \vec{c} be three vectors such that

$\vec{a} \perp (\vec{b} + \vec{c})$, $\vec{b} \perp (\vec{c} + \vec{a})$ and $\vec{c} \perp (\vec{a} + \vec{b})$, if $|\vec{a}| = 1$, $|\vec{b}| = 2$,

$|\vec{c}| = 3$, then $|\vec{a} + \vec{b} + \vec{c}|$ is,

A. $\sqrt{6}$

B. 14

C. $\sqrt{14}$

D. none of these

Answer: C



 Watch Video Solution

14. If two out of the three vectors, \vec{a} , \vec{b} , \vec{c} are unit vectors such that $\vec{a} + \vec{b} + \vec{c} = 0$ and $2\left(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}\right) + 3 = 0$ then the length of the third vector is

- A. 3
- B. 2
- C. 1
- D. 0

Answer: C

 Watch Video Solution

15. Let \vec{a} , \vec{b} , \vec{c} be three unit vectors such that $|\vec{a} + \vec{b} + \vec{c}| = 1$ and $\vec{a} \perp \vec{b}$, if \vec{c} makes angles δ , β with \vec{a} , \vec{b} respectively, then $\cos \delta + \cos \beta$ is equal to

A. 1

B. -1

C. $\frac{3}{2}$

D. 0

Answer: B



Watch Video Solution

16. Let $\vec{a}, \vec{b}, \vec{c}$ be three unit vectors such that angle between \vec{a} and \vec{b} is α , \vec{b} and \vec{c} is β and \vec{c} and \vec{a} is γ . if $|\vec{a} + \vec{b} + \vec{c}| = 1$, then $\cos \alpha + \cos \beta + \cos \gamma =$

A. 1

B. $-\frac{1}{2}$

C. $\frac{3}{2}$

D. $\frac{1}{2}$

Answer: D



Watch Video Solution

17. Let $\vec{a}, \vec{b}, \vec{c}$ be vectors of equal magnitude such that the angle between \vec{a} and \vec{b} is α , \vec{b} and \vec{c} is β and \vec{c} and \vec{a} is γ . Then minimum value of $\cos \alpha + \cos \beta + \cos \gamma$ is

A. $\frac{1}{2}$

B. $-\frac{1}{2}$

C. $\frac{3}{2}$

D. $-\frac{3}{2}$

Answer: D



Watch Video Solution

18. Let \vec{a} , \vec{b} , \vec{c} be three vectors of equal magnitude such that the angle between each pair is $\frac{\pi}{3}$. If $|\vec{a} + \vec{b} + \vec{c}| = \sqrt{6}$, then $|\vec{a}| =$

A. 2

B. -1

C. 1

D. $-\sqrt{\frac{2}{3}}$

Answer: C



Watch Video Solution

19. If $|\vec{a}| = 3$, $|\vec{b}| = 5$ and $|\vec{c}| = 4$ and $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ then the value of $(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c})$ is equal to

A. 0

B. -25

C. 25

D. none of these

Answer: B



[Watch Video Solution](#)

20. Let O be the origin, and OX, OY, OZ be three unit vectors in the direction of the sides QR, RP, PQ , respectively of a triangle PQR . If the triangle PQR varies, then the minimum value of $\cos(P + Q) + \cos(Q + R) + \cos(R + P)$ is: $-\frac{3}{2}$ (b) $\frac{5}{3}$ (c) $\frac{3}{2}$ (d) $-\frac{5}{3}$

A. $-\frac{5}{3}$

B. $-\frac{3}{2}$

C. $\frac{3}{2}$

D. $\frac{5}{2}$

Answer: B



[Watch Video Solution](#)

21. Angle between vectors \vec{a} and \vec{b} where \vec{a} , \vec{b} and \vec{c} are unit vectors satisfying $\vec{a} + \vec{b} + \sqrt{3}\vec{c} = \vec{0}$ is

A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{2}$

Answer: C



Watch Video Solution

22. If $\vec{a} + \vec{b} + \vec{c} = \vec{0}$, $|\vec{a}| = 3$, $|\vec{b}| = 5$ and $|\vec{c}| = 7$, then the angle between \vec{a} and \vec{b} is

A. $\frac{\pi}{2}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{6}$

D. $\frac{\pi}{3}$

Answer: D



Watch Video Solution

23. If \vec{a} , \vec{b} and \vec{c} are unit vectors satisfying $|\vec{a} - \vec{b}|^2 + |\vec{b} - \vec{c}|^2 + |\vec{c} - \vec{a}|^2 = 9$, then $|2\vec{a} + 5\vec{b} + 5\vec{c}|$ is equal to

A. 0

B. 1

C. 2

D. 3

Answer: D



Watch Video Solution

24. The value of 'a' for which the points A, B, C with position vectors $2\hat{i} - \hat{j} + \hat{k}$, $\hat{i} + 2\hat{k}$, $\hat{i} - 3\hat{j} - 5\hat{k}$ and $a\hat{i} - 3\hat{j} + \hat{k}$ respectively are the vertices of a right angled triangle with $C = \pi/2$, are

A. 2 and 1

B. -2 and -1

C. -2 and 1

D. 2 and -1

Answer: A



[Watch Video Solution](#)

25. If R^2 is the magnitude of the projection vector of the vector $\alpha\hat{i} + \beta\hat{j}$ on $\sqrt{3}\hat{i} + \hat{j}$ and if $\alpha = 2 + \sqrt{3}\beta$ then possible value (s) of $|\alpha|$ is /are

A. 1,2

B. 3,4

C. 4,5

D. 3

Answer: D



[Watch Video Solution](#)

26. The vectors $\vec{a} = 3\hat{i} - 2\hat{j} + 2\hat{k}$ and $\vec{b} = -\hat{i} - 2\hat{k}$ are the adjacent sides of a parallelogram. Then, the acute angle between \vec{a} and \vec{b} is

A. $\pi/4$

B. $\pi/3$

C. $3\pi/4$

D. $2\pi/3$

Answer: A



[Watch Video Solution](#)

27. If $\vec{e}_1 = (1, 1, 1)$ and $\vec{e}_2 = (1, 1, -1)$ and \vec{a} and \vec{b} are two vectors that $\vec{e}_1 = 2\vec{a} + \vec{b}$ and $\vec{e}_2 = \vec{a} + 2\vec{b}$ then angle between \vec{a} and \vec{b} is

A. $\cos^{-1}\left(\frac{7}{9}\right)$

B. $\cos^{-1}\left(\frac{7}{11}\right)$

C. $\cos^{-1}\left(-\frac{7}{11}\right)$

D. $\cos^{-1}\left(\frac{6\sqrt{2}}{11}\right)$

Answer: C



Watch Video Solution

28. The values of x for which the angle between the vectors $\vec{a} = x\hat{i} - 3\hat{j} - \hat{k}$ and $\vec{b} = 2x\hat{i} + x\hat{j} - \hat{k}$ is acute, and the angle between the vector \vec{b} and the axis of ordinates is obtuse, are

A. $\frac{1}{2}, 2$

B. $-2, 3$

C. all $x < 0$

D. $x > 0$

Answer: C



Watch Video Solution

29. The value of x for which the angle between $\vec{a} = 2x^2\hat{i} + 4x\hat{j} = \hat{k} + \hat{k}$ and $\vec{b} = 7\hat{i} - 2\hat{j} = x\hat{k}$, is obtuse and the angle between \vec{b} and the z-axis is acute and less than $\pi/6$, are

A. $a < x < 1/2$

B. $1/2 < x < 15$

C. $x > 1/2$ or $x < 0$

D. none of these

Answer: D



Watch Video Solution

30. If \vec{a}, \vec{b} are two unit vectors such that $|\vec{a} + \vec{b}| = 2\sqrt{3}$ and $|\vec{a} - \vec{b}| = 6$ then the angle between \vec{a} and \vec{b} , is

A. $\frac{\pi}{3}$

B. $\frac{2\pi}{3}$

C. $\frac{\pi}{4}$

D. $\frac{\pi}{2}$

Answer: B



Watch Video Solution

31. For any vector \vec{r} ,

$$(\vec{r} \cdot \hat{i})\hat{i} + (\vec{r} \cdot \hat{j})\hat{j} + (\vec{r} \cdot \hat{k})\hat{k} =$$

A. \vec{r}

B. $2\vec{r}$

C. $3\vec{r}$

D. $\vec{0}$

Answer: A

 [Watch Video Solution](#)

32. For any vector \vec{r} , $(\vec{r} \cdot \hat{i})^2 + (\vec{r} \cdot \hat{j})^2 + (\vec{r} \cdot \hat{k})^2$ is equal to

A. 1

B. $|\vec{r}|$

C. \vec{r}

D. $|\vec{r}|^2$

Answer: D

 [Watch Video Solution](#)

33. A vector of magnitude 4 which is equally inclined to the vectors

$\hat{i} + \hat{j}$, $\hat{j} + \hat{k}$ and $\hat{k} + \hat{i}$, is

A. $\frac{4}{\sqrt{3}} (\hat{i} - \hat{j} - \hat{k})$

B. $\frac{4}{\sqrt{3}} (\hat{i} + \hat{j} - \hat{k})$

C. $\frac{4}{\sqrt{3}} (\hat{i} + \hat{j} + \hat{k})$

D. $\frac{4}{\sqrt{3}} (\hat{i} + \hat{j} - \hat{k})$

Answer: C



Watch Video Solution

34. Forces of magnitudes 5 and 3 units acting in the directions

$6\hat{i} + 2\hat{j} + 3\hat{k}$ and $3\hat{i} - \hat{j} + 6\hat{k}$ respectively act on a particle which is

displaced from the point (2,2,-1) to (4,3,1) . The work done by the forces, is

A. 148 unit

B. $\frac{148}{7}$ unit

C. 296 units

D. none of these

Answer: B



Watch Video Solution

35. A groove is in the form of a broken line ABC and the position vectors for the three points are respectively $2\hat{i} - 3\hat{j} + 2\hat{k}$, $3\hat{i} - \hat{k}$, $\hat{i} + \hat{j} + \hat{k}$. A force of magnitude $24\sqrt{3}$ acts on a particle of unit mass kept at the point A and moves it along the groove to the point C. If the line of action of the force is parallel to the vector $\hat{i} + 2\hat{j} + \hat{k}$ all along, the number of units of work done by the force is

A. $144\sqrt{2}$

B. $144\sqrt{3}$

C. $72\sqrt{2}$

D. $72\sqrt{3}$

Answer: C



[Watch Video Solution](#)

36. A particle acted on by constant forces $4\hat{i} = \hat{j} - 3\hat{k}$ and $3\hat{i} + \hat{j} - \hat{k}$ is displaced from the point $\hat{i} + 2\hat{j} + 3\hat{k} \rightarrow 5\hat{i} + 4\hat{j} + \hat{k}$. Find the work done

A. 50 units

B. 20 units

C. 30 units

D. 40 units

Answer: D



[Watch Video Solution](#)

37. If $|\vec{a} \cdot \vec{b}| = |\vec{a} \times \vec{b}|$, then the angle between \vec{a} and \vec{b} is

A. 0°

B. 180°

C. 150°

D. 45°

Answer: D



[Watch Video Solution](#)

38. If \vec{u} and \vec{v} are unit vectors and θ is the acute angle between them, then $2u\vec{u} \times 3\vec{v}$ is a unit vector for

A. no value of θ

B. exactly one value of θ

C. exactly two values of θ

D. more than two values of θ

Answer: B



Watch Video Solution

39. Let A_1, A_2, \dots, A_n ($n < 2$) be the vertices of regular polygon of n sides with its centre at the origin. Let \vec{a}_k be the position vector of the point A_k , $k = 1, 2, \dots, n$

if $\left| \sum_{k=1}^{n-1} (\vec{a}_k \times \vec{a}_{k+1}) \right| = \left| \sum_{k=1}^{n-1} (\vec{a}_k \cdot \vec{a}_{k+1}) \right|$ then the minimum

value of n is

A. 1

B. 2

C. 8

D. 9

Answer: D



Watch Video Solution

40. Let O be the origin , and $\vec{OX}, \vec{OY}, \vec{OZ}$ be three unit vector in the directions of the sides $\vec{OR}, \vec{RP}, \vec{PQ}$ respectively, of a triangle PQR, Then $|\vec{OX} \times \vec{OY}| =$

A. $\sin (P +Q)$

B. $\sin 2R$

C. $\sin (P +R)$

D. $\sin (Q +R)$

Answer: A



Watch Video Solution

41. If the angle between the vectors \vec{a} and \vec{b} is $\frac{\pi}{3}$ and the area of the triangle with adjacent sides parallel to \vec{a} and \vec{b} is 3 is

A. $\sqrt{3}$

B. $2\sqrt{3}$

C. $4\sqrt{3}$

D. $\frac{\sqrt{3}}{2}$

Answer: B



[Watch Video Solution](#)

42. If $\vec{a} = 2\hat{i} - 3\hat{k}$ and $\vec{b} = \hat{i} + 4\hat{j} - 2\hat{k}$ then $\vec{a} \times \vec{b}$ is

A. $10\hat{i} + 2\hat{j} + 11\hat{k}$

B. $10\hat{i} + 3\hat{j} + 11\hat{k}$

C. $10\hat{i} - 3\hat{j} + 11\hat{k}$

D. $10\hat{i} - 3\hat{j} - 10\hat{k}$

Answer: B



[Watch Video Solution](#)

43. For any vector \vec{a}

$|\vec{a} \times \hat{i}|^2 + |\vec{a} \times \hat{j}|^2 + |\vec{a} \times \hat{k}|^2$ is equal to

A. $|\vec{a}|^2$

B. $2|\vec{a}|^2$

C. $3|\vec{a}|^2$

D. $2|\vec{a}|$

Answer: B



Watch Video Solution

44. $(\vec{a} \cdot \hat{i})(\vec{a} \times \hat{i}) + (\vec{a} \cdot \hat{j})(\vec{a} \times \hat{j}) + (\vec{a} \cdot \hat{k})(\vec{a} \times \hat{k})$ is equal to

A. $3\vec{a}$

B. \vec{a}

C. $\vec{0}$

D. $2\vec{a}$

Answer: C

 [Watch Video Solution](#)

45. A unit vector perpendicular to the plane of $\vec{a} = 2\hat{i} - 6\hat{j} - 3\hat{k}$ and $\vec{b} = 4\hat{i} + 3\hat{j} - \hat{k}$ is

A. $\frac{1}{\sqrt{26}}(4\hat{i} + 3\hat{j} - \hat{k})$

B. $\frac{1}{7}(2\hat{i} - 6\hat{j} - 3\hat{k})$

C. $\frac{1}{7}(3\hat{i} + 2\hat{j} + 6\hat{k})$

D. $\frac{1}{7}(2\hat{i} - 3\hat{j} - 6\hat{k})$

Answer: C

 [Watch Video Solution](#)

46. The number of vectors of unit length perpendicular to the vectors

$\vec{a} = \hat{i} + \hat{j}$ and $\vec{b} = \hat{j} + \hat{k}$ is

A. 1

B. 2

C. 4

D. infinite

Answer: B



[Watch Video Solution](#)

47. A unit vector making an obtuse angle with x-axis and perpendicular to the plane containing the points $\hat{i} + 2\hat{j} + 3\hat{k}$, $2\hat{i} + 3\hat{j} + 4\hat{k}$ and $\hat{i} + 5\hat{j} + 7\hat{k}$ also makes an obtuse angle with

A. y-axis

B. z-axis

C. y and z axes

D. x and y axes

Answer: B



Watch Video Solution

48. Let $\vec{u} = \hat{i} + \hat{j}$, $\vec{v} = \hat{i} - \hat{j}$ and $\vec{w} = \hat{i} + 2\hat{j} + 3\hat{k}$. If \hat{n} is a unit vector such that $\vec{u} \cdot \hat{n} = 0$ and $\vec{v} \cdot \hat{n} = 0$, then $|\vec{w} \cdot \hat{n}|$ is equal to

A. 3

B. 0

C. 1

D. 2

Answer: A



Watch Video Solution

49. If $\vec{a} = \hat{i} + \hat{j} - \hat{k}$, $\vec{b} = -\hat{i} + 2\hat{j} + 2\hat{k}$ and $\vec{c} = -\hat{i} + 2\hat{j} - \hat{k}$, then a unit vector normal to the vectors $\vec{a} + \vec{b}$ and $\vec{b} - \vec{c}$, is

A. \hat{i}

B. \hat{j}

C. \hat{k}

D. none of these

Answer: A



Watch Video Solution

50. A unit vector perpendicular to both $\hat{i} + \hat{j}$ and $\hat{j} + \hat{k}$ is

A. $\hat{i} - \hat{j} + \hat{k}$

B. $\hat{i} + \hat{j} + \hat{k}$

C. $\frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$

D. $\frac{\hat{i} - \hat{j} + \hat{k}}{\sqrt{3}}$

Answer: C



Watch Video Solution

51. Let $\vec{a} = 2\hat{i} + \hat{j} - 2\hat{k}$ and $\vec{b} = \hat{i} + \hat{j}$. Let \vec{c} be vector such that $|\vec{c} - \vec{a}| = 3$, $\left| \left(\vec{a} \times \vec{b} \right) \times \vec{c} \right| = 3$ and the angle between \vec{c} and $\vec{a} \times \vec{b}$ be 30° . Then, $\vec{a} \cdot \vec{c}$ is equal to

A. $\frac{25}{8}$

B. 2

C. 5

D. $\frac{1}{8}$

Answer: B



Watch Video Solution

52. If \vec{a} , \vec{b} , \vec{c} are the position vectors of the vertices A,B,C of a triangle ABC. Then the area of triangle ABC is

A. $\frac{1}{2} \left| \vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} \right|$

B. $\frac{1}{2} \left| \vec{a} \times \vec{b} \right|$

C. $\frac{1}{2} \left| \vec{b} \times \vec{c} \right|$

D. $\frac{1}{2} \left| \vec{c} \times \vec{a} \right|$

Answer: A



Watch Video Solution

53. If $\vec{AB} = \vec{b}$ and $\vec{AC} = \vec{c}$ then the length of the perpendicular from A to the line BC is

A. $\frac{\left| \vec{b} \times \vec{c} \right|}{\left| \vec{b} + \vec{c} \right|}$

B. $\frac{\left| \vec{b} \times \vec{c} \right|}{\left| \vec{b} - \vec{c} \right|}$

$$\text{C. } \frac{|\vec{b} \times \vec{c}|}{2|\vec{b} - \vec{c}|}$$

$$\text{D. } \frac{|\vec{b} \times \vec{c}|}{2|\vec{b} + \vec{c}|}$$

Answer: B

 Watch Video Solution

54. The perpendicular distance of the point \vec{c} from the joining \vec{a} and \vec{b} is

$$\text{A. } \frac{|\vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{a} \times \vec{b}|}{|\vec{b} - \vec{a}|}$$

$$\text{B. } \frac{|\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}|}{|\vec{b} - \vec{a}|}$$

$$\text{C. } \frac{|\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}|}{|\vec{a} - \vec{a}|}$$

$$D. \frac{1}{2} \frac{\left| \vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} \right|}{\left| \vec{b} - \vec{a} \right|}$$

Answer: A



Watch Video Solution

55. If the diagonals of a parallelogram are represented by the vectors $3\hat{i} + \hat{j} - 2\hat{k}$ and $\hat{i} + 3\hat{j} - 4\hat{k}$, then its area in square units, is

A. $5\sqrt{3}$

B. $6\sqrt{3}$

C. $\sqrt{42}$

D. $\sqrt{42}$

Answer: C



Watch Video Solution

56. if \vec{a} , \vec{b} , \vec{c} are three vectors such that $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ then

A. $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a}$

B. $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a}$

C. $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{a} \times \vec{c}$

D. $\vec{b} \times \vec{a} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a}$

Answer: B



Watch Video Solution

57. Let \vec{a} , \vec{b} , \vec{c} be unit vectors such that $\vec{a} + \vec{b} + \vec{c} = \vec{0}$, which one of the following is correct ?

A. $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} = \vec{0}$

B. $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} \neq \vec{0}$

C. $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{a} \times \vec{c} = \vec{0}$

D. $\vec{a} \times \vec{b}$, $\vec{b} \times \vec{c}$, $\vec{c} \times \vec{a}$ are mutually perpendicular vectors.

Answer: B



Watch Video Solution

58. Let $\triangle PQR$ be a triangle. Let $\vec{a} = \overrightarrow{QR}$, $\vec{b} = \overrightarrow{RP}$ and $\vec{c} = \overrightarrow{PQ}$. if $|\vec{a}| = 12$, $|\vec{b}| = 4\sqrt{3}$ and $\vec{b} \cdot \vec{c} = 12$, then which of the following is (are) true ?

A. $\frac{1}{2}|\vec{c}|^2 - |\vec{a}| = 12$

B. $\frac{1}{2}|\vec{c}|^2 + |\vec{a}| = 30$

C. $|\vec{a} \times \vec{b} + \vec{c} \times \vec{a}| = 48\sqrt{3}$

D. $\vec{a} \cdot \vec{b} = -72$

Answer: A::C::D



Watch Video Solution

59. Let \vec{a} be a unit vector perpendicular to unit vectors \vec{b} and \vec{c} and if the angle between \vec{b} and \vec{c} is α , then $\vec{b} \times \vec{c}$ is

A. $\pm (\cos \alpha) \vec{a}$

B. $\pm (\cos e\alpha) \vec{a}$

C. $\pm (\sin \alpha) \vec{a}$

D. none of these

Answer: C



Watch Video Solution

60. If the vectors $\vec{a} \cdot \vec{a} = x\hat{i} + y\hat{j} + z\hat{k}$ and $\vec{b} = \hat{j}$ are such that \vec{a} , \vec{b} , and \vec{c} form a right handed system, then \vec{c} is

A. $x\hat{i} - x\hat{k}$

B. $\vec{0}$

C. $y\hat{j}$

D. $-z + \hat{x}\hat{k}$

Answer: A



Watch Video Solution

61. If the area of parallelogram whose diagonals coincide with the following pair of vectors is $5\sqrt{3}$, then vectors are

A. $3\hat{i} + 2\hat{j} - \hat{k}, 3\hat{i} - \hat{j} + 4\hat{k}$

B. $\frac{3}{2}\hat{i} + \frac{1}{2}\hat{j} - \hat{k}, 2\hat{i} - 6\hat{j} + 8\hat{k}$

C. $3\hat{i} + \hat{j} - 2\hat{k}, \hat{i} + 3\hat{j} + 4\hat{k}$

D. none of these

Answer: B



Watch Video Solution

62.

Let

$\vec{a} = 2\hat{i} + \hat{j} + \hat{k}$, $\vec{b} = \hat{i} + 2\hat{j} + \hat{k}$ and $\vec{c} = 2\hat{i} - 3\hat{j} + 4\hat{k}$. A vector \vec{r}

is

A. $-2\hat{i} = 2\hat{j} + 2\hat{k}$

B. $-2\hat{i} + \hat{j} + 3\hat{k}$

C. $3\hat{i} + 2\hat{j} + 4\hat{k}$

D. $\hat{i} - 5\hat{j} + 3\hat{k}$

Answer: D
[Watch Video Solution](#)

63. vectors \vec{a} and \vec{b} are inclined at an angle

$\theta = 60^\circ$. If $|\vec{a}| = 1$, $|\vec{b}| = 2$, then $\left[\left(\vec{a} + 3\vec{b} \right) \times \left(3\vec{a} - \vec{b} \right) \right]^2$

is equal to

A. 225

B. 275

C. 325

D. 300

Answer: D



Watch Video Solution

64. Let \vec{a} , \vec{b} , \vec{c} be unit vectors such that $\vec{a} \cdot \vec{b} = 0 = \vec{a} \cdot \vec{c}$. If the angle between \vec{b} and \vec{c} is $\frac{\pi}{6}$, then \vec{a} equals

A. $\pm 2 \left(\vec{b} \times \vec{c} \right)$

B. $2 \left(\vec{b} \times \vec{c} \right)$

C. $\pm \frac{1}{2} \left(\vec{b} \times \vec{c} \right)$

D. $-\frac{1}{2} \left(\vec{b} \times \vec{c} \right)$

Answer: A



Watch Video Solution

65. Let $\vec{u} = u_1\hat{i} + u_2\hat{j}$ be a unit vector in xy plane and $\vec{w} = \frac{1}{\sqrt{6}}(\hat{i} + \hat{j} + 2\hat{k})$. Given that there exists a vector \vec{c} " in " R_3 " such that $|\vec{u} \times \vec{v}| = 1$ and $\vec{w} \cdot (\vec{u} \times \vec{v}) = 1$, then

A. $|u_1| = |u_2|$

B. $|u_2| = 2|u_1|$

C. $2|u_1| = |u_2|$

D. $|u_1| = 3|u_2|$

Answer: A

 [Watch Video Solution](#)

66. Let $\vec{u} = u_1\hat{i} + u_3\hat{k}$ be a unit vector in xz-plane and $\vec{q} = \frac{1}{\sqrt{6}}(\hat{i} + \hat{j} + 2\hat{k})$. If there exists a vector \vec{v} in such that $|\vec{u} \times \vec{v}| = 1$ and $\vec{w} \cdot (\vec{u} \times \vec{v}) = 1$.Then

A. $|u_1| = |u_3|$

B. $|u_1| = 2|u_3|$

C. $|u_1| = 2|u_3|$

D. $2|u_1| = |u_3|$

Answer: B



Watch Video Solution

67. Let $\vec{u} = u_1\hat{i} + u_2\hat{j} + u_3\hat{k}$ be a unit vector in R^3 and $\vec{w} = \frac{1}{\sqrt{6}}(\hat{i} + \hat{j} + 2\hat{k})$, Given that there exists a vector \vec{v} in R^3 such that $|\vec{u} \times \vec{v}| = 1$ and $\vec{w} \cdot (\vec{u} \times \vec{v}) = 1$ which of the following statements is correct ?

A. There is exactly one choice for such \vec{v}

B. There are exactly two for such \vec{v}

C. There are exactly for such \vec{v}

D. There are infinitely many choices for such \vec{v}

Answer: D



Watch Video Solution

68. IF the force represented by $3\hat{i} + 2\hat{k}$ is acting through the point $5\hat{i} + 4\hat{j} - 3\hat{k}$, then its moment about th point (1,3,1) is

A. $14\hat{i} - 8\hat{j} + 12\hat{k}$

B. $-14\hat{i} + 8\hat{j} - 12\hat{k}$

C. $-6\hat{i} - \hat{j} + 9\hat{k}$

D. $6\hat{i} + \hat{j} - 9\hat{k}$

Answer: A



Watch Video Solution

69. The moment of the couple formed by the forces $5\hat{i} + \hat{j}$ and $-5\hat{i} - \hat{k}$ acting at the points (9, -1, 2) and (3, -2, 1)

respectively, is

A. $-\hat{i} + \hat{j} + 5\hat{k}$

B. $\hat{i} - 11\hat{j} - 5\hat{k}$

C. $-\hat{i} = 11\hat{j} + 5\hat{k}$

D. $\hat{i} - \hat{j} - 5\hat{k}$

Answer: D



Watch Video Solution

70. A force of 39 kg. wt is acting at a point p (-4,2,5) in the direction $12\hat{i} - 4\hat{j} - 3\hat{k}$. The moment of this force about a line through the origin having the direction of $2\hat{i} - 2\hat{j} + \hat{k}$ is

A. 76 units

B. - 76 units

C. $42\hat{i} + 144\hat{j} - 24\hat{k}$

D. none of these

Answer: B



[Watch Video Solution](#)

71. The moment about a line through the origin having the direction of

$$112\hat{i} - 4\hat{j} - 3\hat{k} \text{ is}$$

A. $\frac{760}{13}$

B. $\frac{-760}{13}$

C. $\frac{76}{13}$

D. $\frac{760}{3}$

Answer: B



[Watch Video Solution](#)

72. The moment of the couple consisting of the force through the point

$$2\hat{i} - 3\hat{j} - \hat{k} \text{ is}$$

A. 5

B. $5\sqrt{5}$

C. $\sqrt{5}$

D. 25

Answer: B



[Watch Video Solution](#)

Section I Solved Mcqs

1. The length of the longer diagonal of the parallelogram constructed on

$5\vec{a} + 2\vec{b}$ and $\vec{a} - 3\vec{b}$, if it is given that

$$|\vec{a}| = 2\sqrt{2}, |\vec{b}| = 3 \text{ and } \vec{a} \cdot \vec{b} = \frac{\pi}{4} \text{ is}$$

A. 15

B. $\sqrt{3}$

C. $\sqrt{593}$

D. $\sqrt{369}$

Answer: C



Watch Video Solution

2. If the vectors $\vec{a} = (c \log_2 x) \hat{k}$ make an obtuse angle for any $x \neq (0, \infty)$ then c belongs to

A. $(-\infty, 0)$

B. $(-\infty, -4/3)$

C. $(-4/3, 0)$

D. $(-4/3, \infty)$

Answer: C



Watch Video Solution

3. Let \vec{a} , \vec{b} , \vec{c} represent respectively \vec{BC} , \vec{CA} and \vec{AB} where ABC is a triangle, Then ,

A. $\vec{a} + \vec{b} = \vec{c}$

B. $\vec{b} + \vec{c} = \vec{a}$

C. $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a}$

D. $\left[\begin{matrix} \vec{a} & \vec{b} & \vec{c} \end{matrix} \right] = \left[\begin{matrix} \vec{b} & \vec{c} & \vec{a} \end{matrix} \right] = \left[\begin{matrix} \vec{c} & \vec{a} & \vec{b} \end{matrix} \right] \neq 0$

Answer: C



Watch Video Solution

4. The vector $\hat{i} + x\hat{j} + 3\hat{k}$ is rotated through an angle θ and doubled in magnitude, then it becomes $4\hat{i} + (4x - 2)\hat{j} + 2\hat{k}$. Then values of x are

(A) $-\frac{2}{3}$ (B) $\frac{1}{3}$ (C) $\frac{2}{3}$ (D) 2

A. $-2/3, 2$

B. $1/3, 2$

C. $2/3, -2$

D. $2, -1/3$

Answer: A

 [Watch Video Solution](#)

5. If \vec{a}, \vec{b} are unit vectors such that the vector $\vec{a} + 3\vec{b}$ is perpendicular to $7\vec{a} - \vec{b}$ and $\vec{a} - 4\vec{b}$ is perpendicular to $7\vec{a} - 2\vec{b}$ then the angle between \vec{a} and \vec{b} is

A. $\pi/6$

B. $\pi/4$

C. $\pi/3$

D. $\pi/2$

Answer: C



Watch Video Solution

6. if G is the centroid of $\triangle ABC$ such that \vec{GB} and \vec{GC} are inclined at an obtuse angle, then

A. $5a^2 > b^2 + c^2$

B. $5c^2 > a^2 + b^2$

C. $5b^2 > a^2 + c^2$

D. none of these

Answer:



Watch Video Solution

7. A vector of magnitude 4 which is equally inclined to the vectors $\hat{i} + \hat{j}$, $\hat{j} + \hat{k}$ and $\hat{k} + \hat{i}$, is

A. $\frac{4}{\sqrt{3}}(\hat{i} - \hat{j} - \hat{k})$

B. $\frac{4}{\sqrt{3}}(\hat{i} + \hat{j} - \hat{k})$

C. $\frac{4}{\sqrt{3}}(\hat{i} + \hat{j} + \hat{k})$

D. none of these

Answer: C**Watch Video Solution**

8. Unit vectors equally inclined to the vectors

$$\hat{i}, \frac{1}{3}(-2\hat{i} + \hat{j} + 2\hat{k}) = \pm \frac{4}{\sqrt{3}}(4\hat{j} + 3\hat{k}) \text{ are}$$

A. $\pm \frac{1}{\sqrt{51}}(\hat{i} - 5\hat{j} + 5\hat{k})$

B. $\pm \frac{1}{\sqrt{51}}(\hat{i} - 5\hat{j} - 5\hat{k})$

C. $\pm \frac{1}{\sqrt{t}}51(\hat{i} + 5\hat{j} + 5\hat{k})$

D. none of these

Answer: A



Watch Video Solution

9. If a, b, c are the $p^{\text{th}}, q^{\text{th}}$ and r^{th} terms of G.P then the angle between the _____ vector

$$\vec{u} = (\log a)\hat{i} + (\log b)\hat{j} + (\log c)\hat{k} \text{ and } \vec{v} = (q - r)\hat{i} + (r - p)\hat{j} + (r - p)\hat{k}$$

, is

A. $\frac{\pi}{3}$

B. $\frac{\pi}{6}$

C. π

D. $\frac{\pi}{2}$

Answer: D



Watch Video Solution

10. If a, b, c are the $p^{\text{th}}, q^{\text{th}}$, and r^{th} terms of a HP, then the vectors

$$\vec{u} = a^{-1}\hat{i} + b^{-1}\hat{j} + c^{-1}\hat{k} \text{ and } \vec{v} = (q - r)\hat{i} + (q - r)\hat{j} + (r - p)\hat{k} + (p - r)\hat{k}$$

A. are parallel

B. are orthogonal

C. satisfy $\vec{u} \cdot \vec{v} = 1$

D. satisfy $|\vec{u} \times \vec{v}| = \hat{i} + \hat{j} + \hat{k}$

Answer: B



Watch Video Solution

11. Let a, b, c denote the lengths of the sides of a triangle such that

$$(a - b)\vec{u} + (b - c)\vec{v} + (c - a)(\vec{u} \times \vec{v}) = \vec{0}$$

For any two non-collinear vectors \vec{u} and \vec{v} , then the triangle is

A. right angled

B. equilateral

C. isosceles

D. obtuse angled

Answer: B



Watch Video Solution

12. The vectors \vec{a} , \vec{b} and $\vec{u} \times \vec{u}$ are of the same length and taken pairwise they form equal angles. If $\vec{a} = \hat{i} + \hat{j}$ and $\vec{b} = \hat{j} + \hat{k}$ then \vec{c} is equal to

A. $\hat{i} + \hat{k}, \frac{1}{3}(-\hat{i} + 4\hat{j} - \hat{k})$

B. $\hat{i} + 2\hat{j} + 3\hat{k}, \hat{i} + \hat{j}$

C. $-\hat{i} + \hat{j} + 2\hat{k}, \hat{i} + \hat{k}$

D. $\frac{1}{3}(-\hat{i} + 4\hat{j} - \hat{k}), \hat{j} + \hat{k}$

Answer: A



Watch Video Solution