



# MATHS

BOOKS - OBJECTIVE RD SHARMA

MATHS VOL I (HINGLISH)

TANGENTS AND NORMALS

## Illustration

1. For the curve  $x = t^2 - 1$ ,  $y = t^2 - t$ , the tangent line is perpendicular to  $x$ -axis, then

$$t = \text{(i)} 0 \text{ (ii)} \infty \text{ (iii)} \frac{1}{\sqrt{3}} \text{ (iv)} - \frac{1}{\sqrt{3}}$$

A.  $t = 0$

B.  $t = \infty$

C.  $t = 1/\sqrt{3}$

D.  $t = -1/\sqrt{3}$

**Answer: A**



**Watch Video Solution**

2. The tangent to a given curve is perpendicular to x-axis, if

A.  $\frac{dy}{dx} = 0$

B.  $\frac{dy}{dx} = 1$

C.  $\frac{dx}{dy} = 0$

D.  $\frac{dx}{dy} = 1$

**Answer: C**



**Watch Video Solution**

3. If normal of the curve is parallel to x axis  
then

A.  $\frac{dy}{dx} = 0$

B.  $\frac{dy}{dx} = 1$

C.  $\frac{dx}{dy} = 0$

D.  $\frac{dx}{dy} = 1$

**Answer: C**



**Watch Video Solution**

4. If the tangent to the curve  $xy + ax + by = 0$  at  $(1, 1)$  is inclined at an angle  $\tan^{-1} 2$  with x-axis, then find  $a$  and  $b$  ?

A.  $a = 1, b = 2$

B.  $a = 1, b = -2$

C.  $a = -1, b = 2$

D.  $a = -1, b = -2$

**Answer: B**



**Watch Video Solution**

5. The point on the curve  $y^2 = x$  where tangent makes  $45^\circ$  angle with x-axis, is

A.  $(1/2, 1/4)$

B.  $(1/4, 1/2)$

C.  $(4, 2)$

D.  $(1, 1)$

**Answer: B**



**Watch Video Solution**

6. The point on the curve  $y = 12x - x^2$  where the tangent is parallel to x-axis, is

A. (0, 0)

B. (2, 16)

C. (3, 9)

D. none of these

**Answer: D**



**Watch Video Solution**

7. The slope of the tangent to the curve

$x = t^2 + 3t - 8, y = 2t^2 - 2t - 5$  at the

point  $(2, 1)$  is (A)  $\frac{22}{7}$  (B)  $\frac{6}{7}$  (C)  $\frac{7}{6}$  (D)  $-\frac{6}{7}$

A.  $\frac{22}{7}$

B.  $\frac{6}{7}$

C. -6

D. none of these

**Answer: B**



**Watch Video Solution**



8. For the curve  $x = 3 \cos \theta, y = 3 \sin \theta, 0 \leq \theta \leq \pi$ , the tangent is parallel to the x-axis, where  $\theta =$

A.  $\pi$

B. 0

C.  $\frac{\pi}{3}$

D.  $\frac{\pi}{2}$

**Answer: D**



**Watch Video Solution**

9. The point on the curve  $y = (x - 1)(x - 2)$  at which the tangent makes an angle of  $135^\circ$  with the positive direction of x-axis has coordinates

A. (1, 0)

B. (0, 1)

C. (-1, 0)

D. (0, -1)

**Answer: A**



**Watch Video Solution**

10. If the normal to the curve  $y = f(x)$  at the point  $(3, 4)$  makes an angle  $\frac{3\pi}{4}$  with the positive x-axis, then  $f'(3) =$  (a)  $-1$  (b)  $-\frac{3}{4}$  (c)  $\frac{4}{3}$  (d)  $1$

A.  $-1$

B.  $-\frac{3}{4}$

C.  $-\frac{4}{3}$

D.  $1$

**Answer: C**



**Watch Video Solution**

**11.** The point(s) on the curve  $y^3 + 3x^2 = 12y$  where the tangent is vertical, is(are) ??

$\left( \pm \frac{4}{\sqrt{3}}, -2 \right)$  (b)  $\left( \pm \sqrt{\frac{11}{3}}, 1 \right)$  (0, 0)

(d)  $\left( \pm \frac{4}{\sqrt{3}}, 2 \right)$

A.  $\left( \pm \frac{4}{\sqrt{3}}, -2 \right)$

B.  $\left( \pm \sqrt{\frac{\pi}{3}}, 1 \right)$

C. (0, 0)

D.  $\left( \pm \frac{4}{\sqrt{3}}, 2 \right)$

**Answer: D**



**Watch Video Solution**

**12.** If the slope of tangent to the curve

$y = \frac{ax}{b - x}$  at the point (1, 1) is 2, then

A.  $a = 1, b = -2$

B.  $a = -1, b = 2$

C.  $a = 1, b = 2$

D. none of these

**Answer: C**



**Watch Video Solution**

**13.** The slope of the tangent to the curve

$(y - x^5)^2 = x(1 + x^2)^2$  at the point  $(1, 3)$  is.

A. 4

B. 6

C. 8

D. 2

**Answer: C**



**Watch Video Solution**

**14.** The tangent to the curve  $y = x^3$  at the point  $P(t, t^3)$  cuts the curve again at point Q.

Then, the coordinates of Q are

A. (0, 0)

B.  $(2t, 4t^3)$

C.  $(2t, 8t^3)$

D.  $(-2t, -8t^3)$

**Answer: D**



**Watch Video Solution**

**15.** The point at which the tangent to the curve

$y = x^2 - 4x$  is parallel to x-axis, is

A.  $(0, 4)$



B. (-2, 4)

C. (2,4)

D. (2, -4)

**Answer: D**



**Watch Video Solution**

**16.** The curve  $y - e^{xy} + x = 0$  has a vertical tangent at the point :

A. (1, 1)

B. at no point

C. (0, 1)

D. (1, 0)

**Answer: D**



**Watch Video Solution**

**17.** The angle between the tangents to the curve  $y = x^2 - 5x + 6$  at the point (2, 0) and (3, 0), is

A.  $\pi / 3$

B.  $\pi / 2$

C.  $\pi / 6$

D.  $\pi / 4$

**Answer: B**



**Watch Video Solution**

**18.** The equation of the normal to the curve

$$y = x + \sin x \cos x \quad \text{at} \quad x = \frac{\pi}{2}, \text{ is}$$

A.  $x = 2$

B.  $x = \pi$

C.  $x + \pi = 0$

D.  $2x = \pi$

**Answer: D**



**Watch Video Solution**

**19.** The equation of the normal to the curve  $y = \sin x$  at  $(0, 0)$  is

A.  $x = 0$

B.  $y = 0$

C.  $x + y = 0$

D.  $x - y = 0$

**Answer: C**



**Watch Video Solution**

**20.** The equation of the normal to the curve

$y = x(2 - x)$  at the point  $(2, 0)$  is

A.  $x - 2y = 2$

B.  $x - 2y + 2 = 0$

C.  $2x + y = 4$

D.  $2x + y - 4 = 0$

**Answer: A**



**Watch Video Solution**

**21.** If the equation of the tangent to the curve

$y^2 = ax^3 + b$  at point  $(2, 3)$  is  $y = 4x - 5$ ,

then find the values of  $a$  and  $b$ .

A.  $a = 2, b = 7$

B.  $a = 7, b = 2$

C.  $a = 2, b = -7$

D.  $a = -2, b = 7$

**Answer: C**



**Watch Video Solution**

**22.** The tangent to the curve  $y = e^{2x}$  at the point  $(0,1)$  meets X-axis at

A. (0, 2)

B. (2, 0)

C.  $(-1/2, 0)$

D. none of these

**Answer: C**



**Watch Video Solution**

**23.** If tangent to curve  $2y^3 = ax^2 + x^3$  at point  $(a, a)$  cuts off intercepts  $\alpha, \beta$  on coordinate axes, where  $\alpha^2 + \beta^2 = 61$ , then the



value of 'a' is equal to (A) 20 (B) 25 (C) 30

(D)-30

A.  $\pm 30$

B.  $\pm 5$

C.  $\pm 6$

D.  $\pm 61$

**Answer: A**



**Watch Video Solution**

24. The equation of the tangent to the curve  $y = 1 - e^{x/2}$  at the point of intersection with the y-axis, is

A.  $x + 2y = 0$

B.  $2x + y = 0$

C.  $x - y = 2$

D. none of these

**Answer: A**



Watch Video Solution

25. The normal to the curve  $x = a(1 + \cos \theta)$ ,  $y = a \sin \theta$  at ' $\theta$ ' always passes through the fixed point

A.  $(a, a)$

B.  $(a, 0)$

C.  $(0, a)$

D. none of these

**Answer: B**





26. The area of a triangle formed by a tangent to the curve  $2xy = a^2$  and the coordinate axes, is

A.  $2a^2$

B.  $a^2$

C.  $3a^2$

D. none of these

**Answer: B**



Watch Video Solution

27. If the tangent at a point on the ellipse  $\frac{x^2}{27} + \frac{y^2}{3} = 1$  meets the coordinate axes at  $A$  and  $B$ , and the origin, then the minimum area (in sq. units) of the triangle  $OAB$  is:

A. 9

B.  $\frac{9}{2}$

C.  $9\sqrt{3}$

D.  $3\sqrt{3}$

**Answer: A**



**Watch Video Solution**

**28.** Find the equation of the normal to the curve  $y = (1 + y)^y + \sin^{-1}(\sin^2 x)$  at  $x = 0$ .

A.  $x + y = 2$

B.  $x + y = 1$

C.  $x - y = 1$

D. none of these

**Answer: B**



**Watch Video Solution**

**29.** The normal to the curve  $x^2 + 2xy - 3y^2 = 0$ , at  $(1, 1)$ :

A. meets the curve again in the third quadrant.

B. Meets the curve again the fourth quadrant .

C. does not meet the curve again.

D. meets the curve again in the second quadrant.

**Answer: B**



**Watch Video Solution**

**30.** The area bounded by the coordinate axes and normal to the curve  $y = \log_e x$  at the point  $P(1, 0)$ , is



A. 1 sq. unit

B. 2 sq. units

C.  $\frac{1}{2}$  sq. unit

D. none of these

**Answer: C**



**Watch Video Solution**

**31.**

Consider

$$f(x) = \tan^{-1} \left( \sqrt{\frac{1 + \sin x}{1 - \sin x}} \right), x \in \left( 0, \frac{\pi}{2} \right).$$

A normal to  $y = f(x)$  at  $x = \frac{\pi}{6}$  also passes through the point: (1)  $(0, 0)$  (2)  $\left(0, \frac{2\pi}{3}\right)$  (3)  $\left(\frac{\pi}{6}, 0\right)$  (4)  $\left(\frac{\pi}{4}, 0\right)$

A.  $(0, 0)$

B.  $\left(0, \frac{2\pi}{3}\right)$

C.  $\left(\frac{\pi}{6}, 0\right)$

D.  $\left(\frac{\pi}{4}, 0\right)$

**Answer: B**



**Watch Video Solution**

32. Let  $C$  be a curve given by  $y = 1 + \sqrt{4x - 3}$ ,  $x > \frac{3}{4}$ . If  $P$  is a point on  $C$  such that the tangent at  $P$  has slope  $\frac{2}{3}$ , then a point through which the normal at  $P$  passes, is

A.  $(3, -4)$

B.  $(1, 7)$

C.  $(4, -3)$

D.  $(2, 3)$

**Answer: B**



**Watch Video Solution**

33. The normal to the curve  $y(x - 2)(x - 3) = x + 6$  at the point where the curve intersects the  $y$ -axis, passes

through the point : (1)  $\left(\frac{1}{2}, -\frac{1}{3}\right)$  (2)  $\left(\frac{1}{2}, \frac{1}{3}\right)$

(3)  $\left(-\frac{1}{2}, -\frac{1}{2}\right)$  (4)  $\left(\frac{1}{2}, 1\right)$

A.  $\left(-\frac{1}{2}, -\frac{1}{2}\right)$

B.  $\left(\frac{1}{2}, \frac{1}{2}\right)$

C.  $\left(\frac{1}{2}, -\frac{1}{3}\right)$

D.  $\left(\frac{1}{2}, \frac{1}{3}\right)$

**Answer: B**



**Watch Video Solution**

**34.** If the curves  $y = a^x$  and  $y = e^x$  intersect at an angle  $\alpha$ , then  $\tan \alpha$  equals

A.  $\left| \frac{\log_e a}{1 + \log_e a} \right|$

B.  $\left| \frac{1 + \log_e a}{1 + \log_e a} \right|$

C.  $\left| \frac{\log_e a - 1}{\log_e a + 1} \right|$

D. none of these

**Answer: C**



**Watch Video Solution**

**35.** The angle of intersection of the curves

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ and } x^2 + y^2 = ab, \text{ is}$$

A.  $\tan^{-1} \left( \frac{a - b}{\sqrt{ab}} \right)$

B.  $\tan^{-1} \left( \frac{a + b}{\sqrt{ab}} \right)$

C.  $\tan^{-1} \left( \frac{a - b}{2\sqrt{ab}} \right)$

D. none of these

**Answer: A**



**Watch Video Solution**

**36.** The curves

$$x^3 - 3xy^2 = a \text{ and } 3x^2y - y^3 = b, \text{ where } a$$

and  $b$  are constants, cut each other at an

angle of

A.  $\frac{\pi}{3}$

B.  $\frac{\pi}{4}$

C.  $\frac{\pi}{2}$

D. none of these

**Answer: C**



**Watch Video Solution**

**37.** If the curves  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and  $\frac{x^2}{c^2} + \frac{y^2}{d^2} = 1$  intersect orthogonally, then

A.  $a^2 - b^2 = c^2 - d^2$

B.  $a^2 - c^2 = b^2 - d^2$



$$C. a^2b^2 = c^2d^2$$

$$D. \frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{c^2} + \frac{1}{d^2}$$

**Answer: A**



**Watch Video Solution**

**38.** The curves  $ax^2 + by^2 = 1$  and  $Ax^2 + By^2 = 1$  intersect orthogonally, then

$$A. \frac{1}{a} + \frac{1}{A} = \frac{1}{b} + \frac{1}{B}$$

$$\text{B. } \frac{1}{a} - \frac{1}{A} = \frac{1}{b} - \frac{1}{B}$$

$$\text{C. } \frac{1}{a} + \frac{1}{b} = \frac{1}{B} - \frac{1}{A}$$

$$\text{D. } \frac{1}{a} + \frac{1}{b} = \frac{1}{A} + \frac{1}{B}$$

**Answer: B**



**Watch Video Solution**

**39.** If the curves  $2x^2 + 3y^2 = 6$  and  $ax^2 + 4y^2 = 4$  intersect orthogonally, then  $a =$

A. 2

B. 1

C. 3

D. none of these

**Answer: A**



**Watch Video Solution**

**40.** The curves  $x = y^2$  and  $xy = a^3$  cut orthogonally at a point, then  $a =$

A.  $\frac{1}{3}$

B. 3

C. 2

D.  $\frac{1}{2}$

**Answer: D**



**Watch Video Solution**

41. For a curve  $\frac{(\text{length of normal})}{(\text{length of tangent})}$  is equal to

A. subtangent

B. subnormal

C. slope of tangent

D. slope of normal

**Answer: C**



**Watch Video Solution**

**42.** For a curve  $(\text{length of normal})^2 / (\text{length of tangent})^2$  is equal to

A. (subnormal)/(subtangent )

B. (subtangent)/(subnormal)

C. (tangent)/(normal)

D. constant

**Answer: A**



**Watch Video Solution**

43. At any point of a curve  $\sqrt{\frac{\text{subnormal}}{\text{subtangent}}}$  is equal to

A. the abscissa of that point

B. the ordinate of that point

C. slope of the tangent at that point

D. slope of the normal at that point

**Answer: C**



**Watch Video Solution**

**44.** At any point of a curve ( subtangent ) x  
(subnormal) is equal to the square of the-

A. slope of the tangent at that point

B. slope of the normal at that point

C. abscissa of that point

D. ordinate of that point

**Answer: D**



**Watch Video Solution**

**45.** The subtangent at any point on the curve

$x^m y^n = a^{m+n}$  varies as



A. (abscissae)<sup>2</sup>

B. (ordinate)<sup>2</sup>

C. abscissa

D. ordinate

**Answer: C**



**Watch Video Solution**

**46.** Find the equation of the tangent and the normal at the point 't', on the curve

$$x = a \sin^3 t, y = b \cos^3 t.$$

A.  $4CT^2 = ON^2 = a^2$

B. the length of the tangent =  $\left| \frac{y}{\cos t} \right|$

C. the length of the normal =  $\left| \frac{y}{\sin t} \right|$

D. all the above

**Answer: C**



**Watch Video Solution**

**47.** The length of the normal to the curve

$y = a \left( \frac{e^{-x/a} + e^{x/a}}{2} \right)$  at any point varies as

the

A. abscissa of the point

B. ordinate of the point

C. square of the abscissa of the point

D. square of the ordinate of the point

**Answer: D**



**Watch Video Solution**

**48.** If at any point on a curve the subtangent and subnormal are equal, then the tangent is equal to

A. ordinate

B.  $\sqrt{2}$  ordinate

C.  $\sqrt{2(\text{ordinate})}$

D. none of these

**Answer: B**



**Watch Video Solution**

**49.** Find the length of normal to the curve

$$x = a(\theta + \sin \theta), y = a(1 - \cos \theta) \text{ at } \theta = \frac{\pi}{2}.$$

A.  $2a$

B.  $a\sqrt{2}$

C.  $a/2$

D.  $a/\sqrt{2}$

**Answer: B**



**Watch Video Solution**

## Section I Solved Mcqs

1. The number of possible tangents which can be drawn to the curve  $4x^2 - 9y^2 = 36$ , which are perpendicular to the straight line  $5x + 2y - 10 = 0$ , is zero (b) 1 (c) 2 (d) 4

A.  $5(y - 3) = 2\left(x - \frac{\sqrt{117}}{2}\right)$

B.  $2x - 5y + 10 - 2\sqrt{18} = 0$

C.  $2x - 5y - 10 - 2\sqrt{18} = 0$

D. none of these

**Answer: D**



**Watch Video Solution**

2. Let P be any point on the curve  $x^{2/3} + y^{2/3} = a^{2/3}$ . Then the length of the segment of the tangent between the coordinate axes is of length

A.  $3a$

B.  $4a$

C.  $5a$

D. a

**Answer: D**



**Watch Video Solution**

3. The distance between the origin and the tangent to the curve  $y = e^{2x} + x^2$  drawn at the point  $x = 0$  is

A.  $\frac{1}{\sqrt{5}}$

B.  $\frac{2}{\sqrt{5}}$



C.  $\frac{-1}{\sqrt{5}}$

D.  $\frac{2}{\sqrt{3}}$

**Answer: A**



**Watch Video Solution**

4. Find the point of intersection of the tangents drawn to the curve  $x^2y = 1 - y$  at the points where it is intersected by the curve  $xy = 1 - y$ .

A.  $(0, -1)$

B.  $(1, 1)$

C.  $(0, 1)$

D. none of these

**Answer: C**



**Watch Video Solution**

5. The equation of the tangent to the curve

$y = (2x - 1)e^{2(1-x)}$  at the point of its

maximum, is

A.  $y = -1 = 0$

B.  $x - 1 = 0$

C.  $x + y - 1 = 0$

D.  $x - y + 1 = 0$

**Answer: A**



**Watch Video Solution**

**6.** If the sum of the squares of the intercepts on the axes cut off by tangent to the curve

$x^{\frac{1}{3}} + y^{\frac{1}{3}} = a^{\frac{1}{3}}$ ,  $a > 0$  at  $\left(\frac{a}{8}, \frac{a}{8}\right)$  is 2, then

$a =$  1 (b) 2 (c) 4 (d) 8

A. 1

B. 2

C. 4

D. 8

**Answer: C**



**Watch Video Solution**

7. The point on the curve  $3y = 6x - 5x^3$  the normal at Which passes through the origin, is

A.  $(1, 1/3)$

B.  $(1/3, 1)$

C.  $(2, -28/3)$

D.  $(-1, -1/3)$

**Answer: A**



**Watch Video Solution**

8. If the tangent at any point on the curve  $x^4 + y^4 = c^4$  cuts off intercepts  $a$  and  $b$  on the coordinate axes, the value of  $a^{-\frac{4}{3}} + b^{-\frac{4}{3}}$  is

A.  $c^{-4/3}$

B.  $c^{-1/2}$

C.  $c^{1/2}$

D. none of these

**Answer: A**



**Watch Video Solution**

9. If the tangent at  $(1, 1)$  on  $y^2 = x(2 - x)^2$  meets the curve again at  $P$ , then find coordinates of  $P$ .

A.  $(4, 4)$

B.  $(-1, 2)$

C.  $(9/4, 3/8)$

D. none of these

**Answer: C**



**Watch Video Solution**

10. The two curves  $x^3 - 3xy^2 + 2 = 0$  and  $3x^2y - y^3 - 2 = 0$

- A. cut at right angles
- B. touch each other
- C. cut at an angle  $\pi / 3$
- D. cut at an angle  $\pi / 4$

**Answer: A**



**Watch Video Solution**



11. A curve with equation of the form  $y = ax^4 + bx^3 + cx + d$  has zero gradient at the point  $(0, 1)$  and also touches the  $x$  - axis at the point  $(-1, 0)$  then the value of  $x$  for which the curve has a negative gradient are:  
 $x \geq -1$  b.  $x < 1$  c.  $x < -1$  d.  $-1 \leq x \leq 1$

A.  $x > -1$

B.  $x < 1$

C.  $x < -1$

D.  $-1 \leq x \leq 1$

**Answer: C**



**Watch Video Solution**

**12.** For the curve  $y = ce^{x/a}$ , which one of the following is incorrect?

A. subtangent is constant

B. subnormal varies as the square of the ordinate

C. tangent at  $(x_1, y_1)$  on the curve intersects the x-axis at a distance of  $(x_1 - a)$  from the origin.

D. equation of normal at the point where the curve cuts y-axis is  $cy + ax = c$

**Answer: D**



**Watch Video Solution**

13. If  $m$  is the slope of the tangent to the curve

$$e^y = 1 + x^2, \text{ then}$$

A.  $|m| > 1$

B.  $m < 1$

C.  $|m| < 1$

D.  $|m| \leq 1$

**Answer: D**



**Watch Video Solution**

14. The tangent to the curve  $y = e^x$  drawn at the point  $(c, e^c)$  intersects the line joining  $(c - 1, e^{c-1})$  and  $(c + 1, e^{c+1})$  (a) on the left of  $n = c$  (b) on the right of  $n = c$  (c) at no points (d) at all points

- A. on the left of  $x=c$
- B. on the right of  $x=c$
- C. at no point
- D. at all point

**Answer: A**





Watch Video Solution

15. If  $x + y = k$  is normal to  $y^2 = 12x$ , then  $k$  is 3 (b) 9 (c)  $-9$  (d)  $-3$

A. 3

B. 9

C. -9

D. -3

**Answer: B**



Watch Video Solution

16. If the line  $ax + by + c = 0$  is a tangent to the curve  $xy = 9$ , then

A.  $a > 0, b > 0$

B.  $a > 0, b < 0$

C.  $a < 0, b > 0$

D.  $a < 0, b < 0$

**Answer: A::D**



**Watch Video Solution**

17. The lengths of tangent, subtangent, normal and subnormal for the curve  $y = x^2 + x - 1$  at (1,1) are A,B,C and D respectively, then their increasing order is

A. B,D,A,C

B. B,A,C,D

C. A,B,C,D

D. B,A,D,C

**Answer: D**



**Watch Video Solution**



18. If at each point of the curve  $y = x^3 - ax^2 + x + 1$ , the tangent is inclined at an acute angle with the positive direction of the x-axis, then  $a > 0$  (b)  $a < -\sqrt{3} - \sqrt{3} \leq a \leq \sqrt{3}$  (d) *none of these*

A.  $a > 0$

B.  $a \leq \sqrt{3}$

C.  $|a| \leq \sqrt{3}$

D. none of these

**Answer: C**



**Watch Video Solution**

**19.** If the line  $y = 2x$  touches the curve  $y = ax^2 + bx + c$  at the point where  $x=1$  and the curve passes through the point  $(-1,0)$ , then

A.  $a = \frac{1}{2}, b = 1, c = \frac{1}{2}$

B.  $a = 1, b = \frac{1}{2}, c = \frac{1}{2}$

C.  $a = \frac{1}{2}, c = \frac{1}{2}, b = 1$

D. none of these

**Answer: A**



**Watch Video Solution**

20. If the line joining the points

$(0, 3)$  and  $(5, -2)$  is a tangent to the curve

$y = \frac{C}{x + 1}$ , then the value of  $c$  is 1 (b)  $-2$  (c)

4 (d) none of these

A. 1

B. -2

C. 4

D. none of these

**Answer: C**



**Watch Video Solution**

21. If  $y = f(x)$  be the equation of the line touching the line  $y = 2x + 3$  at  $x = 2$ , then

A.  $f'(2) = 3$

B.  $2f(2) = 7f'(2)$

C.  $f(2) + f'(2) + f''(2) = 2$

D. none of these

**Answer: B**



**Watch Video Solution**

22. The slope of the tangent of the curve

$y = \int_0^x \frac{dx}{1+x^3}$  at the point where  $x = 1$  is

A.  $\frac{1}{2}$

B. 1

C.  $\frac{1}{4}$

D. non-existent

**Answer: A**



**Watch Video Solution**

**23.** Prove that the curve  $y = e^{|x|}$  cannot have a unique tangent line at the point  $x = 0$ .

Find the angle between the one-sided tangents to the curve at the point  $x = 0$ .

A.  $\frac{\pi}{4}$

B.  $\frac{\pi}{6}$

C.  $\frac{\pi}{2}$

D.  $\frac{\pi}{3}$

**Answer: C**



**Watch Video Solution**

**24.** The curve  $y = ax^3 + bx^2 + cx + 5$  touches the x-axis at  $P(-2, 0)$  and cuts the

y-axis at the point  $Q$  where its gradient is 3.

Find the equation of the curve completely.

A.  $a = \frac{1}{2}, b = -\frac{3}{4}, c = 3$

B.  $a = -\frac{1}{2}, b = -\frac{3}{4}, c = 3$

C.  $a = \frac{1}{2}, b = \frac{3}{4}, c = 3$

D. none of these

**Answer: B**



**Watch Video Solution**



25. If the curve  $y = x^2 + bx + c$  touches the line  $y = x$  at the point (1,1), then the set of values of  $x$  for which the curve has a negative gradient is

A.  $(-\infty, 1/2)$

B.  $(1/2, \infty)$

C.  $(-\infty, -1/2)$

D.  $(-1/2, \infty)$

**Answer: A**



Watch Video Solution

26. The sum of the intercepts made on the axes of coordinates by any tangent to the curve  $\sqrt{x} + \sqrt{y} = \sqrt{a}$  is equal to

A.  $2a$

B.  $a$

C.  $\frac{a}{2}$

D. none of these

**Answer: B**





27. The angle at which the curves  $y = \sin x$  and  $y = \cos x$  intersect in  $[0, \pi]$ , is

A.  $\pm \tan^{-1} \sqrt{2}$

B.  $\pm \tan^{-1} 2\sqrt{2}$

C.  $\pm \tan^{-1} \frac{1}{\sqrt{2}}$

D. none of these

**Answer: B**



Watch Video Solution

28. If the tangent at each point of the curve

$$y = \frac{2}{3}x^3 - 2ax^2 + 2x + 5$$

makes an acute angle with the positive direction of x-axis, then

A.  $a \geq 1$

B.  $-1 \leq a \leq 1$

C.  $a \leq -1$

D. none of these

**Answer: B**



**Watch Video Solution**

**29.** If the parabolas  $y = x^2 + ax + b$  and  $y = x(c - x)$  touch each other at the point  $(1,0)$ , then  $a + b + c =$

A. 1

B. -1

C. 0

D. none of these

**Answer: C**



**Watch Video Solution**

**30.** Let  $y = f(x)$  be a parabola, having its axis parallel to the y-axis, which is touched by the line  $y = x$  at  $x = 1$ . Then,  $2f(0) = 1 - f'(0)$

(b)  $f(0) + f'(0) + f^0 = 1$     $f'(1) = 1$    (d)

$f'(0) = f'(1)$

A.  $f'(0) = f'(1)$

B.  $f'(1) = -1$

$$C. f(0) + f'(0) + f''(0) = 1$$

$$D. 2f(0) = 1 - f'(0)$$

**Answer: D**



**Watch Video Solution**

**31.** Find the value of  $n \in \mathbb{N}$  such that the

curve  $\left(\frac{x}{a}\right)^n + \left(\frac{y}{b}\right)^n = 2$  touches the

straight line  $\frac{x}{a} + \frac{y}{b} = 2$  at the point  $(a, b)$ .

A.  $(b, a)$

B.  $(a, b)$

C.  $(1, 1)$

D.  $\left(\frac{1}{b}, \frac{1}{a}\right)$

**Answer: B**



**Watch Video Solution**

**32.** The normal to the curve  $2x^2 + y^2 = 12$  at

the point  $(2, 2)$  cuts the curve again at

$\left(-\frac{22}{9}, -\frac{2}{9}\right)$  (b)  $\left(\frac{22}{9}, \frac{2}{9}\right)$   $(-2, -2)$

(d) none of these



A.  $(-22/9, -2/9)$

B.  $(22/9, 2/9)$

C.  $(-2, -2)$

D. none of these

**Answer: A**



**Watch Video Solution**

**33.** A tangent to the curve  $y = \int_0^x |x| dt$ , which is parallel to the line  $y = x$ , cuts off an

intercept from the y-axis is equal to

A. 1

B.  $-\frac{1}{2}, \frac{1}{2}$

C.  $\frac{1}{2}, 1$

D. -1

**Answer: B**



**Watch Video Solution**

34. The equation of the normal to the curve  $y = e^{-2|x|}$  at the point where the curve cuts the line  $x = -\frac{1}{2}$ , is

A.  $2e(ex + 2y) = 4 - e^2$

B.  $2e(ex - 2y) = e^2 - 4$

C.  $2e(ey - 2x) = e^2 - 4$

D.  $2e(ey + 2x) = e^2 - 4$

**Answer: A**



**Watch Video Solution**

**35.** The equation of the normal to the curve

$y = x^{-x}$  at the point of its maximum is

A.  $x = e$

B.  $x = e^{-1}$

C.  $y = e$

D.  $y = e^{-1}$

**Answer: B**



**Watch Video Solution**

36. The abscissa of a point on the curve  $xy = (a + x)^2$ , the normal which cuts off numerically equal intercepts from the coordinate axes, is  $-\frac{1}{\sqrt{2}}$  (b)  $\sqrt{2}a$  (c)  $\frac{a}{\sqrt{2}}$  (d)  $-\sqrt{2}a$

A.  $\frac{a}{\sqrt{2}}$

B.  $a$

C.  $\sqrt{2}a$

D.  $-\frac{a}{\sqrt{2}}$

**Answer: A::D**



Watch Video Solution

37. Let  $f(x) = \sin x - \tan x, x \in (0, \pi/2)$

then tangent drawn to the curve  $y = f(x)$  at

any point will

A. lie above the curve

B. lie below the curve

C. nothing can be said

D. be parallel to a fixed line.

**Answer: A**



Watch Video Solution

38. If the tangent at a point  $P$  with parameter  $t$ , on the curve  $x = 4t^2 + 3$ ,  $y = 8t^3 - 1$   $t \in R$  meets the curve again at a point  $Q$ , then the coordinates of  $Q$  are

A.  $\left( \frac{35}{9}, \pm \frac{16\sqrt{2}}{27} - 1 \right)$

B.  $\left( \frac{25}{9}, \pm \frac{11}{7} \right)$

C.  $\left( \frac{35}{9}, \pm \frac{16\sqrt{2}}{27} + 1 \right)$

D. none of these

**Answer: A**



**Watch Video Solution**

**39.** If the tangent to the curve  $xy + ax + by = 0$  at  $(1, 1)$  is inclined at an angle  $\tan^{-1} 2$  with x-axis, then find  $a$  and  $b$ ?

A. 0

B.  $\frac{1}{2}$

C.  $-\frac{1}{2}$

D. none of these



**Answer: B**



**Watch Video Solution**

**40.** The slope of the tangent to the curve

$$y = \int_x^{x^2} \cos^{-1} t^2 dt \text{ at } x = \frac{1}{\sqrt[4]{2}} \text{ is}$$

A.  $\left( \frac{\sqrt[4]{8}}{2} - \frac{3}{4} \right) \pi$

B.  $\left( \frac{\sqrt[4]{8}}{3} - \frac{1}{4} \right) \pi$

C.  $\left( \frac{\sqrt[5]{8}}{4} - \frac{1}{3} \right) \pi$

D. none of these

**Answer: B**



**Watch Video Solution**

**41.** The equation of the curve is  $y = f(x)$ . The tangents at  $[1, f(1)]$ ,  $[2, f(2)]$ , and  $[3, f(3)]$  make angles  $\frac{\pi}{6}$ ,  $\frac{\pi}{3}$ , and  $\frac{\pi}{4}$ , respectively, with the positive direction of x-axis. Then the value

of  $\int_2^3 f'(x)f^x dx + \int_1^3 f^x dx$  is equal to  $-\frac{1}{\sqrt{3}}$  (b)  $\frac{1}{\sqrt{3}}$  (e) 0 (d) none of these

A.  $-\frac{1}{\sqrt{3}}$

B.  $\frac{1}{\sqrt{3}}$

C. 0

D. none of these

**Answer: A**



**Watch Video Solution**

**42.** Let  $C$  be the curve  $y^3 - 3xy + 2 = 0$ . If  $H$  is the set of points on the curve  $C$ , where the tangent is horizontal and  $V$  is the set of points

on the curve  $C$ , where the tangent is vertical,  
then  $H = \dots$  and  $V = \dots$ .

A.

$$H = \{(x, y) : y = 0, x \in \mathbb{R}\}, V = \{(1, 1)\}$$

B.

$$H = \{(x, y) : x = 0, y \in \mathbb{R}\}, V = \{(1, 1)\}$$

C.  $H = \emptyset, V = \{(1, 1)\}$

D.

$$H = \{(1, 1)\}, V = \{(x, y) : y = 0, x \in \mathbb{R}\}$$

**Answer: C**



Watch Video Solution

**43.** If  $\sin^{-1} \theta$  is the acute angle between the curves

$$x^2 + y^2 = 4x \quad \text{and} \quad x^2 + y^2 = 8 \quad \text{at} \quad (2, 2),$$

then  $\theta =$

A. 1

B. 0

C.  $1/\sqrt{2}$

D.  $\sqrt{3}/2$

**Answer: C**



**Watch Video Solution**

**44.** If curve  $x^2 = 9a(9 - y)$  and  $x^2 = a(y + 1)$  intersect orthogonally then value of 'a' is

A. 3

B. 4

C. 5

D. 7

**Answer: B**



**Watch Video Solution**

**45.** The equation of the tangent to the curve

$y = x + \frac{4}{x^2}$ , that is parallel to the x-axis, is

A.  $y = 2$

B.  $y = 3$

C.  $y = 0$

D.  $y = 1$

**Answer: B**



**Watch Video Solution**

**46.** The equation of the normal to the parabola,  $x^2 = 8y$  at  $x = 4$  is

A.  $x + y = 6$

B.  $x + 2y = 0$

C.  $3 - 2y = 0$

D.  $x + y = 2$



**Answer: A**



**Watch Video Solution**

**47.** The intercepts on x-axis made by tangents

to the curve,  $y = \int_0^x |t| dt$ ,  $x \in R$ , which are

parallel to the line  $y = 2x$ , are equal to (1)  $\pm 2$

(2)  $\pm 3$  (3)  $\pm 4$  (4)  $\pm 1$

A.  $\pm 1$

B.  $\pm 2$

C.  $\pm 3$

D.  $\pm 4$

**Answer: A**



**Watch Video Solution**

**48.** The complete set of value of 'a' for which there exists at least one line that is tangent to the graph of the curve  $y = x^3 - ax$  at one point and normal to the graph at another point is given by

A.  $(-\infty, -4/3]$

B.  $[-4/3, \infty)$

C.  $[4/3, \infty)$

D.  $(-\infty, 4/3]$

**Answer: C**



**Watch Video Solution**

**49.** If the tangent at a point  $P$  with parameter  $t$ , on the curve  $x = 4t^2 + 3$ ,  $y = 8t^3 - 1$   $t \in \mathbb{R}$  meets the curve again at a point  $Q$ , then the coordinates of  $Q$  are

A.  $(t^2 + 3, -t^3 - 1)$

B.  $(t^2 + 3, t^3 - 1)$

C.  $(16t^2 + 3, -64t^3 - 1)$

D.  $(4t^2 + 3, -8t^3 - 1)$ .

**Answer: A**



**Watch Video Solution**

**Exercise**

1. The equation of the tangents to  $2x^2 + 3y^2 = 36$  which are parallel to the straight line  $x + 2y - 10 = 0$ , are

A.  $x + 2y = 0$

B.  $x + 2y + \sqrt{\frac{288}{15}} = 0$

C.  $x + 2y + \sqrt{\frac{1}{15}} = 0$

D. none of these

**Answer: D**



**Watch Video Solution**

2. If the area of the triangle included between the axes and any tangent to the curve  $x^n y = a^n$  is constant, then find the value of  $n$ .

A. 1

B. 2

C.  $3/2$

D.  $1/2$

**Answer: A**



**Watch Video Solution**

3. Show that the curves  $x = y^2$  and  $xy = k$  cut at right angles; if  $8k^2 = 1$

A.  $2k^2 = 1$

B.  $4k^2 = 1$

C.  $6k^2 = 1$

D.  $8k^2 = 1$

**Answer: D**



**Watch Video Solution**

4. The normal to the curve

$$x = a(\cos \theta - \theta \sin \theta), y = a(\sin \theta - \theta \cos \theta)$$

at any point,  $\theta$ , is such that

- A. makes a constant angle with x-axis
- B. is at a constant distance from the origin
- C. passes through the origin
- D. satisfies all the three conditions

**Answer: B**



**Watch Video Solution**



5. The equation of the tangent to the curve

$x = t \cos t, y = t \sin t$  at the origin, is

A.  $x = 0$

B.  $y = 0$

C.  $x + y = 0$

D.  $x - y = 0$

**Answer: B**



**Watch Video Solution**

6. The equation of the normal to the curve

$$y^4 = ax^3 \text{ at } (a, a) \text{ is}$$

A.  $x + 2y = 3a$

B.  $3x - 4y + a = 0$

C.  $4x + 3y = 7a$

D.  $4x - 3y = a$

**Answer: C**



**Watch Video Solution**

7. Find the angle between the curve

$$y^2 = 4x + 4 \text{ and } y^2 = 36(9 - x).$$

A.  $30^\circ$

B.  $45^\circ$

C.  $60^\circ$

D.  $90^\circ$

**Answer: D**



**Watch Video Solution**

8. The equation(s) of the tangent(s) to the curve  $y = x^4$  from the point  $(2, 0)$  not on the curve is given by

A.  $y = \frac{4098}{81}$

B.  $y - 1 = 5(x - 1)$

C.  $y = \frac{4096}{81} = \frac{2048}{27} \left( x - \frac{8}{3} \right)$

D.  $y - \frac{32}{243} = \frac{80}{81} \left( x - \frac{2}{3} \right)$

**Answer: C**



**Watch Video Solution**

9. The point on the curve  $\sqrt{x} + \sqrt{y} = \sqrt{a}$ , the normal at which is parallel to the x-axis, is

A. (0, 0)

B. (0, a)

C. (a, 0)

D. (a, a)

**Answer: B**



**Watch Video Solution**

10. The length of the Sub tangent at  $(2, 2)$  to the curve  $x^5 = 2y^4$  is

A.  $5/2$

B.  $8/5$

C.  $2/5$

D.  $5/8$

**Answer: B**



**Watch Video Solution**

11. The angle between the curves

$y = \sin x$  and  $y = \cos x$ ,  $0 < x < \frac{x}{2}$ , is

A.  $\tan^{-1}(2\sqrt{2})$

B.  $\tan^{-1}(3\sqrt{2})$

C.  $\tan^{-1}(3\sqrt{3})$

D.  $\tan^{-1}(5\sqrt{2})$

**Answer: A**



**Watch Video Solution**

12. The line, which is parallel to  $X$ -axis and crosses the curve  $y = \sqrt{x}$  at an angle  $45^\circ$ , is

A.  $y = \frac{1}{4}$

B.  $y = \frac{1}{2}$

C.  $y = 1$

D.  $y = 4$

**Answer: B**



**Watch Video Solution**



13. If the normal at the point  $P(at_1, 2at_1)$  meets the parabola  $y^2 = 4ax$  again at  $(at_2, 2at_2)$ , then

A.  $t_1 t_2 = -1$

B.  $t_2 = -t_1 - \frac{2}{t_1}$

C.  $2t_1 = t_2$

D. none of these

**Answer: B**



**Watch Video Solution**

14. If the line  $ax + by + c = 0$  is a normal to the curve  $xy = 1$ , then  $a > 0, b > 0$   
 $a > 0, b < 0$  (c)  $a < 0, b > 0$  (d)  $a < 0, b < 0$  none of these

A.  $(a > 0, b > 0)$  or  $(a < 0, b < 0)$

B.  $(a > 0, b < 0)$  or  $(a < 0, b > 0)$

C.  $(b \leq 0, a \leq 0)$  or  $(a \geq 0, b \leq 0)$

D.  $(a \leq 0, b \leq 0)$  or  $(a \geq 0, b \geq 0)$

**Answer: B**



**Watch Video Solution**

15. The line  $\frac{x}{a} + \frac{y}{b} = 1$  touches the curve  $y = be^{-x/a}$  at the point

A.  $(a, b/a)$

B.  $(-a, b/a)$

C.  $(a, a/b)$

D. none of these

**Answer: D**



**Watch Video Solution**

16. The normal to the curve

$$x = a(\cos \theta + \theta \sin \theta), y = a(\sin \theta - \theta \cos \theta)$$

at any  $\theta$  is such that

A. it makes a constant angle with x-axis

B. it passes through the origin

C. it is at a constant distance from the  
origin

D. none of these

**Answer: C**



Watch Video Solution

17. The point P of the curve  $y^2 = 2x^3$  such that the tangent at P is perpendicular to the line  $4x - 3y + 2 = 0$  is given by

A.  $(2, 4)$

B.  $(1, \sqrt{2})$

C.  $(1/2, -1/2)$

D.  $(1/8, -1/16)$

**Answer: D**



Watch Video Solution

**18.** Find the equation of tangents to the curve  $y = \cos(x + y)$ ,  $-2\pi \leq x \leq 2\pi$  that are parallel to the line  $x + 2y = 0$ .

A.  $x + 2y = 1$

B.  $x + 2y = \frac{\pi}{2}$

C.  $x + 2y = \frac{\pi}{4}$

D. none of these

**Answer: B**



Watch Video Solution

19. The equation of the tangents at the origin to the curve  $y^2 = x^2(1 + x)$  are

A.  $y = \pm x$

B.  $x = \pm y$

C.  $y = \pm 2x$

D. none of these

**Answer: A**



20. The coordinates of the points on the curve

$$x = a(\theta + \sin \theta), y = a(1 - \cos \theta), \quad \text{where}$$

tangent is inclined an angle  $\frac{\pi}{4}$  to the  $x$ -axis

are- (A)  $(a, a)$  (B)  $\left(a\left(\frac{\pi}{2} - 1\right), a\right)$  (C)

$\left(a\left(\frac{\pi}{2} + 1\right), a\right)$  (D)  $\left(a, a\left(\frac{\pi}{2} + 1\right)\right)$

A.  $(a, a)$

B.  $(a(\pi/2 - 1), a)$

C.  $(a(\pi/2 + 1), a)$

D.  $(a, a(\pi/2 + 1))$



**Answer: C**



**Watch Video Solution**

21. The chord joining the points where  $x = p$  and  $x = q$  on the curve  $y = ax^2 + bx + c$  is parallel to the tangent at the point on the curve whose abscissa is

A.  $\frac{1}{2}(p + q)$

B.  $\frac{1}{2}(p - q)$

C.  $\frac{pq}{2}$

D. none of these

**Answer: A**



**Watch Video Solution**

22. All points on the curve

$$y^2 = 4a \left( x + a \sin \frac{x}{a} \right) \quad \text{at which the}$$

tangents are parallel to the axis of x lie on a

A. circle

B. parabola

C. line

D. none of these

**Answer: B**



**Watch Video Solution**

**23.** The fixed point  $P$  on the curve  $y = x^2 - 4x + 5$  such that the tangent at  $P$  is perpendicular to the line  $x + 2y - 7 = 0$  is given by

A. (3, 2)

B. (1, 2)

C. (2, 1)

D. none of these

**Answer: A**



**Watch Video Solution**

**24. 26.** The points of contact of the tangents drawn from the origin to the curve  $y=\sin x$ , lie on the curve

A.  $x^2 - y^2 = xy$

B.  $x^2 + y^2 = x^2y^2$

C.  $x^2 - y^2 = x^2y^2$

D. none of these

**Answer: C**



**Watch Video Solution**

**25.** If the area of the triangle included between the axes and any tangent to the curve  $x^n y = a^n$  is constant, then find the value of  $n$ .

A. -1

B. -2

C. 1

D. 2

**Answer: C**



**Watch Video Solution**

**26.** The tangents to the curve  $x = a(\theta - \sin \theta)$ ,  $y = a(1 + \cos \theta)$  at the points  $\theta = (2k + 1)\pi$ ,  $k \in Z$  are parallel to :

A.  $y = x$

B.  $y = -x$

C.  $y = 0$

D.  $x = 0$

**Answer: C**



**Watch Video Solution**

**27.** The slope of the tangent to the curve

$y = \sin^{-1}(\sin x)$  at  $x = \frac{3\pi}{4}$  is

A. 1

B. -1

C. 0

D. non-existent

**Answer: B**



**Watch Video Solution**

**28.** The slope of the tangent to the curve

$$y = \cos^{-1}(\cos x) \quad \text{at} \quad x = -\frac{\pi}{4}, \text{ is}$$



A. 1

B. 0

C. 2

D. -1

**Answer: D**



**Watch Video Solution**

**29.** The equation of the tangent to the curve  $y = e^{-|x|}$  at the point where the curve cuts the line  $x = 1$ , is

A.  $x + y = e$

B.  $e(x + y) = 1$

C.  $y + ex = 1$

D. none of these

**Answer: D**



**Watch Video Solution**

**30.** The number of points on the curve  $y = x^3 - 2x^2 + x - 2$  where tangents are parallel to x-axis, is

A. 0

B. 1

C. 2

D. 3

**Answer: C**



**Watch Video Solution**

**31.** The angle between the tangents to the curve  $y = x^2 - 5x + 6$  at the point  $(2, 0)$  and  $(3, 0)$ , is

A.  $\pi / 3$

B.  $\pi / 4$

C.  $\pi / 2$

D.  $\pi / 6$

**Answer: C**



**Watch Video Solution**

**32.** The slope of the tangent to the curve

$y = \sqrt{9 - x^2}$  at the point where ordinate and

abscissa are equal, is

A. 1

B. -1

C. 0

D. none of these

**Answer: B**



**Watch Video Solution**

**33.** The slope of the tangent to the curve  $y = x^2 - x$  at the point where the line  $y = 2$  cuts the curve in the first quadrant, is

A. 2

B. 3

C. -3

D. none of these

**Answer: B**



**Watch Video Solution**

**34.** The abscissa of the point on the curve  $ay^2 = x^3$ , the normal at which cuts off equal intercepts from the coordinate axes is

A.  $\frac{2a}{9}$

B.  $\frac{4a}{9}$

C.  $-\frac{4a}{9}$

D.  $-\frac{2a}{9}$

**Answer: B**



**Watch Video Solution**

**35.** The curve given by  $x + y = e^{xy}$  has a tangent parallel to the y-axis at the point

A. (0, 1)

B. (1, 0)

C. (1, 1)

D. none of these

**Answer: B**



**Watch Video Solution**

**36.** The two tangents to the curve

$ax^2 + 2hxy + by^2 = 1, a > 0$  at the points

where it crosses x-axis, are



A. parallel

B. perpendicular

C. inclined at an angle of  $\pi / 4$

D. none of these

**Answer: A**



**Watch Video Solution**

**37.** Let  $P(2, 2)$  and  $Q(1/2, -1)$  be two points on the parabola  $y^2 = 2x$ , The coordinates of the point R on the parabola

$y^2 = 2x$  where the tangent to the curve is parallel to the chord PQ, are

A.  $(2, -1)$

B.  $(1/8, 1/2)$

C.  $(\sqrt{2}, 1)$

D.  $(-\sqrt{2}, 1)$

**Answer: B**



**Watch Video Solution**

38. Any tangent to the curve

$$y = 2x^5 + 4x^3 + 7x + 9$$

- A. is parallel to x-axis
- B. is parallel to y-axis
- C. makes an acute angle with the x-axis
- D. makes an obtuse angle with x-axis

**Answer: C**



**Watch Video Solution**

39. The normal to the curve  $5x^5 - 10x^3 + x + 2y + 6 = 0$  at  $P(0, -3)$  meets the curve again at the point

A.  $(-1, 1), (1, 5)$

B.  $(1, -1), (-1, -5)$

C.  $(-1, -5), (-1, 1)$

D.  $(-1, 5), (1, -1)$

**Answer: B**



**Watch Video Solution**

40. The lines parallel to the normal to the curve  $xy = 1$  is/are  $3x + 4y + 5 = 0$  (b)

$$3x - 4y + 5 = 0 \quad 4x + 3y + 5 = 0 \quad (d)$$

$$3y - 4x + 5 = 0$$

A.  $3x + 4y + 5 = 0$

B.  $3x - 4y + 5 = 0$

C.  $4x + 3y + 5 = 0$

D.  $3y - 4x - 5 = 0$

**Answer: B::D**



**Watch Video Solution**

41. Let P be the point (other than the origin) of intersection of the curves  $y^2 = 4ax$  and  $ay^2 = 4x^3$  such that the normals to the two curves meet x-axis at  $G_1$  and  $G_2$  respectively. Then,  $G_1G_2 =$

A.  $2a$

B.  $4a$

C.  $a$

D. none of these

**Answer: B**



**Watch Video Solution**

**42.** If the sum of the squares of the intercepts on the axes cut off by tangent to the curve

$x^{\frac{1}{3}} + y^{\frac{1}{3}} = a^{\frac{1}{3}}, a > 0$  at  $\left(\frac{a}{8}, \frac{a}{8}\right)$  is 2, then

$a =$  1 (b) 2 (c) 4 (d) 8

A. 1

B. 2

C. 4

D. 8

**Answer: C**



**Watch Video Solution**

## Chapter Test

1. The abscissa of the point on the curve  $ay^2 = x^3$ , the normal at which cuts off equal intercepts from the coordinate axes is



A.  $2a/9$

B.  $4a/9$

C.  $-4a/9$

D.  $-2a/9$

**Answer: B**



**Watch Video Solution**

2. If the curves

$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and  $\frac{x^2}{l^2} - \frac{y^2}{m^2} = 1$  cut each

other orthogonally then.....

A.  $a^2 + b^2 = l^2 + m^2$

B.  $a^2 - b^2 = l^2 - m^2$

C.  $a^2 - b^2 = l^2 + m^2$

D.  $a^2 + b^2 = l^2 - m^2$

**Answer: C**



**Watch Video Solution**

3. The length of normal at any point to the curve,  $y = c \cosh\left(\frac{x}{c}\right)$  is

A. (abscissa)<sup>2</sup>

B. (ordinate)<sup>2</sup>

C. abscissa

D. ordinate

**Answer: B**



**Watch Video Solution**

4. If the sub-normal at any point on  $y^{1-n}x^n$  is of constant length, then find the value of  $n$ .

A. 1

B.  $1/2$

C. 2

D. -2

**Answer: B**



**Watch Video Solution**

5. The angle of intersection of the curves

$y = x^2$ ,  $6y = 7 - x^3$  at  $(1, 1)$ , is

A.  $\pi / 4$

B.  $\pi / 3$

C.  $\pi / 2$

D. none of these

**Answer: C**



**Watch Video Solution**

6. The slope of the tangent to the curve  $x = t^2 + 3t - 8, y = 2t^2 - 2t - 5$  at the point (2, -1), is

A.  $22/7$

B.  $6/7$

C. -6

D. none of these

**Answer: B**



**Watch Video Solution**

7. The two curves  $x^3 - 3xy^2 + 2 = 0$  and  $3x^2y - y^3 - 2 = 0$

A.  $45^\circ$

B.  $60^\circ$

C.  $90^\circ$

D.  $30^\circ$

**Answer: C**



**Watch Video Solution**

8. The tangent and normal at  $P(t)$ , for all real positive  $t$ , to the parabola  $y^2 = 4ax$  meet the axis of the parabola in  $T$  and  $G$  respectively, then the angle at which the tangent at  $P$  to the parabola is inclined to the tangent at  $P$  to the circle passing through the points  $P$ ,  $T$  and  $G$  is

A.  $\tan^{-1} t^2$

B.  $\cot^{-1} t^2$

C.  $\tan^{-1} t$



$$D. \cot^{-1} t$$

**Answer: C**



**Watch Video Solution**

9. If  $y = 4x - 5$  is a tangent to the curve

$y^2 = px^3 + q$  at  $(2, 3)$ , then

A.  $p = 2, q = -7$

B.  $p = -2, q = 7$

C.  $p = -2, q = -7$

D.  $p = 2, q = 7$

**Answer: A**



**Watch Video Solution**

**10.** The curve  $y - e^{xy} + x = 0$  has a vertical tangent at the point :

A. (1, 1)

B. at no point

C. (0, 1)

D. (1, 0)

**Answer: D**



**Watch Video Solution**

**11.** If the parametric of a curve given by  $x = e^t \cos t$ ,  $y = et \sin t$ , then the tangent to the curve at the point  $t = \pi/4$  makes with axis of x the angle

A. 0

B.  $\pi / 4$

C.  $\pi / 3$

D.  $\pi / 2$

**Answer: D**



**Watch Video Solution**

**12.** The length of the normal at  $t$  on the curve

$$x = a(t + \sin t), y = a(1 - \cos t), \text{ is}$$

A.  $a \sin t$

B.  $2a \sin^3 \frac{t}{2} \sec \frac{t}{2}$

C.  $2a \sin \frac{t}{2} \tan \frac{t}{2}$

D.  $2a \sin \frac{t}{2}$

**Answer: C**



**Watch Video Solution**

**13.** For the parabola  $y^2 = 4ax$ , the ratio of the subtangent to the abscissa, is

A. 1:1

B. 2 : 1

C.  $x : y$

D.  $x^2 : y$

**Answer: B**



**Watch Video Solution**

**14.** The length of the subtangent to the curve

$$\sqrt{x} + \sqrt{y} = 3 \text{ at the point } (4, 1), \text{ is}$$

A. 2

B.  $1/2$

C. 3

D. 4

**Answer: A**



**Watch Video Solution**

**15.** The normal to the curve

$$x = a(\cos \theta + \theta \sin \theta), y = a(\sin \theta - \theta \cos \theta)$$

at any  $\theta$  is such that

A. it makes a constant angle with x-axis

B. it passes through the origin

C. it is at a constant distance from the  
origin

D. none of these

**Answer: C**



**Watch Video Solution**



16. Tangents are drawn from the origin to the curve  $y = \cos X$ . Their points of contact lie on

A.  $x^2y^2 = y^2 - x^2$

B.  $x^2y^2 = x^2 + y^2$

C.  $x^2y^2 = x^2 - y^2$

D. none of these

**Answer: C**



**Watch Video Solution**

17. If  $m$  denotes the slope of the normal to the curve  $y = -3\log(9 + x^2)$  at the point  $x \neq 0$ , then,

A.  $n \in [-1, 1]$

B.  $m \in \mathbb{R} - (-1, 1)$

C.  $n \in \mathbb{R} - [-1, 1]$

D.  $m \in (-1, 1)$

**Answer: B**



**Watch Video Solution**

18. If  $m$  be the slope of the tangent to the curve  $e^{2y} = 1 + 4x^2$ , then

A.  $m < 1$

B.  $|m| \leq 1$

C.  $|m| \geq 1$

D. none of these

**Answer: B**



**Watch Video Solution**

19. If the curve  $y = ax^3 + bx^2 + cx$  is inclined at  $45^\circ$  to x-axis at  $(0, 0)$  but touches x-axis at  $(1, 0)$ , then

A.  $a = 1, b = -2, c = 1$

B.  $a = 1, b = 1, c = -2$

C.  $a = -2, b = 1, c = 1$

D.  $a = -1, b = 2, c = 1$

**Answer: A**



**Watch Video Solution**

20. If the curve  $y = ax^2 + bx + c$  passes through the point (1, 2) and the line  $y = x$  touches it at the origin, then

A.  $a = 1, b = -1, c = 0$

B.  $a = 1, b = 1, c = 0$

C.  $a = -1, b = 1, c = 0$

D. none of these

**Answer: B**



**Watch Video Solution**

21. The angle between the tangents to the curve  $y^2 = 2ax$  at the point where  $x = \frac{a}{2}$ , is

A.  $\pi / 6$

B.  $\pi / 4$

C.  $\pi / 3$

D.  $\pi / 2$

**Answer: D**



**Watch Video Solution**

22. The intercepts made by the tangent to the curve  $y = \int_0^x |t| dt$ , which is parallel to the line  $y = 2x$  on y-axis are equal to

A. 1, -1

B. -2, 2

C. 3

D. -3

**Answer: B**



**Watch Video Solution**

23. The line  $\frac{x}{a} + \frac{y}{b} = 2$  touches the curve  $\left(\frac{x}{a}\right)^n + \left(\frac{y}{b}\right)^n = 2$  at the point  $(a, b)$  for

A.  $n = 2$  only

B.  $n = -3$  only

C. any  $n \in R$

D. none of these

**Answer: C**



**Watch Video Solution**



24. The equation of the normal to the curve  $y = e^{-2|x|}$  at the point where the curve cuts the line  $x = 1/2$  is

A.  $2e(ex + 2y) = e^2 - 4$

B.  $2e(ex - 2y) = e^2 - 4$

C.  $2e(ey - 2x) = e^2 - 4$

D. none of these

**Answer: B**



**Watch Video Solution**

25. The length of subtangent to the curve

$$x^2 + xy + y^2 = 7 \text{ at the point } (1, -3) \text{ is}$$

A. 3

B. 5

C. 15

D.  $3/5$

**Answer: C**



**Watch Video Solution**