



PHYSICS

BOOKS - SUNIL BATRA 41 YEARS IITJEE PHYSICS (HINGLISH)

HEAT AND THERMODYNAMICS

Jee Main And Advanced


1. One mole of a mono-atomic ideal gas is mixed with one mole of a diatomic ideal gas. The molar specific heat of the mixture at constant volume is



[Watch Video Solution](#)

2. The variation of temperature of a material as heat is given to it at a constant rate is shown in the figure. The material is in solid state at the

point O. The state of the material at the point P is.....



[Watch Video Solution](#)

3. During an experiment, an ideal gas is found to obey an additional law $VP^2 = \text{constant}$, The gas is initially at a temperature T , and volume V . When it expands to a volume $2V$, the temperature becomes.....

[Watch Video Solution](#)

4. 300 grams of water at 25°C is added to 100 grams of ice at 0°C . The final temperature of the mixture is^{@C}



[Watch Video Solution](#)

5. The earth receives at its surface radiation from the sun at the rate of $1400Wm^{-2}$. The distance of the centre of the sun from the surface of the earth is $1.5 \times 10^{11}m$ and the radius of the sun is 7×10^8m . Treating the sun as a black body, it follows from the above data that its surface temperature isK



[Watch Video Solution](#)

6. A solid copper sphere (density ρ and specific heat c) of radius r at an initial temperature $200K$ is suspended inside a chamber whose walls are at almost $0K$. The time required for the temperature of the sphere to drop to $100K$ is



[Watch Video Solution](#)

7. A point source of heat of power P is placed at the centre of a spherical shell of mean radius R . The material of the shell has thermal conductivity K . If the temperature difference between the outer and inner surface of the shell is not to exceed T , the thickness of the shell should not be less than



[Watch Video Solution](#)

8. A substance of mass M kg requires a power input of P watts to remain in the molten state at its melting point. When the power source is turned off, the sample completely solidifies in time t seconds. The latent heat of fusion of the substance is



[Watch Video Solution](#)

9. A container of volume $1m^3$ is divided into two equal parts by a partition. One part has an ideal gas at $300K$ and the other part is vacuum. The whole system is thermally isolated from the surroundings. When the

partition is removed, the gas expands to occupy the whole volume. Its temperature will now be


 [Watch Video Solution](#)

10. An ideal gas with pressure P , volume V and temperature T is expanded isothermally to a volume $2V$ and a final pressure P_i . If the same gas is expanded adiabatically to a volume $2V$, the final pressure P_a . The ratio of the specific heats of the gas is 1.67. The ratio $\frac{P_a}{P_i}$ is

 [Watch Video Solution](#)

11. Two metal cubes A and B of same size are arranged as shown in Figure. The extreme ends of the combination are maintained at the indicated temperatures. The arrangement is thermally insulated. The coefficients of thermal conductivity of A and B are $300W/m^\circ C$ and $200W/m^\circ C$, respectively. After steady state is reached the temperature t of the

interface will be




[▶ Watch Video Solution](#)

12. A ring shaped tube contain two ideal gases with equal masses and relative molar masses $M_1 = 32$ and $M_2 = 28$.

The gases are separated by one fixed partion and another movable stopper S which can move freely without friction inside the ring. The

angle α as shown in the figure is degrees.



[▶ Watch Video Solution](#)


13. Earth receives 1400 W/m^2 of solar power. If all the solar energy falling on a lens of area 0.2 m^2 is focused on to a block of ice of mass 280 grams, the time taken to melt the ice will be..... Minutes.
(*Latent heat of fusion of ice* = $3.3 \times 10^5 \text{ J/kg.}$)

[▶ Watch Video Solution](#)

14. The root-mean square speeds of the molecules of different ideal gses, maintained at the same temperature are the same.

[▶ Watch Video Solution](#)

15. The volume V versus temperature T graphs for a cetain amount of a perfect gas at two pressure p_1 and p_2 are as shown in Fig. It follows from the graphs that p_1 is greater than p_2 .



[▶ Watch Video Solution](#)

16. Two different gases at the same temperature have equal root mean square velocities.


 [Watch Video Solution](#)

17. The ratio of the velocity of sound in Hydrogen gas $\left(\gamma = \frac{7}{5}\right)$ to that in Helium gas $\left(\gamma = \frac{5}{3}\right)$ at the same temperature is $\sqrt{21/3}$.

 [Watch Video Solution](#)

18. The curves A and B in the figure shown P-V graphs for an isothermal and an adiabatic process for an idea gas. The isothermal process is

represented by the curve A.



[▶ Watch Video Solution](#)

19. At a given temperature, the specific heat of a gas at constant pressure is always greater than its specific heat at constant volume.

[▶ Watch Video Solution](#)

20. The root mean square (rms) speed of oxygen molecules (O_2) at a certain temperature T (degree absolute) is V . If the temperature is

doubled and oxygen gas dissociates into atomic oxygen, the rms speed remains unchanged.

 [Watch Video Solution](#)

21. Two spheres of the same material have radii 1m and 4m and temperatures 4000K and 2000K respectively. The energy radiated per second by the first sphere is greater than that by the second.

 [Watch Video Solution](#)

22. A constant volume gas thermometer works on

A. The Principle of Archimedes

B. Boyel's Law

C. Pascal's Law

D. Charle's Law

Answer: D



Watch Video Solution

23. A metal ball immersed in alcohol weighs W_1 at $0^\circ C$ and W_2 at $50^\circ C$. The coefficient of expansion of cubical the metal is less than that of the alcohol. Assuming that the density of the metal is large compared to that of alcohol, it can be shown that

A. $W_1 > W_2$

B. $W_1 = W_2$

C. $W_1 < W_2$

D. None of these

Answer: C



Watch Video Solution

24. A wall has two layers A and B, each made of different material. Both the layers have the same thickness. The thermal conductivity of the material of A is twice that of B. Under thermal equilibrium, the temperature difference across the wall is 36°C . The temperature difference across the layer A is

A. 6°C

B. 12°C

C. 18°C


D. 24°C

Answer: B



[Watch Video Solution](#)

25. An ideal monatomic gas is taken round the cycle ABCDA as shown in the P-V diagram. The work done during the cycle is



- A. PV
- B. $2PV$
- C. $\frac{1}{2}PV$
- D. zero

Answer: A

[▶ Watch Video Solution](#)

26. If one mole of a monatomic gas ($\gamma = \frac{5}{3}$) is mixed with one mole of a diatomic gas ($\gamma = \frac{7}{5}$), the value of gamma for mixture is

A. 1.40

B. 1.50

C. 1.53

D. 3.07

Answer: B



Watch Video Solution

27. From the following statements concerning ideal gas at any given temperature T , select the correct one (s)

A. The coefficient of volume expansion at constant pressure is the same for all idea gases

B. The average translational kinetic energy per molecule of oxygen gas is $3kT$, k being Boltzmann constant

C. The mean-free path of molecules increases with increases in the pressure

D. In a gaseous mixture, the average translational kinetic energy of the molecules of each component is different

Answer: A

 [Watch Video Solution](#)

28. Three rods of identical area of cross-section and made from the same metal from the sides of an isosceles triangle. ABC, right angled at B. The points A and B are maintained at temperatures T and $\sqrt{2}T$ RESPECTIVELY.

In the steady state the temperature of the point C is T_C .

Assuming that only heat conduction takes place, $\frac{T_C}{T}$ is equal to

A. $\frac{1}{2\sqrt{2} - 1}$

B. $\frac{3}{\sqrt{2} + 1}$

C. $\frac{1}{\sqrt{3}(2\sqrt{2} - 1)}$

D. $\frac{1}{\sqrt{2} + 1}$

Answer: B



Watch Video Solution

29. Two metallic spheres S_1 and S_2 are made of the same material and have got identical surface finish. The mass of S_1 is thrice that of S_2 . Both the spheres are heated to the same high temperature and placed in the same room having lower temperature but are thermally insulated from each other. The ratio of the initial rate of cooling of S_1 to that of S_2 is

A. $\frac{1}{3}$

B. $\frac{1}{\sqrt{3}}$

C. $\frac{\sqrt{3}}{1}$

D. $\left(\frac{1}{3}\right)^{\frac{1}{3}}$

Answer: D



Watch Video Solution

30. The average translational kinetic energy of O_2 (relative molar mass 32) molecules at a particular temperature is 0.048eV. The translational kinetic energy of N_2 (relative molar mass 28) molecules in eV at the same temperature is

- A. 0.0015
- B. 0.003
- C. 0.048
- D. 0.768

Answer: C

[Watch Video Solution](#)

31. A vessel contains 1 mole of O_2 gas (relative molar mass 32) at a temperature T. The pressure of the gas is P. An identical vessel containing

one mole of He gas (relative molar mass 4) at temperature $2T$ has a pressure of

A. $\frac{P}{8}$

B. P

C. $2P$

D. $8P$

Answer: C



[Watch Video Solution](#)

32. A spherical black body with a radius of 12cm radiates 450W power at 500K. If the radius were halved and the temperature doubled, the power radiated in watt would be

A. 225

B. 450

C. 900

D. 1800

Answer: D



[Watch Video Solution](#)

33. A closed compartment containing gas is moving with some acceleration in horizontal direction. Neglect effect of gravity. Then the pressure in the compartment is

- A. same everywhere
- B. lower in the front side
- C. lower in the rear side
- D. lower in the upper side

Answer: B



[Watch Video Solution](#)

34. A gas mixture consists of 2 moles of oxygen and 4 moles of argon at temperature T . Neglecting all vibrational modes, the total internal energy of the system is

A. $4RT$

B. $15RT$

C. $9RT$

D. $11RT$

Answer: D



[Watch Video Solution](#)

35. The ratio of the speed of sound in nitrogen gas to that in helium gas, at 300K is

A. $\left(\sqrt{2/7}\right)$

B. $\left(\sqrt{1/7}\right)$

C. $(\sqrt{3})/5$

D. $(\sqrt{6})/5$

Answer: C



Watch Video Solution

36. A monatomic ideal gas, initially at temperature T_1 , is enclosed in a cylinder fitted with a frictionless piston. The gas is allowed to expand adiabatically to a temperature T_2 by releasing the piston suddenly. If L_1 and L_2 are the length of the gas column before expansion respectively, then $\frac{T_1}{T_2}$ is given by

A. $\left(\frac{L_1}{L_2}\right)^{2/3}$

B. $\left(\frac{L_1}{L_2}\right)$




C. $\left(\frac{L_2}{L_1}\right)$

D. $\left(\frac{L_2}{L_1}\right)^{2/3}$

Answer: D

 Watch Video Solution

37. A block of ice at -10°C is slowly heated and converted to steam at 100°C . Which of the following curves represents the phenomenon qualitatively?







D. \

Answer: A




38. An ideal gas is initially at temperature T and volume V . Its volume is increased by ΔV due to an increase in temperature ΔT , pressure remaining constant. The quantity $\delta = \frac{\Delta V}{V\Delta T}$ varies with temperature as



Answer: C



39. Starting with the same initial conditions, an ideal gas expands from volume $V_1 \rightarrow V_2$ in three different ways. The work done by the gas is W_1 if the process is purely isothermal, W_2 if purely isobaric and W_3 if purely adiabatic. Then




- A. $W_2 > W_1 > W_3$
- B. $W_1 > W_2 > W_3$
- C. $W_2 > W_3 > W_1$
- D. $W_2 > W_1 > W_3$

Answer: A

 Watch Video Solution

40. The plots of intensity versus wavelength for three black bodies at temperature T_1, T_2 and T_3 respectively are as shown. Their temperatures are such that



A. $T_1 > T_2 > T_3$

B. $T_1 > T_3 > T_2$

C. $T_2 > T_3 > T_1$


$$D. T_3 > T_2 > T_1$$

Answer: B



Watch Video Solution

41. Three rods made of same material and having the same cross-section have been joined as shown. In the figure. Each rod is of the same length. The left and right ends are kept at $0^\circ C$ and $90^\circ C$ respectively. The temperature of the junction of the three rods will be



A. $45^\circ C$

B. $60^\circ C$

C. $30^\circ C$

D. $20^\circ C$

Answer: B



[Watch Video Solution](#)

42. In a given process on an ideal gas, $dW = 0$ and $dQ < 0$. Then for the gas

A. the temperature will decrease

B. the volume will increase

C. the pressure will remain constant


D. the temperature will increase

Answer: A



[Watch Video Solution](#)

43. P-V plots for two gases during adiabatic processes are shown in the figure. Plots 1 and 2 should correspond respectively to



A. He and O_2

B. O_2 and He

C. He and Ar

D. O_2 and N_2

Answer: B



[Watch Video Solution](#)

44. When a block of iron in mercury at $0^\circ C$, fraction K_1 of its volume is submerged, while at the temperature $60^\circ C$, a fraction K_2 is seen to be submerged. If the coefficient of volume expansion of iron is γ_{Fe} and that of mercury is γ_{Hg} , then the ratio $(K_1) / (K_2)$ can be expressed as

- A. $\frac{1 + (60\gamma_{Fe})}{1 + (60\gamma_{hg})}$
 B. $\frac{1 - (60\gamma_{Fe})}{1 + (60\gamma_{Hg})}$
 C. $\frac{1 + (60\gamma_{Fe})}{1 - (60\gamma_{Hg})}$
 D. $\frac{1 + (60\gamma_{Hg})}{1 + (60\gamma_{Fe})}$


Answer: A



Watch Video Solution

45. An ideal gas is taken through the cycle $A \rightarrow B \rightarrow C \rightarrow A$, as shown in the figure, If the net heat supplied to the gas in the cycle is 5J, the work

done by the gas in the process CtoA is



- A. $-5J$
- B. $-10J$
- C. $-15J$
- D. $-20J$

Answer: A




[Watch Video Solution](#)


46. Which of the following graphs correctly represents the variation of

$$\beta = - \frac{dV / dP}{V}$$


with P for an ideal gas at constant temperature?




A.



B.



C.



D.

Answer: A



Watch Video Solution

47. An ideal Black-body at room temperature is thrown into a furnace. It is observed that

- A. initially it is the darkest body and at later times the brightest
- B. it is the darkest body at all times
- C. it cannot be distinguished at all times
- D. initially it is the darkest body and at later times it cannot be distinguished


Answer: A



[Watch Video Solution](#)

48. The graph, shown in the adjacent diagram, represents the variation of temperature (T) of two bodies, x and y having same surface area, with time (t) due to the emission of radiation. Find the correct relation

between the emissivity and absorptivity power of the two bodies



- A. $E_x > E_y$ & $a_x < a_y$
- B. $E_x < E_y$ & $a_x > a_y$
- C. $E_x > E_y$ & $a_x > a_y$
- D. $E_x < E_y$ & $a_x < a_y$

Answer: C



[Watch Video Solution](#)

49. Two rods, one of aluminium and the other made of steel, having initial length l_1 and l_2 are connected together to form a single rod of length $l_1 + l_2$. The coefficients of linear expansion for aluminium and steel are α_a and α_s and respectively. If the length of each rod increases by the same amount when their temperature are raised by $t^{\circ}C$, then find the ratio $l_1 / (l_1 + l_2)$

A. $(\alpha_s) / (\alpha_a)$

B. $(\alpha_a) / (\alpha_s)$

C. $(\alpha_s) / ((\alpha_a) / (\alpha_s))$



D. $(\alpha_a) / ((\alpha_a) / (\alpha_s))$

Answer: C




Watch Video Solution

50. The PT diagram for an ideal gas is shown in the figure, where AC is an adiabatic process, find the corresponding PV diagram.




A.



B.

C. 



D.

Answer: B

 **Watch Video Solution**

51. 2kg of ice at 20°C is mixed with 5kg of water at 20°C in an insulating vessel having a negligible heat capacity. Calculate the final mass of water remaining in the container. It is given that the specific heats of water & ice are $1\text{kcal}/\text{kg}/^{\circ}\text{C}$ & 0.5

$\text{kcal}/\text{kg}/^{\circ}\text{C}$ while the latent heat of fusion of ice is $80\text{kcal}/\text{kg}$

A. 7kg

B. 5kg

C. 4kg

D. 2kg

Answer: B



[Watch Video Solution](#)

52. Three discs A, B and C having radii 2, 4 and 6 cm respectively are coated with carbon black. Wavelength for maximum intensity for the three discs

are 300,400 and 500 nm respectively. If Q_A , Q_B and Q_C are power emitted by A,B and D respectively, then

A. Q_A will be maximum

B. Q_B will be maximum

C. Q_C will be maximum


D. $Q_A = Q_B = Q_C$

Answer: B




Watch Video Solution


53. If liquefied oxygen at 1 atmospheric pressure is heated from 50K to 300k by supplying heat at constant rate. The graph of temperature vs time will be




A.



B.



C.



D.

Answer: C



Watch Video Solution

54. Two identical rods are connected between two containers one of them is at $100^{\circ}C$ and another is at $0^{\circ}C$. If rods are connected in parallel then the rate of melting of ice is $q_1 gm/sec$.

If they are connected in series then the rate is q_2 . The ratio q_2/q_1 is

A. 2

B. 4

C. $1/2$

D. $1/4$

Answer: D



Watch Video Solution

55. An ideal gas is initially at P_1, V_1 is expands to P_2, V_2 and then compressed adiabatically to the same volume V_1 and pressure P_3 . If W is the net work done by the gas in complete process which of the following is true.

A. $W > 0, P_3 > P_1$

B. $W < 0, P_3 > P_1$


C. $W > 0, P_3 < P_1$

D. $W < 0, P_3 < P_1$

Answer: B

 Watch Video Solution

56. Variation of radiant energy emitted by sun, filament of tungsten lamp and welding arc as a function of its wavelength is shown in figure. Which of the following option is the correct match?



- A. Sun – T_3 , tungsten filament – T_1 , weld in arc – T_2
- B. Sun – T_2 , tungsten filament – T_1 , weld in arc – T_3
- C. Sun – T_3 , tungsten filament – T_2 , weld in arc – T_1

D. Sun – T_1 , tungsten filament – T_2 , weld \in garc – T_3

Answer: A



[Watch Video Solution](#)

57. In which of the following process, convection does not take place primarily

- A. Sea and land breeze
- B. boiling of water
- C. heating air around a furnace
- D. warming of glass of bulb due to filament

Answer: D



[Watch Video Solution](#)

58. A spherical body of area A , and emissivity $e = 0.6$ is kept inside a black body. What is the rate at which energy is radiated per second at temperature T

A. $0.4AT^4$

B. $0.8AT^4$

C. $6.0AT^4$

D. $1.0AT^4$

Answer: None of these



[Watch Video Solution](#)

59. Calorie is defined as the amount of heat required to raise temperature of 1 g of water by $1^\circ C$ and it is defined under which of the following conditions?

A. From $14.5^\circ C \rightarrow 15.5^\circ C$ at 760mm of Hg.

B. From $98.5^{\circ}C \rightarrow 99.5^{\circ}C$ at 760mm of Hg.

C. From $13.5^{\circ}C \rightarrow 14.5^{\circ}C$ at 76mm of Hg.

D. From $3.5^{\circ}C \rightarrow 4.5^{\circ}C$ at 76mm of Hg.

Answer: A



Watch Video Solution

60. Water of volume 2 litre in a container is heated with a coil of $1kW$ at $27^{\circ}C$. The lid of the container is open and energy dissipates at rate of $160J/s$. In how much time temperature will rise from $27^{\circ}C \rightarrow 77^{\circ}C$ [Given specific heat of water is $4.2kJ/kg$]

A. 7 min

B. 6 min 2s

C. 8 min 20s`

D. 14 min


Answer: C



Watch Video Solution

61. Water is filled up to a height h in a beaker of radius R as shown in the figure. The density of water is ρ , the surface tension of water is T and the atmospheric pressure is P_0 . Consider a vertical section $ABCD$ of the water column through a diameter of the beaker. The force on water on one side on this section by water on the other side of this section has

magnitude



A. $|2P_0RH + \pi R^2 \rho gh - 2RT|$

B. $|2P_0Rgh + R\rho gh^2 + 2RT|$

C. $|P_0\pi R^2 + R\rho gh^2 - 2RT|$

D. $|P_0\pi R^2 + R\rho gh^2 + 2RT|$

Answer: B



[Watch Video Solution](#)

62. An ideal gas is expanding such that $PT^\circ = \text{constant}$. The coefficient of volume expansion of the gas is-

A. $1/T$

B. $1/T$

C. $3/T$

D. $4/T$

Answer: C



[Watch Video Solution](#)

63. A real gas behaves like an ideal gas if its

- A. pressure is temperature are both high
- B. pressure is temperature are both low
- C. pressure is high temperature are both low
- D. pressure is low temperature are both high

Answer: D

 [Watch Video Solution](#)

64. 5.6 liter of helium gas at STP is adiabatically compressed to 0.7 liter.

Taking the initial temperature to be T_1 , the work done in the process is

- A. $\frac{9}{8}RT_1$
- B. $\frac{3}{2}RT_1$
- C. $\frac{15}{8}RT_1$
- D. $\frac{9}{2}RT_1$

Answer: A

 [Watch Video Solution](#)

65. A mixture of 2 moles of helium gas ($a \rightarrow \text{molecular mass} = 4a\mu$) and 1 mole of argon gas ($a \rightarrow \text{molecular mass} = 40a\mu$) is kept at 300K in a container. The ratio of the rms speeds $\left(\frac{v_{rms}(\text{helium})}{v_{rms}(\text{argon})} \right)$ is

A. 0.32

B. 0.45

C. 2.24

D. 3.16

Answer: D

 [Watch Video Solution](#)

66. Two moles of ideal helium gas are in a rubber balloon at $30^\circ C$. The balloon is fully expandable and can be assumed to require no energy in its expansion. The temperature of the gas in the balloon is slowly

changed to $35^{\circ}C$. The amount of heat required in raising the temperature is nearly (take R

$$= 8.31J/mol.K)$$

A. $62J$

B. $104J$

C. $124J$

D. $208J$


Answer: D



[Watch Video Solution](#)

67. Two rectangular blocks, having identical dimensions, can be arranged either in configuration-I or in configuration-II as shown in the figure. One of the blocks has thermal conductivity k and the other $2k$. The temperature difference between the ends along the x-axis is the same in both the configurations. It takes 9s to transport a certain amount of heat from the hot end to the cold end in the configuration-I. The time to

transport the same amount of heat in the configuration-II is



A. $2.0s$

B. $4.5s$

C. $3.0s$

D. $6.0s$

Answer: A



[Watch Video Solution](#)

68. Two non-reactive monoatomic ideal gases have their atomic masses in the ratio 2:3. The ratio of their partial pressures, when enclosed in a vessel kept at a constant temperature, is 4:3. The ratio of their densities is

A. 1:4

B. 1:2

C. 6:9

D. 8:9

Answer: D

 [Watch Video Solution](#)

69. Parallel rays of light of intensity $I = 912 \text{ WM}^{-2}$ are incident on a spherical black body kept in surroundings of temperature 300K. Take Stefan-Boltzmann constant $\sigma = 5.7 \times 10^{-8}$

$\text{Wm}^{-2}\text{K}^{-4}$ and assume that the energy exchange with the

surroundings is only through radiation. The final steady state temperature of the black body is close to

- A. $330K$
- B. $660K$
- C. $990K$
- D. $1550K$


Answer: A



Watch Video Solution

70. A water cooler of storage capacity 120 liters can cool water at a constant rate of P watts. In a closed circulation system (as shown schematically in the figure), the water from the cooler is used to cool an external device that generates constantly 3kW of heat (thermal load). The temperature of water fed into the device cannot exceed 30°C and the entire stored 120 liters of water is initially cooled to 10°C . The entire system is thermally insulated. The minimum value of P (in watts) for

which the device can be operated for 3 hours is



(Specific heat of water is $4.2kJkg^{-1}K^{-1}$ and the density of water is $1000kgm^{-3}$)

- A. 1600
- B. 2067
- C. 2533
- D. 3933

Answer: B

 [Watch Video Solution](#)

71. A gas is enclosed in a cylinder with a movable frictionless piston. Its initial thermodynamic state at pressure $P_i = 10^5$ Pa and volume $V_i = 10^{-3} \text{ m}^3$ changes to a final state at $P_f = (1/32) \times 10^5 \text{ Pa}$ and $V_f = 8 \times 10^{-3} \text{ m}^3$ in an adiabatic quasi-static process, such that $P^3 V^3 = \text{constant}$. Consider another thermodynamic process that brings the system from the same initial state to the same final state in two steps: an isobaric expansion at P_i followed by an isochoric (isovolumetric) process at volume V_f . The amount of heat supplied to the system in the two-step process is approximately

A. 112 J

B. 294 J

C. 588 J

D. 813 J

Answer: C



[Watch Video Solution](#)

72. The ends Q and R of two thin wires, PQ and RS, are soldered (joined) together. Initially each of the of wire has a length of 1m at $10^{\circ}C$. Now the end P is maintained at $10^{\circ}C$, while the ends S is heated and maintained at $400^{\circ}C$. The system is thermally insulated from its surroundings. If the thermal conductivity of wire PQ is twice that of the wire RS and the coefficient of linear thermal expansion of PQ is $1.2 \times 10^{-5} K^{-1}$, the change in length of the wire PQ is

A. $0.78mm$

B. $0.90mm$

C. $1.56mm$

D. $2.34mm$

Answer: A



Watch Video Solution

73. At room temperature the rms speed of the molecules of a certain diatomic gas is found to be 1930 m/sec . The gas is .

A. H_2

B. F_2

C. O_2

D. Cl_2

Answer: A



[Watch Video Solution](#)

74. 70 calories of heat required to raise the temperature of 2 moles of an ideal gas at constant pressure from $30^\circ C \rightarrow 35^\circ C$. The amount of heat required (in calories) to raise the temperature of the same gas through the same range ($30^\circ C \rightarrow 35^\circ C$) at constant volume is:

A. 30

B. 50

C. 70

D. 90

Answer: B



Watch Video Solution

75. Steam at $100^{\circ}C$ is passed into $1.1kg$ of water contained in a calorimeter of water equivalent $0.02kg$ at $15^{\circ}C$ till the temperature of the calorimeter and its content rises to $80^{\circ}C$. What is the mass of steam condensed? Latent heat of steam = $536cal/g$.

A. 0.130

B. 0.065

C. 0.260

D. 0.135

Answer: A



[Watch Video Solution](#)

76. A cylinder of radius R made of a material of thermal conductivity K_1 is surrounded by a cylindrical shell of inner radius R and outer radius $2R$ made of a material of thermal conductivity K_2 . The two ends of the combined system are maintained at two different temperatures. There is no loss of heat across the cylindrical surface and the system is in steady state. The effective thermal conductivity of the system is

A. $K_1 + K_2$

B. $K_1 K_2 / (K_1 + K_2)$

C. $(K_1 + 3K_2) / 4$

D. $(3K_1 + 3K_2) / 4$

Answer: C



[Watch Video Solution](#)

77. For an ideal gas :

- A. the change in internal energy in a constant pressure process from temperature $T_1 \rightarrow T_2$ is equal to $nC_v(T_2 - T_1)$, where C_v is the molar specific heat at constant volume and n the number of moles of the gas.
- B. the change in internal energy of the gas and the work done by the gas are equal in magnitude in an adiabatic process.
- C. the internal energy does not change in an isothermal process.
- D. no heat is added or removed in an adiabatic process.

Answer: A::B::C::D



Watch Video Solution

78. When an ideal diatomic gas is heated at constant pressure, the fraction of the heat energy supplied which increases the internal energy of the gas is

A. $\frac{2}{5}$

B. $\frac{3}{5}$

C. $\frac{3}{7}$

D. $\frac{5}{7}$

Answer: D



[Watch Video Solution](#)

79. Three closed vessels A, B and C are at the same temperature T and contain gasses which obey the Maxwellian distribution of velocities. Vessel A contain only O_2 and N_2 . If the average speed of the O_2 molecules in vessel A is v_1 that of the N_2 molecules in vessel B us v_2 , the average speed of the O_2 molecules in vessel C is

A. $\frac{v_1 + v_2}{2}$

B. v_1

C. $(v_1 v_2)^{\frac{1}{2}}$

D. $\sqrt{\frac{3kT}{M}}$

Answer: B



Watch Video Solution

80. An ideal gas is taken from the state A (pressure P , volume V) to the state B (pressure $p/2$, volume $2V$) along a straight line path in the P-V diagram. Select the correct statement (s) from the following :

- A. The work done by the gas in the process A to B exceeds the work that would be done by it if the system were taken from A to B along the isotherm.
- B. In the T-V diagram, the path AB becomes a part of a parabola
- C. In the P-T diagram, the path AB becomes a part of a hyperbola

D. In going from A to B, the temperature T of the gas first increases to a maximum value and then decreases.

Answer: A::B::D

 [Watch Video Solution](#)

81. Two bodies A and B have thermal emissivities of 0.01 and 0.81 respectively. The outer surface areas of the two bodies are the same. The two bodies emit total radiant power of the same rate. The wavelength λ_B corresponding to maximum spectral radiance in the radiation from B shifted from the wavelength corresponding to maximum spectral radiance in the radiation from A, by $1.00\mu m$. If the temperature of A is 5820K:

A. the temperature of B is 1934K

B. $\lambda_B = 1.5\mu m$

C. the temperature of B is 11604K

D. the temperature of B is 2901K

Answer: A::B



[Watch Video Solution](#)

82. The temperature of an ideal gas is increased from 120K to 480K. If at 120K the root-mean-square velocity of the gas molecules is v , at 480K it becomes

A. $4v$

B. $2v$

C. $v/2$

D. $v/4$

Answer: B



[Watch Video Solution](#)

83. A given quantity of a ideal gas is at pressure P and absolute temperature T . The isothermal bulk modulus of the gas is

A. $\frac{2}{3}P$

B. P

C. $\frac{3}{2}P$

D. $2P$

Answer: B



[Watch Video Solution](#)

84. Two cylinders A and B fitted with pistons contain equal amounts of an ideal diatomic gas at 300K. The piston of A is free to move, while that B is held fixed. The same amount of heat is given to the gas in each cylinder. If the rise in temperature of the gas in A is 30K, then the rise in temperature of the gas in B is

A. $30K$

B. $18K$

C. $50K$

D. $42K$

Answer: D



[Watch Video Solution](#)

85. During the melting of a slab of ice at $273K$ at atmospheric pressure,

A. positive work is done by the ice-water system on the atmosphere.

B. positive work is done on the ice-water system by the atmosphere.

C. the internal energy of the ice-water system increases.

D. the internal energy of the ice-water system decreases.

Answer: B::C



[Watch Video Solution](#)

86. A blackbody is at a temperature of 2880K. The energy of radiation emitted by this object with wavelength between 499nm and 500nm is U_1 , between 999nm and 1000nm is U_2 and between 1499 nm and 1500 nm is U_3 . The Wien constant $b = 2.88 \times 10^6 \text{ nmK}$. Then

A. $U_1 = 0$

B. $U_3 = 0$

C. $U_1 > U_2$

D. $U_2 > U_1$

Answer: D



Watch Video Solution

87. A bimetallic strip is formed out of two identical strips one of copper and the other of brass. The coefficients of linear expansion of the strip goes up by ΔT and the strip bends to form an arc of radius of curvature R. Then R is.

A. proportional to ΔT

B. inversely proportional to ΔT

C. proportional to $|\alpha_B - \alpha_C|$

D. inversely proportional to $|\alpha_B - \alpha_C|$

Answer: B::D

 [Watch Video Solution](#)

88. Two identical containers A and B with frictionless pistons contain the same ideal gas at the same temperature and the same velocity V . The mass of the gas in A is m_A , and that in B is m_B . The gas in each cylinder is now allowed to expand isothermally to the same final volume $2V$. The changes in the pressure in A and B are found to be ΔP and $1.5\Delta P$ respectively. Then

A. $4m_A = 9m_B$

B. $2m_A = 3m_B$

C. $3m_A = 2m_B$

D. $9m_A = 4m_B$

Answer: C



Watch Video Solution

89. Let \bar{v} , v_{rms} and v_p respectively denote the mean speed, Root mean square speed, and most probable speed of the molecules in an ideal monatomic gas at absolute temperature T. The mass of a molecule is m.

Then

A. no molecules can have a speed greater than $\sqrt{2}v_{rms}$

B. no molecule can have a speed less than $v_p / \sqrt{2}$

C. $v_p < \bar{v} < v_{rms}$

D. the average kinetic energy of a molecules is $\frac{3}{4}mv_p^2$.

Answer: C::D



Watch Video Solution

90. A vessel contains a mixture of one mole of oxygen and two moles of nitrogen at 300K. The ratio of the average rotational kinetic energy per O_2 molecules to that per N_2 molecules is

A. 1 : 1

B. 1 : 2

C. 2 : 1


D. depends on the moments of inertia of the two molecules

Answer: A

[Watch Video Solution](#)

91. A black body of temperature T is inside chamber of T_0 temperature initially. Sun rays are allowed to fall from a hole in the top of chamber. If the temperature of black body (T) and chamber (T_0) remains constant,

then



- A. Black body will absorb more radiation
- B. Black body will absorb less radiation
- C. Black body emit more energy
- D. Black body emit energy equal to enrgy absorbed by it

Answer: A::C::D



[Watch Video Solution](#)

92. C_v and C_p denote the molar specific heat capacities of a gas at constant volume and constant pressure, respectively. Then

A. $C_p - C_v$ is larger for a diatomic ideal gas than for a monoatomic ideal gas

B. $C_p + C_v$ is larger for a diatomic ideal gas than for a monoatomic ideal gas

C. C_p / C_v is larger for a diatomic ideal gas than for a monoatomic ideal gas


D. $C_p \cdot C_v$ is larger for a diatomic ideal gas than for a monoatomic ideal gas

Answer: B::D



Watch Video Solution

93. The figure shows the P-V plot of an ideal gas taken through a cycle ABCDA. The part ABC is a semi-circle and CDA is half of an ellipse. Then,




- A. the process during the path $A \rightarrow B$ is isothermal
- B. work done during the path $B \rightarrow C \rightarrow D$
- C. heat flows out of the gas during the path $B \rightarrow C \rightarrow D$
- D. the process during the path $A \rightarrow B$ is isothermal

Answer: B::D

[Watch Video Solution](#)

94. One mole of an ideal gas in initial state A undergoes a cyclic process ABCA, as shown in the figure. Its pressure at A is P_0 . Choose the correct option (s) from the following




- A. Internal energies at A and B are the same
- B. Work done by the gas in process AB is $P_0V_0 \ln 4$
- C. Pressure at C is $\frac{P_0}{4}$
- D. Temperature at C is $\frac{T_0}{4}$

Answer: A::B

[Watch Video Solution](#)

95. A composite block is made of slabs A,B,C,D and E of different thermal conductivities (given in terms of a constant K and sizes (given in terms of length, L) as shown in the figure. All slabs are of same width. Heat ' Q ' flows only from left to right through the blocks. Then in steady state




- A. heat flow through A and E slabs are same.
- B. heat flow through slab E is maximum
- C. temperature difference across slab E is smallest.
- D. heat flow through C = heat flow through B + heat flow through D.

Answer: A::C::D



Watch Video Solution

96. The figure below shows the variation of specific heat capacity (C) of a solid as a function of temperature (T). The temperature is increased continuously from 0 to 500K at a constant rate. Ignoring any volume change, the following statement (s) is (are) correct to a reasonable approximation.



- A. The rate at which heat is absorbed in the range 0-100K varies linearly with temperature T .
- B. Heat absorbed in increasing the temperature from 0-100K is less than the heat required for increasing the temperature from 400-

500K.

C. There is no change in the rate of heat absorption in the range 400-

500K.

D. The rate of heat absorption increases in the range 200-300K.

Answer: A::B::C::D



Watch Video Solution

97. A container of fixed volume has a mixture of a one mole of hydrogen and one mole of helium in equilibrium at temperature T. Assuming the gasses are ideal, the correct statement (s) is (are)

A. The average energy per mole of the gas mixture is $2RT$

B. The ratio of speed of sound in the gas mixture to that in helium gas

is $\sqrt{6/5}$

C. The ratio of the rms speed of helium atoms to that of hydrogen molecules is $1/2$

D. The ratio of the rms speed of helium atoms to that of hydrogen

molecules is $\frac{1}{\sqrt{2}}$


Answer: A::B::C



Watch Video Solution

98. An ideal monoatomic gas is confined in a horizontal cylinder by a spring loaded piston (as shown in the figure). Initially the gas is at temperature T_1 , pressure P_1 and volume V_1 and the spring is in its relaxed state. The gas is then heated very slowly to temperature T_2 , pressure P_2 and volume V_2 . During this process the piston moves out by a distance x . Ignoring the friction between the piston and the cylinder,

the correct statement (s) is (are)



A. If $V_2 = 2V_1$ and $T_2 = 3T_1$, then the energy stored in the

spring is $\frac{1}{4}P_1V_1$

B. If $V_2 = 2V_1$ and $T_2 = 3T_1$, then the change in internal energy is

$3P_1V_1$

C. If $V_2 = 3V_1$ and $T_2 = 4T_1$, then the work done by the gas is

$\frac{7}{3}P_1V_1$

D. If $V_2 = 3V_1$ and $T_2 = 4T_1$, then the heat supplied to the gas is

$\frac{17}{6}P_1V_1$

Answer: B::C



Watch Video Solution


99. A sinker of weight w_0 has an apparent weight w_1 when weighed in a liquid at a temperature t_1 and w_2 when weighed in the same liquid at temperature t_2 . The coefficient of cubical expansion of the material of sinker is β . What is the coefficient of volume expansion of the liquid.



Watch Video Solution

100. Three rods of material X and three rods of material Y are connected as shown in the figure. All the rods are of identical length and cross-sectional area. If the end A is maintained at $60^\circ C$ and the junction E at $10^\circ C$. Calculate the temperature of the junction B, C and D. The thermal conductivity of X is $0.92 \text{ cal/sec} - \text{cm}^\circ C$ and that of Y is

$$0.46 \text{ cal/sec} - \text{cm} - ^\circ \text{C}.$$




[▶ Watch Video Solution](#)

101. Given samples of 1 c.c. of hydrogen and 1 c.c. of oxygen, both at N.T.P. which sample has a larger number of molecules?

[▶ Watch Video Solution](#)

102. A solid material is supplied with heat at a constant rate. The temperature of the material is changing with the heat input as shown in the graph in figure. Study the graph carefully and answer the following

questions:



(i) What do the horizontal regions AB and CD represent?

If CD is equal to 2AB, what do you infer?

What does the slope of DE represents?

The slope of OA is the slope of BC. What does this indicate?

[▶ Watch Video Solution](#)

103. A jar contains a gas and a few drops of water at absolute temperature T_1 . The pressure in the jar is 830mm of mercury. The temperature of the jar is reduced by 1% . The saturation vapour pressures of water at the

two temperatures are 30mm of mercury and 25mm of mercury. Calculate the new pressure in the jar.

 [Watch Video Solution](#)

104. A cyclic process $ABCA$ shown in the $V - T$ diagram is performed with a constant mass of an ideal gas. Show the same process on a $p - V$ diagram. In the figure, CA is parallel to the V -axis and BC is parallel to the T -axis.



 [Watch Video Solution](#)

105. A lead bullet just melts when stopped by an obstacle. Assuming that 25 per cent of the heat is absorbed by the obstacle, find the velocity of the bullet if its initial temperature is 27°C . (Melting point of lead = 327°C , specific heat of lead = $0.03\text{cal or ies/gm}/^\circ\text{C}$, latent heat of fusion of lead = 6cal or ies/gm , $J = 4.2\text{joes/cal or ie}$).

 [Watch Video Solution](#)

 [Watch Video Solution](#)

106. Calculate the work done when one mole of a perfect gas is compressed adiabatically. The initial pressure and volume of the gas are 105 N/m^2 and 6 litres respectively. The final volume of the gas are 2 litre. Molar specific heat of the gas at constant volume is $3R/2$.

 [Watch Video Solution](#)

107. A solid sphere of copper of radius R and a hollow sphere of the same material of inner radius r and outer radius A are heated to the same temperature and allowed to cool in the same environment. Which of them starts cooling faster?

 [Watch Video Solution](#)

108. 1g mole of oxygen at 27°C and 1 atmosphere pressure is enclosed in a vessel.

(a) Assuming the molecules to be moving with v_{rms} , find the number of collisions per second which the molecules make with one square metre area of the vessel wall.

(b) The vessel is next thermally insulated and moves with a constant speed v_0 . It is then suddenly stoppes. The process results in a rise of temperature of the gas by $1^\circ C$. Calculate the speed v_0 . [$k = 1.38 \times 10^{-23} J/K$ and $N_A = 6.02 \times 10^{23} /mol$].




[Watch Video Solution](#)

109. A rectangular box (shown in figure) has a movable and smooth portition which can slide along the length of the box. Both chambers contains 1 mole of monoatomic gas ($\gamma = \frac{5}{3}$) at a pressure walls of box and partition are thermally insulated. Due to heating, gas in left chamber expands until pressure in both chambers become $32P_0$ determine

(a) The final temperature of gas in each chamber

(b) The work done by the gas in the right chamber.



[Watch Video Solution](#)

110. Two glass bulbs of equal volume are connected by a narrow tube and are filled with a gas at $0^{\circ}C$ and a pressure of 76cm of mercury. One of the bulbs is then placed in melting ice and the other is placed in a water bath maintained at $62^{\circ}C$. What is the new value of the pressure inside the bulbs? The volume of the connecting tube is negligible.

[Watch Video Solution](#)

111. A thin tube of uniform cross section is sealed at both ends. It lies horizontally, the middle 5cm containing mercury and the parts on its two sides containing air at the same pressure p . When the tube is held at an angle of 60° with the vertical, the length of the air column above and below the mercury pellet are 46cm and 44.5cm respectively. Calculate the pressure in centimeters of mercury, The temperature of the system is kept at 30°C .



[Watch Video Solution](#)

112. An ideal gas has a specific heat at constant pressure $C_P = \frac{5R}{2}$. The gas is kept in a closed vessel of volume 0.0083m^3 , at a temperature of 300K and a pressure of $1.6 \times 10^6\text{N/m}^2$. An amount of 2.49×10^4 Joules of heat energy is supplied to the gas. Calculate the final temperature and pressure of the gas.



[Watch Video Solution](#)

113. Two moles of helium gas ($\lambda = 5/3$) are initially at temperature $27^\circ C$ and occupy a volume of 20 litres. The gas is first expanded at constant pressure until the volume is doubled. Then it undergoes an adiabatic change until the temperature returns to its initial value.

(i) Sketch the process on a p-V diagram.

(ii) What are the final volume and pressure of the gas?


(iii) What is the work done by the gas ?



[Watch Video Solution](#)

114. An ideal monatomic gas is confined in a cylinder by a spring-loaded piston of cross-section $8.0 \times 10^{-3} m^2$. Initially the gas is at 300K and occupies a volume of $2.4 \times 10^{-3} m^3$ and the spring is in its relaxed (unstretched, un compressed) state, fig. The gas is heated by a small electric heater until the piston moves out slowly by 0.1m. Calculate the final temperature of the gas and the heat supplied (in joules) by the heater. The force constant of the spring is $8000 N/m$, atmospheric pressure is $1.0 \times 10^5 N m^{-2}$. The cylinder and the piston are thermally

insulated. The piston is massless and there is no friction between the piston and the cylinder. Neglect heat loss through lead wires of the heater. The heat capacity of the heater coil is negligible. Assume the spring to be massless.



[Watch Video Solution](#)

115. An ideal gas having initial pressure P , volume V and temperature T is allowed to expand adiabatically until its volume becomes $5.66V$ while its temperature falls to $T/2$.

(i) How many degrees of freedom do the gas molecules have?

Obtain the work done by the gas during the expansion as a function of the initial pressure P and volume V .

 [Watch Video Solution](#)


116. Three moles of an ideal gas $\left(C_p = \frac{7}{2}R\right)$ at pressure, P_A and temperature T_A is isothermally expanded to twice its initial volume. It is then compressed at constant pressure to its original volume. Finally gas is compressed at constant volume to its original pressure P_A .

(a) Sketch P-V and P-T diagrams for the complete process.

(b) Calculate the net work done by the gas, and net heat supplied to the gas during the complete process.

 [Watch Video Solution](#)

117. Two moles of helium gas undergo a cyclic process as shown in Fig. Assuming the gas to be ideal, calculate the following quantities in this process



- The net change in the heat energy
- The net work done
- The net change in internal energy

[▶ Watch Video Solution](#)


118. One mole of a monoatomic ideal gas is taken through the cycle shown in Fig:

$A \rightarrow B$: adiabatic expansion

$B \rightarrow C$: cooling at constant volume

$C \rightarrow D$: adiabatic compression

$D \rightarrow A$: heating at constant volume



The pressure and temperature at A,B,etc. are denoted by P_A, T_A, P_B, T_B etc. respectively. Given that $T_A = 1000K$,

$P_B = (2/3)P_A$ and $P_C = (1/3)P_A$, calculate the following quantities:

- (i) The work done by the gas in the process $A \rightarrow B$
- (ii) The heat lost by the gas in the process $B \rightarrow C$.
- (iii) The temperature T_D . [Given : $(2/3)^{2/5} = 0.85$]

[▶ Watch Video Solution](#)

119. An ideal gas is taken through a cyclic thermodynamic process through four steps. The amounts of heat involved in these steps are $Q_1 = 5960J, Q_2 = -5585J, Q_3 = -2980J$ and $Q_4 = 3645J$,

respectively. The corresponding quantities of work involved are

$W_1 = 2200J$, $W_2 = -825J$, $W_3 = -1100J$ and W_4 respectively.

(1) Find the value of W_4 .

(2) What is the efficiency of the cycle

 [Watch Video Solution](#)

120. A closed container of volume $0.02m^3$ contains a mixture of neon and argon gases, at a temperature of $27^\circ C$ and pressure of $1 \times 10^5 Nm^{-2}$.

The total mass of the mixture is 28g. If the molar masses of neon and argon are 20 and $40gmol^{-1}$ respectively, find the masses of the individual gasses in the container assuming them to be ideal (Universal gas constant $R = 8.314J/mol - K$).

 [Watch Video Solution](#)

121. A gaseous mixture enclosed in a vessel of volume V consists of one mole of a gas A with $\gamma = (C_p/C_v) = 5/3$ and another gas B with $\gamma = 7/5$ at a certain temperature T . The relative molar masses of the

gases A and B are 4 and 32, respectively. The gases A and B do not react with each other and are assumed to be ideal. The gaseous mixture follows the equation $PV^{19/13} = \text{constant}$, in adiabatic processes.

- (a) Find the number of moles of the gas B in the gaseous mixture.
- (b) Compute the speed of sound in the gaseous mixture at $T = 300\text{K}$.
- (c) If T is raised by 1K from 300K , find the % change in the speed of sound in the gaseous mixture.
- (d) The mixture is compressed adiabatically to $1/5$ of its initial volume V . Find the change in its adiabatic compressibility in terms of the given quantities.

 [Watch Video Solution](#)

122. At 27°C two moles of an ideal monoatomic gas occupy a volume V . The gas expands adiabatically to a volume $2V$. Calculate (i) the final temperature of the gas, (ii) change in its internal energy, and (iii) the work done by the gas during this process.

 [Watch Video Solution](#)

123. The temperature of 100g of water is to be raised from $24^{\circ}C$ to $90^{\circ}C$ by adding steam to it. Calculate the mass of the steam required for this purpose.


 [Watch Video Solution](#)

124. One mole of a diatomic ideal gas ($\gamma = 1.4$) is taken through a cyclic process starting from point A. The process $A \rightarrow B$ is an adiabatic compression, $B \rightarrow C$ is isobaric expansion, $C \rightarrow D$ is an adiabatic expansion, and $D \rightarrow A$ is isochoric. The volume ratios are $V_A/V_B = 16$ and $V_C/V_B = 2$ and the temperature at A is $T_A = 300K$. Calculate the temperature of the gas at the points B and D and find the efficiency of the cycle.

 [Watch Video Solution](#)


125. The apparatus shown in the figure consists of four glass columns connected by horizontal section. The height of two central column B and

C are 49 cm each. The two outer columns A and D are open to the atmosphere. A and C are maintained at a temperature of 95°C while the columns B and D are maintained at 5°C . The height of the liquid in A and D measured from the base are 52.8 cm and 51 cm respectively. Determine the coefficient of thermal expansion of the liquid



[▶ Watch Video Solution](#)

126. One mole of an ideal monatomic gas is taken round the cyclic process ABCA as shown in figure. Calculate



- the work done by the gas.
- the heat rejected by the gas in the path CA and the heat absorbed by the gas in the path AB,
- the net heat absorbed by the gas in the path BC,
- the maximum temperature attained by the gas during the cycle.

[▶ Watch Video Solution](#)

127. A solid body X of heat capacity C is kept in an atmosphere whose temperature is $T_A = 300K$. At time $t = 0$ the temperature of X is $T_0 = 400K$. It cools according to Newton's law of cooling. At time t_1 , its temperature is found to be $350K$. At this time (t_1), the body X is

connected to a large box Y at atmospheric temperature is T_4 , through a conducting rod of length L , cross-sectional area A and thermal conductivity K . The heat capacity Y is so large that any variation in its temperature may be neglected. The cross-sectional area A of the connecting rod is small compared to the surface area of X . Find the temperature of X at time $t = 3t_1$.


 [Watch Video Solution](#)

128. Two moles of an ideal monatomic gas, initially at pressure p_1 and volume V_1 , undergo an adiabatic compression until its volume is V_2 . Then the gas is given heat Q at constant volume V_2 .

- (i) Sketch the complete process on a p - V diagram.
- (b) Find the total work done by the gas, the total change in its internal energy and the final temperature of the gas. [Give your answer in terms of p_1 , V_1 , V_2 , Q and R]

 [Watch Video Solution](#)

129. Two moles of an ideal monatomic gas is taken through a cycle ABCA as shown in the P-T diagram. During the process AB, pressure and temperature of the gas vary such that $PT = \text{Constant}$. It $T_1 = 300\text{K}$, calculate



- (a) the work done on the gas in the process AB and
 (b) the heat absorbed or released by the gas in each of the processes.
- Give answer in terms of the gas constant R.


[▶ Watch Video Solution](#)

130. An ice cube of mass 0.1kg at 0°C is placed in an isolated container which is at 227°C . The specific heat S of the container varies with

temperature T according to the empirical relation $S = A + BT$, where $A = 100 \text{ cal/kg} - K$ and $B = 2 \times 10^{-2} \text{ cal/kg} - K^2$. If the final temperature of the container is $27^\circ C$, determine the mass of the container. (Latent heat of fusion of water $= 8 \times 10^4 \text{ cal/kg}$, Specific heat of water $= 10^3 \text{ cal/kg} - K$).

 **Watch Video Solution**

131. A monoatomic ideal gas of two moles is taken through a cyclic process starting from A as shown in figure. The volume ratios are $\frac{V_B}{V_A} = 2$ and $\frac{V_D}{V_A} = 4$. If the temperature T_A at A is $27^\circ C$.



Calculate,

- (a) the temperature of the gas at point B,
- (b) heat absorbed or released by the gas in each process,
- (c) the total work done by the gas during the complete cycle. Express your answer in terms of the gas constant R.

 [Watch Video Solution](#)

132. A cubical box of side 1 meter contains helium gas (atomic weight 4) at a pressure of $100\text{N}/\text{m}^2$. During an observation time of 1 second, an atom travelling with the root-mean-square speed parallel to one of the edges of the cube, was found to make 500 hits with a particular wall, without any collision with other atoms. Take

$$R = \frac{25}{3} \text{ Jmol}^{-1} \text{ K}^{-1} \text{ and } k = 1.38 \times 10^{-23} \text{ J/K}$$


- (a) Evaluate the temperature of the gas.
- (b) Evaluate the average kinetic energy per atom.
- (c) Evaluate the total mass of helium gas in the box.

 [Watch Video Solution](#)

133. An insulated container containing monoatomic gas of molar mass s is moving with a velocity v_0 . If the container is suddenly stopped, find the change in temperature.

 [Watch Video Solution](#)

134. Hot oil is circulated through an insulated container with a wooden lid at the top whose conductivity $K = 0.149 \text{ J}/(\text{m} \cdot ^\circ \text{C} \cdot \text{sec})$, thickness $t = 5 \text{ mm}$, emissivity $= 0.6$. Temperature of the top of the lid is maintaining at $T_l = 127^\circ \text{C}$. If the ambient temperature $T_a = 27^\circ \text{C}$.




Calculate:

(a) rate of heat loss per unit area due to radiation from the lid.

(b) temperature of the oil. $\left(\text{Given } \sigma = \frac{17}{3} \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4} \right)$

 [Watch Video Solution](#)

135. A diatomic gas is enclosed in a vessel fitted with massless movable piston. Area of cross section of vessel is 1 m^2 . Initial height of the piston is 1 m (see the figure). The initial temperature of the gas is 300 K . The temperature of the gas is increased to 400 K , keeping pressure constant, calculate the new height of the piston. The piston is brought to its initial position with no heat exchange. Calculate the final temperature of the gas. You can leave answer in fraction.





[Watch Video Solution](#)


136. A small spherical body of radius r is falling under gravity in a viscous medium. Due to friction the medium gets heated. How does the rate of heating depends on radius of body when it attains terminal velocity?



[Watch Video Solution](#)

137. A cylinder rod of length l , thermal conductivity K and area of cross section A has one end in the furnace at temperature T_1 and the other end in surrounding at temperature T_2 . Surface of the rod exposed to the surrounding has emissivity e . Also $T_2 = T_s + \Delta T$ and $T_s \gg \Delta T$. If

$T_1 - T_s \propto \Delta T$, find the proportionality constant.



[▶ Watch Video Solution](#)

138. A cubical block of co-efficient of linear expansion α_s is submerged partially inside a liquid of co-efficient of volume expansion γ_l . On increasing the temperature of the system by ΔT , the height of the cube inside the liquid remains unchanged. Find the relation between α_s and γ_l .

[▶ Watch Video Solution](#)

139. A cylinder of mass 1kg is given heat of 20,000J at atmospheric pressure. If initially the temperature of cylinder is $20^{\circ}C$, find

- (a) final temperature of the cylinder.
- (b) work done by the cylinder.
- (c) change in internal energy of the cylinder

(Given that specific heat of cylinder= $400Jkg^{-1} \text{ } ^{\circ}C^{-1}$, Atmospheric pressure= $10^5N/m^2$ and Density of cylinder= $9000kg/m^3$)



[Watch Video Solution](#)

140. 0.05 kg steam at 373K and 0.45kg of ice at 253K are mixed in an insulated vessel. Find the equilibrium temperature of the mixture. Given,


$$L_{fusion} = 80cal/g = 336J/g, L_{vap or ization} = 540cal/g = 2268J/g, C_{ice} =$$



[Watch Video Solution](#)

141. A fixed thermally conducting cylinder has a radius R and height L_0 . The cylinder is open at its bottom and has a small hole at its top. A piston

of mass M is held at a distance L from the top surface, as shown in the figure. The atmospheric pressure is P_0 .



The piston is now pulled out slowly and held at a distance $2L$ from the top. The pressure in the cylinder between its top and the piston will then be


- A. P_0
- B. $\frac{P_0}{2}$
- C. $\frac{P_0}{2} + \frac{Mg}{\pi R^2}$
- D. $\frac{P_0}{2} - \frac{Mg}{\pi R^2}$

Answer: A





142. A fixed thermally conducting cylinder has a radius R and height L_0 . The cylinder is open at its bottom and has a small hole at its top. A piston of mass M is held at a distance L from the top surface, as shown in the figure. The atmospheric pressure is P_0 .



While the piston is at a distance $2L$ from the top, the hole at the top is sealed. The piston is then released, to a position where it can stay in equilibrium. In this condition, the distance of the piston from the top is

A. $\left(\frac{2P_0\pi R^2}{\pi R^2 P_0 + Mg}\right)(2L)$


B. $\left(\frac{P_0\pi R^2 - Mg}{\pi R^2 P_0}\right)(2L)$

- C. $\left(\frac{P_0\pi R^2 + Mg}{\pi R^2 P_0}\right)(2L)$
- D. $\left(\frac{P_0\pi R^2}{\pi R^2 P_0 - Mg}\right)(2L)$

Answer: D


 [Watch Video Solution](#)

143. A fixed thermally conducting cylinder has a radius R and height L_0 . The cylinder is open at its bottom and has a small hole at its top. A piston of mass M is held at a distance L from the top surface, as shown in the figure. The atmospheric pressure is P_0 .



The piston is taken completely out of the cylinder. The hole at the top is

sealed. A water tank is brought below the cylinder and put in a position so that the water surface in the tank is at the same level as the top of the cylinder as shown in the figure. The density of the water is ρ . In equilibrium, the height H of the water column in the cylinder satisfies




- A. $\rho g(L_0 - H)^2 + P_0(L_0 - H) + L_0 P_0 = 0$
- B. $\rho g(L_0 - H)^2 - P_0(L_0 - H) - L_0 P_0 = 0$
- C. $\rho g(L_0 - H)^2 + P_0(L_0 - H) - L_0 P_0 = 0$
- D. $\rho g(L_0 - H)^2 - P_0(L_0 - H) + L_0 P_0 = 0$

Answer: C

 [Watch Video Solution](#)

144. A small spherical monoatomic ideal gas bubble ($\gamma = 5/3$) is trapped inside a liquid of density ρ (see figure). Assume that the bubble does not exchange any heat with the liquid. The bubble contains n moles of gas. The temperature of the gas when the bubble is at the bottom is T_0 , the height of the liquid is H and the atmospheric pressure P_0 (Neglect surface tension).



As the bubble moves upwards, besides the buoyancy force the following forces are acting on it

- A. Only the force of gravity

B. The force due to gravity and the force due to the pressure of the liquid

C. The force due to gravity, the force due to the pressure of the liquid and the force due to viscosity of the liquid


D. The force due to gravity and the force due to viscosity of the liquid

Answer: D



[Watch Video Solution](#)

145. A small spherical monoatomic ideal gas bubble ($\gamma = 5/3$) is trapped inside a liquid of density ρ (see figure). Assume that the bubble does not exchange any heat with the liquid. The bubble contains n moles of gas. The temperature of the gas when the bubble is at the bottom is T_0 , the height of the liquid is H and the atmospheric pressure P_0 (Neglect surface tension).



When the gas bubble is at a height y from the bottom, its temperature is-


- A. $T_0 \left(\frac{P_0 + \rho_l g H}{P_0 + \rho_l g y} \right)^{2/5}$
- B. $T_0 \left(\frac{P_0 + \rho_l g (H - y)}{P_0 + \rho_l g H} \right)^{2/5}$
- C. $T_0 \left(\frac{P_0 + \rho_l g H}{P_0 + \rho_l g y} \right)^{3/5}$
- D. $T_0 \left(\frac{P_0 + \rho_l g (H - y)}{P_0 + \rho_l g H} \right)^{3/5}$

Answer: B



Watch Video Solution

146. A small spherical monoatomic ideal gas bubble ($\gamma = 5/3$) is trapped inside a liquid of density ρ (see figure). Assume that the bubble does not exchange any heat with the liquid. The bubble contains n moles of gas. The temperature of the gas when the bubble is at the bottom is T_0 , the height of the liquid is H and the atmospheric pressure P_0 (Neglect surface tension).



The buoyancy force acting on the gas bubble is (Assume R is the universal gas constant)

A. $\rho_l n R g T_0 \frac{(P_0 + \rho_l g H)^{2/5}}{[P_0 + \rho_l g y]^{7/5}}$

B. $\frac{\rho_l n R g T_0}{(P_0 + \rho_l g H)^{2/5} [P_0 \rho_l g (H - y)]^{3/5}}$

$$C. \rho_l n R g T_0 \frac{(P_0 + \rho_l g H)^{3/5}}{[P_0 + \rho_l g y]^{8/5}}$$


$$D. \frac{\rho_l n R g T_0}{(P_0 + \rho_l g H)^{3/5} [P_0 \rho_l g (H - y)]^{2/5}}$$

Answer: B



Watch Video Solution

147. In the figure, a container is shown to have a movable (without friction) piston on top. The container and the piston are all made of perfectly insulated material allowing no heat transfer between outside and inside the container. The container is divided into two compartments by a rigid partition made of a thermally conducting material that allows slow transfer of heat. The lower compartment of the container is filled with 2 moles of an ideal monatomic gas at 700K and the upper compartment is filled with 2 moles of an ideal diatomic gas at 400K. The heat capacities per mole of an ideal monatomic gas are $C_V = \frac{3}{2}R$, $C_P = \frac{5}{2}R$, and those for an ideal diatomic gas are $C_V = \frac{5}{2}R$, $C_P = \frac{7}{2}R$,



Consider the partition to be rigidly fixed so that it does not move. When equilibrium is achieved, the final temperature of the gasses will be


- A. $550K$
- B. $525K$
- C. $513K$
- D. $490K$

Answer: B



[Watch Video Solution](#)

148. In the figure, a container is shown to have a movable (without friction) piston on top. The container and the piston are all made of perfectly insulated material allowing no heat transfer between outside and inside the container. The container is divided into two compartments by a rigid partition made of a thermally conducting material that allows slow transfer of heat. The lower compartment of the container is filled with 2 moles of an ideal monatomic gas at 700K and the upper compartment is filled with 2 moles of an ideal diatomic gas at 400K. The heat capacities per mole of an ideal monatomic gas are $C_V = \frac{3}{2}R$, $C_P = \frac{5}{2}R$, and those for an ideal diatomic gas are $C_V = \frac{5}{2}R$, $C_P = \frac{7}{2}R$,



Now consider the partition to be free to move without friction so that

the pressure of gasses in both compartments is the same. The total work done by the gases till the time they achives equilibrium will be

A. $-200K$

B. $200R$

C. $100R$

D. $-100R$

Answer: D



Watch Video Solution

149. Statement-1: The total translational kenetic energy of fall the molecules of a given mass of an ideal gas is 1.5 times the product of its pressure and its volume because.

Statement-2: The molecules of a gas collide with each other and the velocities of the molecules change due to the collision.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-2 is False, Statement-2 is True

Answer: B

 [Watch Video Solution](#)

150. A metal rod AB of length $10x$ has its one end A in ice at $0^\circ C$, and the other end B in water at $100^\circ C$. If a point P on the rod is maintained at $400^\circ C$, then it is found that equal amounts of water and ice evaporate and melt per unit time. The latent heat of evaporation of water is $540\text{cal}/g$ and latent heat of melting of ice is $80\text{cal}/g$. If the point P is at a distance of λx from the ice end A, find the value λ . [Neglect any heat loss to the surrounding.]



[Watch Video Solution](#)

151. A piece of ice (heat capacity = $2100 \text{ J kg}^{-1} \text{ } ^\circ\text{C}^{-1}$ and latent heat = $3.36 \times 10^5 \text{ J kg}^{-1}$) of mass m grams is at -5°C at atmospheric pressure. It is given 420J of heat so that the ice starts melting. Finally when the ice-water mixture is in equilibrium, it is found that 1gm of ice has melted. Assuming there is no other heat exchange in the process, the value of m is



[Watch Video Solution](#)

152. Two spherical bodies, A(radius 6cm) and B(radius 18cm) are at temperature T_1 and T_2 , respectively. The maximum intensity in the emission spectrum of A is at 500nm and in that of B is at 1500nm. Considering them to be black bodies, what will be the ratio of the rate of total energy radiated by A to that of B?



[Watch Video Solution](#)

153. A diatomic ideal gas is compressed adiabatically to $1/32$ of its initial volume. If the initial temperature of the gas is T_i (in Kelvin) and the final temperature is T_f , the value of T_f/T_i is

 [Watch Video Solution](#)

154. Steel wire of length ' L ' at $40^\circ C$ is suspended from the ceiling and then a mass ' m ' is hung from its free end. The wire is cooled down from $40^\circ C \rightarrow 30^\circ C$ to regain its original length ' L '. The coefficient of linear thermal expansion of the steel is $10^{-5}/^\circ C$, Young's modulus of steel is $10^{11} N/m^2$ and radius of the wire is 1mm. Assume that $L \gg$ diameter of the wire. Then the value of ' m ' in kg is nearly

 [Watch Video Solution](#)

155. A thermodynamic system is taken from an initial state I with internal energy $U_i = -100J$ to the final state f along two different paths iaf and ibf, as schematically shown in the figure. The work done by the system


along the path af, ib and bf are

$W_{af} = 200J$, $W_{ib} = 50J$ and $W_{bf} = 100J$ respectively. The heat

supplied to the system along the path ia, af, ib and bf are Q_{iaf} , Q_{ib} , Q_{bf}

respectively. If the internal energy of the system in the state b is

$U_b = 200J$ and $Q_{iaf} = 500J$, The ratio $\frac{Q_{bf}}{Q_{ib}}$ is



[Watch Video Solution](#)

156. Two spherical stars A and B emit blackbody radiation. The radius of A is 400 times that of B and A emits 10^4 times the power emitted from B.

The ratio $\left(\frac{\lambda_A}{\lambda_B}\right)$ of their wavelengths λ_A and λ_B at which the peaks occur in their respective radiation curves is

 [Watch Video Solution](#)

157. A metal is heated in a furnace where a sensor is kept above the metal surface to read the power radiated (P) by the metal. The sensor has scale that displays $\log_2, (P / P_0)$, where P_0 is constant. When the metal surface is at a temperature of $487^\circ C$, the sensor shows a value 1. Assume that the emissivity of the metallic surface remains constant. What is the value displayed by the sensor when the temperature of the metal surface is raised to $2767^\circ C$?

 [Watch Video Solution](#)

158. Which statement is incorrect ?

- A. all reversible cycles have same efficiency
- B. reversible cycle has more efficiency than an irreversible one
- C. Carnot cycle is reversible one
- D. Carnot cycle has the maximum efficiency in all cycles.

Answer: A



[Watch Video Solution](#)

159. Heat given to a body which raises its temperature by $1^{\circ}C$ is

- A. water equivalent
- B. thermal capacity
- C. specific heat
- D. temperature gradient

Answer: B



[Watch Video Solution](#)

160. Infrared radiation is detected by

- A. spectrometer

B. pyrometer

C. nanometer

D. photometer

Answer: B



Watch Video Solution

161. Which of the following is more close to a black body?

A. black board paint

B. green leaves

C. black holes

D. red roses

Answer: A



Watch Video Solution

162. Cooking gas containers are kept in a lorry moving with uniform speed.

The temperature of the gas molecules inside will

- A. increase
- B. decrease
- C. remain same
- D. decrease for some, while increase for others

Answer: A



[Watch Video Solution](#)

163. If mass-energy equivalence is taken into account, when water is cooled to form ice, the mass of water should

- A. increase
- B. remains unchanged
- C. decrease

D. first increase then decrease

Answer: C



[Watch Video Solution](#)

164. At what temperature is the r.m.s velocity of a hydrogen molecule equal to that of an oxygen molecule at $24^{\circ}C$?

A. $80K$

B. $-73K$

C. $3K$

D. $20K$

Answer: D



[Watch Video Solution](#)

165. Even Carnot engine cannot give 100 % efficiency because we cannot

- A. prevent radiation
- B. find ideal sources
- C. reach absolute zero temperature
- D. eliminate friction.

Answer: C



[Watch Video Solution](#)

166. 1 mole of a gas with $\gamma = 7/5$ is mixed with 1 mole of a gas with $\gamma = 5/3$, then the value of γ for the resulting mixture is

- A. $7/5$
- B. $2/5$
- C. $24/16$
- D. $12/7$

Answer: C



[Watch Video Solution](#)

167. Two spheres of the same material have radii 1m and 4m and temperatures 4000K and 2000K respectively. The ratio of the energy radiated per second by the first sphere to that by the second is

A. 1 : 1

B. 16 : 1

C. 4 : 1

D. 1 : 9

Answer: A



[Watch Video Solution](#)

168. "Heat cannot by itself flow from a body at lower temperature to a body at higher temperature" is a statement or consequence of

- A. second law of thermodynamics
- B. conservation of momentum
- C. conservation of mass
- D. first law of thermodynamics

Answer: A



[Watch Video Solution](#)

169. During an adiabatic process, the pressure of a gas is found to be proportional to the cube of its absolute temperature. The ratio C_P / C_V for the gas is

A. $\frac{4}{3}$

B. 2

C. $\frac{5}{3}$

D. $\frac{3}{2}$

Answer: D



Watch Video Solution

170. Which of the following parameters does not characterize the thermodynamic state of matter?

A. Temperature

B. pressure

C. Work

D. Volume

Answer: C



Watch Video Solution

171. A Carnot engine takes $3 \times 10^6 \text{ cal.}$ of heat from a reservoir at 62° C , and gives it to a sink at 27° C . The work done by the engine is

A. $4.2 \times 10^6 \text{ J}$

B. $8.4 \times 10^6 \text{ J}$

C. $16.8 \times 10^6 \text{ J}$

D. zero

Answer: B



[Watch Video Solution](#)

172. The earth radiates in the infra-red region of the spectrum. The spectrum is correctly given by

A. Rayleigh Jeans law

B. Plank's law of radiation

C. Stefan's law of radiation

D. Wien's law

Answer: D



[Watch Video Solution](#)

173. According to Newton's law of cooling, the rate of cooling of a body is proportional to $(\Delta\theta)^n$, where $\Delta\theta$ is the difference of the temperature of the body and the surroundings, and n is equal to

A. two

B. three

C. four

D. one

Answer: D



[Watch Video Solution](#)

174. One mole of ideal monoatomic gas ($\gamma = 5/3$) is mixed with one mole of diatomic gas ($\gamma = 7/5$). What is γ for the mixture? γ Denotes the ratio of specific heat at constant pressure, to that at constant volume

A. $35/23$

B. $23/15$

C. $3/20$

D. $4/3$

Answer: C



[Watch Video Solution](#)

175. If the temperature of the sun were to increase from T to $2T$ and its radius from R to $2R$, then the ratio of the radiant energy received on earth to what it was previously will be

A. 32

B. 16

C. 4

D. 64

Answer: D



[Watch Video Solution](#)

176. Which of the following statements is correct for any thermodynamic system?

A. The change in entropy can never be zero

B. Internal energy and entropy are state functions

C. The internal energy changes in all processes

D. The work done in an adiabatic process is always zero.

Answer: B



[Watch Video Solution](#)

177. Two thermally insulated vessels 1 and 2 are filled with air at temperature (T_1, T_2) , volume (V_1, V_2) and pressure (P_1, P_2) respectively. If the valve joining the two vessels is opened, the temperature inside the vessel at equilibrium will be

A. $T_1 T_2 (P_1 V_1 + P_2 V_2) / (P_1 V_1 T_2 + P_2 V_2 T_1)$

B. $(T_1 + T_2) / 2$

C. $T_1 + T_2$

D. $T_1 T_2 (P_1 V_1 + P_2 V_2) / (P_1 V_1 T_1 + P_2 V_2 T_2)$


Answer: A



[Watch Video Solution](#)

178. The temperature of the two outer surfaces of a composite slab, consisting of two materials having coefficients of thermal conductivity K and $2K$ and thickness x and $4x$, respectively, are T_2 and T_1 ($T_2 > T_1$). The rate of heat transfer through

the slab, in a steady state is $\left(\frac{A(T_2 - T_1)K}{2}\right) f$, with f equal to



- A. $\frac{2}{3}$
- B. $\frac{1}{2}$
- C. 1
- D. $\frac{1}{3}$

Answer: D



Watch Video Solution

179. Which of the following is incorrect regarding the first law of thermodynamics?

- A. It is a restatement of the principle of conservation of energy
- B. It is not applicable to any cyclic process
- C. It introduces the concept of the internal energy
- D. It introduces the concept of the internal energy


Answer: B::C



[Watch Video Solution](#)

180. The figure shows a system of two concentric spheres of radii r_1 and r_2 are kept at temperature T_1 and T_2 , respectively. The radial rate of flow of heat in a substance between the two concentric spheres is

proportional to



A. $\ln\left(\frac{r_2}{r_1}\right)$

B. $\frac{r_2 - r_1}{r_1 r_2}$

C. $(r_1 - r_2)$


D. $\frac{r_1 r_2}{r_2 - r_1}$

Answer: D



Watch Video Solution

181. A system goes from A and B via two processes. I and II as shown in figure. If ΔU_1 and ΔU_2 are the changes in internal energies in the processes I and II respectively, then




- A. relation between ΔU_1 and ΔU_2 can not be determined
- B. $\Delta U_1 = \Delta U_2$
- C. $\Delta U_2 < \Delta U_1$
- D. $\Delta U_2 > \Delta U_1$

Answer: B



[Watch Video Solution](#)

182. The temperature -entropy diagram of a reversible engine cycle is given in the figure. Its efficiency is



- A. $\frac{1}{4}$
- B. $\frac{1}{2}$
- C. $\frac{2}{3}$
- D. $\frac{1}{3}$

Answer: D



Watch Video Solution

183. A gaseous mixture consists of 16g of helium and 16 g of oxygen. The

ratio $\frac{C_p}{C_v}$ of the mixture is

A. 1.62

B. 1.59

C. 1.54

D. 1.4

Answer: A



[Watch Video Solution](#)

184. Assuming the Sun to be a spherical body of radius R at a temperature of T K, evaluate the total radiant power incident on Earth at a distance r from the sun

where r_0 is the radius of the Earth and σ is Stefan's constant.

A. $4\pi r_0^2 R^2 \sigma \frac{T^4}{r^2}$

$$\text{B. } \pi r_0^2 R^2 \sigma \frac{T^4}{r^2}$$

$$\text{C. } r_0^2 R^2 \sigma \frac{T^4}{4\pi r^2}$$

$$\text{D. } R^2 \sigma \frac{T^4}{r^2}$$

Answer: B



Watch Video Solution

185. Two rigid boxes containing different ideal gases are placed on a table. Box A contains one mole of nitrogen at temperature T_0 , while Box contains one mole of helium at temperature $\left(\frac{7}{3}\right)T_0$. The boxes are then put into thermal contact with each other, and heat flows between them until the gasses reach a common final temperature (ignore the heat capacity of boxes). Then, the final temperature of the gasses, T_f in terms of T_0 is

$$\text{A. } T_f = \frac{3}{7}T_0$$

$$\text{B. } T_f = \frac{7}{3}T_0$$

$$C. T_f = \frac{3}{2}T_0$$

$$D. T_f = \frac{5}{2}T_0$$

Answer: C



Watch Video Solution

186. The work of 142kJ is performed in order to compress one kilo mole of gas adiabatically and in this process the temperature of the gas increases by $7^\circ C$. The gas is ($R = 8.3Jmol^{-1}K^{-1}$)

A. diatomic

B. triatomic

C. a mixture of monoatomic and diatomic


D. monoatomic

Answer: A



Watch Video Solution

187. When a system is taken from state i to state f along the path iaf , it is found that $Q = 50\text{cal}$ and $W = 20\text{cal}$. Along the path ibf $Q = 36\text{cal}$. W along the path ibf is



A. 14cal

B. 6cal

C. 16cal

D. 66cal

Answer: B

 Watch Video Solution

188. A Carnot engine, having an efficiency of $\eta = 1/10$ as heat engine, is used as a refrigerator. If the work done on the system is 10J, the amount of energy absorbed from the reservoir at lower temperature is

A. 100J

B. 99J

C. 90J

D. 1J


Answer: C



[Watch Video Solution](#)

189. One end of a thermally insulated rod is kept at a temperature T_1 and the other at T_2 . The rod is composed of two sections of length l_1 and l_2 and thermal conductivities K_1 and K_2 respectively. The temperature at

the interface of the two section is



- A. $\frac{K_1 l_1 T_1 + K_2 l_2 T_2}{K_1 l_1 + K_2 l_2}$
- B. $\frac{K_2 l_2 T_1 + K_1 l_1 T_2}{K_1 l_1 + K_2 l_2}$
- C. $\frac{K_2 l_1 T_1 + K_1 l_2 T_2}{K_2 l_1 + K_1 l_2}$
- D. $\frac{K_1 l_2 T_1 + K_2 l_1 T_2}{K_1 l_2 + K_2 l_1}$

Answer: D



Watch Video Solution

190. If C_P and C_v denote the specific heats nitrogen per unite mass at constant pressure and constant volume respectively, then

$$(1) C_P - C_v = \frac{R}{28} \quad (2) C_P - C_v = \frac{R}{14}$$

$$(3) C_P - C_v = R \quad (4) C_P - C_v = 28R$$

A. $C_P - C_V = 28R$

B. $C_P - C_V = R/28$

C. $C_P - C_V = R/14$

D. $C_P - C_V = R$

Answer: B



Watch Video Solution

191. The speed of sound in oxygen (O_2) at a certain temperature is $460ms^{-1}$. The speed of sound in helium (He) at the same temperature will be (assume both gases to be ideal)

A. $1421ms^{-1}$

B. $500ms^{-1}$

C. $650ms^{-1}$

D. $330ms^{-1}$

Answer: A

 [Watch Video Solution](#)

192. An insulated container of gas has two chambers separated by an insulating partition. One of the chambers has volume V_1 and contains ideal gas at pressure P_1 and temperature T_1 . The other chamber has volume V_2 and contains ideal gas at pressure P_2 and temperature T_2 . If the partition is removed without doing any work on the gas, the final equilibrium temperature of the gas in the container will be

A. $\left(T_1 T_2 \frac{P_1 V_1 + P_2 V_2}{P_1 V_1 T_2 + P_2 V_2 T_1} \right)$

B. $\frac{P_1 V_1 T_1 + P_2 V_2 T_2}{P_1 V_1 + P_2 V_2}$

$$C. \frac{P_1 V_1 T_2 + P_2 V_2 T_1}{P_1 V_1 + P_2 V_2}$$


$$D. \frac{T_1 T_2 (P_1 V_1 + P_2 V_2)}{P_1 V_1 T_1 + P_2 V_2 T_2}$$

Answer: A




Watch Video Solution



193. A long metallic bar is carrying heat from one end to the other end under steady-state. The variation of temperature θ along the length x of the bar from its hot end is best described by which of the following figures?



A.




B.



Answer: A

 [Watch Video Solution](#)

194. Two moles of helium gas are taken over the cycle ABCDA, as shown in the P-T diagram



Assuming the gas to be ideal the work done on the gas in taking it from A to B is :

A. $300R$

B. $400R$

C. $500R$


D. $200R$

Answer: B



[Watch Video Solution](#)

195. Two moles of helium gas are taken over the cycle ABCDA, as shown in the P-T diagram



The work done on the gas in taking it from D to A is :


- A. $+414R$
- B. $-690R$
- C. $+690R$
- D. $-414R$

Answer: A



Watch Video Solution

196. Two moles of helium gas are taken over the cycle ABCDA, as shown in the P-T diagram



The net work done on the gas in the cycle ABCDA is :

- A. $276R$
- B. $1076R$
- C. $1904R$
- D. zero

Answer: A



[Watch Video Solution](#)

197. One kg of a diatomic gas is at pressure of $8 \times 10^4 \text{ N/m}^2$. The density of the gas is 4 kg/m^3 . What is the energy of the gas due to its thermal motion?

A. $5 \times 10^4 \text{ J}$

B. $6 \times 10^4 \text{ J}$

C. $7 \times 10^4 \text{ J}$

D. $3 \times 10^4 \text{ J}$

Answer: A



[Watch Video Solution](#)

198. Statement-1: The temperature dependence of resistance is usually given as $R = R_0(1 + \alpha\Delta T)$. The resistance of a wire changes from $100\Omega \rightarrow 150\Omega$ when its temperature is increased from $27^\circ \text{C} \rightarrow 227^\circ \text{C}$. This implies that $\alpha = 2.5 \times 10^{-3} / ^\circ \text{C}$.

Statement 2: $R = R_0(1 + \alpha\Delta T)$ is valid only when the change in the temperature ΔT is small and $\Delta R = (R - R_0) \ll R_0$.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is not a correct explanation for Statement-1
- C. Statement-1 is False, Statement-2 is True
- D. Statement-1 is True, Statement-2 is False.

Answer: C

 [Watch Video Solution](#)

199. A diatomic ideal gas is used in a Carnot engine as the working substance. If during the adiabatic expansion part of the cycle the volume of the gas increase from V to $32V$, the efficiency of the engine is

- A. 0.5

B. 0.75

C. 0.99`

D. 0.25

Answer: B



Watch Video Solution

200. A thermally insulated vessel contains an ideal gas of molecular mass M and ratio of specific heats γ . It is moving with speed v and it's suddenly brought to rest. Assuming no heat is lost to the surroundings, its temperature increases by:

A. $\frac{\gamma - 1}{2\gamma R} Mv^2 K$

B. $\frac{\gamma Mv^2}{2R} K$

C. $\frac{\gamma - 1}{2R} Mv^2 K$

D. $\frac{\gamma - 1}{2(\gamma + 1)R} Mv^2 K$

Answer: C



Watch Video Solution

201. Three perfect gases at absolute temperature $T_1, T_2,$ and T_3 are mixed. The masses of molecules are n_1, n_2 and n_3 respectively. Assuming to loss of energy, the final temperature of the mixture is:

A.
$$\frac{n_1 T_1 + n_2 T_2 + n_3 T_3}{n_1 + n_2 + n_3}$$

B.
$$\frac{n_1 T_1^2 + n_2 T_2^2 + n_3 T_3^2}{n_1 T_1 + n_2 T_2 + n_3 T_3}$$

C.
$$\frac{n_1^2 T_1^2 + n_2^2 T_2^2 + n_3^2 T_3^2}{n_1 T_1 + n_2 T_2 + n_3 T_3}$$

D. $(T_1 + T_2 + T_3)/3$

Answer: A



Watch Video Solution

202. A Carnot engine operating between temperature T_1 and T_2 has efficiency $1/6$. When T_2 is lowered by 62K its efficiency increase to $1/3$.

Then T_1 and T_2 are, respectively:

A. 372K and 330K

B. 330K and 268K

C. 310K and 248K

D. 372K and 310K

Answer: D



[Watch Video Solution](#)

203. 100g of water is heated from $30^\circ\text{C} \rightarrow 50^\circ\text{C}$. Ignoring the slight expansion of the water, the change in its internal energy is (specific heat of water is $4184\text{J}/\text{kg}/\text{K}$):

A. 8.4kJ

B. $84kJ$

C. $2.1kJ$

D. $4.2kJ$


Answer: A



[Watch Video Solution](#)

204. A wooden wheel of radius R is made of two semicircular part . The two parts are held together by a ring made of a metal strip of cross sectional area S and length L . L is slightly less than $2\pi R$. To fit the ring on the wheel, it is heated so that its temperature rises by ΔT and it just steps over the wheel. As it cools down to surrounding temperature, it process the semicircle parts together. If the coefficient of linear expansion of the metal is α , and it Young's modulus is Y , the force that

one part of the wheel applies on the other part is :



A. $2\pi SY\alpha\Delta T$

B. $SY\alpha\Delta T$

C. $\pi SY\alpha\Delta T$


D. $2SY\alpha\Delta T$

Answer: D



Watch Video Solution

205. Helium gas goes through a cycle ABCDA (consisting of two isochoric and isobaric lines) as shown in figure Efficiency of this cycle is nearly:
(Assume the gas to be close to ideal gas)







- A. 15.4 %
- B. 9.1 %
- C. 10.5 %
- D. 12.5 %

Answer: A



[Watch Video Solution](#)

206. A liquid in a beaker has temperature $\theta(t)$ at time t and θ_0 is temperature of surroundings, then according to Newton's law of cooling the correct graph between $\log_e(\theta - \theta_0)$ and t is :



Answer: A



Watch Video Solution


207. A Carnot engine, whose efficiency is 40%, takes in heat from a source maintained at a temperature of 500K. It is desired to have an engine of efficiency 60%. Then, the intake temperature for the same exhaust (sink) temperature must be:

- A. efficiency of Carnot engine cannot be made larger than 50%
- B. 1200K
- C. 750K
- D. 600K

Answer: C



Watch Video Solution



208.

The above p-v diagram represents the thermodynamic cycle of an engine, operating with an ideal monatomic gas. The amount of heat, extracted from the source in a single cycle is





- A. p_0v_0
- B. $\left(\frac{13}{2}\right)p_0v_0$
- C. $\left(\frac{11}{2}\right)p_0v_0$
- D. $4p_0v_0$

Answer: B



Watch Video Solution

209. If a piece of metal is heated to temperature θ and the allowed to cool in a room which is at temperature θ_0 , the graph between the temperature T of the metal and time t will be closet to



Answer: C



Watch Video Solution

210. The pressure that has to be applied to the ends of a steel wire of length 10cm to keep its length constant when its temperature is raised by $100^\circ C$ is : (For steel Young's modulus is $2 \times 10^{11} Nm^{-2}$ and coefficient of thermal expansion is $1.1 \times 10^{-5} K^{-1}$)

A. $2.2 \times 10^8 Pa$

B. $2.2 \times 10^9 Pa$

C. $2.2 \times 10^7 Pa$

D. $2.2 \times 10^6 Pa$


Answer: A



Watch Video Solution

211. There is a circular tube in a vertical plane. Two liquids which do not mix and of densities d_1 and d_2 are filled in the tube. Each liquid subtends 90° angle at centre. Radius joining their interface make an

angle α with vertical. Ratio $\frac{d_1}{d_2}$ is :



- A. $\frac{1 + \sin \alpha}{1 - \sin \alpha}$
- B. $\frac{1 + \cos \alpha}{1 - \cos \alpha}$
- C. $\frac{1 + \tan \alpha}{1 - \tan \alpha}$
- D. $\frac{1 + \sin \alpha}{1 - \cos \alpha}$

Answer: C

[▶ Watch Video Solution](#)

212. Three rods of Copper, Brass and Steel are welded together to form a Y shaped structure. Area of cross-section of each rod = 4cm^2 . End of copper rod is maintained at 100°C whereas ends of brass and steel are kept at 0°C . Lengths of the copper, brass and steel rods are 46, 13 and 12 cm respectively. The rods are thermally insulated from surroundings excepts at ends. Thermal conductivities of copper, brass and steel are 0.92, 0.26 and 0.12 CGS units respectively. Rate of heat flow through copper rod is :

A. $1.2\text{cal} / \text{s}$

B. $2.4\text{cal} / \text{s}$

C. $4.8\text{cal} / \text{s}$


D. $6.0\text{cal} / \text{s}$

Answer: C



Watch Video Solution

213. One mole of a diatomic ideal gas undergoes a cyclic process ABC as shown in figure. The process BC is adiabatic. The temperature at A,B and C are 400K, 800K and 600K respectively. Choose the correct statement:



- A. The change in internal energy in whole cyclic process is $250R$.
- B. The change in internal process CA is $700R$.
- C. The change in internal energy in the process AB is $-350R$.
- D. The change in internal energy in the process BC is $-500R$.

Answer: D

 [Watch Video Solution](#)

214. A solid body of constant heat capacity $1J/^\circ C$ is being heated by keeping it contact with reservoirs in two ways:

(i) Sequentially keeping in contact with 2 reservoirs such that each reservoir supplies same amount of heat.

(ii) Sequentially keeping in contact with 8 reservoir such that each reservoir supplies same amount of heat.

In both the cases body is brought from initial temperature $100^\circ C$ to final temperature $200^\circ C$. Entropy change of the body in the tow cases respectively is :

A. $\ln 2, 2 \ln 2$

B. $2 \ln 2, 8 \ln 2$

C. $\ln 2, 4 \ln 2$

D. $\ln 2, \ln 2$

Answer: D



Watch Video Solution

215. Consider a spherical shell of radius R at temperature T . The black body radiation inside it can be considered as an ideal gas of photons with internal energy per unit volume $u = \frac{U}{V} \propto T^4$ and pressure $P = \frac{1}{3} \left(\frac{U}{V} \right)$. If the shell now undergoes an adiabatic expansion the relation between T and R is :

A. $T \propto \frac{1}{R}$

B. $T \propto \frac{1}{R^3}$

C. $T \propto e^{-R}$

D. $T \propto e^{-3R}$

Answer: A

 [Watch Video Solution](#)

216. Consider an ideal gas confined in an isolated closed chamber. As the gas undergoes an adiabatic expansion, the average time of collision

between molecules increase as V^q , where V is the volume of the gas. The

value of q is : $\left(\gamma = \frac{C_p}{C_v}\right)$

A. $\frac{\gamma + 1}{2}$

B. $\frac{\gamma - 1}{2}$

C. $\frac{3\gamma + 5}{6}$


D. $\frac{3\gamma - 5}{6}$

Answer: A



Watch Video Solution

217. n moles of an ideal gas undergoes a process $A \rightarrow B$ as shown in the figure. The maximum temperature of the gas during the process will be:



- A. $\frac{9P_0V_0}{2nR}$
- B. $\frac{9P_0V_0}{nR}$
- C. $\frac{9P_0V_0}{4nR}$
- D. $\frac{3P_0V_0}{2nR}$

Answer: C



Watch Video Solution

218. A pendulum clock loses 12s a day if the temperature is $40^\circ C$ and gains 4s a day if the temperature is $20^\circ C$, The temperature at which the

clock will show correct time, and the co-efficient of linear expansion (α) of the metal of the pendulum shaft are respectively:

A. $30^\circ C: \alpha = 1.85 \times 10^{-3} / ^\circ C$

B. $55^\circ C: \alpha = 1.85 \times 10^{-2} / ^\circ C$

C. $25^\circ C: \alpha = 1.85 \times 10^{-5} / ^\circ C$

D. $60^\circ C: \alpha = 1.85 \times 10^{-4} / ^\circ C$

Answer: C

 [Watch Video Solution](#)

219. An ideal gas under goes a quasi static, reversible process in whichh its molar heat capacity C remains constant. If during this process the relation of pressure P and volume V is given by $PV^n = \text{constant}$, then n is given by (Here C_P and C_V are molar specific heat at constant pressur and constant volume, respectively):

A. $n = \frac{C_P - C}{C - C_V}$

$$\text{B. } n = \frac{C - C_V}{C - C_P}$$

$$\text{C. } n = \frac{C_P}{C_V}$$

$$\text{D. } n = \frac{C - C_P}{C - C_V}$$

Answer: D



Watch Video Solution