



PHYSICS

BOOKS - SUNIL BATRA 41 YEARS IITJEE PHYSICS (HINGLISH)

RAY AND WAVE OPTICS

Jee Main And Advanced

1. A light wave of frequency $5 \times 10^{14} Hz$ enters a medium of refractive index 1.5. In the velocity of the light wave is And its wavelength is



Watch Video Solution

2. A convex lens A of focal length $20cm$ and a concave lens B of focal length $5cm$ are kept along the same axis with a distance d between them.

If a parallel beam of light falling on A and B as a parallel beam, then d is equal tocm



[Watch Video Solution](#)

3. A monochromatic beam of light of wavelength 6000\AA in vacuum enters a medium of refractive index 1.5. In the medium its wavelength is...., its frequency is.....



[Watch Video Solution](#)

4. In Young's double-slit experiment, the two slits act as coherent sources of equal amplitude A and of wavelength λ . In another experiment with the same set-up the two slits are sources of equal amplitude A and wavelength λ , but are incoherent. The ratio of the intensity of light at the midpoint of the screen in the first case to that in the second case is....



[Watch Video Solution](#)

5. A thin lens of refractive index 1.5 has focal length of 15cm in air. When the lens is placed in a medium of refractive index $\frac{4}{3}$, its focal length will becomecm.




[Watch Video Solution](#)

6. A point source emits sound equally in all directions in a non-absorbing medium. Two points P and Q are at the distance of 9meters and 25meters respectively from the source. The ratio of amplitudes of the waves at P and Q is.....



[Watch Video Solution](#)



7.

A slab of a material of refractive index 2 shown in fig. has a curved surface APB of radius of curvature 10 cm and a plane surface CD. On the left of APB is air and on the right CD is water with refractive indices as given in the figure. An object O is placed at a distance of 15 cm from the pole P as shown. The distance of the final image of O from P, as viewed from the left is.....



[Watch Video Solution](#)

8. A thin rod of length $\frac{f}{3}$ is placed along the optic axis of a concave mirror of focal length f such that its image which is real and elongated, just

touches the rod. The magnification is



[Watch Video Solution](#)

9. A ray of light undergoes deviation of 30° when incident on an equilateral prism of refractive index $\sqrt{2}$. The angle made by the ray inside the prism with the base of the prism is



[Watch Video Solution](#)

10. The resolving power of electron microscope is higher than that of an optical microscope because the wavelength of electrons is Than the wavelength of visible light.



[Watch Video Solution](#)

11. If ϵ_0 and μ_0 are, respectively, the electric permittivity and magnetic permeability of free space, ϵ and μ the corresponding quantities in a

medium, the index of refraction of the medium in terms of the above parameters is....



Watch Video Solution

12. A light of wavelength 6000\AA in air, enters a medium with refractive index 1.5 Inside the medium its frequency is....Hz and its wavelength is \AA



Watch Video Solution

13. Two thin lenses, when in contact, produce a combination of power $+10$ diopters. When they are 0.25 m apart, the power reduces to $+6$ diopters. The focal length of the lenses are.... m and ... m .



Watch Video Solution

14. A ray of light is incident normally on one of the faces of a prism of apex angle 30° and refractive index $\sqrt{2}$. The angle of deviation of

the ray is...degrees.



Watch Video Solution

15. The setting sun appears higher in the sky than it really is.



Watch Video Solution

16. The intensity of light at a distance r from the axis of a long cylindrical source is inversely proportional to r .



Watch Video Solution

17. A convex lens of focal length 1 meter and a concave lens of focal length 0.25 meter are kept 0.75 meter apart. A parallel beam of light first passes through the convex lens, then through the concave lens and comes to a focus 0.5 m away from the the concave lens.



Watch Video Solution

18. A beam of white light passing through a hollow prism give no spectrum.



Watch Video Solution

19. The two slits in Young's double slit experiment are illuminated by two different sodium lamps emitting light of the same wavelength. No interference pattern will be observed on the screen.



Watch Video Solution

20. In a Young's double slit experiment performed with a source of white light, only black and white fringes are observed.



Watch Video Solution

21. A parallel beam of white light fall on a combination of a concave and a convex lens, both of the same material. Their focal lengths are 15 cm and 30 cm respectively for the mean wavelength in white light. On the other side of the lens system, one sees coloured patterns with violet colour at the outer edge.



[Watch Video Solution](#)

22. When a ray of light enters a glass slab from air,


- A. its wavelength decreases
- B. its wavelength increases
- C. its frequency decreases
- D. neither its wavelength nor its frequency changes

Answer: C



[Watch Video Solution](#)

23. A glass prism of refractive index 1.5 is immersed in water (refractive index $4/3$). A light beam incident normally on the face AB is totally reflected to reach on the face BC if.



A. $\sin \theta \geq \frac{8}{9}$

B. $\frac{2}{3} < \sin \theta < \frac{8}{9}$

C. $\sin \theta \leq \frac{2}{3}$

D. None of these

Answer: A



Watch Video Solution


24. In Young's double-slit experiment, the separation between the slits is halved and the distance between the slits and the screen is doubled. The fringe width is

- A. unchanged
- B. halved
- C. doubled
- D. quadrupled

Answer: D



Watch Video Solution



25.

A ray of light from a denser medium strike a rarer medium at an angle of incidence i (see Fig). The reflected and refracted rays make as angle of 90 degrees with each other. The angles of reflection and refraction are r and r' The critical angle is

A. $\sin^{-1}(\tan r)$

B. $\sin^{-1}(\tan i)$

C. $\sin^{-1}(\tan r)$

D. $\tan^{-1}(\sin i)$

Answer: A



Watch Video Solution

26. Two coherent monochromatic light beams of intensities I and $4I$ are superposed. The maximum and minimum possible intensities in the resulting beam are

- A. $5I$ and I
- B. $5I$ and $3I$
- C. $9I$ and I
- D. $9I$ and $3I$

Answer: C

[Watch Video Solution](#)

27. Spherical aberration in a thin lens can be reduced by

- A. using a monochromatic light
- B. using a doublet combination

C. using a circular annular mark over the lens

D. increasing the size of the lens

Answer: C



Watch Video Solution

28. A beam of light of wave length 600 nm from a distance source fall on a single slit 1mm wide and a resulting. Diffraction pattern is observed on a screen 2m away. The distance between the first dark fringes on either side of central bright fringe is

A. 1.2 cm

B. 1.2 mm


C. 2.4 cm

D. 2.4 mm

Answer: D



Watch Video Solution



29.

An isosceles prism of angle 120° has a refractive index 1.44. Two parallel monochromatic rays enter the prism parallel to each other in air as shown. The rays emerge from the opposite faces

- A. are parallel to each other
- B. are diverging
- C. make an angle $2[\sin^{-1}(0.72) - 30^\circ]$ with each other
- D. make an angle $2 \sin^{-1}(0.72)$ with each other

Answer: C



[Watch Video Solution](#)

30. A diminished image of an object is to be obtained on a screen 1.0 m from it. This can be achieved by appropriately placing

- A. a concave mirror of suitable focal length
- B. a convex mirror of suitable focal length
- C. a convex lens of focal length less than 0.25 m
- D. a concave lens of suitable focal length

Answer: C



[Watch Video Solution](#)

31. The focal length of the objective and the eye piece of a compound microscope are 2.0 cm and 3.0 cm, respectively. The distance between the objective and the eye piece is 15.0 cm. The final image formed by the eye piece is at infinity. The two lenses are thin. The distance in cm of the

object and the image produced by the objective, measured from the objective lens, are respectively

A. 2.4 and 12.0

B. 2.4 and 15.0

C. 2.0 and 12.0

D. 2.0 and 3.0

Answer: A



Watch Video Solution

32. Consider Fraunhofer diffraction pattern obtained with a single slit illuminated at normal incidence. At the angular position of the first diffraction minimum the phase difference (in radians) between the wavelets from the opposite edges of the slit is

A. $\pi/4$

B. $\pi/2$

C. 2π

D. π


Answer: C



Watch Video Solution

33. In an interference arrangement similar to Young's double-slit experiment, the slits S_1 and S_2 are illuminated with coherent microwave sources, each of frequency 10^6 Hz. The sources are synchronized to have zero phase difference. The slits are separated by a distance $d=150.0$ m. The intensity $I(\theta)$ is measured as a function of θ , where θ is defined as shown. If I_0 is the maximum intensity, then $I(\theta)$ for

Intensity at 90° is given by



A. $I(\theta) = \frac{i_0}{2} f$ or $\theta = 30^\circ$

B. $i(\theta) = \frac{I_0}{4} f$ or $\theta = 90^\circ$

C. $I(\theta) = I_0 f$ or $\theta = 0^\circ$

D. $I(\theta)$ is constant for all

Answer: C



Watch Video Solution

34. A concave lens of glass, refractive index 1.5 has both surfaces of same radius of curvature R . On immersion in a medium of refractive index 1.75, it will behave as a

- A. convergent lens of focal length $3.5 R$
- B. convergent lens of focal length $3.0 R$
- C. divergent lens of focal length $3.5 R$
- D. divergent lens of focal length $3.0 R$

Answer: A



Watch Video Solution

35. Yellow light is used in a single slit diffraction experiment with slit width of 0.6 mm . If yellow light is replaced by X-rays, then the observed pattern will reveal,

- A. that the central maximum is narrower

B. more number of fringes

C. less number fringes


D. no diffraction pattern

Answer: D



Watch Video Solution

36. A thin slice is cut out of a glass cylinder along a plane parallel to its axis. The slice is placed on a flat glass plate as shown in Figure. The observed interference fringes from this combination shall be



- A. straight
- B. circular
- C. equally spaced
- D. having fringe spacing which increases as we go outwards

Answer: A



Watch Video Solution

37. A hollow double concave lens is made of very thin transparent material. It can be filled with air or either of two liquids L_1 or L_2 having refractive indices μ_1 and μ_2 respectively ($\mu_2 > \mu_1 > 1$). The lens will diverge a parallel beam of light if it is filled with


- A. air and placed in air
- B. air and immersed in L_1
- C. L_1 and immersed in L_2
- D. L_2 and immersed in L_1

Answer: D



Watch Video Solution

38. A point source of light B is placed at a distance L in front of the centre of a mirror of width d hung vertically on a wall. A man walks in front of the mirror along a line parallel to the mirror at a distance $2L$ from it as shown in fig. The greatest distance over which he can see the image of the light source in the mirror is



A. $\frac{d}{2}$

B. d


C. $2d$

D. $3d$

Answer: D



Watch Video Solution



39.

A diverging beam of light from a point source S having divergence angle α , falls symmetrically on a glass slab as shown. The angles of incidence of the two extreme rays are equal. If the thickness of the glass slab is t and the refractive index n , then the divergence angle of the emergent beam is

A. zero

B. α

C. $\sin^{-1}\left(\frac{1}{n}\right)$

D. $2 \sin^{-1}\left(\frac{1}{n}\right)$


Answer: B



Watch Video Solution

40. A rectangular glass slab ABCD of refractive index n_1 is immersed in water of refractive index n_2 ($n_1 > n_2$). A ray of light is incident at the surface AB of the slab as shown. The maximum value of the angle of

incidence α_{\max} such that the ray comes out only from the other surface



CD is given by

A. $\sin^{-1} \left[\frac{n_1}{n_2} \cos \left(\sin^{-1} \left(\frac{n_2}{n_1} \right) \right) \right]$

B. $\sin^{-1} \left[n_1 \cos \left(\sin^{-1} \left(\frac{1}{n_2} \right) \right) \right]$

C. $\sin^{-1} \left(\frac{n_1}{n_2} \right)$

D. $\sin^{-1} \left(\frac{n_2}{n_1} \right)$

Answer: A



Watch Video Solution

41. In a double slit experiment instead of taking slits of equal widths, one slit is made twice as wide as the other. Then, in the interference pattern

- A. the intensities of both the maxima and the minima increase
- B. the intensity of the maxima increases and the minima has zero intensity
- C. the intensity of the maxima decreases and that of the minima increases
- D. the intensity of the maxima decreases and the minima has zero intensity

Answer: A



Watch Video Solution

42. In a compound microscope, the intermediate image is

- A. virtual, erect and magnified

- B. real, erect and magnified
- C. real, inverted and magnified
- D. virtual, erect and reduced

Answer: C



Watch Video Solution

43. Two beams of light having intensities I and $4I$ interfere to produce a fringe pattern on a screen. The phase difference between the beams is $\frac{\pi}{2}$ at point A and π at point B. Then the difference between the resultant intensities at A and B is

- A. $2I$
- B. $4I$
- C. $5I$
- D. $7I$

Answer: B



Watch Video Solution

44. In a Young's double slit experiment, 12 fringes are observed to be formed in a certain segment of the screen when light of wavelength 600nm is used. If the wavelength of light is changed to 400nm, number of fringes observed in the same segment of the screen is given by

A. 12

B. 18

C. 24


D. 30

Answer: B



Watch Video Solution

45. A ray of light passes through four transparent media with refractive indices μ_1, μ_2, μ_3 and μ_4 as shown in the figure. The surfaces of all media are parallel. If the emergent ray CD is parallel to the incident ray AB, we must have



A. $\mu_1 = \mu_2$

B. $\mu_2 = \mu_3$

C. $\mu_3 = \mu_4$


D. $\mu_4 = \mu_1$

Answer: D



Watch Video Solution


46. A given ray of light suffers minimum deviation in an equilateral prism P. Additional prism Q and R of identical shape and of the same material as P are now added as shown in the figure. The ray will now suffer



- A. greater deviation
- B. no deviation
- C. same deviation as before
- D. total internal reflection

Answer: C

47. An observer can see through a pin-hole the top end of a thin rod of height h , placed as shown in the figure. The beaker height is $3h$ and its radius h . When the beaker is filled with a liquid up to a height $2h$, he can see the lower end of the rod. Then the refractive index of the liquid is



A. $\frac{5}{2}$

B. $\sqrt{\frac{5}{2}}$





C. $\sqrt{\frac{3}{2}}$

D. $\frac{3}{2}$

Answer: B

 Watch Video Solution

48. Which one of the following spherical lenses does not exhibit dispersion? The radii of curvature of the surfaces of the lenses are as given in the diagrams. `



Answer: C



Watch Video Solution

49. In the ideal double-slit experiment, when a glass-plate(refractive index 1.5) of thickness t is introduced in the path of one of the interfering beams (wave-length λ), the intensity at the position where the central maximum occurred previously remains unchanged. The minimum thickness of the glass-plate is

A. 2λ

B. $\frac{2\lambda}{3}$

C. $\frac{\lambda}{3}$


D. λ

Answer: A



Watch Video Solution

50. Two plane mirrors A and B are aligned parallel to each other, as shown in the figure. A light ray is incident at an angle 30° at a point just inside one end of A. The plane of incidence coincides with the plane of the figure. The maximum number of times the ray undergoes reflections (including the first one) before it emerges out is




- A. 28
- B. 30
- C. 32
- D. 34

Answer: B



51. In the adjacent diagram, CP represents a wavefront and AO & BP, the corresponding two rays. Find the condition on θ for constructive interference at P between the ray BP and reflected ray OP.



A. $\cos \theta = \frac{3\lambda}{2d}$

B. $\cos \theta = \frac{\lambda}{4d}$

C. $\sec \theta - \cos \theta = \frac{\lambda}{d}$

D. $\sec \theta - \cos \theta = \frac{\lambda}{d}$

Answer: B



Watch Video Solution

52. The size of the image of an object, which is at infinity, as formed by a convex lens of focal length 30 cm is 2 cm. If a concave lens of focal length 20 cm is placed between the convex lens and the image at a distance of 26 cm from the convex lens, calculate the new size of the image

A. $\frac{d}{2}$

B. d

C. $2d$


D. $\frac{3}{d}$

Answer: B



Watch Video Solution

53. A ray of light is incident at the glass-water interface at an angle i , it emerges finally parallel to the surface of water, the the value of μ_g would be



A. $\left(\frac{4}{3}\right)$

B. $\frac{1}{\sin i}$

C. $\frac{4}{3}$

D. 1

Answer: B



Watch Video Solution

54. A beam of white light is incident on glass air interface from glass to air such that green light just suffers total internal reflection. The colors of the light which will come out to air are


- A. Violet, Indigok, Blue
- B. All colors except green
- C. Yellow, Orange, Red
- D. White light

Answer: C



Watch Video Solution

55. An equilateral prism is placed on a horizontal surface. A ray PQ is incident onto it. For minumum deviation,



- A. PQ is horizontal
- B. QR is horizontal
- C. RS is horizontal
- D. Any one will be horizontal

Answer: B



Watch Video Solution

56. Monochromatic light of wavelength 400nm and 560nm are incident simultaneously and normally on double slits apparatus whose slit separation is 0.1 mm and screen distance is 1m. Distance between areas of total darkness will be

- A. 4 mm
- B. 5.6mm
- C. 14mm
- D. 28mm

Answer: D



Watch Video Solution

57. A source emits sound of frequency 600Hz inside water. The frequency heard in air will be equal to (velocity of sound in water = $1500 \frac{m}{s}$, velocity of sound in air = $300 \frac{m}{s}$)

A. 3000Hz

B. 120Hz

C. 600Hz

D. 6000Hz

Answer: C



Watch Video Solution

58. A point object is placed at the centre of a glass sphere of radius 6cm and refractive index 1.5. The distance of virtual image from the surface is

A. 6cm

B. 4cm

C. 12cm

D. 9cm

Answer: A

59. In Young's double slit experiment intensity at a point is $\left(\frac{1}{4}\right)$ of the maximum intensity. Angular position of this point is

A. $\sin^{-1}\left(\frac{\lambda}{d}\right)$

B. $\sin^{-1}\left(\frac{\lambda}{2d}\right)$

C. $\sin^{-1}\left(\frac{\lambda}{3d}\right)$

D. $\sin^{-1}\left(\frac{\lambda}{4d}\right)$

Answer: C

60. A convex lens is in contact with a concave lens. The magnitude of the ratio of their focal lengths is $\frac{2}{3}$. Their equivalent focal length is 30 cm.

What are their individual focal lengths?

A. $-15, 10$

B. $-10, 15$

C. $75, 50$

D. $-75, 50$


Answer: A



Watch Video Solution

61. A container is filled with water ($\mu = 1.33$) upto a height of 33.25 cm. A concave mirror is placed 15cm above the water level and the image of an object placed at the bottom is formed 25 cm below the water level. Focal

length of the mirror is




- A. 15 cm
- B. 20 cm
- C. -18.31 cm
- D. 10 cm

Answer: C



Watch Video Solution

62. Focal length of the plano-convex lens is 15 cm. A small object is placed at A as shown in the figure. The plane surface is silvered. The image will form at `




- A. 60 cm to the left of lens
- B. 12 cm to the left of lens
- C. 60 cm to the right of lens
- D. 30 cm to the left of lens

Answer: B



Watch Video Solution

63. The graph shown relationship between object distance and image distance for a equiconvex lens. Then focal length of the lens is `



- A. $0.50 \pm 0.05 \text{ cm}$
- B. $0.50 \pm 0.10 \text{ cm}$
- C. $5.00 \pm 0.05 \text{ cm}$
- D. $5.00 \pm 0.10 \text{ cm}$

Answer: C



Watch Video Solution

64. Rays of light from Sun falls on a biconvex lens of focal length f and the circular image of Sun of radius r is formed on the focal plane of the lens. Then

- A. Area of image is πr^2 and area is directly proportional of f
- B. Area of image is πr^2 and area is directly proportional of f^2
- C. Intensity of image increases if f is increased
- D. If lower half of the lens is covered with black paper area will become half

Answer: B



Watch Video Solution

65. In an experiment to determine the focal length (f) of a concave mirror by the $u - v$ method, a student places the object pin A on the principal axis at a distance x from the pole P. The student looks at the pin and its inverted image from a distance keeping his/her eye in line with PA.

When the student shifts his/her eye towards left, the image appears to the right of the object pin. Then,

A. $x < f$

B. $f < x < 2f$

C. $x = 2f$

D. $x > 2f$

Answer: B



Watch Video Solution

66. A ray of light travelling in water is incident on its surface open to air. The angle of incidence is θ , which is less than the critical angle. Then there will be

A. only a reflected ray and no refracted ray

B. only a refracted ray and no reflected ray

C. a reflected ray and a refracted ray and the angle between them

would be less than $180\text{degree} - 2\theta$

D. a reflected ray and a refracted ray and the angle between them

would be greater than $180\text{degree} - 2\theta$

Answer: C



Watch Video Solution

67. Two beams of red and violet colours are made to pass separately through a prism (angle of the prism is 60degree). In the position of minimum deviation, the angle of refraction will be

A. 30degree for both the colours

B. greater for the violet colour

C. greater for the red colour


D. equal but not 30degree for both the colours

Answer: A



Watch Video Solution

68. A light beam is travelling from Region I to IV (figure). The refractive index in regions I, II, III and IV are $n_0 = \frac{n_0}{2}$ and $\frac{n_0}{8}$ respectively. The angle of incidence θ for which the beam just misses entering region IV is –



A. $\sin^{-1}\left(\frac{3}{4}\right)$

B. $\sin^{-1}\left(\frac{1}{8}\right)$

C. $\sin^{-1}\left(\frac{1}{4}\right)$

D. $\sin^{-1}\left(\frac{1}{3}\right)$

Answer: B



Watch Video Solution

69. A ball is dropped from a height of 20 m above the surface of water in a lake. The refractive index of water is 4.3. A fish inside the lake, in the line of fall of the ball, is looking at the ball. At an instant, when the ball is 12.8 m above the water surface, the fish sees the speed of the ball as

$$\left[Take_g = 10 \frac{m}{s^2} . \right]$$

A. 9 m/s

B. 12 m/s

C. 16 m/s

D. 21.33 m/s

Answer: C



Watch Video Solution


70. A biconvex lens of focal length 15 cm is in front of a plane mirror. The distance between the lens and the mirror is 10 cm. A small object is kept at a distance of 30 cm from the lens. The final image is

- A. virtual and at a distance of 16 cm from the mirror
- B. real and at a distance of 16 cm from the mirror
- C. virtual and at a distance of 20 cm from the mirror
- D. real and at a distance of 20 cm from the mirror


Answer: B

[Watch Video Solution](#)


71. A light ray travelling in glass medium is incident of glass- air interface at an angle of incidence θ . The reflected (R) and transmitted (T) intensities, both as function of θ , are plotted The correct sketch is




A.



B.



C.



D.

Answer: C



Watch Video Solution

72. A bi-convex lens is formed with two thin plano-convex lenses as shown in the figure. Refractive index n of the first lens is 1.5 and that of the second lens is 1.2. Both the curved surface are of the same radius of

curvature $R=14$ cm. For this bi-convex lens, for an object distance of 40 cm, the image distance will be `



A. -280.0cm

B. 40.0 cm

C. 21.5 cm

D. 13.3 cm

Answer: B



Watch Video Solution

73. Young's double slit experiment is carried out by using green, red and blue light, one color at a time. The fringe width recorded are b_G, b_R and b_B respectively. Then,

A. $b_G > b_B > b_R$

B. $b_B > b_G > b_R$

C. $b_R > b_B > b_G$

D. $b_R > b_G > b_B$

Answer: D



Watch Video Solution

74. A ray of light travelling in the direction $\frac{1}{2}(\hat{i} + \sqrt{3}\hat{j})$ is incident on a plane mirror. After reflection, it travels along the direction $\frac{1}{2}(\hat{i} - \sqrt{3}\hat{j})$. The angle of incidence is

A. 30degree

B. 60degree

C. 45degree

D. 75degree

Answer: A



Watch Video Solution

75. In the Young's double slit experiment using a monochromatic light of wavelength λ , the path difference (in terms of an integer n) corresponding to any point having half the peak

A. $(2n + 1) \frac{\lambda}{2}$

B. $(2n + 1) \frac{\lambda}{4}$

C. $(2n + 1) \frac{\lambda}{8}$

D. $(2n + 1) \frac{\lambda}{16}$


Answer: B



Watch Video Solution

76. A point source S is placed at the bottom of a transparent block of height 10mm and refractive index 2.72. It is immersed in a lower refractive index liquid as shown in the figure. It is found that the light emerging from the block to the liquid forms a circular bright spot of diameter 11.54

mm on the top of the block. The refractive index of the liquid is `



A. 1.21

B. 1.3

C. 1.36


D. 1.42

Answer: C



Watch Video Solution

77. A parallel beam of light is incident from air at an angle α on the side PQ of a right angled triangular prism of refractive index $n = \sqrt{2}$. Light undergoes total internal reflection at the face PR. The minimum value of α is 45 degree. The $\angle \theta$ of the prism is



- A. 15 degree
- B. 22.5 degree
- C. 30 degree
- D. 45 degree


Answer: A



Watch Video Solution

78. A small object is placed 50 cm to the left of a thin convex lens of focal length 30 cm. A convex spherical mirror of radius of curvature 100 cm is placed to the right of the lens at a distance of 50 cm. The mirror is tilted such that the axis of the mirror is at an angle $\theta = 30^\circ$ to the axis of the lens, as shown in the figure

If the origin of the coordinate system is taken to be at the centre of the lens, the coordinates (in cm) of the point (x, y) at which the image is formed are



A. $(0, 0)$

B. $(50 - 25\sqrt{3}, 25)$

C. $(25, 25\sqrt{3})$

D. $\left(\frac{125}{3}, 25\sqrt{3}\right)$

Answer: C



Watch Video Solution

79. In the Young's double slit experiment, the interference pattern is found to have as intensity ratio between the bright and dark fringes as 9.

This implies that

A. the intensities at the screen due to the two slits are 5 units and 4 units respectively

B. the intensity at the screen due to the two slits are 4 units respectively

C. the amplitude ratio is 3

D. the amplitude ratio is 2

Answer: B::D



Watch Video Solution

80. A convex lens of focal length 40 cm is in contact with a concave lens of focal length 25 cm. The power of the combination is

- A. -1.5 diopters
- B. -6.5 diopters
- C. $+6.5$ diopters
- D. $+6.67$ diopters

Answer: A



Watch Video Solution

81. White light is used to illuminate the two slits in a Young's double slit experiment. The separation between the slits is b and the screen is at a

distance d (gtb) from the slits. At a point on the screen directly in front of one of the slits, certain wavelength are missing. Some of these missing wavelength are

A. $\lambda = \frac{b^2}{d}$

B. $\lambda = \frac{2b^2}{d}$

C. $\lambda = \frac{b^2}{3d}$

D. $\lambda = \frac{2b^2}{3d}$

Answer: A::C



Watch Video Solution

82. A converging lens is used to form an image on a screen. When the upper half of the lens is covered by an opaque screen

A. half the image will disappear.

B. complete image will be formed.

C. intensity of the image will increase

D. intensity of the image will decrease.

Answer: B::D



Watch Video Solution

83. A short linear object of length b lies along the axis of a concave mirror of focal length f at a distance u from the pole of the mirror. The size of the image is approximately equal to

A. $b \left(\frac{u - f}{f} \right)^{\frac{1}{2}}$

B. $b \left(\frac{f}{u - f} \right)^{\frac{1}{2}}$

C. $b \left(\frac{u - f}{f} \right)$


D. $b \left(\frac{f}{u - f} \right)^2$

Answer: D



Watch Video Solution

84. A beam of light consisting of red, green and blue colours is incident on right angled prism. The refractive indices of the material of the prism for the above red, green and blue wavelengths are 1.39, 1.44 and 1.47 respectively. The prism will.



- A. separate part of the red colour from the green and blue colours
- B. separate part of the blue colour from the red and green colours
- C. separate all the three colours from one another
- D. not separate even partially any colour from the other two colours.

Answer: A



Watch Video Solution

85. An astronomical telescope has an angular magnification of magnitude 5 for distant object. The separation between the objective and the eyepiece is 36 cm and the final image is formed at infinity. The focal length f_0 of the objective and the focal length f_e of the eyepiece are

A. $f_0 = 45\text{cm}$ and $f_e = -9\text{cm}$

B. $f_0 = 50\text{cm}$ and $f_e = 10\text{cm}$

C. $f_0 = 7.2\text{cm}$ and $f_e = 5\text{cm}$

D. $f_0 = 30\text{cm}$ and $f_e = 6\text{cm}$.

Answer: D



Watch Video Solution

86. A thin prism P_1 with angle 4degree and made from glass of refractive index 1.54 is combined with another thin prism P_2 made from glass of refractive index 1.72 to produce dispersion without deviation. The angle of the prism P_2 is

A. 5.33°

B. 4°

C. 3°

D. 2.6°

Answer: C



Watch Video Solution

87. A planet is observed by astronomical refracting telescope having an objective of focal length 16cm and an eyepiece of focal length 2cm.

A. The distance between the objective and the eyepiece is 16.02 m

B. The angular magnification of the planet is -800


C. The image of the planet is inverted

D. The objective is larger than the eyepiece

Answer: A::B::C::D

88. Two thin convex lenses of focal lengths f_1 and f_2 are separated by a horizontal distance d (where $d < f_1, d < f_2$) and their centres are displaced by a vertical separation Δ as shown in the fig.

Taking the origin of coordinates O , at the centre of the first lens the x and y coordinates of the focal point of this lens system, for a parallel beam of rays coming from the left, are given by:



A. $x = \frac{f_1 f_2}{f_1 + f_2}, y = \Delta$

B. $x = \frac{f_1(f_2 + d)}{f_1 + f_2 - d}, y = \frac{\Delta}{f_1 + f_2 - d}$

C. $x = \frac{f_1 f_2 + d(f_1 - d)}{f_1 + f_2 - d}, y = \frac{\Delta(f_1 - d)}{f_1 + f_2 - d}$

$$\text{D. } x = \frac{f_1 f_2 + d(f_1 - d)}{f_1 + f_2 - d}, y = 0$$

Answer: C



Watch Video Solution

89. Which of the following form(s) a virtual and erect image for all position of the object?

- A. Convex lens
- B. Concave lens
- C. Convex mirror
- D. Concave mirror

Answer: B::C



Watch Video Solution

90. A real image of a distant object is formed by a plano-convex lens of its principal axis. Spherical aberration

- A. is absent.
- B. is smaller if the curved surface of the lens faces the object.
- C. is smaller if the plane surface of the lens faces the object.
- D. is the same whichever side of the lens faces the object

Answer: B



Watch Video Solution

91. A ray of light travelling in a transparent medium falls on a surface separating the medium from air at an angle of incidence of 45° . The ray undergoes total internal reflection. If n is the refractive index of the medium with respect to air, select the possible value (s) of n from the following:

A. 1.3

B. 1.4

C. 1.5

D. 1.6

Answer: C::D



Watch Video Solution

92. A parallel monochromatic beam of light is incident normally on a narrow slit. A diffraction pattern is formed on a screen placed perpendicular to the direction of the incident beam. At the first minimum of the diffraction pattern, the phase difference between the rays coming from the two edges of the slit is

A. 0

B. $\frac{\pi}{2}$

C. π

D. 2π

Answer: D



Watch Video Solution

93. A concave mirror is placed on a horizontal table, with its axis directed vertically upwards. Let O be the pole of the mirror and C its centre of curvature. A point object is placed at C. It has a real image, also located at C. If the mirror is now filled with water, the image will be.

- A. real, and will remain at C.
- B. real, and located at a point between C and ∞ .
- C. virtual, and located at a point between C and O.
- D. real, and located at a point between C and O

Answer: D



Watch Video Solution

94. A spherical surface of radius of curvature R separates air (refractive index 1.0) from glass (refractive index 1.5). The centre of curvature is in the glass. A point object P placed in air is found to have a real image Q in the glass. The line PQ cuts the surface at a point O , and $PO = OQ$. The distance PO

A. $5R$

B. $3R$

C. $2R$

D. $1.5R$

Answer: A



Watch Video Solution

95. In a Young's double slit experiment, the separation between the two slits is d and the wavelength of the light is λ . The intensity of light falling

on slit 1 is four times the intensity of light falling on slit 2. Choose the correct choice (s).

- A. If $d = \lambda$, The screen will contain only one maximum
- B. If $\lambda < d < 2\lambda$, at least one more maximum (besides the central maximum) will be observed on the screen
- C. If the intensity of light falling on slit 1 is reduced so that it becomes equal to that of slit 2, the intensities of the observed dark and bright fringes will increase
- D. If the intensity of light falling on slit 2 is increased so that it becomes equal to that of slit 1, the intensities of the observed dark and bright fringes will increase

Answer: A::B::C::D



Watch Video Solution

96. A student performed the experiment of determination of focal length of a concave mirror by $u - v$ method using an optical bench of length 1.5 meter. The focal length of the mirror used is 24 cm. The maximum error in the location of the image can be 0.2 cm. The 5 sets of (u, v) values recorded by the student (in cm) are: $(42, 56)$, $(48, 48)$, $(60, 40)$, $(66, 33)$, $(78, 39)$. The data set (s) that cannot come from experiment and is (are) incorrectly recorded, is (are)

A. $(42, 56)$

B. $(48, 48)$

C. $(66, 33)$


D. $(78, 39)$

Answer: C::D



Watch Video Solution

97. A ray OP of monochromatic light is incident on the face AB of prism ABCD near vertex B at an incident angle of 60° (see figure). If the refractive index of the material of the prism is $\sqrt{3}$, which of the following is (are) are correct? `




- A. The ray gets totally internally reflected at face CD
- B. The ray comes out through face AD
- C. The angle between the incident ray and the emergent ray is 90°
- D. The angle between the incident ray and the emergent ray is 120°

Answer: A::B::C::D



Watch Video Solution

98. A transparent thin film of uniform thickness and refractive index $n_1 = 1.4$ is coated on the convex spherical surface of radius R at one end of a long solid glass cylinder of refractive index $n_2 = 1.5$, as shown in the figure. Rays of light parallel to the axis of the cylinder traversing through the film from air to glass get focused at distance f_1 from the film, while rays of light traversing from glass to air get focused at distance f_2 from the film, Then `



A. $|f_1| = 3R$

B. $|f_1| = 2.8R$

C. $|f_2| = 1.4R$

D.

Answer: A::C



Watch Video Solution

99. A light source, which emits two wavelength $\lambda_1 = 400nm$ and $\lambda_2 = 600nm$, is used in a Young's double slit experiment. If recorded fringe width for λ_1 and λ_2 are β_1 and β_2 and the number of fringes for them within a distance y on one side of the central maximum are m_1 and m_2 respectively, then

A. $\beta_2 > \beta_1$

B. $m_1 > m_2$

- C. From the central maximum, 3^{rd} maximum of λ_2 overlaps with 5^{th} minimum of λ_1
- D. The angular separation of fringes for λ_1 is greater than λ_2 .


Answer: A::B::C



Watch Video Solution

100. Two identical glass rods S_1 and S_2 (refractive index=1.5) have one convex end of radius of curvature 10 cm. They are placed with the curved surfaces at a distance d as shown in the figure, with their axes (shown by the dashed line) aligned. When a point source of light P is placed inside rod S_1 on its axis at a distance of 50 cm from the curved face, the light rays emanating from it are found to be parallel to the axis inside S_2 . The

distance d is



A. 60cm

B. 70cm

C. 80 cm

D. 90 cm

Answer: B



Watch Video Solution

101. A plano-convex lens is made of a material of refractive index n . When a small object is placed 30cm away in front of the curved surface of the lens, an image of double the size of the object is produced. Due to reflection from the convex surface of the lens, another faint image is observed at a distance of 10 cm away from the lens. Which of the following statement (s) is (are) true?

- A. The refractive index of the lens 2.5
- B. The radius of curvature of the convex surface is 45 cm
- C. The faint image is erect and real
- D. The focal length of the lens is 20 cm

Answer: A::D




Watch Video Solution

102. A transparent slab of thickness d has a refractive index $n(z)$ that increases with z . Here z is the vertical distance inside the slab, measured

from the top. The slab is placed between two media with uniform refractive indices n_1 and $n_2 (> n_1)$, θ_i , from medium 1 and emerges in medium 2 with refraction angle θ_f with a lateral displacement l .

Which of the following statement(s) is (are) true?




- A. $n_1 \sin \theta_i = n_2 \sin \theta_f$
- B. $n_1 \sin \theta_i = (n_2 - n_1) \sin \theta_f$
- C. l is independent of n_2
- D. l is dependent of $n(z)$

Answer: A::C::D



Watch Video Solution

103. While conducting the Young's double slit experiment, a student replaced the two slits with a large opaque plate in the x - y plane containing two small holes that act as two coherent point sources (S_1, S_2) emitting light of wavelength 600nm . The student mistakenly placed the screen parallel to the x - z plane (f or $z > 0$) at a distance $D=3\text{ m}$ from the mid-point of S_1, S_2 , as shown schematically in the figure. The distance between the sources $d = 0.6003\text{mm}$. The origin O is at the intersection of the screen and the line joining S_1, S_2 . Which of the following is (are) true of the intensity pattern of the screen?



A. Straight bright and dark bands parallel to the x -axis

- B. The region very close to the point O will be dark
- C. Hyperbolic bright and dark bands with foci symmetrically placed about O in the x-direction
- D. Semi circular bright and dark bands centered at point.


Answer: B::D



Watch Video Solution

104. A pin is placed 10cm in front of a convex lens of focal length 20cm, made of a material having refractive index 1.5 . The surface of lens farther away from the pin is silvered and has a radius of curvature 22cm.

Determine the position of the final image. Is the image real or virtual?




[Watch Video Solution](#)

105. A ray of light is incident at an angle of 60° on the face of a prism having refracting angle 30° . The ray emerging out of the prism makes an angle 30° with the incident ray. Show that the emergent ray is perpendicular to the face through which it emerges and calculate the refractive index of the material of prism.





[Watch Video Solution](#)

106. A rectangular block of refractive index μ is placed on a printed page lying on a horizontal surface as shown in Fig. , Find the minimum value of μ so that the letter L on the page is not visible from any of the vertical sides.



Watch Video Solution

107. What is the relation between the refractive indices μ , μ_1 and μ_2 if the behaviour of light rays is shown in Figure.



[▶ Watch Video Solution](#)

108. An object is placed 21 cm in front of a concave mirror of radius of curvature 10cm. A glass slab of thickness 3cm and refractive index 1.5 is then placed close to the mirror in the space between the object and the mirror. The distance of the near surface of the slab from the mirror is 1cm. The final image from the mirror will be formed at

[▶ Watch Video Solution](#)

109. The convex surface of a thin concave-convex lens of glass of refractive index 1.5 has a radius of curvature 20 cm. The concave surface has a


radius of curvature 60 cm. The convex side is silvered and placed on a horizontal surface as shown in figure. (a) Where should a pin be placed on the axis so that its image is formed at the same place ? (b) If the concave part is filled with water ($\mu = 4/3$), find the distance through which the pin should be moved so that the image of the pin again coincides with the pin.



Watch Video Solution

110. Screen S is illuminated by two point sources A and B. Another source C sends a parallel beam of light towards point P on the screen (see figure). Line AP is normal to the screen and the lines AP, BP and CP are in one plane. The distance AP, BP and CP are 3m, 1.5m and 1.5m respectively. The radiant powers of sources A and B are 90 W and 180 W respectively. The beam from C is of intensity $\frac{20\text{ W}}{\text{m}^2}$. Calculate the

intensity at P on the screen.



[Watch Video Solution](#)

111. A plano-convex lens has thickness 4cm. When placed on a horizontal table with the curved surface in contact with it, the apparent depth of the bottom-most point of the lens is found to be 3cm. If the lens is inverted such that the plane face is in contact with the table, the apparent depth of the center of the plane face of the lens is found to be $25/8$ cm. Find the focal length of the lens.



[Watch Video Solution](#)

112. A beam of light consisting of two wavelengths $650nm$ and $520nm$ is used to obtain interference fringes in a Young's double slit experiment.

(a) Find the distance of the third bright fringe on the screen from the central maximum for the wavelength $650nm$.

(b) What is the least distance from the central maximum where the bright fringes due to both the wavelengths coincide? The distance between the slits is $2mm$ and the distance between the plane of the slits and screen is $120cm$.




Watch Video Solution

113. A monochromatic light is incident on the plane interface AB between two media of refractive indices μ_1 and $(\mu_2 > \mu_1)$ at an angle of incidence θ as shown in Fig.


The angle θ is infinitesimally greater than the critical angle for the two media so that total internal reflection takes place. Now, if a transparent slab DEFG of uniform thickness and of refractive index μ_3 is introduced on the interface (as shown in the figure), show that for any value of μ_3 all

light will ultimately be reflected back into medium II.



[Watch Video Solution](#)

114. A right angled prism is to be made by selecting a proper material and the angles A and B ($B \leq A$), as shown in figure. It is desired that a ray of light incident on the face AB emerges parallel to the incident direction after two internal reflections.



(a) What should be the minimum refractive index n for this to be possible?

(b) For $n = \frac{5}{3}$ is it possible to achieve this with the angle B equal to 30 degrees?




Watch Video Solution

115. A parallel beam of light travelling in water (refractive index $= \frac{4}{3}$) is refracted by a spherical bubble of radius 2 mm situated in water. Assuming the light rays to be paraxial. i. find the position of the image due to refraction at the first surface and the position of the final image, and ii draw a ray diagram showing the positions of both the images.



116. In a modified Young's double-slit experiment, a monochromatic uniform and parallel beam of light of wavelength 6000\AA and intensity $(10/\pi) \text{ W m}^{-2}$ is incident normally on two circular apertures A and B of radii 0.001 m and 0.002 m , respectively. A perfectly transparent film of thickness 2000\AA and refractive index 1.5 for the wavelength of 6000\AA is placed in front of aperture A (see the figure). Calculate the power (in mW) received at the focal spot F of the lens. Then lens is symmetrically placed with respect to the aperture. Assume that 10% of the power received by each aperture goes in the original direction and is brought to the focal


spot.



Watch Video Solution

117. A narrow monochromatic beam of light of intensity 1 is incident on a glass plate as shown in figure. Another identical glass plate is kept close to the first one and parallel to it. Each glass plate reflects 25 % of the light incident on it and transmits intensities in the interference pattern

formed by two beams obtained after one reflection at each plate.




Watch Video Solution

118. Two parallel beams of light P and Q (separation d) containing radiation of wavelengths 4000\AA and 5000\AA (which are mutually coherent in each wavelength separately) are incident normally on a prism as shown in fig. The refractive index of the prism as a function of wavelength is given by the relation. $\mu(\lambda) = 1.20 + \frac{b}{\lambda^2}$ Where λ is in \AA and b is positive constant. The value of b is such that the condition wave length

and is not satisfied for the other.

(a) Find the value of b .


(b) find the deviation of the beams transmitted through the face AC. (c) A convergent lens is used to bring these transmitted beams into focus. If the intensities of transmission from the face AC, are 41 and I respectively, find the resultant intensity at the focus. `



Watch Video Solution

119. Light is incident at an angle α on one planar end of a transparent cylindrical rod of refractive index μ . Determine the least value of μ so that the light entering the rod does not emerge from the curved surface of

rod irrespective of the value of α




Watch Video Solution

120. In figure S is a monochromatic point source emitting light of wavelength $\lambda = 500nm$. A thin lens of circular shape and focal length $0.10m$ is cut into two identical halves L_1 and L_2 by a plane passing through a diameter. The two halves are placed symmetrically about the central axis SO with a gap of $0.5mm$. The distance along the axis from A to L_1 and L_2 is $0.15m$, while that from L_1 and L_2 to O is $1.30m$. The screen at O is normal to SO .

(a) If the 3rd intensity maximum occurs at point P on screen, find distance


OP .

(b) If the gap between L_1 and L_2 is reduced from its original value of 0.5 mm , will the distance OP increase, decrease or remain the same?




[Watch Video Solution](#)

121. An image Y is formed of a point object X by a lens whose optic axis is AB as shown in Figure. Draw a ray diagram to locate the lens and its focus. If the image Y of object X is formed by a concave mirror (having the same optic axis AB) instead of lens, draw another ray diagram to locate the mirror and its focus. Write down the steps of construction of the ray diagrams.





122. A ray of light travelling in air is incident at grazing angle (incident angle = 90°) on a long rectangular slab of a transparent medium of thickness $t = 1.0$ (see figure). The point of incidence is the origin $A(O, O)$. The medium has a variable index of refraction $n(y)$ given by : $n(y) = [ky^{3/2} + 1]^{1/2}$, where $k = 1.0 \text{ m}^{-3/2}$. the refractive index of air is 1.0.



- (i) Obtain a relation between the slope of the trajectory of the ray at a point $B(x, y)$ in the medium and the incident angle at that point
- (ii) obtain an equation for the trajectory $y(x)$ of the ray in the medium.

(ii) Determine the coordinates (x_1, y_1) of the point P where the ray intersects upper surface of the slab -air boundary.

Indicate the path of the ray subsequently.




Watch Video Solution

123. A right angles prism $(45^\circ, 90^\circ, 45^\circ)$ of refractive index n has a plate of refractive index $(n_1 < n)$ cemented to its diagonal face. The assembly is in air. A ray is incident on AB.

a. Calculate the angle of incidence at AB for which the ray strikes the diagonal face at the critical angle.

b. Assuming $n = 1.351$, calculate the angle of incidence at AB for which

the refracted ray passes through the diagonal face undeviated.



Watch Video Solution


124. A double slit apparatus is immersed in a liquid of refractive index 1.33. It has slit and the screen 1 mm. The slits are illuminated by a parallel beam of light whose wavelength in air is 6300\AA

- calculate the fringe width.
- One of the slits of the apparatus is covered by a thin glass sheet of refractive index 1.53. Find the smallest thickness of the sheet to bring the adjacent minima on the axis.



Watch Video Solution

125. A thin plano-convex lens of focal length f is split into two halves. One of the halves is shifted along the optical axis as shown in figure. The separation between object and image planes is 1.8 m . The magnification of the image, formed by one of the ball lens is 2 . Find the focal length of the lens and separation between the two halves. Draw the ray diagram for image formation.




Watch Video Solution

126. In Young's experiment the upper slit is covered by a thin glass plate of refractive index 1.4 while the lower slit is covered by another glass plate, having the same thickness as the first one but having refractive


index 1.7 interference pattern is observed using light of wavelength 5400\AA

It is found that point P on the screen where the central maximum ($n = 0$) fell before the glass plates were inserted now has $3/4$ the original intensity. It is further observed that what used to be the fourth maximum earlier, lies below point P while the fifth minimum lies above P. Calculate the thickness of glass plate. (Absorption of light by glass plate may be neglected.)



Watch Video Solution

127. A prism of refractive index n_1 & another prism of reactive index n_2 are stuck together without a gap as shown in the figure. The angle of the prisms are as shown. n_1 & n_2 depend on λ , the wavelength of light according to $n_1 = 1.20 + \frac{10.8 \times 10^4}{\lambda^2}$ & $n_2 = 1.45 + \frac{1.80 \times 10^4}{\lambda^2}$ where λ is in nm.




- (i) Calculate the wavelength λ_0 for which rays incident at any angle on the interface BC pass through without bending at that interface.
- (ii) for light of wavelength λ_0 , find the angle of incidence i on face AC such that the deviation produced by the combination of prism is minimum.



Watch Video Solution


128. A coherent parallel beam of microwaves of wavelength $\lambda = 0.5\text{mm}$ falls on a Young's double-slit apparatus. The separation between the slits is 1.0 mm . The intensity of microwaves is measured on a screen placed parallel to the plane of the slits at a distance of 1.0 m from it as shown in Fig. 2.42.

If the incident beam makes an angle of 30° with the x -axis (as in the dotted arrow shown in the figure), find the y -coordinates of the first minima on either side of the central maximum.



Watch Video Solution

129. The young's double slit experiment is done in a medium of refractive index $4/3$. A light of 600nm wavelength is falling on the slits having 0.45 mm separation. The lower slit S_2 is covered by a thin glass sheet of thickness $10.4\mu\text{m}$ and refractive index 1.5 . The interference pattern is observed on a screen placed 1.5m from the slits as shown



- Find the location of the central maximum (bright fringe with zero path difference) on the y -axis.
- Find the light intensity at point O relative to the maximum fringe intensity.

(c) Now, if 600 nm light is replaced by white light of range 400 to 700 nm find the wavelength of the light that from maxima exactly point O . [All wavelengths in this problem are for the given medium of refractive index $4/3$. Ignore dispersion]




[Watch Video Solution](#)

130. The XY plane is the boundary between two transparent media. Medium 1 with $z \geq 0$ has a refractive index of $\sqrt{2}$ and medium 2 with $z \leq 0$ has a refractive index of $\sqrt{3}$. A ray of light in medium 1 given by the vector $6\sqrt{3}\hat{i} + 8\sqrt{3}\hat{j} - 10\hat{k}$ is incident on the plane of separation. Find the unit vector in the direction of the refracted ray in medium 2.



[Watch Video Solution](#)



131.


A quarter cylinder of radius R and refractive index 1.5 is placed on a table. A point object P is kept at a distance of mR from it. Find the value of m for which a ray from P will emerge parallel to the table as shown in the figure.



Watch Video Solution

132. A convex lens of focal length 15 cm and a concave mirror of focal length 30 cm are kept with their optic axis PQ and RS parallel but separated in vertical direction by 0.6 cm as shown. The distance between the lens and mirror is 30 cm. An upright object AB of height 1.2 cm is placed on the optic axis PQ of the lens at a distance of 20 cm from the lens. If AB' is the image after refraction from the lens and the reflection


from the mirror, find the distance of AB' from the pole of the mirror and obtain its magnification. Also locate positions of A and B' with respect to the optic axis RS .



Watch Video Solution

133. The refractive indices of the crown glass for violet and red lights are 1.51 and 1.49 respectively and those of the flint glass are 1.77 and 1.73 respectively. A prism of angle 6° is made of crown glass. A beam of white light is incident at a small angle on this prism. The other thin flint glass prism is combined with the crown glass prism such that the net mean deviation is 1.5° anticlockwise.

(i) Determine the angle of the flint glass prism.




(ii) A screen is placed normal to the emerging beam at a distance of 2 m from the prism combination. Find the distance between red and violet spot on the screen. Which is the topmost colour on screen.



Watch Video Solution

134. A vessel ABCD of 10 cm width has two small slits S_1 and S_2 sealed with identical glass plates of equal thickness. The distance between the slits is 0.8 mm . POQ is the line perpendicular to the plane AB and passing


through O , the middle point of S_1 and S_2 . A monochromatic light source is kept at S , 40 cm below P and 2 m from the vessel, to illuminate the slits as shown in the figure. Calculate the position of the central bright fringe on the other wall CD with respect of the line OQ . Now, a liquid is poured into the vessel and filled up to OQ . The central bright fringe is found to be at Q . Calculate the refractive index of the liquid.



Watch Video Solution

135. A thin equiconvex lens of refractive index $\frac{3}{2}$ is placed on a horizontal plane mirror as shown in figure. The space between the lens and the mirror is filled with a liquid of refractive index $\frac{4}{3}$. It is found that when a point object is placed 15 cm above the lens on its principal

axis, the object coincides with its own image.



Q. If another liquid is filled instead of water, the object and the image coincide at a distance 25 cm from the lens.


Calculate the refractive index of the liquid.



Watch Video Solution

136. A point source S emitting light of wavelength $600nm$ is placed at a very small height h above the flat reflecting surface AB (see figure). The intensity of the reflected light is 36 % of the intensity. Interference fringes are observed on a screen placed parallel to the reflecting surface a very large distance D from it.

(A) What is the shape of the interference fringes on the screen?




(B) Calculate the ratio of the minimum to the maximum to the maximum intensities in the interference fringes formed near the point P (shown in the figure) (c) if the intensities at point P corresponds to a maximum, calculate the minimum distance through which the reflecting surface AB should be shifted so that the intensity at P again becomes maximum.



Watch Video Solution


137. Find the focal length of the lens shown in Fig . The radii of curvature of both the surfaces are equal to R .



Watch Video Solution

138. A prism ($\mu_P = \sqrt{3}$) has an angle of prism $A = 30^\circ$. A thin film ($\mu_f = 2.2$) is coated on face AC as shown in the figure. Light of wavelength $550nm$ is incident on the face AB at 60° angle of incidence, find

(i) the angle of its emergence from the face AC and




(ii) the minimum thickness (in nm) of the film for which the emerging light is of maximum possible intensity.



[Watch Video Solution](#)

139. Figure shows an irregular block of material of refractive index $\sqrt{2}$. A ray of light strikes the face AB as shown. After refraction, it is incident on a spherical surface CD of radius of curvature 0.4 m and enters a medium

of refractive index 1.514 to meet PQ at E. Find the distance OE up to two places of decimal.



Watch Video Solution

140. In YDSE, two wavelengths of $500nm$ and $700nm$ are used. What is the minimum their maxima coincide ? Take $D/d = 10^3$, symbols have standard meaning.




Watch Video Solution

141. An object is approaching a convex lens of focal length 0.3m with a speed of 0.01ms^{-1} . Find the magnitudes of the ratio of change of position and lateral magnification of image when the object is at a distance of 0.4m from the lens



Watch Video Solution

142. What will be the minimum angle of incidence such that the total internal reflection occurs on both the surfaces?




Watch Video Solution

143. Two identical prism of index $\sqrt{3}$ are kept as shown in the figure. A light ray strikes the first prism at face AB. Find,

(a) the angle of incidence, so that the emergent ray from the first prism has minimum deviation.


(b) through what angle the prism DCE should be rotated about C so that the final emergent ray also has minimum deviation.



Watch Video Solution

144. Fig. shows a surface XY separating two transparent media, medium 1 and medium 2. Lines ab and cd represent wavefronts of a light wave travelling in medium 1 and incident on XY. Line ef and gh represent

wavefront of the light wave in medium 2 after refraction.



Light travel as a


- A. parallel beam in each medium
- B. convergent beam in each medium
- C. divergent beam in each medium
- D. divergent beam in one medium and convergent beam in the other medium.

Answer: A



Watch Video Solution

145. Fig. shows a surface XY separating two transparent media, medium 1 and medium 2. Lines ab and cd represent wavefronts of a light wave travelling in medium 1 and incident on XY. Line ef and gh represent wavefront of the light wave in medium 2 after refraction.



The phase of the light wave at c, d, e, and f are ϕ_c , ϕ_d , ϕ_e and ϕ_f , respectively. It is given that $\phi_c \neq \phi_f$. Then

- A. ϕ_c cannot be equal to ϕ_d
- B. ϕ_d can be equal to ϕ_c
- C. $(\phi_d - \phi_f)$ is equal to $(\phi_c - \phi_e)$


D. $(\phi_d - \phi_c)$ is not equal to $(\phi_c - \phi_e)$

Answer: C



Watch Video Solution

146. Fig. shows a surface XY separating two transparent media, medium 1 and medium 2. Lines ab and cd represent wavefronts of a light wave travelling in medium 1 and incident on XY. Line ef and gh represent wavefront of the light wave in medium 2 after refraction.



Speed of light

is

- A. the same in medium- 1 and medium- 2
- B. larger in medium- 1 than in medium- 2
- C. larger in medium- 2 than in medium- 1
- D. different at b and d.

Answer: B




Watch Video Solution

147. Most materials have the refractive index, $n > 1$. So, when a light ray from air enters a naturally occurring material, then by Snell's law, $\frac{\sin \theta_1}{\sin \theta_2} = \frac{n_1}{n_2}$, it is understood that the refracted ray bends towards the normal. But it never emerges on the same side of the normal as the incident ray. According to electromagnetism, the refractive index of the medium is given by the relation, $n = (c/v) = \pm \sqrt{\epsilon_r, \mu_r}$, where c is the speed of the electromagnetic waves in vacuum, v its speed in the medium, ϵ_r and μ_r are negative, one must choose the negative root of n . Such negative refractive index materials can now be artificially prepared


and are called meta-materials. They exhibit significantly different optical behaviour, without violating any physical laws. Since n is negative, it results in a change in the direction of propagation of the refracted light. However, similar to normal materials, the frequency of light remains unchanged upon refraction even in meta-materials.

Answer the following questions :


For light incident from air on a meta-material, the appropriate ray diagram is




A.



B.



C.



D.

Answer: C



Watch Video Solution

148. Most materials have the refractive index, $n > 1$. So, when a light ray from air enters a naturally occurring material, then by Snell's law, $\frac{\sin \theta_1}{\sin \theta_2} = \frac{n_1}{n_2}$, it is understood that the refracted ray bends towards the normal. But it never emerges on the same side of the normal as the incident ray. According to electromagnetism, the refractive index of the medium is given by the relation, $n = (c/v) = \pm \sqrt{\epsilon_r, \mu_r}$, where c is the speed of the electromagnetic waves in vacuum, v its speed in the medium, ϵ_r and μ_r are negative, one must choose the negative root of n . Such negative refractive index materials can now be artificially prepared and are called meta-materials. They exhibit significantly different optical behaviour, without violating any physical laws. Since n is negative, it results in a change in the direction of propagation of the refracted light. However, similar to normal materials, the frequency of light remains unchanged upon refraction even in meta-materials.

Answer the following questions :

Choose the correct statement.

A. the speed of light in the meta-material is $v = c|n|$

B. the speed of light in the meta-material is $v = \frac{c}{|n|}$

C. the speed of light in the meta-material is $v = c$.

D. The wavelength of the light in the meta-material(λ_m) is given by

$\lambda_m = \lambda_{air}|n|$, where λ_{air} is wavelength of the light in air.

Answer: B




Watch Video Solution

149. Light guidance in an optical fibre can be understood by considering a structure comprising of thin solid glass cylinder of refractive index n_1 surrounded by a medium of lower refractive index n_2 . The light guidance in the structure takes place due to successive total internal reflections at the interface of the media n_1 and n_2 as shown in the figure. All rays with the angle of incidence i less than a particular value i_m are confined in the

medium of refractive index n_1 . The numerical aperture (NA) of the structure is defined as $\sin i_m$

For two structure namely S_1 with $n_1 = \frac{\sqrt{45}}{4}$ and $n_2 = \frac{3}{2}$, and S_2 with $n_1 = \frac{8}{5}$ and $n_2 = \frac{7}{5}$ and taking the refractive index of water to be $\frac{4}{3}$ and that of air to be 1, the correct option (s) is (are) ?



A. NA of S_1 immersed in water is the same as that of S_2 immersed in a

liquid of refractive index $\frac{16}{3\sqrt{15}}$

B. NA of S_1 immersed in liquid of refractive index $\frac{6}{\sqrt{15}}$ is the same as

the of S_2 immersed in water

C. NA of S_1 placed in air is the same as that of S_2 immersed in liquid

of refractive index $\frac{4}{\sqrt{15}}$

D. NA of S_1 placed in air is the same as that of S_2 placed in water

Answer: A::C




Watch Video Solution

150. Light guidance in an optical fibre can be understood by considering a structure comprising of thin solid glass cylinder of refractive index n_1 surrounded by a medium of lower refractive index n_2 . The light guidance in the structure takes place due to successive total internal reflections at the interface of the media n_1 and n_2 as shown in the figure. All rays with the angle of incidence i less than a particular value i_m are confined in the medium of refractive index n_1 . The numerical aperture (NA) of the structure is defined as $\sin i_m$

If two structure of same cross-sectional area, but different numerical apertures NA_1 and NA_2 ($NA_2 < NA_1$) are joined longitudinally, the

numerical aperture of the combined structure is `



A. $\frac{NA_1 NA_2}{NA_1 NA_2}$

B. $NA_1 + NA_2$

C. NA_1

D. NA_2

Answer: D



Watch Video Solution

151. In each of the questions, assertion(A) is given by corresponding statement of reason (R) of the statemens. Mark the correct answer.

Q. Statement I: The formula connecting u, v and f for a spherical mirror is valid only for mirrors whose sizes are very small compared to their radii of curvature.

Statement II: Laws of reflection are strictly valid for plane surfaces, but not for large spherical surfaces.

- A. Statement- 1 is true, Statement- 2 is True, Statement- is a correct explanation for Statement- 1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement- 1
- C. Statement- 1 is True, Statement- 2 is False
- D. Statement- 1 is False, Statement- 2 is True.

Answer: C



Watch Video Solution

152. The focal length of a thin biconvex lens is 20cm . When an object is moved from a distance of 25cm in front of it to 50cm , the magnification of its image changes from $m_{25} \rightarrow m_{50}$. The ratio $\frac{m_{25}}{m_{50}}$ is.



Watch Video Solution

153. A large glass slab ($\mu = 5/3$) of thickness 8cm is placed over a point source of light on a plane surface. It is seen that light emerges out of the top surface of the slab from a circular area of radius R cm. What is the value of R ?




Watch Video Solution

154. Image of an object approaching a convex mirror of radius of curvature 20m along its optical axis is observed to move from $\frac{25}{3}\text{m}$ to $\frac{50}{7}\text{m}$ in 30 seconds. What is the speed of the object in km per hour?



Watch Video Solution

155. Water (with refractive index $= 4/3$) in a tank is 18cm deep. Oil of refraction index $7/4$ lies on water making a convex surface of radius of curvature $R = 6\text{cm}$ as shown in Fig. Consider oil to act as a thin lens. An object S is placed 24cm above water surface. The location of its image is at $x\text{cm}$ above the bottom of the tank. Then x is.



Watch Video Solution


156. A Young's double slit interference arrangement with slits S_1 and S_2 is immersed in water (refractive index $= 4/3$) as shown in the figure. The positions of maxima on the surface of water are given by $x^2 = p^2 m^2 \lambda^2 - d^2$, where λ is the wavelength of light in air (refractive index $= 1$), $2d$ is the separation between the slits and m is an integer. The value of P is



Watch Video Solution

157. Consider a concave mirror and a convex lens (refractive index 1.5) of focal length 10cm each separated by a distance of 50cm in air (refractive index $= 1$) as shown in the Fig. An object is placed at a distance of 15cm from the mirror. Its erect image formed by this combination has magnification M_1 . When this set up is kept in a medium of refractive

index $7/6$, the magnification becomes M_2 . The magnitude $\left(\frac{M_2}{M_1}\right)$ is :



[▶ Watch Video Solution](#)

158. A monochromatic beam of light is incident at 60° on one face of an equilateral prism of refractive index n and emerges from the opposite face making an angle θ with the normal. For $n = \sqrt{3}$, the value of θ is 60° and $\frac{d\theta}{dn} = m$. The value of m is.

[▶ Watch Video Solution](#)

159. An astronomical telescope has large aperture to

- A. reduce spherical aberration
- B. have high resolution
- C. increase span of observation
- D. have low dispersion

Answer: B



Watch Video Solution

160. If two mirrors are kept at 60° to each other, then the number of images formed by them is

- A. 5
- B. 6
- C. 7
- D. 8

Answer: A



Watch Video Solution

161. Electrimagnetic waves are transverse is nature is evident by

- A. polarization
- B. interference
- C. reflection
- D. differaction

Answer: A



Watch Video Solution

162. Wavelength of ligh used in an optical instrument are $\lambda = 4000\text{\AA}$ and

$\lambda_2 = 5000\text{\AA}$

, then ratio of their respective resolvable powers (or respond $\in g \rightarrow$

λ_1 and λ_2)

A. 16: 25

B. 9: 1

C. 4: 5

D. 5: 4

Answer: D



Watch Video Solution

163. Which of the following is used in optical fibres?

A. total internal reflection

B. scattering

C. diffraction

D. refracton

Answer: A



Watch Video Solution

164. Consider telecommunication through optical fibres. Which of the following statements is not true?

- A. Optical fibres can be of graded refractive index
- B. Optical fibres are subject to electromagnetic interference from outside
- C. Optical fibres have extremely low transmission loss
- D. Optical fibre may have homogeneous core with a suitable cladding

Answer: B



Watch Video Solution

165. To demonstrate the phenomenon of interference, we require two sources which emit radiation

- A. of nearly the same frequency

B. of the same frequency

C. of different wavelengths

D. of the same frequency and having a definite phase relationship

Answer: D



Watch Video Solution

166. The image formed by an objective of a compound microscope is

A. virtual and diminished

B. real and diminished

C. real and enlarged

D. virtual and enlarged

Answer: C



Watch Video Solution

167. To get three images of a single object, one should have two plane mirrors at an angle of

A. 60°

B. 90°

C. 120°

D. 30°


Answer: B



Watch Video Solution

168. A light ray is incident perpendicularly to one face of a 90° prism and is totally internally reflected at the glass-air interface. If the angle of

reflection is 45° , we conclude that the refractive index n



A. $n > \frac{1}{\sqrt{2}}$

B. $n > \sqrt{2}$

C. $n < \frac{1}{\sqrt{2}}$

D. $n < \sqrt{2}$

Answer: B



Watch Video Solution

169. A plano convex lens of refractive index 1.5 and radius of curvature 30cm. Is silvered at the curved surface. Now this lens has been used to form the image of an object. At what distance from this lens an object be placed in order to have a real image of size of the object.

A. 60 cm

B. 30 cm

C. 20 cm

D. 80 cm

Answer: C



Watch Video Solution

170. The angle of incidence at which reflected light is totally polarized for reflection from air to glass (refractive index n),

A. $\tan^{-1}\left(\frac{1}{n}\right)$

B. $\sin^{-1}\left(\frac{1}{n}\right)$

C. $\sin^{-1}(n)$

D. $\tan^{-1}(n)$

Answer: D



Watch Video Solution

171. The maximum number of possible interference maxima for slit-separation equal to twice the wavelength in Young's double-slit experiment is

A. three

B. five

C. infinite

D. zero

Answer: B

[Watch Video Solution](#)

172. An electromagnetic wave of frequency $\nu = 3.0\text{MHz}$ passes from vacuum into a dielectric medium with permittivity $\epsilon = 4.0$. Then

- A. wave length is halved and frequency remains unchanged
- B. wave length is doubled and frequency becomes half
- C. wave length is doubled and the frequency remains unchanged
- D. wave length and frequency both remain unchanged.

Answer: A

[Watch Video Solution](#)

173. A fish looking up through the water sees the outside world contained in a circular horizon. If the refractive index of water is $\frac{4}{3}$ and the fish is 12 cm below the surface, the radius of this circle is cm is

A. $\frac{36}{\sqrt{7}}$

B. $36\sqrt{7}$

C. $4\sqrt{5}$

D. $36\sqrt{5}$

Answer: A



Watch Video Solution

174. Two point white dots are 1mm apart on a black paper. They are viewed by eye of pupil diameter 3mm. Approximately, what is the maximum distance at which these dots can be resolved by the eye? [Take wavelength of light = 500nm]

A. 1m

B. 5m

C. 3m

D. 6m

Answer: B



Watch Video Solution

175. A thin glass (refractive index 1.5) lens has optical power of $-5D$ in air.

Its optical power in a liquid medium with refractive index 1.6 will be

A. $-1D$

B. $1D$

C. $-25D$

D. $25D$

Answer: B



Watch Video Solution

176. A Young's double slit experiment uses a monochromatic source. The shape of the interference fringes formed on a screen is

- A. circle
- B. hyperbola
- C. parabola
- D. straight line

Answer: D



Watch Video Solution

177. If I_0 is the intensity of the principal maximum in the single slit diffraction pattern. Then what will be its intensity when the slit width is doubled?

- A. $4I_0$
- B. $2I_0$
- C. $\frac{I_0}{2}$
- D. I_0

Answer: A



Watch Video Solution

178. When an unpolarized light of intensity I_0 is incident on a polarizing sheet, the intensity of the light which does not get transmitted is

A. $\frac{1}{4}I_0$

B. $\frac{1}{2}I_0$

C. I_0

D. zero

Answer: B



Watch Video Solution

179. The refractive index of a glass is 1.520 for red light and 1.525 for blue light. Let D_1 and D_2 be angles of minimum deviation for red and blue

light respectively in a prism of this glass. Then,

A. $D_1 < D_2$

B. $D_1 = D_2$

C. D_1 can be less than or greater than D_2 depending upon the angle of prism

D. $D_1 > D_2$

Answer: A



Watch Video Solution

180. In a Young's double slit experiment the intensity at a point where the path difference is $\frac{\lambda}{6}$ (λ being the wavelength of light used) is I . If I_0 denotes the maximum intensity, $\frac{I}{I_0}$ is equal to

A. $\frac{3}{4}$

B. $\frac{1}{\sqrt{2}}$

C. $\frac{\sqrt{3}}{2}$

D. $\frac{1}{2}$

Answer: A



Watch Video Solution

181. Two lenses of power $-15D$ and $+5D$ are in contact with each other.

The focal length of the combination is

A. $+10cm$

B. $-20cm$

C. $-10cm$


D. $+20cm$

Answer: C






Watch Video Solution

182. In an experiment, electrons are made to pass through a narrow slit of width d comparable to their de Broglie wavelength. They are detected on a screen at a distance D from the slit (see figure).




Which of the following graphs can be expected to represent the number of electrons N detected as a function of the detector position y ($y=0$ corresponds to the middle of the slit).





C.




D.

Answer: D




Watch Video Solution

183. A student measures the focal length of a convex lens by putting an object pin at a distance u from the lens and measuring the distance v of the image pin. The graph between u and v plotted by the student should look like




A.

(b)




B.

(c)



C.

(d)



D.

Answer: C



Watch Video Solution

184. An experiment is performed to find the refractive index of glass using a travelling microscope. In this experiment distances are measured by

- A. a vernier scale provided on the microscope
- B. a standard laboratory scale
- C. a meter scale provided on the microscope

D. a screw gauge procided on the microscope

Answer: A



Watch Video Solution

185. A micture of light, consisting of wavelength 590nm and an unknown wavelength, illuminates Young's double slit and gives rise to two overlapping interference patterns on the scree. The central maximum of both lights coincide. Further, it is obseved that the third bright fringe of known light coincides with the 4th bright fringe of the unknown light. From this data, the wavelength of the unknown light is:

A. 885.5 nm

B. 442.5 nm


C. 776.8nm

D. 393.4 nm

Answer: B



186. A transparent solid cylindrical rod has a refractive index of $\frac{2}{\sqrt{3}}$. It is surrounded by air. A light ray is incident at the mid-point of one end of the rod as shown in the figure. The incident angle θ for which the light ray grazes along the wall of the rod is:



A. $\sin^{-1}\left(\frac{\sqrt{3}}{2}\right)$

B. $\sin^{-1}\left(\frac{2}{\sqrt{3}}\right)$

C. $\sin^{-1}\left(\frac{1}{\sqrt{3}}\right)$

D. $\sin^{-1}\left(\frac{1}{2}\right)$

Answer: C



Watch Video Solution

187. In an optics experiment, with the position of the object fixed, a student varies the position of a convex lens and for each position, the screen is adjusted to get a clear image of the object. A graph between the object distance u and the image distance v , from the lens, is plotted using the same scale for the two axes. A straight line passing through the origin and making an angle of 45° with x-axis meets the experimental curve at P. The coordinates of P will be.

A. $\left(\frac{f}{2}, \frac{f}{2}\right)$

B. (f, f)

C. $(4f, 4f)$

D. $(2f, 2f)$

Answer: D



Watch Video Solution

188. An initially parallel cylindrical beam travels in a medium of refractive index $\mu(I) = \mu_0 + \mu_2 I$, where μ_0 and μ_2 are positive constants and I is the intensity of the light beam. The intensity of the beam is decreasing with increasing radius.

30 . At the beam enters the medium, it will

A. diverge

B. converge

C. diverge near the axis and converge near the periphery

D. travel as a cylindrical beam

Answer: B



Watch Video Solution

189. An initially parallel cylindrical beam travels in a medium of refractive index $\mu(I) = \mu_0 + \mu_2 I$, where μ_0 and μ_2 are positive constants and I is the intensity of the light beam. The intensity of the beam is decreasing with increasing radius.

31. The initial shape of the wavefront of the beam is

- A. convex
- B. concave
- C. convex near the axis and concave near the priphery
- D. planar

Answer: D



Watch Video Solution

190. An initially parallel cylindrical beam travels in a medium of refractive index $\mu(I) = \mu_0 + \mu_2 I$, where μ_0 and μ_2 are positive constants and I is the intensity of the light beam. The intensity of the beam is decreasing

with increasing radius.

32. The speed of light in the medium is

- A. minimum on the axis of the beam
- B. the same everywhere in the beam
- C. directly proportional to the intensity I
- D. maximum on the axis of the beam

Answer: A



Watch Video Solution

191. Let the x - z plane be the boundary between two transparent media.

Medium 1 in $z \geq 0$ has a refractive index of $\sqrt{2}$ and medium 2 with $z < 0$

has a refractive index of $\sqrt{3}$. A ray of light in medium 1 given by the vector

$\vec{A} = 6\sqrt{3}\hat{i} + 8\sqrt{3}\hat{j} - 10\hat{k}$ is incident on the plane of separation. The

angle of refraction in medium 2 is:

- A. 45°

B. 60°

C. 74°

D. 30°

Answer: A



Watch Video Solution

192. This question has a paragraph followed by two statements, Statement - 1 and Statement - 2. Of the given four alternatives after the statements, choose the one that describes the statements. A thin air film is formed by putting the convex surface of a plane-convex lens over a plane glass plate. With monochromatic light, this film gives an interference pattern due to light reflected from the top (convex) surface and the bottom (glass plate) surface of the film.

Statement - 1: When light reflects from the air-glass plate interface, the reflected wave suffers a phase change of π .

Statement - 2 : The centre of the interference pattern is dark.

- A. Statement - 1 is true, Statement - 2 is true, Statement - 2 is the correct explanation of Statement -1.
- B. Statement - 1 is true, Statement - 2 is true, Statement - 2 is not the correct explanation of Statement -1.
- C. Statement - 1 is false, Statement -2 is true.
- D. Statement -1 is true, Statement -2 is false.

Answer: B



Watch Video Solution

193. A car is fitted with a convex side-view mirror of focal length 20 cm. A second car 2.8m behind the first car is overtaking the first car at a relative speed of $15 \frac{m}{s}$. The speed of the image of the second car as seen in the mirror of the first one is:

A. $\frac{1}{15} \frac{m}{s}$

B. $10 \frac{m}{s}$

C. $15 \frac{m}{s}$

D. $\frac{1}{10} \frac{m}{s}$

Answer: A



Watch Video Solution

194. An electromagnetic wave in vacuum has the electric and magnetic field \vec{E} and \vec{B} , which are always perpendicular to each other. The direction of polarization is given by \vec{X} and that of wave propagation by \vec{K} . Then

A. $\vec{X} \parallel \vec{B}$ and $\vec{k} \parallel \vec{B} \times \vec{E}$

B. $\vec{X} \parallel \vec{E}$ and $\vec{k} \parallel \vec{E} \times \vec{B}$

C. $\vec{X} \parallel \vec{B}$ and $\vec{k} \parallel \vec{E} \times \vec{B}$

D. $\vec{X} \parallel \vec{E}$ and $\vec{k} \parallel \vec{B} \times \vec{E}$

Answer: B



195. In Young's double slit experiment, one of the slit is wider than other, so that amplitude of the light from one slit is double of that from other slit. If I_m be the maximum intensity, the resultant intensity I when they interfere at phase difference ϕ is given by:

A. $\frac{I_m}{9}(4 + 5 \cos \phi)$

B. $\frac{I_m}{3}\left(1 + 2 \cos^2\left(\frac{\phi}{2}\right)\right)$

C. $\frac{I_m}{5}\left(1 + 4 \cos^2\left(\frac{\phi}{2}\right)\right)$

D. $\frac{I_m}{9}\left(1 + 8 \cos^2\left(\frac{\phi}{2}\right)\right)$

Answer: D

[Watch Video Solution](#)

196. An object 2.4 m in front of a lens forms a sharp image on a film 12 cm behind the lens. A glass plate 1 cm thick, of refractive index 1.50 is

interposed between lens and film with its plane faces parallel to film. At what distance (from lens) should object shifted to be in sharp focus of film?

A. 7.2 m

B. 2.4 m

C. 3.2 m

D. 5.6 m

Answer: D



Watch Video Solution

197. The diameter of a plano convex lens is 6 cm and thickness at the centre is 3 mm . If the speed of light in the material of the lens is $2 \times 10^8\text{ m/s}$, what is the focal length of the lens ?

A. 15 cm

B. 20 cm

C. 30 cm

D. 10 cm

Answer: C



Watch Video Solution

198. A beam of unpolarised light of intensity I_0 is passed through a polaroid A and then through another polaroid B which is oriented so that its principal plane makes an angle of 45° relative to that of A. The intensity of the emergent light is

A. I_0

B. $\frac{I_0}{2}$

C. $\frac{I_0}{4}$


D. $\frac{I_0}{8}$

Answer: C



Watch Video Solution

199. Two coherent point sources S_1 and S_2 are separated by a small distance d as shown. The fringes obtained on the screen will be




- A. points
- B. straight lines
- C. semi-circles
- D. concentric circles



Answer: D



200. The graph between angle of deviation (δ) and angle of incidence (i) for a triangular prism is represented by



B. 



Answer: C



Watch Video Solution

201. A thin convex lens made from crown glass $\left(\mu = \frac{3}{2}\right)$ has focal length f . When it is measured in two different liquids having refractive indices $\frac{4}{3}$ and $\frac{5}{3}$, it has the focal lengths f_1 and f_2 respectively. The correct relation between the focal lengths is ,

- A. $f_1 = f_2 < f$
- B. $f_1 > f$ and f_2 becomes negative
- C. $f_2 > f$ and f_1 becomes negative
- D. f_1 and f_2 both become negative

Answer: B



Watch Video Solution

202. A green light is incident from the water to the air - water interface at the critical angle (θ) . Select the correct statement.

- A. The entire spectrum of visible light will come out of the water at an angle of 90° to the normal.
- B. The spectrum of visible light whose frequency is less than that of green light will come out to the air medium.
- C. The spectrum of visible light whose frequency is more than that of green light will come out to the air medium.
- D. The entire spectrum of visible light will come out of the water at various angles to the normal

Answer: B



Watch Video Solution

203. Two beams A and B, of plane polarized light with mutually perpendicular planes of polarization are seen through a polaroid. From the position when the beam a has maximum intensity (and beam B has zero ntensity), a rotation of polaroid through 30° makes the two beams

appear equally bright. If the initial intensities of the two beams are I_A and I_B respectively, then $\frac{I_A}{I_B}$ equals:

A. 3

B. $\frac{3}{2}$

C. 1

D. $\frac{1}{3}$

Answer: D



Watch Video Solution

204. Assuming human pupil to have a radius of 0.25 cm and a comfortable viewing distance of 25 cm, the minimum separation between two objects than human eye can resolve at 500nm wavelength is :

A. $100\mu m$

B. $300\mu m$

C. $1\mu m$

D. $30\mu m$

Answer: D



Watch Video Solution

205. On a hot summer night, the refractive index of air is smallest near the ground and increases with height from the ground. When a light beam is directed horizontally, the Huygens' principle leads us to conclude that as it travels, the light beam:


- A. bends downwards
- B. bends upwards
- C. becomes narrower
- D. goes horizontally without any deflection

Answer: B



Watch Video Solution

206. Monochromatic light is incident on a glass prism of angle A . If the refractive index of the material of the prism is μ , a ray, incident at an angle θ , on the face AB would get transmitted through the face AC of the prism provided:



- A. $\theta > \cos^{-1} \left[\mu \sin \left(A + \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$
- B. $\theta < \cos^{-1} \left[\mu \sin \left(A + \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$
- C. $\theta > \sin^{-1} \left[\mu \sin \left(A - \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$
- D. $\theta < \sin^{-1} \left[\mu \sin \left(A - \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$

Answer: C



Watch Video Solution

207. The box of a pin hole camera, of length L , has a hole of radius a . It is assumed that when the hole is illuminated by a parallel beam of light of wavelength λ the spread of the spot (obtained on the opposite wall of the camera) is the sum of its geometrical spread and the spread due to diffraction. The spot would then have its minimum size (say b_{\min}) when:

A. $a = \sqrt{\lambda L}$ and $b_{\min} = \sqrt{4\lambda L}$

B. $a = \frac{\lambda^2}{L}$ and $b_{\min} = \sqrt{4\lambda L}$

C. $a = \frac{\lambda^2}{L}$ and $b_{\min} = \left(\frac{2\lambda^2}{L} \right)$

D. $a = \sqrt{\lambda L}$ and $b_{\min} = \left(\frac{2\lambda}{L} \right)$

Answer: A



Watch Video Solution

208. An observer looks at a distant tree of height 10m with a telescope of magnifying power of 20. to the observer the tree appears:

- A. 20 times taller
- B. 20 times nearer
- C. 10 times taller
- D. 10 times nearer

Answer: B



Watch Video Solution

209. In an experiment for determination of refractive index of glass of a prism by $i - \delta$, plot it was found that a ray incident at angle 35° , suffers a deviation of 40° and that it emerges at angle 79° . In that case which of the following is closest to the maximum possible value of the refractive index?

A. 1.7

B. 1.8

C. 1.5

D. 1.6

Answer: C



Watch Video Solution