



PHYSICS

BOOKS - CENGAGE PHYSICS (HINGLISH)

LINEAR AND ANGULAR SIMPLE HARMONIC MOTION

Illustration

1. What is the time - period of $x = A \sin(\omega t + \alpha)$?



Watch Video Solution

2. Find time period of the function,

$$y = \sin \omega t + \sin 2\omega t + \sin 3\omega t$$



[Watch Video Solution](#)

3. Identify which of the following function represent simple harmonic motion.

(i) $Y = Ae^{I\omega t}$ (ii) $Y = ae^{-\omega t}$

(iii) $y = a \sin^2 \omega t$ (iv) $y = a \sin \omega t + b \cos \omega t$

(v) $y = \sin \omega t + b \cos 2\omega t$



[Watch Video Solution](#)

4. A particle executes SHM with an angular frequency $\omega = 4\pi \text{ rad/s}$. If it is at its extreme position initially, then find the time when it is at a distance $\frac{\sqrt{2}}{2}$ times its amplitude from the mean position.



[Watch Video Solution](#)

5. A particle executes SHM with an amplitude 8 cm and a frequency 4 cm initially, in the positive direction, determine its displacement equation and the maximum velocity and acceleration.



[Watch Video Solution](#)

6. A particle executing SHM oscillates between two fixed points separated by 20cm . If its maximum velocity be 30cm/s . Find its velocity when its displacement is 5cm from its mean position.






[Watch Video Solution](#)

7. A particle executing SHM with time period of 2s : Find the time taken by it to move from one amplitude to half the amplitude position.



[Watch Video Solution](#)

8. Write the equation of SHM for the situations show below:



 Watch Video Solution

9. A partical of $mass\ m = 1\text{kg}$ oscillates simple harmonically with angular frequency 1rad/s . Find the

phase of the particle at $t = 1s$ and $2s$. Start calculating time when the particle moves up passing through the mean position.


 [Watch Video Solution](#)

10. If $x = A \sin(\omega t + \alpha)$, Find phase constant α in $x = A \sin(\omega t + \alpha)$, a particle executing SHM is going along

 [Watch Video Solution](#)

11. Shows the displacement time graph of a particle executing SHM with a time period T . Four points 1, 2, 3 and 4 are marked on the graph where the displacement is

half that of the amplitude.



- a. Identify the point of same displacement but with opposite direction of motion . Find the time difference between them.
- b. Identify the point where the particles move in the same difference. Find the time difference between them.



Watch Video Solution

12. A particle executes SHM with same frequency and amplitude along the same straight line. They cross each other, at a point midway between the mean and the extreme position. Find the Phase difference between them.



[Watch Video Solution](#)

13. A particle executes SHM with amplitude A and angular frequency ω . At an instant the particle is at a distance $A/5$ from the mean position and is moving away from it. Find the time after which it will come back to this position again and also find the time after which it will pass through the mean position.



Watch Video Solution

14. two particle executing SHM with same frequency and amplitudes A and $2A$ on same straight line with same mean position cross each other in opposite direction at a distance $A/3$ from mean position. Find the phase difference in the two SHM s.



Watch Video Solution

15. A particle of mass 0.50 kg executes a simple harmonic motion under a force $F = - (50Nm^{-1})x$. If it crosses the centre of oscillation with a speed of $10ms^{-1}$, find the amplitude of the motion.



Watch Video Solution



Watch Video Solution

16. A $20g$ partical is oscillating simple barmonically with a period of 2 second and maximum kinetic energy $2J$. The total machanical energy of the partical is zero , find
- Amplitude of oscillation
 - potential energy as a punction of displacement x relative to mean position.



Watch Video Solution

17. A body is executing SHM under action of the a force of whose maximum is $50N$. magnitude of force acting on the particle at the time when its energy is half kinetic

energy and half potential is (Assume potential energy to be zero at mean position).

 [Watch Video Solution](#)

18. A point particle of mass 0.1kg is executing SHM of amplitude 0.1m . When the particle passes through the mean position, its kinetic energy is $8 \times 10^{-3}\text{J}$. Write down the equation of motion of this particle when the initial phase of oscillation is 45° .

 [Watch Video Solution](#)

19. If a particle moves in a potential energy field $U = U_0 - ax + bx^2$, where a and b are particle constants

obtain an expression for the force acting on it as a function of position. At what point does the force vanish?

Is this a point of stable equilibrium ?

Calculate the force constant and frequency of the particle.

 [Watch Video Solution](#)

20. A particle executes SHM with an amplitude of 10cm and frequency 2Hz , at $t = 0$, the particle is at point where potential energy and kinetic energy are same. Find the equation of displacement of particle.

 [Watch Video Solution](#)

21. A particle of mass 0.2kg undergoes SHM according to the equation $x(t) = 3 \sin(\pi t + \pi/4)$.

i. What is the total energy of the particle if potential energy of zero at mean position ?

ii. What are the kinetic and potential energies of particle at time $t = 1\text{s}$?

iii. At what time instants is the particles energies purely kinetic?



Watch Video Solution

22. A particle of mass 0.2kg executes simple harmonic motion along a path of length 0.2m at the rate of 600 oscillations per minute. Assume at $t = 0$. The particle

start SHM in positive direction. Find the kinetic potential energies in joules when the displacement is $x = A/2$ where, A stands for the amplitude.

 [Watch Video Solution](#)

23. A point particle of mass 0.1kg is executing SHM of amplitude 0.1m . When the particle passes through the mean position, its kinetic energy is $8 \times 10^{-3}\text{J}$. Write down the equation of motion of this particle when the initial phase of oscillation is 45° .

 [Watch Video Solution](#)

24. A particle of mass m is located in a unidimensional potential field where potential energy of the particle depends on the coordinates x as: $U(x) = U_0(1 - \cos Ax)$, U_0 and A constants.

Find the period of small oscillation that the particle performs about the equilibrium position.

 [Watch Video Solution](#)

25. Find the amplitude of the simple harmonic motion obtained by combining the motions

$$x_1 = (2.0\text{cm})\sin \omega t$$

$$\text{and } x_2 = (2.0\text{cm})\sin\left(\omega t + \frac{\pi}{3}\right)$$

 [Watch Video Solution](#)

26. $x_1 = 3 \sin \omega t \Rightarrow x_2 = 4 \cos \omega t$. Find (i) amplitude of resultant SHm, (ii) equation of the resultant SHm.

 [Watch Video Solution](#)

27. Two particles A and B execute simple harmonic motion according to the equation $y_1 = 3 \sin \omega t$ and $y_2 = 4 \sin[\omega t + (\pi/2)] + 3 \sin \omega t$. Find the phase difference between them.

 [Watch Video Solution](#)

28. If the displacement of a moving point at any time is given by an equation of the form $y(t) = a \cos \omega t + b \sin \omega t$, shown that the motion is simple harmonic . If $a = 3m$, $b = 4m$ and $\omega = 2$: determine the period , amplitude, maximum velocity and maximum acceleration.

 [Watch Video Solution](#)

29. If two SHMs are represented by $y_1 = 10 \sin(4\pi t + \pi/2)$ and $y_2 = 5(\sin 2\pi t + \sqrt{8} \cos 2\pi t)$, compare their amplitudes .

 [Watch Video Solution](#)

30. A force $F = -10x + 2$ acts on a particle of mass 0.1kg where x is in m and F in newton. If is released from rest at $x = 0$, find:

a. Amplitude:

b. Time period:

c. Equation of motion.



Watch Video Solution

31. A person normally weighing 60kg stands on a platform which oscillates up and down harmonically at a frequency 2.0sec^{-1} and an amplitude 5.0cm . If a machine on the platform gives the person's weight

against time deduce the maximum and minimum reading


it will shown, $T_{akeg} = 10m / \text{sec}^2$.




[Watch Video Solution](#)

32. A spring of stiffness constant k and natural length l is cut into two parts of length $3l/4$ and $l/4$, respectively, and an arrangement is made as shown in figure . If the mass is slightly displaced , find the time period of

oscillation.



(a)





(b)

[▶ Watch Video Solution](#)


33. A particle of mass m is attached with three springs A , B and C of equal force constants k as shown in figure. The particle is pushed slightly against the spring C

and released. Find the time period of oscillation.



 Watch Video Solution


34. Two light spring of force constants k_1 and k_2 and a block of mass m are in the line AB on a smooth horizontal table such that one end of each spring is fixed on right supports and the other end is free as shown in figure



The distance CD between the free ends of the spring is 60cm . If the block ($k_1 = 1.8\text{N/m}$, $k_2 = 3.2\text{N/m}$ and $m = 200\text{g}$). Is the motion simple harmonic?

 Watch Video Solution

35. figure shown a partical mass $m = 100g$ attaches with four identical spring , each of length $l = 10cm$. Inital tension in each spring is $F_0 = 25N$. Neglecting gravity , Calculate the period of small obscillation of the article along a line perpenducular to the plane of the figure .



[▶ Watch Video Solution](#)

36. Find the time period of m if pulley P is light and small.





Fig. 4.43


 Watch Video Solution

37. A solid cylinder of mass m is attached to a horizontal spring with force constant k . The cylinder can roll without slipping along the horizontal plane. (See the accompanying figure.) Show that the center of mass of the cylinder executes simple harmonic motion with a period $T = 2\pi\sqrt{\frac{3m}{2k}}$, if displaced from mean position.



Watch Video Solution

38. figure (a) and (b) represent spring- block system. If m is displacement slightly , find the time period of oscillation of the system.



[▶ Watch Video Solution](#)

39. Two identical balls A and B each of mass 0.1kg are attached to two identical mass less is springs. The spring-mass system is constrained to move inside a right smooth pipe bent in the form of a circle as shown in

figure . The pipe is fixed in a horizontal plane. The center of the balls can move in a circle of radius $0.06m$. Each spring has a natural length of $0.06\pi m$ and force constant $0.1N/m$. Initially, both the balls are displaced by an angle $\theta = \pi/6$ radians with respect to diameter PQ of the circle and released from rest.

- a. Calculate the frequency of oscillation of the ball B.
- b. What is the total energy of the system ?
- c. Find the speed of the ball A when A and B are at the two

ends of the diameter PQ.





Fig. 1.26



Watch Video Solution

40. The system shown in the figure can move on a smooth surface. They are initially compressed by 6cm and then released, then choose the correct options.

$$k = 800 \text{ N/m}$$



- (a) The system performs, SHM with time period $\frac{\pi}{10} \text{ s}$
- (b) The block of mass 3 kg perform SHM with amplitude 4 cm
- (c) The block of mass 6 kg will have maximum momentum of $2.40 \text{ kg} - \text{m} / \text{s}$
- (d) The time periods of two blocks are in the ratio of $1 : \sqrt{2}$

 [Watch Video Solution](#)

41. A simple pendulum of length 40 cm oscillates with an angular amplitude of 0.04 rad . Find a. the time period b.

the linear amplitude of the bob, c. The speed of the bob when the string makes 0.02 rad with the vertical and d. the angular acceleration when the bob is in momentary rest.


Take $g = 10\text{ms}^{-2}$.



[Watch Video Solution](#)

42. A ball is suspended by a thread of length l at the point O on an incline wall as shown. The inclination of the wall with the vertical is θ . (a) the thread is displaced through a small angle away from the vertical and (b) the ball is released. Find the period of oscillation of pendulum.

Consider $\theta < 90^\circ$ for both cases



a. $\alpha > \beta$

b. $\alpha < \beta$

Assuming that any impact between the wall and the ball is elastic.

 Watch Video Solution

43. figure show the identical simple pendulums of length l . One is tilted at an angle α and imparted an initial velocity v_1 towards mean position and at a velocity v_2 at an initial angular displacement β . Find the phase difference in oscillations of these two pendulums





Fig 4



Watch Video Solution

44. In previous question, if the second pendulum bob is thrown at velocity v_2 at an angle β from mean position, but on other side of mean position, find the phase difference in the two SHMs now as show in the figure.



[Watch Video Solution](#)

45. Derive an expression for the angular frequency of small oscillation of the bob of a simple pendulum when it

is immersed in a liquid of density ρ . Assume the density of the bob as σ and length of the string as l .

 [Watch Video Solution](#)

46. What is the period of a pendulum formed by pivoting a meter stick so that is free to rotate about a horizontal axis passing through the 75cm mark?

 [Watch Video Solution](#)

47. A uniform disc of radius 5.0cm and mass 200g is fixed at its centre to a metal wire, the other end of which is fixed to a ceiling. The hanging disc is rotated about the wire through an angle and is released. If the disc makes torsional

oscillations with time period 0.20s , find the torsional constant of the wire.




Fig. 4.72



Watch Video Solution

48. Assume that a tunnel is dug across the earth (radius= R) passing through its centre. Find the time a particle takes to cover the length of the tunnel if (a) it is projected into the tunnel with a speed of $\sqrt{(gR)}$ (b) it is released from a height R above the tunnel (c) it is thrown

vertically upward along the length of tunnel with a speed of \sqrt{gR} .

[Watch Video Solution](#)

49. Consider a solid cylinder of the density ρ_s cross section area A and h floating in a liquid of density ρ_l as shown in figure ($\rho_l > \rho_s$). It is depressed slightly and allowed to oscillation.




Fig. 4.76

[Watch Video Solution](#)

50. A V-shaped glass tube of uniform cross section is kept in a vertical plane as shown. A liquid is poured in the tube. In equilibrium the level of liquid in both limbs of tube are equal. Find the angular frequency of small oscillation of liquid.




Fig. 4.77

- A. $\pi \sqrt{\frac{2h}{g \sin \alpha \cdot \sin \beta}}$
- B. $2\pi \sqrt{\frac{h}{g \sin \alpha \cdot \sin \beta}}$
- C. $2\pi \sqrt{\frac{h}{2g \sin \alpha \cdot \sin \beta}}$

D. None of these

Answer: B

 [Watch Video Solution](#)

51. Find the amplitude and initial phase of a particle in SHM, whose motion equation is given as

$$y = A \sin \omega t + B \cos \omega t$$

 [Watch Video Solution](#)

52. Two simple harmonic motions are represented by the following equations

$$y_1 = 10 \sin(\pi/4)(12t + 1)$$

$$y_2 = 5(\sin 3\theta t + \sqrt{3} \cos 3\theta t)$$

Here t is in seconds.

Find out the ratio of their amplitudes. What are the time period of the two motion?

 [Watch Video Solution](#)

Solved Example

1. A uniform horizontal plank is resting symmetrically in a horizontal position on two cylindrical drums, which are spinning in opposite directions about their horizontal axes with equal angular velocity. The distance between the axes is $2L$ and the coefficient of friction


between the plank and cylinder is μ . If the plank is displaced slightly from the equilibrium position along its length and released, show that it performs simple horizontal motion. Calculate also the time period of motion.



[Watch Video Solution](#)

2. A uniform plank of mass m , free to move in the horizontal direction only, is placed at the top of a solid cylinder of mass $2m$ and radius R . The plank is attached to a fixed wall by means of a light spring constant k . There is no slipping between the cylinder and the plank, and between the cylinder and the ground. Find the time


period of small oscillation of the system.



[Watch Video Solution](#)


3. A block of mass m hangs by means of a string which goes over a pulley of mass m and moment of inertia I , as shown in the diagram. The string does not relative to the

pulley. Find the frequency of small oscillations.



[View Text Solution](#)

4. The pulley shown in figure has a moment of inertias I about its axis and mass m . find the time period of vertical oscillation of its centre of mass. The spring has spring constant k and the string does not slip over the pulley.




 Watch Video Solution

5. An L-shaped bar of mass M is pivoted at one of its end so that it can freely rotate in a vertical plane, as shown in the figure

a. Find the value of θ_0 at equilibrium

b. If it is slightly displacement from its equilibrium position, find the frequency of oscillation.



[Watch Video Solution](#)

6. A certain of a perfect gas is enclosed in a cylinder of volume V_0 fitted with a smooth heavy piston of mass m and area a . The piston is displaced through a small distance downwards so as to compress the gas isothermally, and find its period. Take the atmospheric pressure as $P_{a \rightarrow m}$.




[Watch Video Solution](#)

7. A spherical ball of mass m and radius r rolls without slipping on a rough concave surface of large radius R . It makes small oscillations about the lowest point. Find the time period.



[Watch Video Solution](#)

8. A simple pendulum of length L and mass m has a spring of force constant k connected to it at a distance h below its point of suspension. Find the frequency of vibrations of the system for small values of amplitude.



 Watch Video Solution

9. One end of an ideal spring is fixed to a wall at origin O and axis of spring is parallel to x -axis. A block of mass $m = 1\text{kg}$ is attached to free end of the spring and it is performing SHM. Equation of position of the block in coordinate system shown in figure is $x = 10 + 3\sin(10t)$. Here, t is in second and x in cm . Another block of mass $M = 3\text{kg}$, moving towards the origin with velocity 30cm/s collides with the block performing SHM at $t = 0$ and gets stuck to it. Calculate



- (a) new amplitude of oscillations,
- (b) new equation for position of the combined body,
- (c) loss of energy during collision. Neglect friction.



[Watch Video Solution](#)

10. A circular spring of natural length l_0 is cut and weided with two beads of masses m_1 and m_2 each that the such that the ratio of the original spring is k find the frequency of oscillation of the heads in a smooth horizontal rigid tube. Assume $m_{(1)} = m$ and $m_{(2)} = 3m$.





Fig. 4.110



Watch Video Solution

11. A uniform cylinder of mass m and radius R is in equilibrium on an inclined by the action of a light spring of stiffness, gravity and reaction force acting on it .If the angle of inclination of the plane is ϕ , find the angular frequency of small oscillation of the cylinder



[Watch Video Solution](#)

12. A 2kg mass is attached to a spring of force constant $600\text{N}/\text{m}$ and rests on a smooth horizontal surface. A second mass of 1kg slides along the surface toward the first at $6\text{m}/\text{s}$.

(a) Find the amplitude of oscillation if the masses make a perfectly inelastic collision and remain together on the spring. what is the period of oscillation ?


(b) Find the amplitude and period of oscillation if the collision is perfectly elastic.

(c) For each case , write down the position x as a function of time t for the mass attached to the spring, assuming that the collision occurs at time $t = 0$. What is the impulse given to the 2kg mass in each case ?




[Watch Video Solution](#)

13. A spring block pendulum is shown in figure . The system is hanging in equilibrium. A bullet of mass $m/2$ moving at a speed u bites the block from downwards direction and gets embedded in it. Find the amplitude of oscillation of the block now. Also find the time taken by the block to reach its upper extreme position after hit by bullet.



14. Figure shown a spring block system hanging in equilibrium. The block of system is pulled down by the distance x and imparted a velocity v in downward direction as shown in figure. Find the time it will take to reach its mean position. Also find the maximum distance to which it will move before returning back towards mean


position.



[Watch Video Solution](#)

15. Figure shown a block P of mass m resting on a smooth horizontal surface, attached to a spring of force constant k which is rigidly fixed on the wall on left side, shown in the figure . At a distance l to the right of the block there is a rigid wall. If block is pushed towards left so that spring

is compressed by a distance $5l/3$ and when released, it will start its oscillations. If collision of block with the wall is considered to be perfectly elastic. Find the time period of oscillation of the block.




[▶ Watch Video Solution](#)

Exercise 4.1

1. i. The acceleration versus time graph of a particle in SHM is shown in the figure. Plot the displacement versus time

graph



ii. The frequency of oscillation is.....

iii. The displacement amplitude is

iv. At $t = 0$, the velocity of the particle is

v. The kinetic energy of the particle is maximum at $t = \dots\dots$

and $t = \dots\dots$

vi. The potential energy is maximum at $t = \dots\dots$: $t = \dots\dots$ and t

=

 Watch Video Solution

2. A particle slides back and forth between two inclined frictionless planes.




Fig. 4.28

a. If h is the initial height of the particle, the period of oscillation.....

b. Is the motion oscillatory? Is it SHM?



Watch Video Solution

3. The equation of displacement of two waves are given as

$$y_1 = 10 \sin\left(3\pi t + \frac{\pi}{3}\right), y_2 = 5[\sin 3\pi t + \sqrt{3} \cos 3\pi t]$$

Then what is the ratio of their amplitudes

 [Watch Video Solution](#)

4. Suppose a tunnel is dug along a diameter of the earth.

A particle is dropped from a point a distance h directly above the tunnel. The motion of the particle as seen from the earth is

 [Watch Video Solution](#)


5. The equation of motion of a particle started at $t = 0$ is given by $x = 5 \sin(20t + \pi/3)$ where x is in centimeter and t in second. When does particle

- (a) first come to rest
- (b) first have zero acceleration
- (c) first have maximum speed?



[Watch Video Solution](#)

6. A particle starts SHM from mean position O executing SHM A and B are the two points at which its velocity is zero. It passes through a certain point O at time $t_1 = 0.5$ and $t_2 = 1.5s$ with a speed of $3m/s$.



i. The maximum speed

ii. ratio AP / PB

[Watch Video Solution](#)

7. If the maximum speed and acceleration of a particle executing SHM is 20cm/s and 100cm/s^2 , find the time period of oscillation.

[Watch Video Solution](#)

8. A particle is performing SHM of amplitude 'A' and time period 't'. Find the time taken by the particle to go from $0 \rightarrow A/2$.



[Watch Video Solution](#)

9. A particle of mass 2kg is moving of a straight line under the action force $F = (8 - 2x)\text{N}$. It is released at rest from $x = 6\text{m}$.

- Is the particle moving simple harmonically.
- Find the equilibrium position of the particle.
- Write the equation of motion of the particle.
- Find the time period of SHM.



[Watch Video Solution](#)

10. A particle executing simple harmonic motion has amplitude of 1m and time period 2s . At $t = 0$, net force on the particle is zero. Find the equation of displacement of the particle.



[Watch Video Solution](#)

11. In the previous question, find maximum velocity and maximum acceleration.



[Watch Video Solution](#)

12. A particle in SHM has a period of 4s. It takes time t_1 to start from mean position and reach half the amplitude. In another case it takes a time t_2 to start from mean position and reach half the amplitude. Find the ratio t_1/t_2



[Watch Video Solution](#)

13. A particle is subjected to two simple harmonic motions in the same direction having equal amplitudes and equal frequency. If the resultant amplitude is equal to the amplitude of the individual motions. Find the phase difference between the individual motions.



[Watch Video Solution](#)

14. A particle executes SHM of period $1.2s$ and amplitude $8cm$. Find the time it takes to travel $3cm$ from the positive extremity of its oscillation. $[\cos^{-1}(5/8) = 0.9rad]$



Watch Video Solution


15. A cylinder of mass M and radius R is resting on a horizontal platform (which is parallel to the x - y plane) with its axis fixed along the y -axis and free to rotate about its axis. The platform is given a motion in the x -direction given by $x = A \cos(\omega t)$. There is no slipping between the cylinder and platform. The maximum torque acting on the cylinder during its motion is



Watch Video Solution

 Watch Video Solution

16. The figure shows the displacement-time graph of a particle executing *SHM*. If the time period of oscillation is $2s$, then the equation of motion is given by



 Watch Video Solution

17. A body executing SHM has its velocity $10\text{cm}/\text{sec}$ and $7\text{cm}/\text{sec}$ when its displacement from

the mean position are 3cm and 4cm , respectively.

Calculate the length of the path.


 [Watch Video Solution](#)

18. A body undergoing *SHM* about the origin has its equation is given by $X = 0.2 \cos 5\pi t$. Find its average speed from $t = 0 \rightarrow t = 0.7\text{sec}$.

 [Watch Video Solution](#)

19. The acceleration-displacement ($a - X$) graph of a particle executing simple harmonic motion is shown in

the figure. Find the frequency of oscillation.



[▶ Watch Video Solution](#)

20. A block is kept on a horizontal table. The table is undergoing simple harmonic motion of frequency 3 Hz in a horizontal plane. The coefficient of static friction between block and the table surface is 0.72 . Find the

maximum amplitude of the table at which the block does not slip on the surface.



[Watch Video Solution](#)

21. A linear harmonic oscillator has a total mechanical energy of $200J$. Potential energy of it at mean position is $50J$. Find

(i) the maximum kinetic energy,

(ii) the minimum potential energy,

(iii) the potential energy at extreme positions.



[Watch Video Solution](#)

22. The potential energy of a particle oscillating along x -axis is given as

$$U = 20 + (x - 2)^2$$

Here, U is in joules and x in meters. Total mechanical energy of the particle is $36J$.

(a) State whether the motion of the particle is simple harmonic or not.

(b) Find the mean position.

(c) Find the maximum kinetic energy of the particle.



[Watch Video Solution](#)

23. $x_1 = 3 \sin \omega t$

$$x_2 = 5 \sin(\omega t + 53^\circ)$$

$$x_3 = -10 \cos \omega t$$

Find amplitude of resultant SHM.


 [Watch Video Solution](#)

Exercise 4.2

1. A block of mass m is suspended from the ceiling of a stationary standing elevator through a spring of spring constant k . Suddenly, the cable breaks and the elevator starts falling freely. Show that the block now executes a simple harmonic motion of amplitude $m \frac{g}{k}$ in the elevator

 [Watch Video Solution](#)


2. The left block in figure collides inelastically with the right block and sticks to it. Find the amplitude of the resulting simple harmonic motion.



[Watch Video Solution](#)

3. A ball of a mass m is connected to two rubber bands of length L , each under tension T as shown in figure . The ball is displaced by a small distance y perpendicular to the length of the rubber band. Assuming the tension does not change , shown that (a) the restoring force is

$-(2T/L)y$ and (b) the system exhibits simple harmonic motion with an angular frequency $\omega = \sqrt{2T/mLs}$.



[Watch Video Solution](#)

4. A mass M attached to a spring oscillation with a period of $2s$. If the mass is increased by $2kg$, the period increases by $1s$, find the initial mass m assuming that Hooke's law is obeyed.

[Watch Video Solution](#)




Fig. 14.11

5.


A horizontal rod of mass m and length L is pivoted at one end. The rod's other end is supported by a spring of force constant k . The rod is displaced by a small angle θ from its horizontal equilibrium position and released. The angular frequency of the subsequent simple harmonic motion is



[Watch Video Solution](#)

6. A pendulum has a period T for small oscillations. An obstacle is placed directly beneath the pivot, so that only the lowest one - quarter of the string can follow the pendulum bob when it swings to the left of its resting position. The pendulum is released from rest at a certain point. How long will it take to return to that point again ? In answering this question, you may assume that the angle between the moving string and the vertical stays

small throughout the motion.



[Watch Video Solution](#)

7. A horizontal spring block system of mass M executes simple harmonic motion. When the block is passing

through its equilibrium position, an object of mass m is put on it and the two move together. Find the new amplitude and frequency of vibration. Given, k is the spring constant of the system.




[Watch Video Solution](#)

8. A spring of spring constant $200N/m$ has a block of mass $1kg$ hanging at its one end and from the other end the spring is attached to a ceiling of an elevator. The elevator rises upwards with an acceleration of $g/3$.

When acceleration is suddenly ceased, then what should be the angular frequency and elongation during the time


when the elevator is accelerating?



[Watch Video Solution](#)

9. With the assumption of no slipping, determine the mass m of the block which must be placed on the top of a 6 kg cart in order that the system period is 0.75 s . What is

the minimum coefficient of static friction μ_s for which the block will not slip relative to the cart is displaced 50mm from the equilibrium position and released? Take $(g = 9.8\text{m/s}^2)$.




[▶ Watch Video Solution](#)

10. A simple pendulum of length l swings from a small angle θ . Its swinging is constrained by the smooth inclined planes OP and PC . Assuming elastic collision of the bob with the plane PC , find


a. angular amplitude for the motion of the bob in the left hand side of its mean position.

b. time period for a complete cycle of motion of the bob.



[Watch Video Solution](#)

11. A uniform rod of length l is pivoted distance x from the top of the rod. Neglecting friction find the (a) value of x for minimum period of oscillation, (b) minimum period of oscillation of the rod.





[Watch Video Solution](#)

12. The period of oscillation of a spring pendulum is T . If the spring is cut into four equal parts, then find the time period corresponding to each part.



[Watch Video Solution](#)

13. A uniform stick of length l is hinged so as to rotate about a horizontal axis perpendicular to the stick, at a distance from the center x . Find the value of x , for which the time period is minimum.



[Watch Video Solution](#)

14. A ball is released in a smooth dimetrical tunnel of earth

a. After how much time will it pass through the center of earth?

b. With what speed will the ball pass the center earth?



[Watch Video Solution](#)

15. A body is in *SHM* with period T when oscillated from a freely suspended spring. If this spring is cut in two parts of length ratio 1:3 & again oscillated from the two the two parts separatedly, then the periods are T_1 & T_2 then find T_1/T_2 .



[Watch Video Solution](#)


16. A point mass m is suspended at the end of a massless wire of length l and cross section. If Y is the Young's modulus for the wire, obtain the frequency of oscillation for the simple harmonic motion along the vertical line.



Watch Video Solution

17. In the figure shown, the block A of mass m collides with the identical block B and after collision they stick


together. Calculate the amplitude of resultant vibration.



[▶ Watch Video Solution](#)

18. Figure shown a block P of mass m resting on a smooth floor at a distance l from a rigid wall. Block is pushed towards right by a distance $3/2$ and released. When block passes from its mean position another block of mass m_1


so that the combined block just collides with the left wall.



[Watch Video Solution](#)

19. Figure shown a block P of mass m resting on a smooth horizontal ground, attached to one end of a spring of force constant k in its natural length. If another block of same mass and moving with a velocity u towards right is placed on the block and sticks to it due to friction, find the time it will take to reach its extreme position. Also find the


amplitude of oscillation of the combined mass $2m$.



[▶ Watch Video Solution](#)

20. Figure shown a spring block system hanging in equilibrium. If a velocity v_0 is imparted to the block in downwards direction . Find the amplitude of SHM of the block and the time after which it will reach a point at half

the amplitude of block



[Watch Video Solution](#)

21. Find the amplitude of the harmonic motion obtained by combining the motions

$$x_1 = (2.0\text{cm})\sin \omega t$$

and $x_2 = (2.0\text{cm})\sin(\omega t + \pi/3)$.



[Watch Video Solution](#)

22. $x_1 = 3 \sin \omega t, x_2 = 4 \cos \omega t$

Find (i) amplitude of resultant SHM, (ii) equation of the resultant SHM.



[Watch Video Solution](#)

23. A particle is subjected to two simple harmonic motions

$$x_1 = A_1 \sin \omega t$$

and $x_2 = A_2 \sin(\omega t + \pi/3)$

Find (a) the displacement at $t = 0$, (b) the maximum speed of the particle and (c) the maximum acceleration of the particle.




[Watch Video Solution](#)

Subjective

1. A rigid rod of mass m with a ball of mass M attached to the free end is restrained to oscillate in a vertical plane as shown in the figure. Find the natural frequency of

oscillation.



[Watch Video Solution](#)

2. A rectangular tank having base $15\text{cm} \times 20\text{cm}$ is filled with water (density $\rho = 1000\text{kg}/\text{m}^3$) up to 20cm and force constant $K = 280\text{N}/\text{m}$ is fixed to the bottom of the tank so that the spring remains vertical.

This system is in an elevator moving downwards with

acceleration $a = 2m/s^2$. A cubical block of side $l = 10\text{cm}$ and mass $m = 2\text{kg}$ is gently placed over the spring and released gradually, as shown in the figure.




Fig. 4.124

- Calculate compression of the spring in equilibrium position.
- If the block is slightly pushed down from equilibrium position and released, calculate frequency of its vertical oscillations.

 [Watch Video Solution](#)

3. A body A mass $m_1 = 1\text{kg}$ and body B of mass $m_2 = 4.1\text{kg}$. The body A perform free vertical harmonic oscillations with the amplitude 1.6cm and frequency 25Hz . Neglecting the mass of the spring , find the maximum and minimum value of force that the system exerts on the bearing surface.





Fig. 4.125



Watch Video Solution


4. In the arrangement shown in figure the sleeve M of mass $m = 0.20\text{kg}$ is fixed between two identical springs whose combined stiffness is equal to $x = 20\text{N}/\text{m}$. The sleeve can slide without friction over a horizontal bar AB . The arrangement rotates with a constant angular velocity $\omega = 4.4\text{rad}/\text{s}$ about a vertical axis passing through the middle of the bar. Find the period of small oscillations of the sleeve. At what values of ω will there be no oscillations of the sleeve?






Watch Video Solution

5. A vertical pole of length l , density ρ , area of cross section A floats in two immiscible liquids of densities ρ_1 and ρ_2 . In equilibrium position the bottom end is at the interface of the liquids. When the cylinder is displaced vertically, find the time period of oscillation.



Watch Video Solution

6. In the shown arrangement, both the spring are in their natural lengths. The coefficient of friction between m_2 and m_1 is μ . There is no friction between m_1 and the surface. If the blocks are displaced slightly, they together perform simple harmonic motion. Obtain



- Frequency of such oscillations.
- The condition if the friction force on block m_2 is to act in the direction of its displacement from mean position.
- If the condition obtained in (b) is met, what can be maximum of their oscillations ?



7. A uniform disc of mass m and radius R is connected with two light springs 1 and 2. The springs are connected at the highest point M and the CM 'N' of the disc. The other ends of the springs are rigidly attached with vertical walls. If we shift the CM in horizontal by a small distance, the disc oscillates simple harmonically. Assuming a perfect rolling of the disc on the horizontal surface, find the angular frequency of oscillation.




Fig 6.10



Watch Video Solution

8. Consider a liquid which fills a uniform U - tube uniform U - tube, as shown in figure , up to a height h . Find angular the frequency of small oscillation of th eliquid in the U - tube.




Fig. 4.130



Watch Video Solution

9. A particle of mass m is located in a unidimensional potential field where potential energy of the particle depends on the coordinates x as $U(x) = \frac{A}{x^2} - \frac{B}{x}$ where A and B are positive constant.

Find the time period of small oscillation that the particle perform about equilibrium position.



[Watch Video Solution](#)

10. A particle of mass $2kg$ is moving of a straight line under the actin force $F = (8 - 2x)N$. It is released at rest from $x = 6m$.

a. Is the partical moving simple harmonically.

b. Find the equilibrium position of the particle.

c. Write the equation of motion of the particle.


d. Find the time period of SHM.



[Watch Video Solution](#)


11. A body of mass m hangs from a smooth fixed pulley P_1 by the inextensible string fitted with the springs of stiffness k_1 and k_2 . The string passes over the smooth light pulley P_2 which is connected with another ideal spring of stiffness k_2 . Find the period of oscillation of the

body.



Watch Video Solution


12. A block of mass m connected with a smooth prismatic wedge of mass M is released from rest when the spring is relaxed. Find the angular frequency of oscillation.



[Watch Video Solution](#)

13. A stepped pulley having mass m radius of gyration k is connected with two ideal springs of stiffnesses k_1 and k_2 as shown in figure. If the pulley shown in the figure. If the

pulley rolls without sliding, find the angular frequency of its oscillation.




[▶ Watch Video Solution](#)

14. A stepped dies of mass M and radius R is pivoted at its center C smoothly. An inextensible string connected with a light spring of stiffness k passes over the pulley. One end of the string is rigidly connected with the ground and the other end is attached to a body of mass

m . If the string does not slide on the pulley, find the angular frequency of oscillation of the system


of the system.



[▶ Watch Video Solution](#)

15. A disc of mass m hanged by a string is attached at P and a spring of stiffness k is attached at O . Find the frequency of small angular oscillation of the disc if the

string does not slide over the pulley. Assume $I_0 = MR^2$ of the disc about O .



 Watch Video Solution


16. A uniform cylinder of length (L) and mass (M) having cross sectional area (A) is suspended, with its length vertical, from a fixed point by a massless spring, such that it is half - submerged in a liquid of density (ρ) at equilibrium position. When the cylinder is given a small downward push and released it starts oscillating vertically with small amplitude. If the force constant of the spring is (k), the frequency of oscillation of the cylinder is.



Watch Video Solution

17. Disregarding gravity, find the period of oscillation of the particle connected with four springs as shown in the figure.


(given: $\theta = 45^\circ$. $B\eta = 30^\circ$)



[▶ Watch Video Solution](#)

18. A smooth of mass m_1 is lying on a rigid horizontal string A bob of mass m_2 hangs from, the ring by an inextensible light string of length l . Find angular

frequency of oscillation of the system.



[▶ Watch Video Solution](#)

19. A smooth piston of mass m area of cross - section A is in equilibrium in a gas jar when the pressure of the gas is P_0 . Find the angular frequency of oscillation of the piston,

assuming adiabatic Change of state of the gas.





Fig. 4.1

[▶ Watch Video Solution](#)

20. If velocity of a particle moving along a straight line changes sinusoidally with time as shown in given graph. Find the average speed over time interval $t = 0$ to

$t = 2(2n - 1)$ second, n being any positive interget.



[Watch Video Solution](#)

21. In the figure shown, mass $2m$ connected with a spring of force constant k is at rest and in equilibrium. A partical of mass m is released from height $4.5mg/k$ from $2m$. The partical stick to the block. Neglecting the duration of collision find time from the release of m to the moment

when the spring has maximum compression.




Fig. 4.141



Watch Video Solution

1. While a particle executes linear simple harmonic motion
- A. its linear velocity and acceleration pass through their maximum and minimum values once in each oscillation.
 - B. Its linear velocity and acceleration pass through their maximum and minimum values twice in each oscillation.
 - C. its linear velocity and acceleration pass through their maximum and minimum values four times in each oscillation.

D. its linear velocity and acceleration always attain their peak values simultaneously.

Answer: B

 [Watch Video Solution](#)

2. While a particle executes simple harmonic motion, the rate of change of acceleration is maximum and minimum respectively at

- A. the mean position and extreme positions
- B. the extreme positions and mean position
- C. the mean position alternatively

D. the extreme positions alternatively.

Answer: A



Watch Video Solution

3. A hollow sphere filled with water through a small hole at the top. It is then hung by a long thread and made to oscillate. As the water slowly flows out of the hole at the bottom, the period of oscillation will

- A. go on increasing
- B. go on decreasing
- C. first increases and then decreases
- D. first decreases and then increases

Answer: D



Watch Video Solution

4. A simple pendulum oscillates slightly above a large horizontal metal plate. The bob is given a charge. The time period

- A. has no effect, whatever be the nature of charge
- B. always decreases, whatever be the nature of charge
- C. always increases, whatever be the nature of charge
- D. a increases or decreases depending upon the nature of charge.

Answer: B



Watch Video Solution

5. A block is resting on a piston which executes simple harmonic motion in vertical plain with a period of $2.0s$ in vertical plane at an amplitude just sufficient for the block to separate from the piston. The maximum velocity of the piston is

A. $\frac{5}{\pi} \frac{m}{s}$

B. $\frac{10}{\pi} \frac{m}{s}$

C. $\frac{\pi}{2} \frac{m}{s}$

D. $\frac{20}{\pi} \frac{m}{s}$

Answer: B



Watch Video Solution

6. The number of independent constituent simple harmonic motions yielding a resultant displacement equation of the periodic motion as

$$y = 8 \sin^2\left(\frac{t}{2}\right) \sin(10t) \text{ is}$$

A. 8


B. 6

C. 4

D. 3

Answer: D

 Watch Video Solution



7.

The diagram below shows a sinusoidal curve. The equation of the curve will be

A. $y = 10 \sin\left(16t + \frac{\pi}{4}\right) \text{ cm}$


B. $y = 10 \sin\left(16t + \frac{\pi}{3}\right) \text{ cm}$

C. $y = 10 \sin\left(16t - \frac{\pi}{4}\right) \text{cm}$

D. $y = 10 \cos\left(16t + \frac{\pi}{4}\right) \text{cm}$

Answer: C

 Watch Video Solution



8.

Fig. 4.143

The following figure shows the displacement versus time graph for two particles A and B executing simple

harmonic motions. The ratio of their maximum velocities is

A. 3:1

B. 1:3

C. 1:9


D. 9:1

Answer: A



Watch Video Solution

at one of the extreme positions will be



9.

The variation of velocity of a particle executing SHM with time is shown in fig. The velocity of the particle when a phase change of $\frac{\pi}{6}$ takes place from the instant it is at one of the extreme positions will be

A. $3.53 \frac{m}{s}$

B. $2.5 \frac{m}{s}$

C. $4.330 \frac{m}{s}$

D. none of these

Answer: B



Watch Video Solution

10. In the previous problem, the displacement of the particle from the mean position corresponding to the instant mentioned is

A. $\frac{5}{\pi}m$

B. $\frac{5\sqrt{3}}{\pi}m$

C. $\frac{10\sqrt{3}}{\pi}m$

D. $\frac{5\sqrt{3}}{2\pi}m$

Answer: B



Watch Video Solution

11. In problem 9 , the acceleration of the particle is

A. $\frac{5\sqrt{3}\pi}{2} \frac{m}{s}$

B. $\frac{5\pi^2}{2} \frac{m}{s^2}$

C. $\frac{5\sqrt{3}\pi}{4} \frac{m}{s^2}$

D. $5\sqrt{3} \frac{m}{s^2}$

Answer: C



Watch Video Solution

12. In problem 9 the acceleration of the particle is

A. $\frac{10}{\pi}m$ and $5\pi \frac{m}{s^2}$

B. $\frac{5}{\pi}m$ and $\frac{5\pi}{2} \frac{m}{s^2}$


C. $\frac{10}{\pi}m$ and $\frac{5\pi}{2} \frac{m}{s^2}$

D. $\frac{5}{\pi}m$ and $\frac{5\pi}{4} \frac{m}{s^2}$

Answer: C



Watch Video Solution



13.

Figure. Shows the variation of force acting on a particle of mass 400 g executing simple harmonic motion. The frequency of oscillation of the particle is

A. 4 s^{-1}

B. $\left(\frac{5}{2\pi}\right)\text{ s}^{-1}$

C. $\left(\frac{1}{8\pi}\right)\text{ s}^{-1}$

D. $\left(\frac{1}{2\pi}\right)\text{ s}^{-1}$

Answer: B



Watch Video Solution

14. A block of mass 1kg hangs without vibrating at the end of a spring whose force constant is $200\frac{N}{m}$ and which is attached to the ceiling of an elevator. The elevator is rising with an upward acceleration of $\frac{g}{3}$ when the acceleration suddenly ceases. The angular frequency of the block after the acceleration ceases is

A. $13\frac{rad}{s}$

B. $14\frac{rad}{s}$

C. $15\frac{rad}{s}$

D. none of these

Answer: B

 [Watch Video Solution](#)

15. A vertical spring carries a 5kg body and is hanging in equilibrium an additional force is applied so that the spring is further stretched. When released from this position. It performs 50 complete oscillation in 25 s, with an amplitude of 5 cm. The additional force applied is

A. 80 N

B. $80\pi^2 N$

C. $4\pi^2 N$

D. $4N$

Answer: C

 [Watch Video Solution](#)

16. A particle performs SHM in a straight line. In the first second, starting from rest, it travels a distance a and in the next second it travels a distance b in the same direction. The amplitude of the SHM is

A. $\frac{2a^2}{3b - a}$

B. $\frac{3a^2}{3a - b}$

C. $\frac{2a^2}{3a - b}$

D. $\frac{3a^2}{3b - a}$

Answer: C



Watch Video Solution

17. A particle free to move along the (x - axis) has potential energy given by

$$U(x) = k[1 - \exp(-x^2)] \text{ or } -\infty \leq x \leq +\infty,$$

where (k) is a positive constant of appropriate dimensions. Then.

A. for small displacement from $x = 0$, the motion is simple harmonic.

B. if its total mechanical energy is $\frac{k}{2}$, it has its minimum kinetic energy at the origin

C. for any finite non zero value of x , there is a force directed away from the origin

D.

Answer: A

 [Watch Video Solution](#)

18. Two simple harmonic motion are represented by equations

$$y_1 = 4 \sin(10t + \phi) \text{ and } y_2 = 5 \cos 10t$$

What is the phase difference between their velocities ?

A. ϕ

B. $-\phi$

C. $\left(\phi + \frac{\pi}{2}\right)$

D. $\left(\phi - \frac{\pi}{2}\right)$

Answer: D

 [Watch Video Solution](#)

19. The metallic bob of a simple pendulum has the relative density ρ . The time period of this pendulum is T if the metallic bob is immersed in water the new time period is given by

A. $2\pi\sqrt{\frac{l}{ng}}$

B. $2\pi\sqrt{\frac{l}{\left(1 - \frac{1}{n}\right)g}}$

C. $2\pi\sqrt{\frac{\ln}{g}}$

D. $2\pi\sqrt{\frac{l}{(n-1)g}}$

Answer: B



Watch Video Solution

20. Two particles move parallel to x - axis about the origin with the same amplitude and frequency. At a certain instant they are found at distance $\frac{A}{3}$ from the origin on opposite sides but their velocities are found to be in the same direction. What is the phase difference between the two ?

A. $\cos^{-1}\left(\frac{7}{9}\right)$

B. $\cos^{-1}\left(\frac{5}{9}\right)$

C. $\cos^{-1}\left(\frac{4}{9}\right)$

D. $\cos^{-1}\left(\frac{1}{9}\right)$

Answer: A



Watch Video Solution

21. The potential energy of a particle executing SHM along the x-axis is given by $U = U_0 - U_0 \cos ax$. What is the period of oscillation?

A. $2\pi\sqrt{\frac{ma}{U_0}}$

B. $2\pi\sqrt{\frac{U_0}{ma}}$

C. $\frac{2\pi}{a} \sqrt{\frac{m}{U_0}}$

D. $2\pi \sqrt{\frac{m}{aU_0}}$

Answer: C



Watch Video Solution

22. A particle executing SHM of amplitude a has displacement $\frac{a}{2}$ at $t = \frac{T}{4}$ and a negative velocity. The epoch of the particle is

A. $\frac{\pi}{3}$

B. $\frac{2\pi}{3}$

C. π

D. $\frac{5\pi}{3}$

Answer: A



Watch Video Solution

23. A block of mass 4 kg hangs from a spring constant $k = 400 \frac{N}{m}$. The block is pulled down through 15 cm below and released. What is its kinetic energy when the block is 10 above the equilibrium position.?

A. $5J$

B. $2.5J$

C. $1J$

D. $1.9J$

Answer: B

 [Watch Video Solution](#)

24. A body of mass 100 g attached to a spring executed SHM of period 2 s and amplitude 10 cm. How long a time is required for it to move from a point 5 cm below its equilibrium position to a point 5 cm above it, when it makes simple harmonic vertical oscillation (take $g = 10 \frac{m}{s^2}$)?

A. 0.6 s

B. $\frac{1}{3}$ s

C. 1.5s

D. 2.2s

Answer: B



Watch Video Solution

25. A particle executing SHM has velocities u and v and $a \leq$ *rationa* and b in two of its position. Find the distance between these two positions.

A. $\frac{u^2 - v^2}{a + b}$

B. $\frac{v^2 + u^2}{a - b}$

C. $\frac{v^2 + u^2}{a + b}$

D. $\frac{v^2 - u^2}{a - b}$

Answer: A



Watch Video Solution

26. Two particles are executing identical simple harmonic motions described by the equations $x_1 = a \cos\left(\omega t + \left(\frac{\pi}{6}\right)\right)$ and $x_2 = a \cos\left(\omega t + \frac{\pi}{3}\right)$. The minimum interval of time between the particles crossing the respective mean positions is $\frac{\pi}{2\omega}$

- A. $\frac{\pi}{2\omega}$
- B. $\frac{\pi}{3\omega}$
- C. $\frac{\pi}{4\omega}$
- D. $\frac{\pi}{6\omega}$

Answer: D



Watch Video Solution

27. The KE and PE , at is a particle executing SHM with amplitude A will be equal when its displacement is

A. $A\sqrt{2}$

B. $\frac{A}{2}$

C. $\frac{A}{\sqrt{2}}$



D. $A\sqrt{\frac{2}{3}}$




Answer: C



Watch Video Solution

28. A body is performing simple harmonic motion with amplitude a and line period T variation of its acceleration (f) with time(t) is shown in figure If a line x velocity of the body is x which of the following graph is correct?










Answer: A

 **Watch Video Solution**

29. A body is performing simple harmonic motion with amplitude a and time period T . The variation of its acceleration (f) with time (t) is shown in figure. If a line x velocity of

the body is x which of the following graph is correct?





D.

Answer: D

 Watch Video Solution





Fig. 4.148

30.

A particle is performing SHM. Its kinetic energy K varies with time t as shown in the figure. Then



A.

Period of oscillation of the particle is equal to T .

B. excess potential energy U of the particle varies with time t as shown in Fig.

C. excess potential energy U of the particle varies with time t as shown in fig.

D. none of these

Answer: B



Watch Video Solution

31. Two particle P and Q describe $S.H.M.$ of same amplitude a same frequency f along the same straight line .The maximum distance between the two particles is $a\sqrt{2}$ The phase difference between the two particle is

A. zero

B. $\frac{\pi}{2}$

C. $\frac{\pi}{6}$


D. $\frac{\pi}{3}$

Answer: B



Watch Video Solution

32. Two masses (m_1) and (m_2) are suspended together by a massless spring of spring constant (k). When the masses are in equilibrium, (m_1) is removed without disturbing the system. Find the angular frequency and amplitude of oscillation of (m_2).



A. $\frac{m_1 g}{k}$

B. $\frac{m_2 g}{k}$

C. $\frac{(m_1 + m_2)g}{k}$

D. $\frac{(m_2 - m_1)g}{k}$

Answer: A



33. A body of mass m is released from a height h to a scale pan hung from a spring. The spring constant of the spring is k , the mass of the scale pan is negligible and the body does not bounce relative to the pan, then the amplitude of vibration is

A. $\frac{mg}{k} \sqrt{1 - \frac{2hk}{mg}}$

B. $\frac{mg}{k}$

C. $\frac{mg}{k} + \frac{mg}{k} \sqrt{1 + \frac{2hk}{mg}}$

D. $\frac{mg}{k} - \frac{mg}{k} \sqrt{1 - \frac{2hk}{mg}}$

Answer: C



[Watch Video Solution](#)

34. Frequency of a particle executing SHM is 10 Hz. The particle is suspended from a vertical spring. At the highest point of its oscillation the spring is unstretched.

Maximum speed of the particle is $\left(g = 10 \frac{m}{s}\right)$

A. $2\pi \frac{m}{s}$

B. $\pi \frac{m}{s}$

C. $\frac{1}{\pi} \frac{m}{s}$

D. $\frac{1}{2\pi} / \frac{m}{s}$

Answer: D

[Watch Video Solution](#)

35. The potential energy of a particle of mass 1kg in motion along the x - axis is given by: $U = 4(1 - \cos 2x)$, where x in meters. The period of small oscillation (in sec) is

A. 2π

B. π

C. $\frac{\pi}{2}$

D. $\sqrt{2}\pi$

Answer: C



Watch Video Solution

36. An object of mass 0.2kg executes simple harmonic oscillation along the $x - a\xi$ with a frequency of $(25/\pi)\text{Hz}$. At the position $x = 0.04$, the object has Kinetic energy of 0.5J and potential energy 0.4J . The amplitude of oscillations is.....m.

A. 0.05m

B. 0.06m

C. 0.01m

D. none of these

Answer: B



Watch Video Solution

37. The string of a simple pendulum replaced by a uniform rod of length L and mass M while the bob has a mass m .

It is allowed to make small oscillation. Its time period is

A. $2\pi \sqrt{\left(\frac{2M}{3m}\right) \frac{L}{g}}$

B. $2\pi \sqrt{\frac{2(M + 3m)L}{3(M + 2m)g}}$


C. $2\pi \sqrt{\left(\frac{M + m}{M + 3m}\right) \frac{L}{g}}$

D. $2\pi \sqrt{\left(\frac{2m + M}{3(M + 2m)}\right) \frac{L}{g}}$

Answer: B



Watch Video Solution



38.

A uniform semicircular ring having mass m and radius r is hanging at one of its ends freely as shown in Fig. The ring is slightly disturbed so that it oscillates in its own plane. The time period of oscillation of the ring is

A. $2\pi \sqrt{\frac{r}{g\left(1 + \frac{1}{\pi^2}\right)}}$

B. $2\pi \sqrt{\frac{r}{g\left(1 + \frac{4}{\pi^2}\right)^{\frac{1}{2}}}}$


C. $2\pi \sqrt{\frac{r}{g\left(1 - \frac{2}{\pi^2}\right)^{\frac{1}{2}}}}$

D. $2\pi \sqrt{\frac{2r}{g\left(1 + \frac{4}{\pi}\right)^{\frac{1}{2}}}}$

Answer: D



Watch Video Solution



39.

15

Two springs with negligible masses and force constants $k_1 = 200 \frac{N}{m}$ and $k_2 = 160 \frac{N}{m}$ are attached to the block of mass $m = 10kg$ as shown in the fig. Initially the block is at rest, at the equilibrium position which both springs are neither stretched nor compressed. At time $t = 0$, sharp impulse of $50Ns$ is given to the block with a hammer along the spring.

A. Period of oscillations for the mass m is $\frac{\pi}{6}$ s.

B. Maximum velocity of the mass m during its

oscillation is $10\frac{m}{s}$

C. Data are insufficient to determine maximum velocity.

D. Amplitude of oscillation is $0.83m$.

Answer: D



Watch Video Solution

The frequency of small os




Fig. 4.153

40.

A thin uniform vertical rod of mass m and length l pivoted at point O is shown in Fig. The combined stiffness of the springs is equal to k . The mass of the spring is negligible.

The frequency of small oscillation is .

A. $\sqrt{\frac{3k}{2m} + \frac{g}{l}}$

B. $\sqrt{\frac{3k}{2m} + \frac{3g}{l}}$

C. $\sqrt{\frac{3k}{m} + \frac{3g}{2l}}$

D. $\sqrt{\frac{3k}{m} + \frac{2g}{3l}}$

Answer: C



Watch Video Solution

41. A particle executes SHM with time period 8 s. Initially, it is at its mean position. The ratio of distance travelled by it in the 1st second to that in the 2nd second is

A. $\sqrt{2} : 1$

B. $1 : (\sqrt{2} - 1)$

C. $(\sqrt{2} + 1) : \sqrt{2}$

D. $(\sqrt{2} - 1) : 1$

Answer: B



Watch Video Solution

42. A particle executed S.H.M. starting from its mean position at $t = 0$, If its velocity is $\sqrt{3}b\omega$, when it is at a distance b from the mean position, when $\omega = \frac{2\pi}{T}$, the time taken by the particle to move from b to the extreme position on the same side is

A. $\frac{5\pi}{6\omega}$

B. $\frac{\pi}{3\omega}$

C. $\frac{\pi}{2\omega}$

D. $\frac{\pi}{4\omega}$

Answer: B



43. In a certain oscillatory system (particle is performing SHM), the amplitude of motion is 5 m and the time period is 4 s. the minimum time taken by the particle for passing between points, which are at distances of 4 m and 3 m from the centre and on the same side of it will approximately be

A. $\frac{16}{45} s$

B. $\frac{7}{45} s$

C. $\frac{8}{45} s$

D. $\frac{13}{45} s$

Answer: C



44. A particle of mass m moving along x-axis has a potential energy $U(x) = a + bx^2$ where a and b are positive constant. It will execute simple harmonic motion with a frequency determined by the value of

- A. b alone
- B. b and a alone
- C. b and m alone
- D. b, a and m alone

Answer: C

45. The instantaneous displacement x of a particle executing simple harmonic motion is given by $x = a_1 \sin \omega t + a_2 \cos \left(\omega t + \frac{\pi}{6} \right)$. The amplitude A of oscillation is given by

A. $\sqrt{a_1^2 + a_2^2 + 2a_1a_2 \cos \left(\frac{\pi}{6} \right)}$

B. $\sqrt{a_1^2 + a_2^2 + 2a_1a_2 \cos \left(\frac{\pi}{3} \right)}$

C. $\sqrt{a_1^2 + a_2^2 - 2a_1a_2 \cos \left(\frac{\pi}{6} \right)}$

D. $\sqrt{a_1^2 + a_2^2 - 2a_1a_2 \cos \left(\frac{\pi}{3} \right)}$

Answer: D



Watch Video Solution

46. A simple harmonic motion along the x-axis has the following properties: amplitude = $0.4m$ the time to go from one extreme position to other is $2s$ and $x = 0.3m$ at $t = 0.5s$. the general equation of the simple harmonic motion is

A. $x = (0.5m)\sin\left[\frac{\pi t}{2} + 8^\circ\right]$

B. $x = (0.5m)\sin\left[\frac{\pi t}{2} - 8^\circ\right]$

C. $x = (0.5m)\cos\left[\frac{\pi t}{2} + 8^\circ\right]$

D. $x = (0.5m)\cos\left[\frac{\pi t}{2} - 8^\circ\right]$

Answer: B



Watch Video Solution

47. A spring balance has a scale that can read from 0 to 50 kg. The length of the scale is 20 cm. a body suspended from this balance when displaced and released oscillated harmonically with a time period of 0.6 s. The mass of the body is (take $g = 10 \frac{m}{s^2}$)

A. 10 kg

B. 25 kg


C. 18 kg

D. 22.8 kg

Answer: D



Watch Video Solution



48.

A soil cylinder of mass M and radius R is connected to a spring as shown in fig. The cylinder is placed on a rough horizontal surface. All the parts except the cylinder shown in the figure are light. If the cylinder is displaced slightly from its mean position and released, so that it performs pure rolling back and forth about its equilibrium position, determine the time period of oscillation?

A. $2\pi\sqrt{\frac{M}{k}}$


B. $2\pi\sqrt{\frac{3M}{2k}}$

C. $2\pi\sqrt{\frac{3M}{k}}$

D. none of these

Answer: B

 Watch Video Solution



49.

A block A is connected to spring and performs simple harmonic motion with a time period of 2 s. Another block B rests on A. The coefficient of static friction between A and B is $\mu_s = 0.6$. The maximum amplitude of oscillation

which the system can have so that there is no relative motion between A and B is (take $\pi^2 = g = 10$)

A. 0.3 m

B. 0.6 m

C. 0.4 m

D. 0.52 m

Answer: B



[Watch Video Solution](#)

50. A block of mass m is suspended from the ceiling of a stationary standing elevator through a spring of spring constant k . Suddenly, the cable breaks and the elevator

starts falling freely. Show that the block now executes a simple harmonic motion of amplitude $m\frac{g}{k}$ in the elevator

A. the block executes simple harmonic motion with

time period $2\pi\sqrt{\frac{m}{k}}$

B. the block executes simple harmonic motion with

amplitude $\frac{mg}{k}$

C. the block executes simple harmonic motion about

its mean position and the mean position is the position when the spring acquires its natural length.

D. all of the above.

Answer: D

51. A mass m attached to a spring of spring constant k is stretched a distance x_0 from its equilibrium position and released with no initial velocity. The maximum speed attained by mass in its subsequent motion and the time at which this speed would be attained are, respectively.


A. $\sqrt{(k)(m)}x_0, \pi\sqrt{\frac{m}{k}}$

B. $\sqrt{\frac{k}{m}}\frac{x_0}{2}, \frac{\pi}{2}\sqrt{\frac{m}{k}}$

C. $\sqrt{\frac{k}{m}}x_0, \frac{\pi}{2}\sqrt{\frac{m}{k}}$

D. $\sqrt{\frac{k}{m}}\frac{x_0}{2}, \pi\sqrt{\frac{m}{k}}$

Answer: C



52.

A plank of mass 12 kg is supported by two identical springs as shown in Fig. The plank always remains horizontal. When the plank is pressed down and released it performs simple harmonic motion with time period 3 s. When a block of m is attached to the plank the time period changes to 6 s. The mass of the block is

A. 48 kg

B. 36 kg

C. 24 kg

D. 12 kg

Answer: B



Watch Video Solution

53. The time taken by a particle performing *SHM* on a straight line to pass from point *A* to *B* where its velocities are same is 2 seconds .After another 2 seconds it return to *B* The time period of oscillation is (in seconds)

A. 2 s

B. 4 s

C. 6 s

D. 8 s

Answer: D



Watch Video Solution

54. Two springs are made to oscillate simple harmonically due to the same mass individually. The time periods obtained are T_1 and T_2 . If both the springs are connected in series and then made to oscillate by the same mass, the resulting time period will be

A. $T_1 + T_2$

B. $\frac{T_1 T_2}{T_1 + T_2}$


C. $\sqrt{T_1^2 + T_2^2}$

D. $\frac{T_1 + T_2}{2}$

Answer: C

 Watch Video Solution

is



55.

A thin-walled tube of mass m and radius R has a rod of

mass m and very small cross section soldered on its inner surface. The side-view of the arrangement is as shown in the following figure. The entire arrangement is placed on a rough horizontal surface. The system is given a small angular displacement from its equilibrium position, as a result, the system performs oscillations. The time period of resulting oscillations if the tube rolls without slipping is

A. $2\pi\sqrt{\frac{4R}{g}}$

B. $2\pi\sqrt{\frac{2R}{g}}$

C. $2\pi\sqrt{\frac{R}{g}}$

D. none of these

Answer: B

56. A thin uniform rod of mass 1 kg and length 12 cm is suspended by a wire that passes through its centre and is perpendicular to its length. The wire is twisted and the rod is set oscillating. Time period of oscillation is found to be 3 s. When a flat triangular plate is suspended in same way through its centre of mass, the time period is found to be 6 s. The moment of inertia of the triangular plate about this axis is

A. $0.12 \text{ kg} - \text{m}^2$

B. $0.24 \text{ kg} - \text{m}^2$

C. $0.48 \text{ kg} - \text{m}^2$

D. information insufficient

Answer: C

 **Watch Video Solution**

57. A particle performs SHM about $x = 0$ such that at $t = 0$ it is at $x = 0$ and moving towards positive extreme. The time taken by it to go from $x = 0$ to $x = \frac{A}{2}$ is time the time taken to go from $x = \frac{A}{2}$ to a. The most suitable option for the blank space is

A. 2

B. $\frac{1}{2}$

C. $\frac{3A}{T}$

D. $\frac{A}{2T}$

Answer: B

 [Watch Video Solution](#)

58. A particle performs SHM with a period T and amplitude a . The mean velocity of particle over the time interval during which it travels $a/2$ from the extreme position is

A. $\frac{A}{T}$

B. $\frac{2A}{T}$

C. $\frac{3A}{T}$

D. $\frac{A}{2T}$

Answer: C



Watch Video Solution

59. A particle performs simple harmonic motion about O with amplitude A and time period T . The magnitude of its acceleration at $t = \frac{T}{8}$ s after the particle reaches the extreme position would be

A. $\frac{4\pi^2 A}{\sqrt{2T^2}}$

B. $\frac{4\pi^2 A}{T^2}$

C. $\frac{2\pi^2 A}{\sqrt{2T^2}}$

D. none of these

Answer: A



Watch Video Solution

60. In the previous question, the magnitude of velocity of particle at the mentioned instant is

A. $\frac{\pi A}{T}$

B. $\frac{\sqrt{2}\pi A}{T}$

C. zero

D. $\sqrt{\frac{7}{8}} \times \frac{2\pi A}{T}$

Answer: B



Watch Video Solution

61. An object of mass 4 kg is attached to a spring having spring constant $100\frac{N}{m}$. It performs simple harmonic motion on a smooth horizontal surface with an amplitude of 2 m. A 6 kg object is dropped vertically onto the 4 kg object when it crosses the mean position, and sticks to it. the change in amplitude of oscillation due to collision is

A. $1m$

B. *zero*

C. $2\left[1 - \sqrt{\frac{2}{5}}\right]$

D. $2\left[1 - \frac{1}{\sqrt{5}}\right]$

Answer: C



Watch Video Solution

62. A cork floating on the pond water executes a simple harmonic motion, moving up and down over a range of 4 cm. The time period of the motion is 1 s. at $t = 0$, the cork is at its lowest position of oscillation, the position and velocity of the cork at $t = 10.5s$, would be

A. 2 cm above the mean position, $0 \frac{m}{s}$

B. 2 cm below the mean position $0 \frac{m}{s}$

C. 1 cm above the mean position $2\sqrt{3}\pi \frac{m}{s}$ up

D. 1 cm below the mean position, $2\sqrt{3}\pi \frac{m}{s}$ up

Answer: A



Watch Video Solution

63. A spring is placed in vertical position by suspending it from a hook at its top. A similar hook on the bottom of the spring is at 11 cm above a table top. A mass of 75 g and of negligible size is then suspended from the bottom hook, which is measured to be 4.5 cm above the table top. The mass is then pulled down a distance of 4 cm and released. Find the approximate. Position of the bottom hook after 5 s? Take $g = 10 \frac{m}{s^2}$ and hook's mass to be negligible.

- A. 5 cm above the table top
- B. 4.5 cm above the table top
- C. 9 cm above the table top
- D. 0.5 cm above the table top

Answer: D



Watch Video Solution

64. A particle is performing *SHM* according to the equation $x = (3\text{cm})\sin\left(\frac{2\pi}{18}t + \frac{\pi}{6}\right)$, where t is in seconds. The distance travelled by the particle in 39 s is

A. 24cm

B. 1.5cm


C. 25.5cm

D. none of these

Answer: C



Watch Video Solution



65.

Two springs, each of unstretched length 20 cm but having different spring constant $k_1 = 1000 \frac{N}{m}$ and $k_2 = 3000 \frac{N}{m}$ are attached to two opposite faces of a small block of mass $m = 100g$ kept on a smooth horizontal surface as shown in fig. The outer ends of the two springs are now attached to two pins P_1 and P_2 whose locations are shown in the figure. as a result of this, the block acquires a new equilibrium position. The block has been displaced by small amount from its equilibrium position and released to perform simple harmonic motion: then

A. new equilibrium position is at 35 cm from P_1 and

time period of simple harmonic motion is $\frac{\pi}{100}$ s.

B. New equilibrium position is at 20 cm from P_1 and

time period of simple harmonic motion is $\frac{\pi}{100}$ s

C. new equilibrium position is at 35 cm from P_1 and

time period of simple harmonic motion is $\frac{\pi}{25}$ s

D. new equilibrium position is at 30 cm from P_1 and

time period of simple harmonic motion is $\frac{\pi}{26}$ s

Answer: A



Watch Video Solution

66. A particle of mass m is present in a region where the potential energy of the particle depends on the x -coordinate according to the expression $U = \frac{a}{x^2} - \frac{b}{x}$, where a and b are positive constant. The particle will perform.

A. oscillatory motion but not simple harmonic motion about its mean position for small displacements

B. simple harmonic motion with time period $2\pi\sqrt{\frac{8a^2m}{b^4}}$ about its mean position for small displacements

C. neither simple harmonic motion nor oscillatory about its mean position for small displacements

D. none of the above

Answer: D

 [Watch Video Solution](#)

67. A particle performing simple harmonic motion having time period 3 s is in phase with another particle which also undergoes simple harmonic motion at $t = 0$. The time period of second particle is T (less than 3s). If they are again in the same phase for the third time after 45 s, then the value of T will be

A. 2.8 s

B. 2.7 s

C. 2.5 s

D. none of these

Answer: C

 [Watch Video Solution](#)

68. A particle performs *SHM* on x- axis with amplitude A and time period T . The time taken by the particle to travel a distance $A/5$ from rest is

A. $\frac{T}{20}$


B. $\frac{T}{2\pi} \cos^{-1} \left(\frac{4}{5} \right)$

C. $\frac{T}{2\pi} \cos^{-1} \left(\frac{1}{5} \right)$

D. $\frac{T}{2\pi} \sin^{-1} \left(\frac{1}{5} \right)$

Answer: B

 Watch Video Solution



69.

The coefficient of friction between block of mass m and $2m$ is $\mu = 2 \tan \theta$. There is no friction between block of mass $2m$ and inclined plane. The maximum amplitude of

the two block system for which there is no relative motion between both the blocks is

A. $g \sin \theta \sqrt{\frac{k}{m}}$

B. $\frac{mg \sin \theta}{k}$


C. $\frac{3mg \sin \theta}{k}$

D. none of these

Answer: C



Watch Video Solution



70.

A block of mass m is suspended from a spring and executes vertical SHM of time period T as shown in Fig.

The amplitude of the SHM is A and spring is never in compressed state during the oscillation. The magnitude of minimum force exerted by spring on the block is

A. $mg - \frac{4\pi^2}{T^2}mA$

B. $mg + \frac{4\pi^2}{T^2}mA$

C. $mg - \frac{\pi^2}{T^2}mA$

D. $mg + \frac{\pi^2}{T^2}mA$

Answer: A



Watch Video Solution

71. A particle performs *SHM* of amplitude A along a straight line .When it is at a distance of $\frac{\sqrt{3}}{2}A$ from mean position its kinetic energy gets increased by an amount of $\frac{1}{2}m\omega^2A^2$ due to an impulsive force. Then its new amplitude becomes

A. $\frac{\sqrt{5}}{2}A$

B. $\frac{\sqrt{3}}{2}A$

C. $\sqrt{2}A$

D. $\sqrt{5}A$

Answer: C



Watch Video Solution

72. A horizontal spring-block system of mass $2kg$ executes $S.H.M$ when the block is passing through its equilibrium position an object of mass $1kg$ is put on it the two move together The new amplitude of vibration is (A being its initial amplitude)

A. $\sqrt{\frac{2}{3}}A$

B. $\sqrt{\frac{3}{2}}A$


C. $\sqrt{2}A$

D. $\frac{A}{\sqrt{2}}$

Answer: A



Watch Video Solution



73.

A metre stick swinging in vertical plane about a fixed horizontal axis passing through its one end undergoes small oscillation of frequency f_0 . If the bottom half of the stick were but off, then its new frequency of small oscillation would become.

A. f_0

B. $\sqrt{2}f_0$

C. $2f_0$

D. $2\sqrt{2}f_0$

Answer: B



Watch Video Solution

74. A physical pendulum is positioned so that its centre of gravity is above the suspension point. When the pendulum is released it passes the point of stable equilibrium with an angular velocity ω . The period of small oscillations of the pendulum is

A. $\frac{4\pi}{\omega}$

B. $\frac{2\pi}{\omega}$

C. $\frac{\pi}{\omega}$

D. $\frac{\pi}{2\omega}$

Answer: A

 **Watch Video Solution**

75. A particle executing harmonic motion is having velocities v_1 and v_2 at distances x_1 and x_2 from the equilibrium position. The amplitude of the motion is

A. $\sqrt{\frac{v_1^2 x_2 - v_2^2 x_1}{v_1^2 - v_2^2}}$


B. $\sqrt{\frac{v_1^2 x_1^2 - v_2^2 x_2^2}{v_1^2 + v_2^2}}$

C. $\sqrt{\frac{v_1^2 x_2^2 - v_2^2 x_1^2}{v_1^2 - v_2^2}}$

D. $\sqrt{\frac{v_1^2 x_2^2 + v_2^2 x_1^2}{v_1^2 + v_2^2}}$

Answer: C

 Watch Video Solution



76.

A wire is bent at an angle θ . A rod of mass m can slide along the bended wire without friction as shown in Fig. A soap film is maintained in the frame kept in a vertical position and the rod is in equilibrium as shown in the figure. If rod is displaced slightly in vertical direction, then the time period of small oscillation of the rod is

A. $2\pi\sqrt{\frac{l}{g}}$

B. $2\pi\sqrt{\frac{l \cos \theta}{g}}$


C. $2\pi\sqrt{\frac{l}{g \cos \theta}}$

D. $2\pi\sqrt{\frac{l}{g \tan \theta}}$

Answer: A



Watch Video Solution



77.

A solid right circular cylinder of weight 10 kg and cross sectional area 100cm^2 is suspended by a spring, where $k = 1\frac{\text{kg}}{\text{cm}}$, and hangs partially submerged in water of density $1000\frac{\text{kg}}{\text{m}^3}$ as shown in Fig. What is its period when it makes simple harmonic vertical oscillations? (Take $g = 10\frac{\text{m}}{\text{s}^2}$)

A. 0.6 s

B. 1 s

C. 1.5 s

D. 2.2 s

Answer: A

 **Watch Video Solution**




Fig. 4.164

78.

A block A of mass m is placed on a smooth horizontal

platform P and between two elastic massless springs S_1 and S_2 fixed horizontally to two fixed vertical walls. The elastic constants of the two springs are equal to k and the equilibrium distance between the two springs both in relaxed states is d . The block is given a velocity v_0 initially towards one of the springs and it then oscillated between the springs. The time period T of oscillations and the minimum separation d_m of the spring will be

A. $T = 2\left(\frac{d}{v} + \pi\sqrt{\frac{m}{k}}\right), d_m = d$

B. $T = 2\left(\frac{d}{v} + 2\pi\sqrt{\frac{m}{k}}\right), d_m = d - v\sqrt{\frac{m}{k}}$

C. $T = 2\left(\frac{d}{v} + 2\pi\sqrt{\frac{m}{k}}\right), d_m = d - 2v\sqrt{\frac{m}{k}}$

D. $T = 2\pi\sqrt{\frac{m}{k}}, d_m = d$

Answer: A

79. A certain simple harmonic vibrator of mass 0.1 kg has a total energy of 10 J. Its displacement from the mean position is 1 cm when it has equal kinetic and potential energies. The amplitude A and frequency n of vibration of the vibrator are

A. $A = \sqrt{2} \text{ cm}, n = \frac{500}{\pi} \text{ Hz}$

B. $A = \sqrt{2} \text{ cm}, n = \frac{1000}{\pi} \text{ Hz}$

C. $A = \frac{1}{\sqrt{2}} \text{ cm}, n = \frac{500}{\pi} \text{ Hz}$

D. $A = \frac{1}{\sqrt{2}} \text{ cm}, n = \frac{1000}{\pi} \text{ Hz}$

Answer: A

80. A simple pendulum of length l and mass m is suspended in a car that is moving with constant speed v around a circle of radius r . Find the period of oscillation and equilibrium position of the pendulum.

A. $\frac{1}{2\pi} \sqrt{\frac{g}{l}}$

B. $\frac{1}{2\pi} \sqrt{\frac{g}{R}}$


C. $\frac{1}{2\pi} \sqrt{\frac{\left(g^2 + \frac{v^4}{R^2}\right)}{l}}$

D. $\frac{1}{2\pi} \sqrt{\frac{v^2}{Rl}}$

Answer: C



Watch Video Solution



81.

One end of a spring of force constant K is fixed to a vertical wall and the other to a body of mass m resting on a smooth horizontal surface. There is another wall at a distance x_0 from the body. The spring is then compressed by $3x_0$ and released. The time taken to strike the wall

from the instant of release is (given $\sin^{-1}\left(\frac{1}{3}\right) = \left(\frac{\pi}{9}\right)$)

A. $\frac{\pi}{6} \sqrt{\frac{m}{K}}$

B. $\frac{2\pi}{3} \sqrt{\frac{m}{K}}$

C. $\frac{\pi}{4} \sqrt{\frac{m}{K}}$

D. $\frac{11\pi}{18} \sqrt{\frac{m}{K}}$


Answer: D



Watch Video Solution

82. A block P of mass m is placed on horizontal frictionless plane. A second block of same mass m is placed on it and is connected to a spring of spring constant k , the two blocks are pulled by distance A . Block Q oscillates without slipping. What is the maximum value of frictional force

between the two blocks.



A. zero

B. k

C. $\frac{kA}{2}$

D. μg


Answer: C



Watch Video Solution

83. A uniform stick of mass M and length L is pivoted at its center. Its ends are attached to two springs each of the constant K . In the position shown in figure the stick is horizontal. The spring is displayed through a small length when the spring is displayed through a

small angle of and released. The stick :



A. executes non periodic motion

B. executes periodic motion which is not simple harmonic .


C. executes SHM of frequency $\frac{1}{2\pi} \sqrt{\frac{6K}{M}}$.

D. executes SHM of frequency $\frac{1}{2\pi} \sqrt{\frac{K}{2M}}$.

Answer: C



Watch Video Solution



84.

A mass m is suspended from a spring of force constant k and just touches another identical spring fixed to the floor as shown in the fig. The time period of small oscillations is

A. $2\pi\sqrt{\frac{m}{k}}$

B. $\pi\sqrt{\frac{m}{k}} + \pi\sqrt{\frac{m}{\frac{k}{2}}}$

C. $\pi\sqrt{\frac{m}{\frac{3k}{2}}}$

D. $\pi\sqrt{\frac{m}{k}} + \pi\sqrt{\frac{m}{2k}}$

Answer: D



Watch Video Solution

85. A street car moves rectilinearly from station A to the next station B (from rest to rest) with an acceleration varying according to the law $f = a - bx$, where a and b are constants and x is the distance from station A. The

distance between the two stations and the maximum velocity are

A. $x = \frac{2a}{b}, v_{\max} = \frac{a}{\sqrt{b}}$

B. $x = \frac{b}{2a}, v_{\max} = \frac{a}{b}$

C. $x = \frac{a}{2b}, v_{\max} = \frac{b}{\sqrt{a}}$

D. $x = \frac{a}{b}, v_{\max} = \frac{\sqrt{a}}{b}$

Answer: A



Watch Video Solution

86. Two particle P and Q describe $S.H.M.$ of same amplitude a same frequency f along the same straight

line .The maximum distance between the two particles is

$a\sqrt{2}$ The phase difference between the two particle is

A. zero

B. $\frac{\pi}{2}$

C. $\frac{\pi}{6}$

D. $\frac{\pi}{3}$

Answer: B



Watch Video Solution

87. The velocity v of a particle of mass is moving along a straight line change with time t as $\frac{d^2v}{dt^2} = -Kv$ where

K is a particle constant which of the following statement is correct?

A. the particle does not perform SHM.

B. The particle performs SHM with time period

$$2\pi\sqrt{\frac{m}{K}}$$


C. The particle performs SHM with frequency $\frac{\sqrt{K}}{2\pi}$

D. The particle performs SHM with time period $\frac{2\pi}{K}$.

Answer: C



Watch Video Solution



88.

The oscillations represented by curve 1 in the graph are expressed by equation $x = A \sin \omega t$. The equation for the oscillation represented by curve 2 is expressed as

A. $x = 2A \sin\left(\omega t - \frac{\pi}{2}\right)$

B. $x = 2A \sin\left(\omega t + \frac{\pi}{2}\right)$

C. $x = -2A \sin\left(\omega t - \frac{\pi}{2}\right)$

D. $x = A \sin\left(\omega t - \frac{\pi}{2}\right)$

Answer: A

 Watch Video Solution




Fig. 4.170

89.

Graph shows the $x(t)$ curves for the three experiments involving a particular spring block system oscillating in SHM. The kinetic energy of the system is maximum at $t = 4$ s. For the situation.

A. 1

B. 2

C. 3

D. same in all

Answer: A

 [Watch Video Solution](#)

90. The acceleration of a particle moving along x-axis is $a = -100x + 50$. It is released from $x = 2$. Here a and x are in S.I units. The motion of particle will be:

A. periodic, oscillatory but not SHM.

B. periodic but not oscillatory

C. oscillatory but not periodic.

D. simple harmonic.

Answer: D

 [Watch Video Solution](#)

91. In the above question, the speed of the particle at origin will be:

A. $10\sqrt{2}\frac{m}{s}$

B. $1.5\frac{m}{s}$

C. $10\frac{m}{s}$

D. none of these

Answer: A



Watch Video Solution

92. A particle performs *SHM* of amplitude A along a straight line. When it is at a distance of $\frac{\sqrt{3}}{2}A$ from mean position its kinetic energy gets increased by an amount of $\frac{1}{2}m\omega^2A^2$ due to an impulsive force. Then its new amplitude becomes


A. $\frac{\sqrt{5}}{2}A$

B. $\frac{\sqrt{3}}{2}A$

C. $\sqrt{2}A$

D. $\sqrt{5}A$

Answer: C



93.

Fig. 1.17

A horizontal rod of mass m and length L is pivoted at one end. The rod's other end is supported by a spring of force constant k . The rod is displaced by a small angle θ from its horizontal equilibrium position and released. The angular frequency of the subsequent simple harmonic motion is

- A. $\sqrt{\frac{3k}{m}}$
- B. $\sqrt{\frac{k}{3m}}$

C. $\sqrt{\frac{3k}{m} + \frac{3g}{2L}}$

D. $\sqrt{\frac{k}{m}}$

Answer: A

 [Watch Video Solution](#)

94. A small mass executes linear *SHM* about O with amplitude a and period T . Its displacement from O at time $T/8$ after passing through O is:

A. $\frac{a}{8}$

B. $\frac{a}{2\sqrt{2}}$

C. $\frac{a}{2}$

D. $\frac{a}{\sqrt{2}}$

Answer: D

 [Watch Video Solution](#)

95. Time period of a particle executing *SHM* is 8 sec. At $t = 0$ it is at the mean position. The ratio of the distance covered by the particle in the *1st* second to the *2nd* second is:

A. $\frac{1}{\sqrt{2} + 1}$

B. $\sqrt{2}$

C. $\frac{1}{\sqrt{2}}$

D. $\sqrt{2} + 1$

Answer: D



Watch Video Solution

96. A particle performs SHM with a period T and amplitude a . The mean velocity of particle over the time interval during which it travels $a/2$ from the extreme position is

A. $\frac{a}{T}$

B. $\frac{2A}{T}$





C. $\frac{3A}{T}$

D. $\frac{A}{2T}$

Answer: C



97. A graph of the square of the velocity against the square of the acceleration of a given simple harmonic motion is



Answer: D



Watch Video Solution

98. A plank with a small block on top of it is under going vertical *SHM*. Its period is 2 sec. The minimum amplitude at which the block will separate from plank is :

A. $\frac{10}{\pi^2}$

B. $\frac{\pi^2}{10}$

C. $\frac{20}{\pi^2}$

D. $\frac{20}{\pi^2}$

Answer: A



Watch Video Solution

99. The potential energy of a simple harmonic oscillator of mass 2 kg in its mean position is 5 J. If its total energy is 9 J and its amplitude is 0.01 m, its time period would be

A. $\frac{\pi}{10} s$

B. $\frac{\pi}{20} s$

C. $\frac{\pi}{50} s$

D. $\frac{\pi}{100} s$

Answer: D



Watch Video Solution

100. A spring mass system performs $S. H. M$ if the mass is doubled keeping amplitude same, then the total energy of $S. H. M$ will become :

- A. double
- B. half
- C. unchanged
- D. 4 times

Answer: C



Watch Video Solution

101. A particle of mass m moves in a one dimensional potential energy $U(x) = -ax^2 + bx^4$, where a and b are positive constant. The angular frequency of small oscillation about the minima of the potential energy is equal to

A. $\pi \sqrt{\frac{a}{2b}}$

B. $2\sqrt{\frac{a}{m}}$

C. $\sqrt{\frac{2a}{m}}$

D. $\sqrt{\frac{a}{2m}}$

Answer: B



Watch Video Solution

102. A particle of mass m moves in the potential energy U shown above. The period of the motion when the particle has total energy E is

A. $2\pi\sqrt{\frac{m}{k}} + 4\sqrt{\frac{2E}{mg^2}}$

B. $2\pi\sqrt{\frac{m}{k}}$

C. $\pi\sqrt{\frac{m}{k}} + 2\sqrt{\frac{2E}{mg^2}}$

D. $2\sqrt{\frac{2E}{mg^2}}$

Answer: C



Watch Video Solution

103. The displacement of a body executing *SHM* is given by $x = A \sin(2\pi t + \pi/3)$. The first time from $t = 0$ when the velocity is maximum is

A. 0.33 s

B. 0.16 s

C. 0.25 s

D. 0.5 s

Answer: A



Watch Video Solution

104. Two particles are in *SHM* in a straight line about same equilibrium position. Amplitude A and time period T of both the particles are equal. At time $t = 0$, one particle is at displacement $y_1 = +A$ and the other at $y_2 = -A/2$, and they are approaching towards each other. after what time they cross each other?

A. $\frac{T}{3}$

B. $\frac{T}{4}$

C. $\frac{4T}{6}$


D. $\frac{T}{6}$

Answer: D



Watch Video Solution

105. A system of two identical rods (L-shaped) of mass m and length l are resting on a peg P as shown in the figure. If the system is displaced in its plane by a small angle θ , find the period of oscillations:



A. $2\pi\sqrt{\frac{\sqrt{2}l}{3g}}$

B. $2\pi\sqrt{\frac{2\sqrt{2}l}{3g}}$

C. $2\pi\sqrt{\frac{2l}{3g}}$

D. $3\pi\sqrt{\frac{l}{3g}}$

Answer: C



Watch Video Solution

106. A particle is subjected to two mutually perpendicular simple harmonic motions such that its X and y coordinates are given by $X = 2\sin\omega t$,
 $y = 2\sin\left(\omega + \frac{\pi}{4}\right)$

The path of the particle will be:

- A. an ellipse
- B. a straight line
- C. a parabola

D. circle

Answer: A



Watch Video Solution

107. Two simple harmonic motions $y_1 = A \sin \omega t$ and $y_2 = A \cos \omega t$ are superimposed on a particle of mass m .

A. $\frac{1}{2}m\omega^2 A^2$

B. $m\omega^2 A^2$

C. $\frac{1}{4}m\omega^2 A^2$

D. zero

Answer: B



Watch Video Solution

Multiple Correct

1. A coin is placed on a horizontal platform which undergoes vertical simple harmonic motion of angular frequency ω . The amplitude of oscillation is gradually increased. The coin will leave contact with the platform for the first time

A. at the highest position of the platform

B. at the mean position of the platform

C. for an amplitude of $\frac{g}{\omega^2}$

D. for an amplitude of $\sqrt{\frac{g}{\omega}}$

Answer: A::C



Watch Video Solution

2. For a simple harmonic motion with given angular frequency ω , two arbitrary initial conditions are necessary and sufficient to determine the motion completely. These initial conditions may be

- A. initial position and initial velocity
- B. amplitude and initial phase
- C. total energy of oscillation and amplitude
- D. total energy of oscillation and initial phase.

Answer: A::B::D



Watch Video Solution

3. The potential energy U of a body of unit mass moving in one dimensional conservative force field is given by $U = x^2 - 4x + 3$. All units are in SI. For this situation mark out the correct statement (s).

- A. The body will perform simple harmonic motion about $x = 2$ units.
- B. The body will perform oscillatory motion but not simple harmonic motion.
- C. The body will perform simple harmonic motion with time period $\sqrt{2}\pi s$.


D. If speed of the body at equilibrium position is $4\frac{m}{s}$,

then the amplitude of oscillation would be $2\sqrt{2}m$

Answer: A::C::D



Watch Video Solution



4.

For the spring pendulum shown in fig. the value of spring

constant is $3 \times 10^4 \frac{N}{m}$ and amplitude of oscillation is $0.1m$. The total mechanical energy of oscillating system is 200 J. Mark out the correct option (s).

- A. Minimum PE of the oscillating system is 50 J
- B. Maximum PE of the oscillating system is 200J
- C. maximum KE of the oscillating system is 200 J
- D. minimum KE of the oscillating system is 150 J.

Answer: A::B

 [Watch Video Solution](#)

ent(s).




Fig. 17.17

5.

An object of mass m is performing simple harmonic motion on a smooth horizontal surface as shown in Fig. just as the oscillating object reaches its extreme position, another object of mass $2m$ is dropped on to oscillating object, which sticks to it. For this situation mark out the correct statement (s).


- A. Amplitude of oscillations remains unchanged.
- B. time period of oscillation remains unchanges.

C. The total mechanical energy of the system does not change.

D. The maximum speed of oscillating object changes.

Answer: A::C::D

 Watch Video Solution



6.

A simple pendulum consists of a bob of mass m and a

light string of length l as shown in the fig. another identical ball moving with the small velocity v_0 collides with the pendulum's bob and sticks to it. For this new pendulum of mass $2m$, mark out the correct statements (s).

A. Time period of the pendulum is $2\pi\sqrt{\frac{l}{g}}$.

B. The equation of motion for this pendulum is

$$\theta = \frac{v_0}{2\sqrt{gl}} \sin \left[\sqrt{\frac{g}{l}} t \right]$$

C. The equation of motion for this pendulum is

$$\theta = \frac{v_0}{2\sqrt{gl}} \cos \left[\sqrt{\frac{g}{l}} t \right].$$

D. Time period of the pendulum is $2\pi\sqrt{\frac{2l}{g}}$.

Answer: A:B



7. A particle performing simple harmonic motion undergoes initial displacement of $\frac{A}{2}$ (where A is the amplitude of simple harmonic motion) in 1 s. At $t = 0$, the particle may be at the extreme position or mean position. The time period of the simple harmonic motion can be

- A. 6s
- B. 2.4s
- C. 12s
- D. 1.2s

Answer: A::C



8. A particle is subjected to two simple harmonic motions along x and y directions according to $x = 3 \sin 100\pi t$,
 $y = 4 \sin 100\pi t$.

A. Motion of particle will be on ellipse travelling in clockwise direction.

B. Motion of particle will be on a straight line with slope $\frac{4}{3}$

C. Motion will be simple harmonic motion with amplitude 5.

D. Phase difference between two motions is $\frac{\pi}{2}$

Answer: B::C



Watch Video Solution

9. The speed v of a particle moving along a straight line, when it is at a distance (x) from a fixed point of the line is given by $v^2 = 108 - 9x^2$ (all equation are in CGS units):

A. The motion is uniformly accelerated along the straight line

B. the magnitude of the acceleration at a distance 3 cm from the point is $27 \frac{cm}{s^2}$

C. the motion is simple harmonic about the given fixed point

D. the maximum displacement from the fixed point is 4
cm

Answer: B::C

 [Watch Video Solution](#)

10. A horizontal plank has a rectangular block placed on it. The plank starts oscillating vertically and simple harmonically with an amplitude of 40 cm. The block just loses contact with the plank when the latter is at momentary rest Then.

A. the period of oscillation is $\left(\frac{2\pi}{5}\right)$

- B. the block weighs double its actual weight, then the plank is at one of the positions of momentary rest.
- C. the block weighs 1.5 times its weight on the plank halfway down
- D. the block weighs its true weight on the plank when the later moves fastest.

Answer: A::B::C::D

 [Watch Video Solution](#)

11. A 20g particle is subjected to two simple harmonic motions $x_1 = 2 \sin 10t, x_2 = 4 \sin\left(10t + \frac{\pi}{3}\right)$. Where x_1 and x_2 are in metres and t is in seconds.

A. the displacement of the particle at $t = 0$ will be

$$2\sqrt{3} \text{ m}$$

B. maximum speed of the particle will be $20\sqrt{7} \frac{m}{s}$

C. Magnitude of maximum acceleration of the particle

$$\text{will be } 200\sqrt{7} \frac{m}{s^2}$$

D. Energy of the result motion will be 28 J

Answer: A::B::C::D



Watch Video Solution




Fig. 4.176

12.

A spring block system undergoes SHM on a smooth horizontal surface, the block is now given some charge and a uniform horizontal electric field E is switched on as shown in Fig. As a result

- A. Time period of oscillation will increase
- B. Time period of oscillation will decrease
- C. Time period of oscillation will remain unaffected
- D. the mean position of SHM will shift to the right.

Answer: C::D



Watch Video Solution

13. The potential energy of a particle of mass 0.1 kg, moving along the x -axis, is given by $U = 5x(x - 4)J$, where x is in meter. It can be concluded that

A. the particle is acted upon by a constant force

B. the speed of the particle is maximum at $x = 2m$

C. the particle executes SHM

D. the period of oscillation of the particle $\left(\frac{\pi}{5}\right)$ s

Answer: B::C::D



Watch Video Solution



14. The time period of a particle in simple harmonic motion is T . Assume potential energy at mean position to be zero. After a time of $\frac{T}{6}$ it passes its mean position its,

- A. velocity will be half its maximum velocity
- B. displacement will be half its amplitude
- C. acceleration will be nearly 86 % of its maximum
- D. $KE = PE$

Answer: A:C



Watch Video Solution



15.

Figure. (a) shows a spring of force constant k fixed at one end and carrying a mass m at the other end placed on a horizontal frictionless surface. The spring is stretched by a force F . Figure. (b) shows the same spring with both ends free and a mass m fixed at each free end. Each of the spring is stretched by the same force F . The mass in case (a) and the masses in case (b) are then released. Which of the following statements are true?

- A. While oscillating the maximum extension of the spring is more in case (a) than in case (b).
- B. The maximum extension of the spring is same in both cases.
- C. The time period of oscillation is the same in both cases.
- D. The time period of oscillation in case (a) is $\sqrt{2}$ time that in case (b).

Answer: B::D



Watch Video Solution

16. When the point of suspension of pendulum is moved, its period of oscillation

A. decreases when it moves vertically upwards with an acceleration a

B. decreases when it moves vertically downwards with acceleration greater than $2g$.

C. Increases when it moves horizontally with acceleration a

D. all of the above.

Answer: A::B



Watch Video Solution

17. The displacement time relation for a particle can be expressed as $y = 0.5[\cos^2(n\pi t) - \sin^2(n\pi t)]$ This relation shows that

- A. the particle executing a SHM with amplitude 0.5m
- B. the particle is executing a SHM with frequency n time that of a second's pendulum
- C. the particle is executing a SHM and the velocity in its mean position is $(n\pi) \frac{m}{s}$
- D. the particle is not executing a SHM at all

Answer: A::C



Watch Video Solution


18. At two particular closest instant of time t_1 and t_2 the displacements of a particle performing SHM are equal. At these instant

- A. instantaneous speed are equal
- B. instantaneous acceleration are equal
- C. phase of the motion are unequal
- D. kinetic energies are equal

Answer: A::B::C::D



Watch Video Solution



19.

Two blocks connected by a spring rest on a smooth horizontal plane as shown in Fig. A constant force F start acting on block m_2 as shown in the figure. Which of the following statements are not correct?

A. Length of the spring increases continuously if

$$m_1 > m_2.$$

B. Blocks start performing SHM about centre of mass of the system, which moves rectilinearly with constant acceleration.

C. Blocks start performing oscillation about centre of mass of the system with increasing amplitude.

D. Acceleration of m_2 is maximum at initial moment of time only.

Answer: A::C::D



Watch Video Solution

20. A block of mass m is suspended by a rubber cord of natural length $l = \frac{mg}{k}$, where k is force constant of the cord. The block is lifted upwards so that the cord becomes just tight and then block is released suddenly. Which of the following will not be true?

- A. Block performs periodic motion with amplitude greater than l .
- B. Block performs SHM with amplitude equal to l .
- C. Blocks will never return to the position from where it was released.
- D. Angular frequency ω is equal to $1 \frac{\text{rad}}{\text{s}}$

Answer: A::C::D

 [Watch Video Solution](#)

21. The displacement (x) of a particle as a function of time (t) is given by

$$x = a \sin(bt + c)$$

Where a, b and c are constant of motion. Choose the correct statements from the following.

- A. The motion repeats itself in a time interval of $\frac{2\pi}{b}$
- B. The energy of the particle remains constant.
- C. The velocity of the particle zero at $x = \pm a$
- D. The acceleration of the particle is zero at $x = \pm a$

Answer: A::B::C

 [Watch Video Solution](#)

22. A simple pendulum is oscillating between extreme position P and Q about the mean position O. Which of the

following statements are true about the motion of pendulum?

A. At point O, the acceleration of the bob is different from zero.

B. The acceleration of the bob is constant throughout the oscillation.


C. The tension in the string is constant throughout the oscillation.

D. The tension is maximum at O and minimum at P or Q.

Answer: A:D



Watch Video Solution



23.

A cylindrical block of density d stays fully immersed in a beaker filled with two immiscible liquids of different densities d_1 and d_2 . The block is in equilibrium with half of it in liquid 1 and the other half in liquid 2 as shown in the Fig. If the block is given a displacement downwards released, then neglecting friction study the following statements.

A. It executes simple harmonic motion.

B. Its motion is periodic but not simple harmonic.

C. The frequency of oscillation is independent of the size of the cylinder.

D. The displacement of the centre of the cylinder is symmetric about its equilibrium position.

Answer: A::D



Watch Video Solution

24. A mass of 0.2 hg is attached to the lower end of a massless spring of force constant $200 \frac{N}{m}$ the upper end of which is fixed to a rigid support. Study the following statements.

- A. In equilibrium the spring will be stretched by 1 cm.
- B. If the mass is raised till the spring becomes unstretched and then released, it will go down by 2 cm before moving upwards.
- C. The frequency of oscillation will be nearly 5 Hz.
- D. If the system is taken to the moon, the frequency of oscillation will be the same as that on the earth.

Answer: A::B::C::D



Watch Video Solution

25. A spring of spring constant k stores 5 J of energy when stretched by 25 cm. It is kept vertical with one end fixed. A

mass m is attached to the other end. It makes 5

Oscillation per second. Then

A. $m = 0.16\text{kg}$

B. $m = 0.32\text{kg}$

C. $k = 160\frac{N}{m}$

D. $k = 320\frac{N}{m}$

Answer: A:C



Watch Video Solution

26. A particle is subjected to SHM as given by equation

$x_1 = A \sin \omega t$ and $x_2 = A_2 \sin\left(\omega t + \frac{\pi}{3}\right)$. The maximum

acceleration and amplitude of the resultant motion are

a_{\max} and A , respectively Then.

A. $a_{\max} = \omega^2 \sqrt{A_1^2 + A_2^2 + A_1 A_2}$

B. $a_{\max} = \omega^2 \sqrt{A_1 A_2}$

C. $A = A_1 + A_2$

D. $A = \sqrt{A_1^2 + A_2^2 + A_1 A_2}$

Answer: A:D



Watch Video Solution

27. Three simple harmonic motions in the same direction each of amplitude a and periodic time T , are superposed.

The first and second and the second and third differ in

phase from each other by $\frac{\pi}{4}$, with the first and third not being identical . Then.

A. the resultant motion is not simple harmonic

B. the resultant amplitude is $(\sqrt{2} + 1)a$


C. the phase difference between the second SHM and the resultant motion is zero.

D. the energy in the resultant motion is three times the energy in each separate SHM.

Answer: B::C



Watch Video Solution



28.

A horizontal platform with a mass m placed on it is executing SHM along y -axis. If the amplitude of oscillation is 2.5 cm, the minimum period of the motion for the mass not to be detached from the platform $\left(g = 10 \frac{m}{s^2}\right)$

A. $\frac{10}{\pi} s$

B. $\frac{\pi}{10} s$

C. $\frac{\pi}{\sqrt{10}} s$

D. $\frac{1}{\sqrt{10}} s$

Answer: B::D



Watch Video Solution

29. The speed v of a particle moving along a straight line, when it is at a distance (x) from a fixed point of the line is given by $v^2 = 108 - 9x^2$ (all equation are in CGS units):

A. The motion is uniformly accelerated along the straight line

B. the magnitude of the acceleration at a distance 3 cm from the point is $27 \frac{cm}{s^2}$

C. the motion is simple harmonic about the given fixed point

D. the maximum displacement from the fixed point is 4

cm

Answer: B::C

 **Watch Video Solution**

30. For a body executing SHM with amplitude A , time period T , maximum velocity v_{\max} and phase constant zero, which of the following statements are correct for $0 \leq t \leq \frac{T}{4}$ (y is displacement from mean position)?

A. At $y = \left(\frac{A}{2}\right), v > \left(\frac{v_{\max}}{2}\right)$

B. For $v = \left(\frac{v_{\max}}{2}\right), y = < \left(\frac{A}{2}\right)$

C. For $t = \left(\frac{T}{8}\right), y > \left(\frac{A}{2}\right)$

D. for $y = \left(\frac{A}{2}\right), y < \left(\frac{T}{8}\right)$.

Answer: A::B::C::D

 **Watch Video Solution**

31. If y , v and a represent displacement velocity and acceleration at any instant for a particle executing SHM, which of the following statements are true?

- A. v and y may have same direction.
- B. v and a may have same direction.
- C. a and y may have same direction.
- D. a and v may have opposite direction.

Answer: A::B::D



Watch Video Solution

32. The time period of a particle in simple harmonic motion is T . Assume potential energy at mean position to be zero. After a time of $\frac{T}{6}$ it passes its mean position its,

A. velocity will be one half its mean position, its

B. displacement will be one half its amplitude

C. acceleration will be nearly 85% of its maximum

acceleration

D. $KE = PE$

Answer: A::C



Watch Video Solution

33. The potential energy of a particle of mass 0.1 kg, moving along the x -axis, is given by $U = 5x(x - 4)J$, where x is in meter. It can be concluded that

A. the particle is acted upon by a constant force

B. the speed of the particle is maximum at $x = 2m$

C. the particle executes SHM

D. the period of oscillation of the particle $\left(\frac{\pi}{5}\right) s$

Answer: B::C::D



Watch Video Solution

34. A horizontal plank has a rectangular block placed on it. The plank starts oscillating vertically and simple harmonically with an amplitude of 40 cm. The block just loses contact with the plank when the latter is at momentary rest. Then.

A. the period of oscillation is $\left(\frac{2\pi}{5}\right)$

B. the block weight double its weight, when the plank is at one of the positions of momentary rest.

C. the block weighs 0.1 times its weight on the plank halfway up.

D. the block weighs 1.5 times its weight on the plank
halfway down.

Answer: A::B::C::D

 [Watch Video Solution](#)

35. A particle moves in the X-Y plane according to the equation $\vec{r} = \left(\vec{i} + 2\vec{j} \right) A \cos \omega t$. The motion of the particle is

A. on a straight line

B. on an ellipse

C. periodic

D. simple harmonic.

Answer: A::C::D



Watch Video Solution

Assertion Reasoning

1. Q. Statement I: The total energy of a particle performing simple harmonic motion could be negative. Statement II: Potential energy of a system could be magnetic.

A. Statement I is true statement II is true, Statement II is a correct explanation for Statement I.

- B. Statement I is true statement II is true, Statement II is NOT a correct explanation for Statement I.
- C. Statement I is true, Statement II is false
- D. Statement I is false, Statement II is true

Answer: A



Watch Video Solution




Fig. 4.181

2.

Q. Statement I: Two cubical blocks of same material and of sides a and $2a$, respectively are attached rigidly and symmetrically to each other as shown. The system of two blocks is floating in water in such a way that upper surface of bigger blocks is just submerged in the water. If the system of blocks is displaced slightly in vertical directions, then the amplitude of oscillation on either side of equilibrium position would be different. Statement II:

The force constant on two sides of equilibrium position in the above described situation is different.

A. Statement I is true statement II is true, Statement II is a correct explanation for Statement I.

B. Statement I is true statement II is true, Statement II is NOT a correct explanation for Statement I.

C. Statement I is true, Statement II is false

D. Statement I is false, Statement II is true

Answer: A



Watch Video Solution




Fig. 4.182

3.

Q. Statement I Three pendulums are suspended from ceiling as shown in Fig. These three pendulums are set to oscillate as shown by arrows, and it is found that all three have same time period. Now, all three are taken to a place where acceleration due to gravity changes to $\frac{4}{9}$ th of its value at the first place. If spring pendulum makes 60 cycles in a given time at this place, then torsion pendulum and simple pendulum will also make 60 oscillation in same

(given) time interval. Statement II: Time period of torsion pendulum is independent of acceleration due to gravity.

A. Statement I is true statement II is true, Statement II is a correct explanation for Statement I.

B. Statement I is true statement II is true, Statement II is NOT a correct explanation for Statement I.

C. Statement I is true, Statement II is false

D. Statement I is false, Statement II is true

Answer: D



Watch Video Solution

4. A. Statement I is true statement II is true, Statement II is a correct explanation for Statement I.

b. Statement I is true, Statement II is true, Statement II is NOT a correct explanation for Statement I.

c. Statement I is true, Statement II is false.

d. Statement I is false, Statement II is true.

Q. Statement I: A circular metal hoop is suspended on the edge by a hook. the hoop can oscillate side to side in the plane of the hoop, or it can oscillate back and forth in a direction perpendicular to the plane of the hoop. The time period of oscillation would be more when oscillation are carried out in the plane of hoop.

Statement II: Time period of physical pendulum is more if moment of inertial of the rigid body about corresponding axis passing through the pivoted point is more.

- A. Statement I is true statement II is true, Statement II is a correct explanation for Statement I.
- B. Statement I is true statement II is true, Statement II is NOT a correct explanation for Statement I.
- C. Statement I is true, Statement II is false
- D. Statement I is false, Statement II is true

Answer: A

 [Watch Video Solution](#)

5. STATEMENT-1 : In simple pendulum performing $S. H. M$, net acceleration is always between tangential and radial acceleration except at lowest point.

STATEMETN-2 : At lowest point tangential acceleration is zero.

A. Statement I is true statement II is true, Statement II is a correct explanation for Statement I.

B. Statement I is true statement II is true, Statement II is NOT a correct explanation for Statement I.

C. Statement I is true, Statement II is false

D. Statement I is false, Statement II is true

Answer: D



Watch Video Solution

6. Statement I: If the amplitude of a simple harmonic oscillator is doubled, its total energy becomes four times.

Statement II: The total energy is directly proportional to the square of the amplitude of vibration of the harmonic oscillator.

A. Statement I is true statement II is true, Statement II is a correct explanation for Statement I.

B. Statement I is true statement II is true, Statement II is NOT a correct explanation for Statement I.

C. Statement I is true, Statement II is false

D. Statement I is false, Statement II is true

Answer: A

Comprehension

1. One end of an ideal spring is fixed to a wall at origin O and axis of spring is parallel to x -axis. A block of mass $m = 1\text{kg}$ is attached to free end of the spring and it is performing SHM. Equation of position of the block in co-ordinate system shown in figure is $x = 10 + 3\sin(10t)$. Here, t is in second and x in cm . Another block of mass $M = 3\text{kg}$, moving towards the origin with velocity 30cm/s collides with the block performing SHM at $t = 0$ and gets stuck to it. Calculate



(a) new amplitude of oscillations,

(b) new equation for position of the combined body,

(c) loss of energy during collision. Neglect friction.

A. $20 \frac{\text{rad}}{\text{s}}$

B. $5 \frac{\text{rad}}{\text{s}}$

C. $100 \frac{\text{rad}}{\text{s}}$

D. $50 \frac{\text{rad}}{\text{s}}$


Answer: B



Watch Video Solution

2. One end of an ideal spring is fixed to a wall at origin O and axis of spring is parallel to x-axis. A block of mass

$m = 1\text{kg}$ is attached to free end of the spring and it is performing SHM. Equation of position of the block in coordinate system shown in figure is $x = 10 + 3\sin(10t)$. Here, t is in second and x in cm . Another block of mass $M = 3\text{kg}$, moving towards the origin with velocity 30cm/s collides with the block performing SHM at $t = 0$ and gets stuck to it. Calculate



- (a) new amplitude of oscillations,
- (b) new equation for position of the combined body,
- (c) loss of energy during collision. Neglect friction.

A. 3 cm

B. 20 cm

C. 10 cm

D. 100 cm

Answer: A



Watch Video Solution

3. One end of an ideal spring is fixed to a wall at origin O and axis of spring is parallel to x -axis. A block of mass $m = 1\text{kg}$ is attached to free end of the spring and it is performing SHM. Equation of position of the block in coordinate system shown in figure is $x = 10 + 3\sin(10t)$. Here, t is in second and x in cm . Another block of mass $M = 3\text{kg}$, moving towards the origin with velocity 30cm/s collides with the block performing SHM at $t = 0$

and gets stuck to it. Calculate



- (a) new amplitude of oscillations,
(b) new equation for position of the combined body,
(c) loss of energy during collision. Neglect friction.

A. $(10 + 3 \sin 5t) \text{ cm}$

B. $(10 - 3 \sin 5t) \text{ cm}$


C. $(10 + 3 \cos 10t) \text{ cm}$

D. $(10 - 3 \cos 10t) \text{ cm}$

Answer: B



Watch Video Solution



4.

A block of mass m is connected to a spring constant k and is at rest in equilibrium as shown. Now, the block is displaced by h below its equilibrium position and imparted a speed v_0 towards down as shown in the Fig. As a result of the jerk, the block executes simple harmonic motion about its equilibrium position. Based on this information, answer the following question.

Q. The amplitude of oscillation is

A. h


B. $\sqrt{\frac{mv_0^2}{k} + h^2}$

C. $\sqrt{\frac{m}{k}v_0} + h$

D. none of these

Answer: B

 Watch Video Solution



5.

A block of mass m is connected to a spring constant k and

is at rest in equilibrium as shown. Now, the block is displaced by h below its equilibrium position and imparted a speed v_0 towards down as shown in the Fig. As a result of the jerk, the block executes simple harmonic motion about its equilibrium position. Based on this information, answer the following question.

Q. The equation for the simple harmonic motion is

A. $y = -A \sin \left[\sqrt{\frac{k}{m}} t + \sin^{-1} \left(\frac{h}{A} \right) \right]$

B. $y = -A \cos \left[\sqrt{\frac{k}{m}} t + \sin^{-1} \left(\frac{h}{A} \right) \right]$


C. $y = A \sin \left[\sqrt{\frac{k}{m}} t + \cos^{-1} \left(\frac{h}{A} \right) + \frac{\pi}{2} \right]$

D. $y = A \sin \left[\sqrt{\frac{k}{m}} t + \cos^{-1} \left(\frac{h}{A} \right) + \frac{\pi}{4} \right]$

Answer: A



Watch Video Solution



6.

A block of mass m is connected to a spring constant k and is at rest in equilibrium as shown. Now, the block is displaced by h below its equilibrium position and imparted a speed v_0 towards down as shown in the Fig. As a result of the jerk, the block executes simple harmonic motion about its equilibrium position. Based on this information, answer the following question.

Q. Find the time taken by the block to cross the mean position for the first time.

A.
$$\frac{2\pi - \cos^{-1}\left(\frac{h}{A}\right)}{\sqrt{\frac{k}{m}}}$$

B.
$$\frac{\frac{\pi}{2} - \cos^{-1}\left(\frac{h}{A}\right)}{\sqrt{\frac{k}{m}}}$$


C.
$$\frac{\pi - \sin^{-1}\left(\frac{h}{A}\right)}{\sqrt{\frac{k}{m}}}$$

D.
$$\frac{\pi - \sin^{-1}\left(\frac{h}{A}\right)}{2\sqrt{\frac{k}{m}}}$$

Answer: C



Watch Video Solution



7.

A block of mass m is connected to a spring of spring constant k as shown in Fig. The block is found at its equilibrium position $t = 1\text{ s}$ and it has a velocity of $+0.25\frac{m}{s}$ at $t = 2\text{ s}$. The time period of oscillation is 6 s .

Based on the given information answer the following question: Q. the amplitude of oscillation is

A. $\frac{3}{2\pi} \text{ cm}$


B. $3m$

C. $\frac{1}{\pi}m$

D. $1.5m$

Answer: A

 Watch Video Solution



8.

A block of mass m is connected to a spring of spring constant k as shown in Fig. The block is found at its equilibrium position $t = 1s$ and it has a velocity of

$+0.25 \frac{m}{s}$ at $t = 2s$. The time period of oscillation is 6 s.

Based on the given information answer the following question:

Q. Determine the velocity of particle at $t = 5s$.

A. $-0.4 \frac{m}{s}$

B. $0.5 \frac{m}{s}$


C. $-0.25 \frac{m}{s}$

D. none of these

Answer: C



Watch Video Solution



9.

In physical pendulum, the time period for small oscillation

is given by $T = 2\pi \sqrt{\frac{I}{Mgd}}$ where I is the moment of

inertial of the body about an axis passing through a

pivoted point O and perpendicular to the plane of

oscillation and d is the separation point between centre

of gravity and the pivoted point. In the physical pendulum a special point exists where if we concentrate the entire mass of body, the resulting simple pendulum (w.r.t. pivot point O) will have the same time period as that of physical pendulum. This point is termed centre of oscillation.

$$T = 2\pi \sqrt{\frac{I}{Mgd}} = 2\pi \sqrt{\frac{L}{g}}$$

Moreover, this point

possesses two other important remarkable properties:

Property I: Time period of physical pendulum about the centre of oscillation (if it would be pivoted) is same as in the original case.

Property II: If an impulse is applied at the centre of oscillation in the plane of oscillation, the effect of this impulse at pivoted point is zero. Because of this property, this point is also known as the centre of percussion. From the given information answer the following question:

Q. A uniform rod of mass M and length L is pivoted about point O as shown in Figgt It is slightly rotated from its mean position so that it performs angular simple harmonic motion. For this physical pendulum, determine the time period oscillation.

A. $2\pi\sqrt{\frac{L}{g}}$

B. $\pi\sqrt{\frac{7L}{3g}}$

C. $2\pi\sqrt{\frac{2l}{3g}}$

D. none of these

Answer: B



Watch Video Solution

10. In physical pendulum, the time period for small oscillation is given by $T = 2\pi\sqrt{\frac{I}{Mgd}}$ where I is the moment of inertial of the body about an axis passing through a pivoted point O and perpendicular to the plane of oscillation and d is the separation point between centre of gravity and the pivoted point. In the physical pendulum a speacial point exists where if we concentrate the entire mass of body, the resulting simple pendulum (w.r.t. pivot point O) will have the same time period as that of physical pendulum This point is termed centre of oscillation.

$$T = 2\pi\sqrt{\frac{I}{Mgd}} = 2\pi\sqrt{\frac{L}{g}} \quad \text{Moreover, this point}$$


possesses two other important remarkable properties:

Property I: Time period of physical pendulum about the

centre of oscillation (if it would be pivoted) is same as in the original case.

Property II: If an impulse is applied at the centre of oscillation in the plane of oscillation, the effect of this impulse at pivoted point is zero. Because of this property, this point is also known as the centre of percussion. From the given information answer the following question:

Q. For the above question, locate the centre of oscillation.



- A. $\frac{L}{4}$ from O (down)
- B. $\frac{L}{4}$ from O (up)
- C. $\frac{2L}{3}$ from O (down)
- D. $\frac{7L}{12}$ from O (down)

Answer: D



Watch Video Solution

11. In physical pendulum, the time period for small oscillation is given by $T = 2\pi \sqrt{\frac{I}{Mgd}}$ where I is the moment of inertial of the body about an axis passing through a pivoted point O and perpendicular to the plane of oscillation and d is the separation point between centre of gravity and the pivoted point. In the physical pendulum a special point exists where if we concentrate the entire mass of body, the resulting simple pendulum (w.r.t. pivot point O) will have the same time period as that of physical pendulum This point is termed centre of

oscillation.

$$T = 2\pi \sqrt{\frac{I}{Mgd}} = 2\pi \sqrt{\frac{L}{g}}$$

Moreover, this point


possesses two other important remarkable properties:

Property I: Time period of physical pendulum about the centre of oscillation (if it would be pivoted) is same as in the original case.

Property II: If an impulse is applied at the centre of oscillation in the plane of oscillation, the effect of this impulse at pivoted point is zero. Because of this property, this point is also known as the centre of percussion. From the given information answer the following question:

Q. If an impulse J is applied at the centre of oscillation in the plane of oscillation, then angular velocity of the rod

will be .





- A. $\frac{4J}{ML}$
- B. $\frac{2J}{ML}$
- C. $\frac{3J}{2ML}$
- D. $\frac{J}{ML}$

Answer: A



Watch Video Solution



12.

A block of mass m is suspended from one end of a light spring as shown. The origin O is considered at distance

equal to natural length of the spring from the ceiling and vertical downwards direction as positive y -axis. When the system is in equilibrium a bullet of mass $\frac{m}{3}$ moving in vertical up wards direction with velocity v_0 strikes the block and embeds into it. As a result, the block (with bullet embedded into it) moves up and start oscillating. Based on the given information, answer the following question:

Q. Mark out the correct statements (s).

A. the block bullet system performs SHM about

$$y = \frac{mg}{k}.$$

B. The block bullet system performs oscillatory motion

but not SHM about $y = \frac{mg}{k}$.

C. The block bullet system performs SHM about

$$y = \frac{4mg}{3k}.$$



D. The block bullet system performs oscillatory motion

but not SHM about $y = \frac{4mg}{3k}$.

Answer: C



Watch Video Solution



13.

A block of mass m is suspended from one end of a light spring as shown. The origin O is considered at distance

equal to natural length of the spring from the ceiling and vertical downwards direction as positive y-axis. When the system is in equilibrium a bullet of mass $\frac{m}{3}$ moving in vertical up wards direction with velocity v_0 strikes the block and embeds into it. As a result, the block (with bullet embedded into it) moves up and start oscillating. Based on the given information, answer the following question:

Q. The amplitude of oscillation would be

A. $\sqrt{\left(\frac{4mg}{3k}\right)^2 + \frac{mv_0^2}{12k}}$

B. $\sqrt{\frac{mv_0^2}{12k} + \left(\frac{mg}{3k}\right)^2}$



C. $\sqrt{\frac{mv_0^2}{6k} + \left(\frac{mg}{k}\right)^2}$

D. $\sqrt{\frac{mv_0^2}{6k} + \left(\frac{4mg}{3k}\right)^2}$

Answer: B



Watch Video Solution



14.

A block of mass m is suspended from one end of a light spring as shown. The origin O is considered at distance

equal to natural length of the spring from the ceiling and vertical downwards direction as positive y-axis. When the system is in equilibrium a bullet of mass $\frac{m}{3}$ moving in vertical up wards direction with velocity v_0 strikes the block and embeds into it. As a result, the block (with bullet embedded into it) moves up and start oscillating. Based on the given information, answer the following question:

Q. The time taken by the block bullet system to move from $y = \frac{mg}{k}$ (initial equilibrium position) to $y = 0$ (natural length of spring) is (A represents the amplitude of motion)

A. $\sqrt{\frac{4m}{3k}} \left[\cos^{-1} \left(\frac{mg}{3kA} \right) - \cos^{-1} \left(\frac{4mg}{3kA} \right) \right]$


B. $\sqrt{\frac{3k}{4m}} \left[\cos^{-1} \left(\frac{mg}{3kA} \right) - \cos^{-1} \left(\frac{4mg}{3kA} \right) \right]$

C. $\sqrt{\frac{4m}{6k}} \left[\sin^{-1} \left(\frac{4mg}{3kA} \right) - \sin^{-1} \left(\frac{mg}{3kA} \right) \right]$

D. none of these

Answer: A

 **Watch Video Solution**



188

15.

Two identical blocks A and B, each of mass $m = 3 \text{ kg}$, are connected with the help of an ideal spring and placed on a smooth horizontal surface as shown in Fig. Another

identical blocks C moving velocity $v_0 = 0.6 \frac{m}{s}$ collides with A and sticks to it, as a result, the motion of system takes place in some way

Based on this information answer the following questions:

Q. After the collision of C and A, the combined body and block B would

A. oscillate about centre of mass of system and centre of mass is at rest.


B. oscillate about centre of mass of system and centre of mass is moving.

C. oscillate but about different location other than the centre of mass.

D. not oscillate.

Answer: B

 Watch Video Solution



188

16.

Two identical blocks A and B, each of mass $m = 3\text{kg}$, are connected with the help of an ideal spring and placed on a smooth horizontal surface as shown in Fig. Another identical blocks C moving velocity $v_0 = 0.6 \frac{m}{s}$ collides

with A and sticks to it, as a result, the motion of system takes place in some way

Based on this information answer the following questions:

Q. Oscillation energy of the system i.e., part of the energy which is oscillation (changing) between potential and kinetic forms is

A. 0.27 J

B. 0.09 J


C. 0.18 J

D. 0.45 J

Answer: B



Watch Video Solution



188

17. Two identical blocks A and B, each of mass $m = 3\text{kg}$, are connected with the help of an ideal spring and placed on a smooth horizontal surface as shown in Fig. Another identical blocks C moving velocity $v_0 = 0.6 \frac{m}{s}$ collides with A and sticks to it, as a result, the motion of system takes place in some way

Based on this information answer the following questions:

Q. The maximum compression of the spring is

A. $3\sqrt{30}\text{mm}$

B. $3\sqrt{20}\text{mm}$

C. $3\sqrt{10}\text{mm}$

D. $3\sqrt{50}\text{mm}$

Answer: C



Watch Video Solution




Fig. 4.189

18.

A small block of mass m is fixed at upper end of a massive vertical spring of spring constant $k = \frac{2mg}{L}$ and natural length $10L$. The lower end of spring is free and is at a height L from fixed horizontal floor as shown. The spring is initially unstressed and the spring block system is released from rest in the shown position.

Q. At the instant the speed of block is maximum the magnitude of force exerted by the spring on the block is

A. $\frac{mg}{2}$

B. mg

C. *zero*

D. none of these

Answer: B

 **Watch Video Solution**




Fig. 4.189

19.

A small block of mass m is fixed at upper end of a massive

vertical spring of spring constant $k = \frac{2mg}{L}$ and natural length $10L$. The lower end of spring is free and is at a height L from fixed horizontal floor as shown. The spring is initially unstressed and the spring block system is released from rest in the shown position.

Q. As the block is coming down, the maximum speed attained by the block is

A. \sqrt{gL}

B. $\sqrt{3gL}$

C. $\frac{3}{2}\sqrt{gL}$

D. $\sqrt{\frac{3}{2}gL}$

Answer: C



Watch Video Solution




Fig. 4.189

20.

A small block of mass m is fixed at upper end of a massive vertical spring of spring constant $k = \frac{2mg}{L}$ and natural length $10L$. The lower end of spring is free and is at a height L from fixed horizontal floor as shown. The spring is initially unstressed and the spring block system is released from rest in the shown position.

Q. Till the blocks reaches its lowest position for the first time, the time duration for which the spring remains compressed is

A. $\pi \sqrt{\frac{L}{2g}} + \sqrt{\frac{L}{4g} \frac{\sin^{-1}(1)}{3}}$

B. $\frac{\pi}{4} \sqrt{\frac{L}{g}} + \sqrt{\frac{L}{4g} \frac{\sin^{-1}(1)}{3}}$



C. $\pi \sqrt{\frac{L}{2g}} + \sqrt{\frac{L}{4g} \frac{\sin^{-1}(2)}{3}}$

D. $\frac{\pi}{2} \sqrt{\frac{L}{2g}} + \sqrt{\frac{L}{4g} \frac{\sin^{-1}(2)}{3}}$

Answer: B



Watch Video Solution



21.

A 100 g block is connected to a horizontal massless spring of force constant $25.6 \frac{N}{m}$. As shown in Fig. the block is free to oscillate on a horizontal frictionless surface. The block is displaced 3 cm from the equilibrium position and, at $t = 0$, it is released from rest at $x = 0$. It executes simple harmonic motion with the positive x -direction indicated in Fig. The position-time ($x-t$)

graph of motion of the block is as shown in Fig.



Q. When the block is at position A on the graph, its

- A. position and velocity both are negative
- B. position is positive and velocity is negative
- C. position is negative and velocity is positive
- D. position and velocity both the positive

Answer: B



Watch Video Solution



22.

A 100 g block is connected to a horizontal massless spring of force constant $25.6 \frac{N}{m}$. As shown in Fig. the block is free to oscillate on a horizontal frictionless surface. The block is displaced 3 cm from the equilibrium position and, at $t = 0$, it is released from rest at $x = 0$. It executes simple harmonic motion with the positive x -direction indicated in Fig. The position time ($x - t$) graph of motion of the

block is as shown in Fig.



Q. When the block is at position B on the graph its.

- A. position and velocity both are positive.
- B. position is positive and velocity is negative
- C. position is negative and velocity is positive
- D. position and velocity are negative.

Answer: C



Watch Video Solution



23.

A 100 g block is connected to a horizontal massless spring of force constant $25.6 \frac{N}{m}$. As shown in Fig. the block is free to oscillate on a horizontal frictionless surface. The block is displaced 3 cm from the equilibrium position and, at $t = 0$, it is released from rest at $x = 0$. It executes simple harmonic motion with the positive x -direction indicated in Fig. The position time ($x - t$) graph of motion of the

block is as shown in Fig.



When the block is at position C on the graph, its

- A. velocity is maximum and acceleration is zero
- B. velocity is minimum and acceleration is zero
- C. velocity is zero and acceleration is negative
- D. velocity is zero and acceleration is positive.

Answer: C



Watch Video Solution



24.

A 100 g block is connected to a horizontal massless spring of force constant $25.6 \frac{N}{m}$. As shown in Fig. the block is free to oscillate on a horizontal frictionless surface. The block is displaced 3 cm from the equilibrium position and, at $t = 0$, it is released from rest at $x = 0$. It executes simple harmonic motion with the positive x -direction indicated in Fig. The position-time ($x - t$) graph of motion of the block is as shown in Fig.

Position of the block as a function of time can now be expressed as

A. $x = 3 \cos\left(16t + \frac{\pi}{2}\right) \text{cm}$

B. $x = 3 \cos\left(16t + \frac{\pi}{3}\right) \text{cm}$


C. $x = 3.5 \cos\left(16t + \frac{\pi}{6}\right) \text{cm}$

D. $x = 3.2 \cos\left(16t + \frac{\pi}{4}\right) \text{cm}$


Answer: C



Watch Video Solution



(a)



(b)

25.

A 100 g block is connected to a horizontal massless spring of force constant $25.6 \frac{N}{m}$. As shown in Fig. the block is free to oscillate on a horizontal frictionless surface. The block is displaced 3 cm from the equilibrium position and, at $t = 0$, it is released from rest at $x = 0$. It executes simple harmonic motion with the positive x -direction indicated in Fig. The position-time ($x-t$) graph of motion of the block is as shown in Fig.

Velocity of the block as a function of time can be expressed as

A. $v = -\sin 48 \left(16t \frac{\pi}{2} \right) \frac{cm}{s}$

B. $v = -48 \sin \left(16t \frac{\pi}{3} \right) \frac{cm}{s}$


C. $v = -56 \sin \left(16t \frac{\pi}{4} \right) \frac{cm}{s}$

D. $v = -56 \left(16t \frac{\pi}{6} \right) \frac{cm}{s}$

Answer: D



Watch Video Solution



26.

A spring having a spring constant k is fixed to a vertical wall as shown in Fig. A block of mass m moves with velocity v towards the spring from a parallel wall opposite to this wall. The mass hits the free end of the spring compressing it and is decelerated by the spring force and comes to rest and then turns the spring is decelerated by the spring force and comes to rest and then turns back till the spring acquires its natural length and contact with the spring is broken. In this process, it regains its angular speed in the opposite direction and makes a perfect elastic collision on the opposite left wall and starts

moving with same speed as before towards right. The above processes are repeated and there is periodic oscillation.

Q. What is the maximum compression produced in the spring?

A. $v\sqrt{\frac{m}{k}}$

B. $\sqrt{\frac{m}{k}}$


C. $v\sqrt{mk}$

D. $v\sqrt{\frac{k}{m}}$

Answer: A



Watch Video Solution



27.

A spring having a spring constant k is fixed to a vertical wall as shown in Fig. A block of mass m moves with velocity v towards the spring from a parallel wall opposite to this wall. The mass hits the free end of the spring compressing it and is decelerated by the spring force and comes to rest and then turns the spring is decelerated by the spring force and comes to rest and then turns back till the spring acquires its natural length and contact with the spring is broken. In this process, it regains its angular speed in the opposite direction and makes a perfect elastic collision on the opposite left wall and starts

moving with same speed as before towards right. The above processes are repeated and there is periodic oscillation.

Q. What is the time period of oscillation ?

A. $\pi \sqrt{\frac{m}{k}}$

B. $\sqrt{\frac{\pi m}{k}} + \frac{L}{m}$


C. $\pi \sqrt{\frac{m}{k}} + \frac{2L}{v}$

D. $\pi \sqrt{\frac{m}{k}} + \frac{L}{v}$

Answer: C



Watch Video Solution



28.

A and B are two fixed points at a distance $3l$ apart. A particle of mass m placed at a point P experiences the force $2\left(\frac{mg}{l}\right)\vec{P}A$ and the force $\left(\frac{mg}{l}\right)\vec{P}B$ simultaneously. Initially at $t = 0$, the particle is projected from A towards B with speed $3\sqrt{gl}$.

Q. The particle moves simple harmonically with period T and amplitude A .

A. $A = 2l, T = 2\pi\sqrt{\frac{l}{g}}$


B. $A = 3l, T = 2\pi\sqrt{\frac{l}{2g}}$

C. $A = 2l, T = 2\pi\sqrt{\frac{l}{3g}}$

D. $A = l, T = 2\pi\sqrt{\frac{l}{g}}$

Answer: C

 Watch Video Solution



29.

A and B are two fixed points at a distance $3l$ apart. A particle of mass m placed at a point P experiences the force $2\left(\frac{mg}{l}\right)\vec{P}A$ and the force $\left(\frac{mg}{l}\right)\vec{P}B$ simultaneously. Initially at $t = 0$, the particle is projected from A towards B with speed $3\sqrt{gl}$.

Q. The instant t at which the particle arrives at B in terms of the periodic time T will be

A. $t = \frac{T}{2}$


B. $t = \frac{T}{3}$

C. $t = \frac{T}{4}$

D. $t = \frac{2T}{3}$

Answer: B

 Watch Video Solution



30.

A and B are two fixed points at a distance $3l$ apart. A particle of mass m placed at a point P experiences the force $2\left(\frac{mg}{l}\right)\vec{P}A$ and the force $\left(\frac{mg}{l}\right)\vec{P}B$ simultaneously. Initially at $t = 0$, the particle is projected

from A towards B with speed $3\sqrt{gl}$.

Q. The velocity of the particle when it reaches B will be

A. zero

B. $3\sqrt{gl}$


C. $2\sqrt{gl}$

D. \sqrt{gl}

Answer: A



Watch Video Solution



31.

A body of mass m is attached by an inelastic string to a suspended spring of spring constant k . Both the string

and the spring have negligible mass and the string is inextensible and of length L . Initially, the mass m is at rest.

Q. If the mass m is now raised up to point A (the top end of the string see fig. and allowed to fall from rest, the maximum extension of the spring in the subsequent motion will be

A. L

B. $\frac{mg}{k}$

C. $\frac{mg}{k} \sqrt{1 + \frac{2kL}{mg}}$

D. $\frac{mg}{k} \left[1 + \sqrt{1 + \frac{2kL}{mg}} \right]$

Answer: D



Watch Video Solution

32. A body of mass m is attached by an inelastic string to a suspended spring of spring constant k . Both the string and the spring have negligible mass and the string is inextensible and of length L . Initially, the mass m is at rest.

If the mass m , from the initial position of rest is pulled down a distance A and then released, assuming that the string remains taut throughout the motion, the maximum (downwards) acceleration of the oscillating body will be

A. $\frac{kA}{m}$

B. $\frac{kA}{2m}$

C. $\frac{g}{2}$

D. g

Answer: A

 [Watch Video Solution](#)

33. A body of mass m is attached by an inelastic string to a suspended spring of spring constant k . Both the string and the spring have negligible mass and the string is inextensible and of length L . Initially, the mass m is at rest.

The largest amplitude A_{\max} , for which the string will remain taut throughout the motion is

A. $\frac{mg}{2k}$

B. $\frac{mg}{k}$

C. $\frac{2mg}{3k}$


D. L

Answer: B



Watch Video Solution

Integer



1.

Two uniform ropes having linear mass densities m and $4m$ are joined to form a closed loop. The loop is hanging over a fixed frictionless small pulley with the lighter rope above as shown in the fig. (in the figure equilibrium

position is shown). Now if point A (joint) is slightly displaced in downward direction and released, it is found that the loop performs SHM with the period of oscillation equal to N . Find the value of N (take $l = \frac{150m}{4\pi^2}$, $g = 10\frac{m}{s^2}$).



[Watch Video Solution](#)

2. In the figure shown, mass $2m$ connected with a spring of force constant k is at rest and in equilibrium. A particle of mass m is released from height $4.5mg/k$ from $2m$. The particle stick to the block. Neglecting the duration of collision find time from the release of m to the moment

when the spring has maximum compression.





Fig. 4.141



Watch Video Solution



3.

Two simple pendulums A and B having lengths l and $\frac{l}{4}$ respectively are released from the position as shown in Fig. Calculate the time (in seconds) after which the two strings become parallel for the first time. (Take $l = \frac{90}{\pi^2}$ m and $g = 10 \frac{m}{s^2}$).

 [Watch Video Solution](#)

value of λ .





Fig. 4.207

4.

A weightless rigid rod with a small iron bob at the end is hinged at point A to the wall so that it can rotate in all directions. The rod is kept in the horizontal position by a vertical inextensible string of length 20 cm , fixed at its midpoint. The bob is displaced slightly perpendicular to the plane of the rod and string. Find period of small oscillations of the system in the form $\frac{\pi X}{10} \text{ s}$ and fill the value of X.

 Watch Video Solution



5.


A rod of mass m and length l hinged at one end is connected by two springs of spring constant k_1 and k_2 so that it is horizontal at equilibrium. What is the angular

frequency of the system? (in $\frac{rad}{s}$) (Take $l = 1m, b = \frac{1}{4}m,$

$$K_1 = 16 \frac{N}{m}, K_2 = 61 \frac{N}{m}.$$



Watch Video Solution




6.

A block of mass m is tied to one end of a string which passes over a smooth fixed pulley A and under a light smooth movable pulley B . The other end of the string is attached to the lower end of a spring of spring constant K_2 . Find the period of small oscillation of mass m about

its equilibrium position (in second). (Take $m = \pi^2 kg$,

$$K_2 k = 4K_1, K_1 = 17 \frac{N}{m}.)$$

 Watch Video Solution




7.

A uniform disc of mass m and radius R is pivoted smoothly at its centre of mass. A light spring of stiffness k is attached with the disc tangentially as shown in the Fig.

Find the angular frequency in $\frac{rad}{s}$ of torsional oscillation of the disc. (Take $m = 5kg$ and $K = 10 \frac{N}{m}$.)



Watch Video Solution




8.

A uniform disc of mass m and radius $R = \frac{80}{23\pi^2}m$ is pivoted smoothly at P . If a uniform disc of mass m and radius R is welded at the lowest point of the disc, find the period of *SHM* of the system (disc + ring). (in seconds)



Watch Video Solution




9.

In the arrangement shown in Fig. Pulleys are small and light and spring are ideal and $K_1 = 25\left(\pi^2\right) \frac{N}{m}$, $K_2 = 2K_1$, $K_3 =$ and $K_4 = 4K_1$ are the force constant of the spring. Calculate the period of small vertical oscillation of block of mass $m = 3\text{kg}$.



Watch Video Solution

of the figure? Neglect effects of gravity.



10.

A small body of mass m is connected to two horizontal spring of elastic constant k , natural length $\frac{3d}{4}$. In the equilibrium position both springs are stretched to length d , as shown in Fig. What will be the ratio of period of the motion $\left(\frac{T_b}{T_a}\right)$ if the body is displaced horizontally by a small distance where T_a is the time period when the particle oscillates along the line of spring T_b is time plane of the figure? Neglect effect of gravity.



Watch Video Solution

Subjective type

1. Describe the motion of a particle acted upon by the force:

A. $F = 3x + 3$

B. $F = -3x - 3$

C. $F = -3x + 3$

D. $F = -3x - 3$

Answer: A::B::C::D



Watch Video Solution

2. The potential energy (U) of a body of unit mass moving in a one-dimension force field is given by

$$U = (x^2 - 4x + 3) . \text{ All units are in S.I.}$$

A. Find the equilibrium position of the body

B. Show that oscillations of the body about this equilibrium position is simple harmonic motion and find its time period

C. Find the amplitude of oscillations if speed of the body at equilibrium position is $2\sqrt{6}m / s$

D. Find the amplitude of oscillators if speed of the body at equilibrium position is $2\sqrt{6}m / s$.

Answer: A::B::C



Watch Video Solution

Single correct Answer Type

1. A boby is moving in a room with a velocity of $20m/s$ perpendicular to the two walls separated by 5 meters .There is no friction and the collisionnn with the walls are elastic.

- A. Not periodic
- B. Periodic but not smple harmonic
- C. Periodic and simple harmonic
- D. Periodic with variable time period

Answer: B



Watch Video Solution

2. The equation of motion of a particle is $x = a\cos(\alpha t)^2$

.The motion is

- A. periodic but not oscillatory
- B. periodic and oscillatory
- C. oscillatory but not periodic
- D. neither periodic nor oscillatory

Answer: C



Watch Video Solution

3. As the expression involving sine function, which of the following equations does not represent a simple harmonic motion?

A. $y = a \sin \omega t$

B. $y = a \tan \omega t$

C. $y = a \cos \omega t$

D. $y = a \sin \omega t + b \cos \omega t$

Answer: D



Watch Video Solution

4. The displacement of a particle is represented by the equation $y = \sin^3 \omega t$. The motion is

- A. non-periodic
- B. periodic but not simple harmonic
- C. simple harmonic with period $2\pi / \omega$
- D. simple harmonic with period π / ω

Answer: B



Watch Video Solution

5. Value of the acceleration A of a particle moving in simple harmonic motion as a function of its displacement

x are given in the table below .

$A(\text{mm s}^{-2})$	16	8	0	- 8	- 16
$x(\text{mm})$	- 4	- 2	0	2	4

The period of the motion is

A. $\frac{1}{\pi} s$

B. $\frac{2}{\pi} s$

C. $\frac{\pi}{2} s$

D. πs

Answer: D



Watch Video Solution

6. The time taken by a particle performing SHM on a straight line to pass from point A to B where its velocities are same is 2 seconds . After another 2 second it returns to B .The time period of oscillation is (in seconds):

A. 2

B. 4

C. 6

D. 8

Answer: D



Watch Video Solution

7. A particle is performing simple harmonic motion along x – axis with amplitude 4 cm and time period 1.2 sec. The minimum time taken by the particle to move from $x=2$ cm to $x = +4$ cm and back again is given by

A. 0.6 sec

B. 0.4 sec

C. 0.3 sec

D. 0.2 sec

Answer: B



Watch Video Solution

8. Two particles P and Q describe $S.H.M.$ of same amplitude and same frequency f along the same straight line. The maximum distance between the two particles is $a\sqrt{2}$. The phase difference between the two particles is

A. zero

B. $\frac{\pi}{2}$

C. $\frac{\pi}{6}$

D. $\frac{\pi}{3}$

Answer: B



Watch Video Solution

9. A particle executing SHM of amplitude 4 cm and $T = 4s$. The time taken by it to move from position to half the amplitude is

A. 1 sec

B. 1/sec

C. $2/3$ sec


D. $\sqrt{3/2}$ sec

Answer: C



Watch Video Solution

10. Two particles undergo *SHM* along parallel lines with the same time period (T) and equal amplitude. At a particular instant, one particle is at its extreme position while the other is at its mean position. They move in the same direction. They will cross each other after a further time



- A. $T/8$
- B. $3T/8$
- C. $T/6$
- D. $4T/3$

Answer: B



Watch Video Solution

11. Two particles execute *SHM* of same amplitude of 20cm with same period along the same line about the same equilibrium position. If phase difference is $\pi/3$ then the maximum distance between these two will be

A. 10cm

B. 20cm

C. $10\sqrt{2}\text{cm}$

D. $20\sqrt{2}\text{cm}$

Answer: A



Watch Video Solution

12. The maximum acceleration of a particle in SHM is made two times keeping the maximum speed to be constant. It is possible when .

- A. amplitude of oscillation is double while frequency remains constant
- B. amplitude is doubled while frequency is halved
- C. frequency is double while amplitude is halved
- D. frequency is double while amplitude remains constant

Answer: C



Watch Video Solution

13. A particles is SHM is described by the displacement function $x(t) = a \cos(\omega t + \theta)$. If the initial ($t = 0$) position of the particle is 1 cm and its initial velocity is $\pi \text{ cm/s}$. The angular frequency of the particle is $\pi \text{ rad/s}$.

Then its amplitude is

A. 1 cm

B. $\sqrt{2} \text{ cm}$

C. 2 cm

D. 2.5 cm

Answer: B





14. Two particles P and Q describe harmonic motions of same period, same amplitude along the same line about the same equilibrium position O . When P and Q are on opposite sides of O at the same distance from O , they have the same speed of 1.2 m/s in the same direction. When their displacements are the same, they have the same speed at 1.6 m/s in opposite directions. The maximum velocity in m/s of either particle is

A. 2.8

B. 2.5

C. 2.4

D. 2

Answer: D



Watch Video Solution

15. A particle executes simple harmonic motion with an amplitude of 4 cm. At the mean position, the velocity of the particles is 10 cm/s. The distance of the particles from the mean position when its speed becomes 5 cm/s is

A. $\sqrt{3}$ cm

B. $\sqrt{5}$ cm

C. $2(\sqrt{3})$ cm

D. $2(\sqrt{5})$ cm

Answer: C



Watch Video Solution

16. A particle is executing SHM according to the equation $x=A \cos \omega t$. Average speed of the particle during the interval $0 \leq t \leq \frac{\pi}{6\omega}$

A. $\frac{\sqrt{3}A\omega}{2}$

B. $\frac{\sqrt{3}A\omega}{4}$

C. $\frac{3A\omega}{\pi}$

D. $\frac{3A\omega}{\pi} (2 - \sqrt{3})$

Answer: D



Watch Video Solution

17. The total energy of the body executing S.H.M is E . Then .
The kinetic energy when the displacement half of the amplitude is

A. $\frac{E}{2}$

B. $\frac{E}{4}$

C. $\frac{3E}{4}$

D. $\frac{\sqrt{3}}{4}E$

Answer: C



Watch Video Solution

18. A body is executing simple Harmonic Motion . At displacement x its potential energy is E_1 and at a displacement y its potential energy is E_2 . The potential energy E at displacement $(x + y)$ is

A. $\sqrt{E} = \sqrt{E_1} - \sqrt{E_2}$

B. $\sqrt{E} = \sqrt{E_1} + \sqrt{E_2}$

C. $E = E_1 + E_2$

D. $E = E_1 - E_2$

Answer: B



Watch Video Solution

19. An object of mass 0.2kg executes Simple harmonic along X-axis with frequency of $\frac{25}{\pi}\text{Hz}$. At the position $x=0.04\text{ m}$, the object has kinetic energy of oscillation in meter is equal to

A. 0.05

B. 0.06

C. 0.01

D. None of these

Answer: B



Watch Video Solution

20. A particle of mass (m) is executing oscillations about the origin on the (x) axis. Its potential energy is $V(x) = k|x|^3$ where (k) is a positive constant. If the amplitude of oscillation is a , then its time period (T) is.


- A. Proportional to $\frac{1}{\sqrt{a}}$
- B. Independent of a .
- C. Proportional to \sqrt{a}
- D. Proportional to $a^{3/2}$

Answer: A



Watch Video Solution


21. The variation of potential energy of harmonic oscillator is as show in figure . The spring constant is




- A. $1 \times 10^2 \text{ N/m}$
- B. 150 N/m
- C. $0.667 \times 10^2 \text{ N/m}$
- D. $3 \times 10^2 \text{ N/m}$

Answer: B

22. A particle is executing SHM between points $-X_m$ and X_m , as shown in figure(i). The velocity $V(t)$ of the particle is partially graphed and shown in figure (ii). Two points A and B corresponding to time t_1 and time t_2 respectively are marked on the $V(t)$ curve.



(i)



(ii)

- A. At time t_1 , it is going towards X_m .
- B. At time t_1 , its speed is decreasing
- C. At time t_2 , its position lies in between $-X_m$ and O

D. The phase difference $\Delta\phi$ between points A and B

must be expressed as $90^\circ < \Delta\phi < 180^\circ$.


Answer: B::C



Watch Video Solution

23. One end of a spring of force constant k is fixed to a vertical wall and the other to a block of mass m resting on a smooth horizontal surface. There is another wall at distance x_0 from the block. The spring is then compressed

by $2x_0$ and released. The time taken to strike the wall is



A. $\frac{1}{6}\pi\sqrt{\frac{k}{m}}$

B. $\sqrt{\frac{k}{m}}$

C. $\frac{2\pi}{3}\sqrt{\frac{k}{m}}$

D. $\frac{\pi}{4}\sqrt{\frac{k}{m}}$


Answer: C



Watch Video Solution

24. Three masses 700g, 500g, and 400 g are suspended at the end of a spring as shown and are in equilibrium. When the 700 g mass is removed, the system oscillates with a period of 3 seconds, when the 500 gm mass is also

removed, it will oscillate with a period of



A. 1 s

B. 2 s

C. 3 s

D. $\sqrt{\frac{12}{5}} s$


Answer: B



Watch Video Solution

25. Four massless springs whose force constants are $2k, 2k, k$ and $2k$, respectively, are attached to a mass M kept on a frictionless plane (as shown in figure), If the mass M is displaced in the horizontal direction, then the frequency

of oscillation of the system is



A. $\frac{1}{2\pi} \sqrt{\frac{k}{4M}}$

B. $\frac{1}{2\pi} \sqrt{\frac{4k}{M}}$

C. $\frac{1}{2\pi} \sqrt{\frac{k}{7M}}$

D. $\frac{1}{2\pi} \sqrt{\frac{7k}{M}}$

Answer: B



Watch Video Solution

26. A body at the end of a spring executes SHM with a period t_1 , while the corresponding period for another spring is t_2 , If the period of oscillation with the two spring in series is T , then

A. $T = t_1 + t_2$

B. $T^2 = t_1^2 + t_2^2$

C. $\frac{1}{T} = \frac{1}{t_1} + \frac{1}{t_2}$

D. $\frac{1}{T^2} = \frac{1}{t_1^2} + \frac{1}{t_2^2}$


Answer: B



Watch Video Solution

27. Two identical springs are attached to a small block P. The other ends of the springs are fixed at A and B, When P is in equilibrium the extension of bottom spring is 10 cm. The period of small vertical oscillations of P about is

equilibrium position is ($useg = 9.8m / s^2$)



A. $\frac{2\pi}{7}\text{sec}$

B. $\frac{\pi}{7}\text{sec}$

C. $\frac{2\pi}{5}\text{sec}$

D. none of these

Answer: B



Watch Video Solution

28. Figure shows a system consisting of a massless pulley, a spring of force constant k and a block of mass m . If the block is slightly displaced vertically down from its equilibrium position and then released, the period of its vertical oscillation is





A. $2\pi\sqrt{\frac{m}{K}}$

B. $\pi\sqrt{\frac{m}{4K}}$

C. $\pi\sqrt{\frac{m}{K}}$

D. $4\pi\sqrt{\frac{m}{K}}$


Answer: D




Watch Video Solution

29. A block of mass m is at rest on the another block of same mass as shown in figure. Lower block is attached to

the spring then the maximum amplitude of motion so that both the block will remain in contact is





A. $\frac{mg}{2k}$

B. $\frac{mg}{k}$

C. $\frac{2mg}{k}$

D. $\frac{3mg}{2k}$

Answer: C



Watch Video Solution

30. The displacement of a particle from its mean position
(in metre) is given by

$$y = 0.2 \sin(10\pi t + 1.5\pi) \cos(10\pi t + 0.5\pi).$$

The motion of particle is

- A. Periodic but not S.H.M.
- B. Non-periodic
- C. Simple harmonic motion with period 0.1 s
- D. Simple harmonic motion with period 0.2 s

Answer: C



[View Text Solution](#)

31. The displacement of a particle varies with time as

$$x = 12 \sin \omega t - 16 \sin^2 \omega t \quad (\text{in cm}) \quad \text{it is motion is}$$

S. H. M. then its maximum acceleration is

A. $12 \omega^2$

B. $36 \omega^2$

C. $144 \omega^2$

D. $\sqrt{192} \omega^2$

Answer: B



Watch Video Solution

32. A particle is acted simultaneously by mutually perpendicular simple harmonic motions

$x = a \cos \omega t$ and $y = a \sin \omega t$. The trajectory of motion of the particle will be

A. an ellipse

B. a parabola

C. a circle

D. a straight line

Answer: C



Watch Video Solution

33. A disc of radius R and mass M is pivoted at the rim and it set for small oscillations. If simple pendulum has to have the same period as that of the disc, the length of the simple pendulum should be

A. $\frac{5}{4}R$

B. $\frac{2}{3}R$

C. $\frac{3}{4}R$

D. $\frac{3}{2}R$


Answer: D



Watch Video Solution

34. A particle of mass $m = 2\text{kg}$ executes *SHM* in xy -plane between point A and B under action of force $\vec{F} = F_x \hat{i} + F_y \hat{j}$. Minimum time taken by particle to move from A to B is 1 sec. At $t = 0$ the particle is at $x = 2$

and $y = 2$. Then F_x as function of time t is



A. $-4\pi^2 \sin \pi t$

B. $-4\pi^2 \cos \pi t$


C. $4\pi^2 \cos \pi t$

D. None of these

Answer: B

Multiple Correct Answer Type

1. A particles starts from point $x = -\frac{\sqrt{3}}{2}A$ and move towards negative extreme as shown . If the time period of oscillation is T , them



- A. The equation of the SHM is $x = A \sin\left(\frac{2\pi}{T} + \frac{4\pi}{3}\right)$
- B. The time taken by the particle to go directly from its initial position to negative extreme is $\frac{T}{12}$

C. The time taken by the particle to reach at mean

position is $\frac{T}{3}$

D. The equation of the SHM is $x = A \sin\left(\frac{2\pi}{T}t + \frac{\pi}{3}\right)$

Answer: A::B::C

 [Watch Video Solution](#)

2. A particle is in linear simple harmonic motion between two points A and B 10cm apart (figure). Take the direction from A to B as positive direction and choose the correct statements.

$$AO = OB = 5\text{cm}$$

$$BC = 8\text{cm}$$




- A. The sign of velocity , acceleration and force on the particle when it is 3 cm away from A going towards B are positive
- B. The sign of velocity of the particle at C given towards B is negative
- C. The sign of velocity , acceleration and force on the particle when it is 4 cm away from B going towards A are negative
- D. The sign of acceleration and force on the particle when it is at point B is negative

Answer: A::C::D

 Watch Video Solution

3. The displacement versus time curves for a particle executing SHM is shown in figure. Choose the correct statement (s)



A. Phases of the oscillator is same at $t = 0s$ and

$$t = 2s$$

B. Phases of the oscillator is same at $t = 2s$ and

$$t = 6s$$

C. Phases of the oscillator is same at $t = 1s$ and

$$t = 7s$$

D. Phases of the oscillator is same at $t = 1s$ and

$$t = 5s$$

Answer: B::D



Watch Video Solution

4. Which of the following statements is/are true for a simple harmonic oscillator?


- A. Force acting is directly proportional to displacement from the mean position and opposite to it
- B. motion is periodic
- C. Acceleration of the oscillator is constant
- D. the velocity is periodic

Answer: A::B::D

 [Watch Video Solution](#)

5. The displacement -time graph of a particle executing *SHM* is shown in figure. Which of the following

statements is/are true ?



- A. The force is zero at $t = \frac{3T}{4}$
- B. The acceleration is maximum at $t = \frac{4T}{4}$
- C. The velocity is maximum at $t = \frac{T}{4}$
- D. The PE is equal to KE of oscillation at $t = \frac{T}{2}$

Answer: A::B::C

 **Watch Video Solution**

6. A linear harmonic oscillator of force constant $2 \times 10^6 \text{ N/m}$ and amplitude 0.01 m has a total mechanical energy of 160 Joules. Its

A. Maximum P.E is 100 J.

B. Maximum K.E is 100J

C. Maximum P.E is 160 J

D. Minimum P.E is zero


Answer: B::C



Watch Video Solution

Multiple Correct Answer Type

1. The springs shown in the figure are all upstretched in the beginning when a man starts pulling the block. The man exerts a constant force F on the block.



A. the amplitude of oscillation is

$$\frac{(K_2 + K_3)F}{K_1K_2 + K_2K_3 + K_3K_1}$$

B. the oscillation frequency is

$$\frac{1}{2\pi} \left[\frac{K_1K_2 + K_2K_3 + K_3K_1}{M(K_2 + K_3)} \right]^{1/2}$$

C. the amplitude of oscillation is

$$\frac{2(K_2 + K_3)F}{K_1K_2 + K_2K_3 + K_3K_1}$$

D. the oscillation frequency is

$$\frac{1}{\pi} \left[\frac{K_1K_2 + K_2K_3 + K_3K_1}{M(K_2 + K_3)} \right]^{1/2}$$

Answer: A:B

 [Watch Video Solution](#)

2. A spring has natural length 40cm and spring constant $500\text{N}/\text{m}$. A block of mass 1kg is attached at one end of the spring and other end of the spring is attached to a ceiling. The block is released from the position, where the spring has length 45cm .

A. the block will perform SHM of amplitude 5 cm.

B. the block will have maximum velocity

$$30\sqrt{5} \text{ cm/sec.}$$

C. the block will have maximum acceleration 15 m/s^2 ,


D. the minimum potential energy of the spring will be zero.

Answer: B::C::D

 [Watch Video Solution](#)

3. A 1kg block is executing simple harmonic motion of amplitude 0.1 m on a smooth horizontal surface under the restoring force of a spring of spring constant

100Nm^{-1} . A block of mass 3 kg is gently placed on it at the instant it passes through the mean position. Assuming that the two blocks move together, find the frequency and the amplitude of the motion.



- A. amplitude of the motion is 5 cm
- B. the oscillation frequency is $\frac{5}{2\pi}\text{ Hz}$
- C. amplitude of the motion is 10 cm
- D. the oscillation frequency is $\frac{5}{\pi}\text{ Hz}$


Answer: A:B



Watch Video Solution

4. A mass M is in static equilibrium on a massless vertical spring as shown in the figure. A ball of mass m dropped from certain height sticks to the mass M after colliding with it.

The oscillations they perform reach to height 'a' above the original level of scales & depth 'b' below it.



- (a) Find the constant of force of the spring.,
- (b) Find the oscillation frequency.

(c) What is the height above the initial level from which the mass m was dropped?

A. the force constant of the spring is $\frac{2mg}{b-a}$

B. the oscillation frequency is $\frac{1}{2\pi} \sqrt{\frac{2mg}{(b-a)(M+m)}}$

C. the force constant of the spring is $\frac{mg}{b-a}$


D. the oscillation frequency is $\frac{1}{2\pi} \sqrt{\frac{mg}{(b-a)(M+m)}}$

Answer: A:B

 [Watch Video Solution](#)

5. Two blocks A (5kg) and B (2kg) attached to the ends of a spring constant 1120 N/m are placed on a smooth horizontal plane with the spring unreformed.

Simultaneously velocities of 3 m/s and 10 m/s along the line of the spring in the same direction are imparted to A and B then



A. the maximum extension of the spring is 0.25 m

B. the first maximum compression occurs after start at

$$\frac{3\pi}{56} \text{ s}$$

C. the maximum extension of the spring is 0.50 m

D. Time period of oscillation is $\frac{\pi}{14} \text{ s}$

Answer: B::D



Watch Video Solution

6. The displacement of a particle varies with time according to the relation $y = a \sin \omega t + b \cos \omega t$.

A. The motion is SHM

B. The motion is SHM with amplitude $a+b$

C. The motion is SHM with amplitude $a^2 + b^2$

D. The motion is SHM with amplitude $\sqrt{a^2 + b^2}$

Answer: A::D



7. Three simple harmonic motions in the same direction having each of amplitude "a" and the same period are superposed. If each differs in phase from the next by $\pi/4$ then

A. the resultant amplitude is $(1 + \sqrt{2})a$.

B. the phase of the resultant motion relative to first is 90° .

C. The energy associated with the resultant motion is $(3 + 2\sqrt{2})$ times the energy associated with any single motion.

D. the resulting motion is not simple harmonic.

Answer: A::C



[Watch Video Solution](#)

8. Function $x = A \sin^2 \omega t + B \cos^2 \omega t + C \sin \omega t \cos \omega t$

Represents SHM.

A. For any value of A, B and C (except C = 0).

B. If $A=B, C=2B$ and amplitude = $|B\sqrt{2}|$

C. If $A=B, C=0$

D. If $A=B, C=2B$ amplitude = $|B|$.

Answer: B::D



[View Text Solution](#)

9. The x-coordinate of a particle moving on x-axis is given by $x = 3 \sin 100t + 8 \cos^2 50t$, where x is in cm and t is time in seconds. Which of the following is/are correct about this motion

A. the motion of the particle is not S.H.M.

B. the amplitude of the S.H.M. of the particle is 5 units

C. the amplitude of the resultant S.H.M. is $\sqrt{73}$ units

D. the maximum displacement of the particle from the origin is 9 units.

Answer: B::D



Watch Video Solution

Subjective type

1. Consider a pair of identical pendulums, which oscillate with equal amplitude independently such that when one pendulum is at its extreme position making an angle of 2° to the right with the vertical, the other pendulum makes an angle of 1° to the left of the vertical. What is the phase difference between the pendulums?



[Watch Video Solution](#)

2. A uniform square plate of side a is hinged at one of its corners. It is suspended such that it can rotate about a horizontal axis. The time period of small oscillation about its equilibrium position.



Watch Video Solution

3. A particle of mass 'm' is moving in the x-y plane such that its x and y coordinate vary according to the law $x = a \sin \omega t$ and $y = a \cos \omega t$ where 'a' and ' ω ' are positive constants and 't' is time. Find

(a) equation of the path. Name the trajectory (path).

(b) whether the particle moves in clockwise or anticlockwise direction


(c) magnitude of the force on the particle at any time t.



Watch Video Solution

Single correct answer type

1. The period of a simple pendulum whose bob is hollow metallic sphere is T . The period is T_1 when the bob is filled with sand, T_2 where it is filled with mercury and T_3 when it is half filled with mercury. Which of the following is true?



A. $T = T_1 = T_2 > T_3$

B. $T_1 = T_1 = T_3 > T$

C. $T > T_3 > T_1 = T_2$

D. $T = T_1 = T_2 < T_3$

Answer: D



Watch Video Solution

2. A pendulum has time period T in air when it is made to oscillate in water it acquired a time period $T = \sqrt{2}T$ The density of the pendulum bob is equal to (density) of water = 1)

A. $\sqrt{2}$

B. 2

C. $2\sqrt{2}$

D. none of these

Answer: B



Watch Video Solution

3. A sphere of radius r is kept on a concave mirror of radius of curvature R . The arrangement is kept on a horizontal surface (the surface of concave mirror is friction less and sliding not rolling). If the sphere is displaced from its equilibrium position and left, then it executes $S. h. M$. The period of oscillation will be

A. $2\pi\sqrt{\left(\frac{(R-r)1.4}{g}\right)}$

B. $2\pi\sqrt{\left(\frac{R-r}{g}\right)}$

C. $2\pi\sqrt{\left(\frac{rR}{a}\right)}$

D. $2\pi\sqrt{\left(\frac{R}{gr}\right)}$

Answer: B



Watch Video Solution

4. Two simple pendulums of length $0.5m$ and $0.2m$ respectively are given small linear displacement in one direction at the same time. They will again be in the same phase when the pendulum of shorter length has completed oscillations

A. 5

B. 1

C. 2

D. 3

Answer: C

 [Watch Video Solution](#)

5. The bob of a simple pendulum is displaced from its equilibrium position 'O' to a position 'Q' which is at a height 'h' above 'O' and the bob is then released. Assuming the mass of the bob to be 'm' and time period of oscillation to be $2.0s$, the tension in the string when the bob passes through 'O' is

A. $m(g + \pi\sqrt{2gh})$

B. $m\left(g + \sqrt{\pi^2 gh}\right)$

C. $m\left(g + \sqrt{\frac{\pi^2}{2} gh}\right)$

$$D. m \left(g + \sqrt{\frac{\pi^2}{3} gh} \right)$$

Answer: A



Watch Video Solution

6. Two simple pendulum whose lengths are 100cm and 121cm are suspended side by side. Then bobs are pulled together and then released. After how many minimum oscillations of the longer pendulum will two be in phase again. ?

A. 11

B. 10

C. 21

Answer: B



Watch Video Solution

7. Two pendulums having time period T and $\frac{5T}{4}$. They start $S.H.M.$ at the same time from mean position. what will be the phase difference between them after the bigger pendulum has completes one oscillation ?

A. 45°

B. 90°

C. 60°

D. 30°

Answer: B



Watch Video Solution

8. Two simple pendulum first of bob mass M_1 and length L_1 second of bob mass M_2 and length L_2 $M_1 = M_2$ and $L_1 = 2L_2$. if the vibrational energy of both is same which is correct?


- A. Amplitude of B greater than A
- B. Amplitude of B smaller than A
- C. Amplitudes will be same
- D. none of these

Answer: B




Watch Video Solution


9. In case of a simple pendulum, time period versus length is depicted by




A.



B.



C.



D.

Answer: B

 **Watch Video Solution**

10. A U tube of uniform bore of cross sectional area A has been set up vertically with open ends facing up. Now m gm of a liquid of density d is poured into it. The column of liquid in this tube will oscillate with a period T such that

$$\text{A. } T = 2\pi \sqrt{\frac{M}{g}}$$

$$\text{B. } T = 2\pi \sqrt{\frac{MA}{gd}}$$

$$\text{C. } T = 2\pi \sqrt{\frac{M}{gdA}}$$

$$\text{D. } T = 2\pi \sqrt{\frac{M}{2Agd}}$$

Answer: D



Watch Video Solution

11. A horizontal platform with an object placed on it is executing S.H.M. in the vertical direction. The amplitude of oscillation is 3.92×10^{-3} m. what must be the least period of these oscillations. So that the object is not detached from the platform

A. 0.1256 sec

B. 0.1356 sec

C. 0.1456 sec

D. 0.1556 sec

Answer: A



Watch Video Solution

12. The metallic bob of a simple pendulum has the relative density ρ . The time period of this pendulum is T if the metallic bob is immersed in water the new time period is given by

A. $T \frac{\rho - 1}{\rho}$

B. $T \frac{\rho}{\rho - 1}$

$$C. T \sqrt{\frac{\rho - 1}{\rho}}$$

$$D. T \sqrt{\frac{\rho}{\rho - 1}}$$

Answer: D

 [Watch Video Solution](#)

13. A brass cube of side a and density σ is floating in mercury of density ρ . If the cube is displaced a bit vertically, it executes S.H.M. Its time period will be

$$A. 2\pi \sqrt{\frac{\sigma a}{\rho g}}$$

$$B. 2\pi \sqrt{\frac{\rho a}{\sigma g}}$$

$$C. 2\pi \sqrt{\frac{\rho g}{\sigma a}}$$

$$D. 2\pi \sqrt{\frac{\sigma g}{\rho a}}$$


Answer: A



Watch Video Solution

14. A man weighing $60kg$ stands on the horizontal platform of a spring balance. The platform starts executing simple harmonic motion of amplitude $0.1m$ and frequency $\frac{2}{\pi}$. its which of the following statement is

correct ?



A. The spring balance reads the weight of man as 60

kg

B. The spring balance reading fluctuates between 60

kg and 70 kg.

C. The spring balance reading fluctuates between 50 kg and 60 kg.

D. The spring balance reading fluctuates between 50 kg and 70 kg.

Answer: D



Watch Video Solution