



# PHYSICS

## BOOKS - CENGAGE PHYSICS (HINGLISH)

### PROPERTIES OF SOLIDS AND FLUIDS

#### Illustration

1. Two rods  $A$  and  $B$ , each of equal length for different materials are suspended from a

common support as shown in the figure. The rods  $A$  and  $B$  can support a maximum load of  $W_1 = 600N$  and  $W_2 = 6000N$  respectively. If their cross sectional area are  $A_1 = 10mm^2$  and  $A_2 = 1000mm^2$ , respectively then identify the stronger material.



[Watch Video Solution](#)

2. Assume that if the shear stress in steel exceeds about  $4.00 \times 10^8 N/m^2$  the steel

reptures. Determine the shearing force necessary to *a* shear a steel bolt  $1.00\text{cm}$  in diameter and *b* punch a  $1.00\text{cm}$  diameter hole in a steel plate  $0.500\text{cm}$  thick.



[Watch Video Solution](#)

**3.** A bar of cross section  $A$  is subjected to equal and opposite tensile force at its ends. Consider a plane section of the bar whose normal makes an angle  $\theta$  with the axis of the bar.



- a. What is the tensile stress on the plane?
- b. What is the shearing stress on the this plane?
- c. For what value of  $\theta$  is the tensile stress maximum?
- d. For what value of  $\theta$  is the shearing stress maximum?



**Watch Video Solution**

4. a. Estimate the force with which a karate master strikes a board, assuming the hand's speed at the moment of impact is  $10.0\text{ms}^{-1}$ , decreasing to  $1.00\text{ms}^{-1}$  during a  $0.002\text{s}$  time interval of contact between the hand and the board. The mass of his hand and arm is  $1.00\text{kg}$ .

b Estimate the shear, assuming this force is exerted on a  $1.00\text{cm}$  thick pine board that is  $10.0\text{m}$  wide. c. If the maximum shear stress a pine board can support before breaking is  $3.60 \times 10^6 \text{N}/\text{m}^2$ , will the board break?



**Watch Video Solution**

5. A steel wire  $2m$  long is suspended from the ceiling. When a mass is hung from its lower end, the increase in length recorded is  $1cm$ . Determine the strain in the wire.

A. 0.002

B. 0.005

C. 0.008

D. 0.002

**Answer: B**



[Watch Video Solution](#)

6. A brass rod of length  $1m$  is fixed to a vertical wall at one end, with the other end keeping free to expand. When the temperature of the rod is increased by  $120^{\circ}C$ , the length increases by  $3cm$ . What is the strain?



[Watch Video Solution](#)

7. A  $30.0\text{kg}$  hammer, moving with speed  $20.0\text{ms}^{-1}$  strikes a steel spike  $2.30\text{cm}$  in diameter. The hammer rebounds with speed  $10.0\text{ms}^{-1}$  after  $0.110\text{s}$ . What is the average strain in the spike during the impact.?



[Watch Video Solution](#)

8. The shear modulus for a metal is  $50000\text{Mpa}$ . Suppose that a shear force of  $200\text{N}$  is applied on the upper surface of a cube of this metal



that is  $3.0\text{cm}$  on each edge. How far will the top surface be displaced?

A.  $2.67 \times 10^{-7} m$

B.  $4.5 \times 10^{-7} m$

C.  $6.0 \times 10^{-7} m$

D.  $1.33 \times 10^{-7} m$

**Answer: D**



**Watch Video Solution**

9. When a weight  $W$  is hung from one end of a wire of length  $L$  (other end being fixed), the length of the wire increases by  $l$ . If the same wire is passed over a pulley and two weights  $W$  each are hung at the two ends, what will be the total elongation in the wire?



**Watch Video Solution**

**10.** A load of  $10kN$  is supported from a pulley which in turn is supported by a rope of sectional area  $1 \times 10^3 mm^2$  and modulus of elasticity  $10^3 Nmm^{-2}$ , as shown in figure. Neglecting the friction at the pulley, determine the deflection of the load.



**Watch Video Solution**

**11.** A uniform heavy rod of weight  $W$ , cross sectional area  $a$  and length  $L$  is hanging from

fixed support. Young modulus of the material of the rod is  $Y$ . Neglect the lateral contraction. Find the elongation of the rod.



[Watch Video Solution](#)

**12.** A uniform rod of length  $L$  and mass  $M$  is pulled horizontally on a smooth surface with a force  $F$ . Determine the elongation of rod if Young's modulus of the material is  $Y$ .



[Watch Video Solution](#)

**13.** A  $0.05m$  cube has its upper face displaced by  $0.2cm$  by a tangential force of  $8N$ . Calculate the shearing strain, shearing stress and modulus of rigidity of the material of the cube.



**Watch Video Solution**

**14.** A cube of sponge rubber with edge length  $5cm$  has a force of  $2N$  applied horizontally to the top face (parallel to an edge) while the bottom face is held fixed. If the top face is

displaced horizontally through a distance of  $1\text{mm}$ , find the shear modulus for the sponge rubber. (in  $N/m^2$ )

A.  $2 \times 10^4$

B.  $3 \times 10^4$

C.  $4 \times 10^4$

D.  $5 \times 10^4$

**Answer: C**



**Watch Video Solution**

**15.** Two parallel and opposite forces, each of magnitude  $4000N$ , are applied tangentially to the upper and lower faces of a cubical metal block  $25cm$  on a side. Find the angle of shear and the displacement of the upper surface relative to the lower surface. The shear modulus for the metal is  $80GPa$ .



**Watch Video Solution**

**16.** A horizontal aluminium rod of diameter  $4.8cm$  projects  $5.3cm$  from a wall. A  $1200kg$

object is suspended from the end of the rod.

The shear modulus of aluminium is

$3.0 \times 10^{10} \text{ N/m}^2$ . Neglecting the mass of the

rod find a shearing stress on the rod and b the

vertical deflection of the end of the rod.



[Watch Video Solution](#)

**17.** A metal cube of side  $10\text{cm}$  is subjected to a

shearing stress of  $10^6 \text{ N/m}^2$ . Calculate the

modulus of rigidity if the of the cube is



displaced by  $0.05\text{cm}$  with respect to its bottom.

A.  $1 \times 10^8 \text{ N/m}^2$

B.  $2 \times 10^8 \text{ N/m}^2$

C.  $3 \times 10^8 \text{ N/m}^2$

D.  $4 \times 10^8 \text{ N/m}^2$

**Answer: B**



**Watch Video Solution**

**18.** A hydraulic press contains  $0.25m^3$  ( $250L$ ) of oil. Find the decrease in volume of the oil when it is subjected to a pressure increase  $\Delta p = 1.6 \times 10^7 Pa$ . The bulk modulus of the oil is  $B = 5.0 \times 10^9 Pa$ .



**Watch Video Solution**

**19.** Compressibility of water is  $5 \times 10^{-10} m^2 / N$ . Find the decrease in volume

of  $100\text{mL}$  of water when subjected to a pressure of  $15\text{MPa}$ .



[Watch Video Solution](#)

**20.** Bulk modulus for rubber is  $9.8 \times 10^8 \text{Nm}^{-2}$ . To what depth should a rubber ball be taken in a take so that its volume is decreased by  $0.1\%$



[Watch Video Solution](#)

21. A steel bar  $ABCD$   $40\text{cm}$  long is made up of three parts  $AB$ ,  $BC$  and  $CD$ , as shown in figure. The rod is subjected to a pull of  $25\text{kN}$ . Determine the stress in the three parts and the total extension of the rod. Young's modulus for steel  $= 2 \times 10^{11}\text{Nm}^{-2}$ .



[Watch Video Solution](#)

22. Three elastic wires  $PQ$ ,  $PR$  and  $PS$  support a body  $P$  of mass  $M$ , as shown in figure. The wires are of the same material and cross sectional area, the middle one being vertical. Find the loads by each wire.



[Watch Video Solution](#)

23. A composite tube is made by striking a thin steel tube on a brass tube. If  $A_S$  and  $A_B$

are the respective sectional areas of the steel and brass tubes and  $Y_S$  and  $Y_B$  their Young's moduli, then find the Young's modulus of single tube of the same length and total sectional area, which would be have in the same facition as that of the composite tube.



**Watch Video Solution**

**24.** Two vertical rods of equal lengths, one of steel and the other of copper, are suspended from the ceiling at a distance  $l$  apart and are

connected rigidly to a rigid horizontal bar at their lower ends. If  $A_S$  and  $A_C$  be their respective cross-sectional areas, and  $Y_S$  and  $Y_C$ , their respective Young's moduli of elasticities, where should a vertical force  $F$  be applied to the horizontal bar in order that the bar remains horizontal?



**Watch Video Solution**

**25.** A metallic wire is stretched by suspending a weight of it. If  $\alpha^2$  is the longitudinal strain and  $Y$  is its Young's modulus of elasticity, then show that the elastic potential energy per unit volume is given by  $\frac{1}{2}Y\alpha^2$ .



**Watch Video Solution**

**26.** Calculate the elastic potential energy per unit volume of water at a depth of  $1\text{km}$ .



Compressibility ( $\alpha$ ) of water =  $5 \times 10^{-10}$  SI  
units. Density of water =  $10^3 \text{ kg/m}^3$



[Watch Video Solution](#)

27. A catapult consists of two parallel rubber strings each of lengths,  $10\text{cm}$  and cross sectional area  $10\text{mm}^2$ . When stretched by  $5\text{cm}$ , it can throw a stone of mass  $10\text{gm}$  to a vertical height of  $25\text{m}$ . Determine Young's modulus of elasticity of rubber.



[Watch Video Solution](#)

**28.** A plate of area  $100\text{cm}^2$  is placed on the upper surface of castor oil,  $2\text{mm}$  thick. Taking the coefficient of viscosity to be  $15.5$  poise, calculate the horizontal force necessary to move the plate with a velocity  $3\text{cm}^{-1}$ .



**Watch Video Solution**

**29.** A metal plate of area  $0.10\text{m}^2$  is connected to a  $0.01\text{kg}$  mass via a string that passes over an ideal pulley (considered to be frictionless),

as shown in the figure. A liquid with a film thickness of  $3.0\text{mm}$  is placed between the plate and the table. When released the plate moves to the right with a constant speed of  $0.085\text{m.s}^{-1}$ . Find the coefficient of viscosity of the liquid.



[Watch Video Solution](#)

**30.** A sliding fit cylindrical body of mass of  $1\text{kg}$  drops vertically down at a constant velocity of

$5\text{cm s}^{-1}$ . Find the viscosity of the oil.



[Watch Video Solution](#)

**31.** A powder comprising particle of various sizes is stirred up in a vessel filled to a height of  $10\text{cm}$  with water. Assuming the particle to be spherical, find the size of the largest particle that will remain in suspension after  $1\text{h}$  (density of powder  $= 4\text{g}/\text{cm}^3$ , viscosity of water  $= 0.01\text{poise}$ ).



Watch Video Solution

32. Spherical particles of pollen are shaken up in water and allowed to settle. The depth of water is  $2 \times 10^{-2} m$ . What is the diameter of the largest particles remaining in suspension one hour later? Density of pollen  $= 1.8 \times 10^3 kgm^{-3}$  viscosity of water  $= 1 \times 10^{-2}$  poise and density of water  $= 1 \times 10^{-5} kgm^{-3}$



Watch Video Solution

**33.** A small sphere falls from rest in a viscous liquid. Due to friction, heat is produced. Find the relation between the rate of production of heat and the radius of the sphere at terminal velocity.



**Watch Video Solution**

**34.** A cube of mass  $m = 800g$  floats on the surface of water. Water wets it completely. The cube is  $10cm$  on each edge. By what additional distance is it buoyed up or down by surface

tension? Surface tension of water

$$= 0.07 \text{Nm}^{-1}$$



[Watch Video Solution](#)

**35.** Find the maximum possible mass of a greased needle floating on water surface.



[Watch Video Solution](#)

**36.** A film of water is formed between two straight parallel wires each  $10\text{cm}$  long and at

separation  $0.5\text{cm}$ . Calculate the work required to increase  $1\text{mm}$  distance between the wires.

Surface tension of water =  $72 \times 10^{-3}\text{N/m}$ .



[Watch Video Solution](#)

**37.** Calculate the work done against surface tension in blowing a soap bubble from a radius of  $10\text{cm}$  to  $20\text{cm}$ , if the surface tension of soap solution is  $25 \times 10^{-3}\text{N/m}$ .



[Watch Video Solution](#)



**38.** Calculate the difference  $h$  in water levels in two communicating capillary tube of radius  $1\text{mm}$  and  $1.5\text{mm}$ . Surface tension of water  $= 0.07\text{Nm}^{-1}$



**Watch Video Solution**

**39.** A vessel filled with air under pressure  $p_0$  contains a soap bubble of diameter  $d$ . The air pressure have been reduced  $n$ -fold, and the bubbled diameter increased  $r$ -fold is

othermally. Find the surface tension of the soap water solution.



[Watch Video Solution](#)

**40.** What should be the pressure inside a small air bubble of  $0.1\text{mm}$  radius situated just below the surface of water? Surface tension of water  $= 72 \times 10^{-3}\text{N/m}$  and atmospheric pressure  $= 1.013 \times 10^5\text{N/m}^2$



[Watch Video Solution](#)

**41.** Two separate air bubbles (radii  $0.002\text{cm}$  and  $0.004$ ) formed of the same liquid (surface tension  $0.07\text{N}/\text{m}$ ) come together to form a double bubble. Find the radius and the sense of curvature of the internal film surface common to both the bubbles.



**Watch Video Solution**

**42.** A vertical capillary with inside diameter  $0.50\text{mm}$  is submerged into water so that the length of its part emerging outside the water

surface is equal to  $25\text{mm}$ . Find the radius of curvature of the meniscus. Surface tension of water is  $73 \times 10^{-3}\text{N/m}$ .



[Watch Video Solution](#)

**43.** A glass rod of diameter  $d_1 = 1.5\text{mm}$  is inserted symmetrically into a glass capillary with inside diameter  $d_2 = 2.0\text{mm}$ . Then the whole arrangement is vertically oriented and brought in contact with the surface of water. To what height will the liquid rise in the

capillary?

Surface tension of water =  $73 \times 10^{-3} \text{ N/m}$



[Watch Video Solution](#)

**44.** The end of a capillary tube with a radius  $r$  is immersed in water. Is mechanical energy conserved when the water rises in the tube? The tube is sufficiently long. If not calculate the energy change.



[Watch Video Solution](#)

**45.** A glass capillary tube of internal radius  $r = 0.25\text{mm}$  is immersed in water. The top end of the tube projects by  $2\text{cm}$  above the surface of water. At what angle does the liquid meet the tube? Surface tension of water  $= 0.7\text{Nm}^{-1}$ .



**Watch Video Solution**

**46.** Derive an expression for the height of capillary rise between two parallel plates dipping in a liquid of density  $\sigma$  separated by a

distance  $d$ . The surface tension of the liquid is  $T$ .



[Watch Video Solution](#)

## Solved Examples

1. A boy has a catapult made of a rubber cord of length  $42\text{cm}$  and diameter  $6.0\text{mm}$ . The boy stretches the cord by  $20\text{cm}$  to catapult a stone of mass  $20\text{g}$ . The stone flies off with a speed of  $20\text{ms}^{-1}$ . Find Young's modulus for

rubber. Ignore the change in the cross section of the cord in stretching.



[Watch Video Solution](#)

2. A steel wire of cross-sectional area  $0.5\text{mm}^2$  is held between two fixed supports. If the wire is just taut at  $20^\circ\text{C}$ , determine the tension when the temperature falls to  $0^\circ\text{C}$ . Coefficient of linear expansion of steel is  $1.2 \times 10^{-5} \text{ }^\circ\text{C}^{-1}$  and its Young's modulus is  $2.0 \times 10^{11} \text{ Nm}^{-2}$ .





[Watch Video Solution](#)

3. A sphere of radius  $10\text{cm}$  and mass  $25\text{kg}$  is attached to the lower end of a steel wire which is suspended from the ceiling of a room. The point of support is  $521\text{cm}$  above the floor. When the sphere is set swinging as a simple pendulum, its lowest point just grazes the floor. Calculate the velocity of the ball at its lowest position.



[Watch Video Solution](#)

4. A steel bolt is inserted into a copper tube as shown in the figure. Find the forces induced in the bolt and in the tube when the nut is turned through one revolution. Assume that the length of the tube is  $l$ , the pitch of the bolt thread is  $h$  and the cross sectional areas of the steel bolt and the copper tube are  $A_s$  and  $A_c$  respectively.



[Watch Video Solution](#)

5. A circular ring of radius  $R$  and mass  $m$  made of a uniform wire of cross sectional area  $A$  is rotated about a stationary vertical axis passing through its centre and perpendicular to the plane of the ring. If the breaking stress of the material of the ring is  $\sigma_b$ , then determine the maximum angular speed  $\omega_{\max}$  at which the ring may be rotated without failure.



[Watch Video Solution](#)

6. A glass plate of length  $10\text{cm}$ , breadth  $1.54\text{cm}$  and thickness  $0.20\text{cm}$  weight  $8.2\text{g}$  in air. It is held vertically with the long side horizontal and the lower half under water. Find the apparent weight of the plate. Surface tension of water  $= 7.3 \times 10^{-2}\text{N/m}$  and  $= 9.8\text{ms}^{-12}$



**Watch Video Solution**

7. A barometer contains two uniform capillaries of radii  $1.4 \times 10^{-3}m$  and  $7.2 \times 10^{-4}m$ . If the height of liquid in narrow tube is  $0.2m$  more than that in wide tube, calculate the true pressure difference. Density of liquid  $= 10^3 kg/m^3$ , surface tension  $= 72 \times 10^{-3} N/m$  and  $g = 9.8ms^{-2}$ .



**Watch Video Solution**

8. A glass capillary sealed at the upper end is of length  $0.11\text{m}$  an internal diameter  $2 \times 10^{-5}\text{m}$ . The tube is immersed vertically into a liquid of surface tension  $5.06 \times 10^{-N} / \text{m}$ . To what length the capillary has to be immersed so that the liquid level inside and outside the capillary becomes the same. What will happen to water level inside the capillary if the seal is now broken?



[Watch Video Solution](#)

9. An open capillary tube contains a drop of water. When the tube is in its vertical position, the drop forms a column with a length of a  $2\text{cm}$ , b.  $4\text{cm}$ , and c.  $2.98\text{cm}$ . The internal diameter of the capillary tube is  $1\text{mm}$ . Determine the radii of curvature of the upper and lower menisci in each case. Consider the wetting to be complete. Surface tension of water =  $0.0075\text{N}/\text{m}$



**Watch Video Solution**

**10.** Two spherical soap bubble coalesce. If  $V$  is the consequent change in volume of the contained air and  $S$  the change in total surface area, show that

$$3PV + 4ST = 0$$

where  $T$  is the surface tension of soap bubble and  $P$  is


Atmospheric pressure



**Watch Video Solution**



11. A uniform wire having mass per unit length  $\lambda$  is placed over a liquid surface. The wire causes the liquid to depress by  $y$  ( $y \ll a$ ) as shown in figure. Find surface tension of liquid. Neglect end effect.



[Watch Video Solution](#)


## Exercise 5.1

1. Stress and pressure are both forces per unit area.  $T$  in what respect does stress differ from Pressure?



[Watch Video Solution](#)

2. The stress and strain graphs for two materials,  $A$  and  $B$  are shown here. Answer the following:



give

reasons

A. Which material has greater Young's

modulus?

B. Which material is more ductile?

C. Which material is more brittle?

D. Which material has greater tensile strength?



[Watch Video Solution](#)

**3.** State whether the following statements are true or false with reasons.

- a. Elastic forces are always conservative.
- b. Elastic forces are strictly conservative only when Hooke's law is obeyed.
- c. When a wire is loaded beyond the elastic


limit and then reloaded, the work done disappears completely as heat.



[Watch Video Solution](#)

4. A horizontal force of magnitude  $F$  acts at the end  $P$  of a uniform rigid rod which is welded at point  $Q$ . In each case 1 and 2, as shown in Fig., find the reaction force acting at a point  $C$  at a distance  $x$  from the fixed end  $Q$  of the

rod.




[Watch Video Solution](#)

5. A rubber ball of bulk modulus  $B$  is taken to a depth  $h$  of a liquid of density  $\rho$ . Find the fractional change in the radius of the ball.



[Watch Video Solution](#)


6. Find out longitudinal stress and tangential stress on a fixed block.



[Watch Video Solution](#)

7. Find out bulk stress on the spherical object of radius  $10/\pi$  cm Piston if area and mass of

piston are  $50\text{cm}^2$  and  $50\text{kg}$ , respectively, for a cylinder filled with gas as Gas shown in figure.




[Watch Video Solution](#)



8. Three rods of uniform area of cross section

$A = 10^{-7} \text{ m}^2$  are arranged as shown in Fig



Find out

the shift in point  $B$ ,  $C$  and  $D$ .



Watch Video Solution

9. Find the depth of lake at which density of water is 1 % greater than that at the surface.

Given compressibility  $k = 50 \times 10^{-6} atm^{-1}$ .



[Watch Video Solution](#)

10. A rubber cube of side  $5cm$  has one side fixed while a tangential force equal to  $1800N$  is applied to opposite face. Find the shearing strain and the lateral displacement of the

strained face. Modulus of rigidity for rubber is


$$2.4 \times 10^6 \text{ N/m}^2.$$



**Watch Video Solution**

**11.** A mass ' $m$ ' is attached with rods as shown in Fig. This mass is slightly stretched and released. Find out whether the motion of mass is *S. H. M.* If yes then find out the time


period.



Watch Video Solution

**12.** A ball of mass ' $m$ ' drops from a height which sticks to a massless hanger after striking it. Neglecting overturning. Find out the maximum extension in rod, assuming that

the rod is massless.



Watch Video Solution

Exercise 5.2


1. A man is rowing a boat with a constant velocity  $v_0$  in a river. The contact area of boat is ' $A$ ' and coefficient of viscosity is  $\eta$ . The depth of river is ' $D$ '. Find the force required to row the boat.



[Watch Video Solution](#)

2. A cubical block (of side  $2m$ ) of mass  $20kg$  slides on inclined plane lubricated with the oil of viscosity  $\eta = 10^{-1}$  with constant velocity of

$10ms^{-1}$ . Find out the thickness of the layer of liquid (take  $g = 10ms^{-2}$ ).



**Watch Video Solution**



3. A drop of water of radius  $0.0015\text{mm}$  is falling in air. If the coefficient of viscosity of air is  $1.8 \times 10^{-3}\text{kg/m}^3$ , what will be the terminal velocity of the drop? Density of water  $= 1.0 \times 10^3\text{kg/m}^3$  and  $g = 9.8\text{N/kg}$ . Density of air can be neglected.



[Watch Video Solution](#)

4. A metallic sphere of radius  $1.0 \times 10^{-3}\text{m}$  and density  $1.0 \times 10^4\text{kg/m}^3$  enters a tank of

water, after a free fall through a distance of  $h$  in the earth's gravitational field. If its velocity remains unchanged after entering water, determine the value of  $h$ . Given: coefficient of viscosity of water  $= 1.0 \times 10^{-3} \text{Ns/m}^2$ ,  $g = 10 \text{ms}^{-2}$  and density of water  $= 1.0 \times 10^3 \text{kg/m}^3$ .



[Watch Video Solution](#)

5. Find the minimum force required to drag a hard polythene plate of area  $2 \text{m}^2$  on a thin

film of oil of thickness  $0.25\text{cm}$  and  $\eta = 15$  poise. Assume the speed of the plate is  $10\text{cm s}^{-1}$ .



[Watch Video Solution](#)

6. A force of  $3.14\text{N}$  is required to drag a sphere of radius  $4\text{cm}$  with a speed of  $5\text{m s}^{-1}$  in a medium in gravity free space. Find the coefficient of the viscosity of the medium.



[Watch Video Solution](#)

## Exercise 5.3

1. Why is moisture retained longer in the soil if it is harrowed?



[Watch Video Solution](#)

2. A capillary tube is dipped in water vertically. It is long enough for the water to rise to the maximum height  $h$  in the tube. The length of portion immersed in water is  $I < h$ . The lower end of the tube is closed and then the tube is


taken out and opened again. Will all the water flow out of tube? Explain.



[Watch Video Solution](#)

**3.** Two soap bubbles  $A$  and  $B$  of different diameters are blown at the two ends of a bent tube. By opening the stopcock  $S$ , the two bubbles are put in communication. What will

happen?



[Watch Video Solution](#)

4. A mercury drop of radius  $R$  is sprayed into  $n$  droplets of equal size. Calculate the energy

expanded if surface tension of mercury is  $T$ .



[Watch Video Solution](#)

5. If a number of little droplets of water, each of radius  $r$ , coalesce to form a single drop of radius  $R$ , show that the rise in temperature will be given by  $\frac{3T}{J} \left( \frac{1}{r} - \frac{1}{R} \right)$  where  $T$  is the surface tension of water and  $J$  is the mechanical equivalent of heat.



[Watch Video Solution](#)

6. A liquid of specific gravity 1.5 is observed to rise  $3.0\text{cm}$  in a capillary tube of diameter  $0.50\text{mm}$  and the liquid wets the surface of the tube. Calculate the excess pressure inside a spherical bubble of  $1.0\text{cm}$  diameter blown from the same liquid.



[Watch Video Solution](#)

7. A drop of water of volume  $0.05\text{cm}^3$  is pressed between two glass plates, as a consequence of which, it spreads and occupies



an area of  $40\text{cm}^2$ . If the surface tension of water is  $70\text{dyne/cm}$ , find the normal force required to separate out the two glass plates in newton.



[Watch Video Solution](#)

8. A glass tube of circular cross section is closed at one end. This end is weighted and the tube floats vertically in water, heavy end down. How far below the water surface is the end of the tube? Given: outer radius of the

tube is  $0.14\text{cm}$ , mass of weighted tube is  $0.2\text{g}$ , surface tension of water  $73\text{dyn}/\text{cm}$  and  $g = 980\text{cm}\cdot\text{s}^{-2}$ .



[Watch Video Solution](#)

9. If a  $5\text{cm}$  long capillary tube with  $0.1\text{mm}$  internal diameter, open at both ends, is slightly dipped in water having surface tension  $75\text{dyn}/\text{cm}$ , state whether: water will rise halfway in the capillary, (ii) water will rise up to the upper end of capillary, (iii) water will

overflow out of the upper end of capillary.

Explain your answer.




[Watch Video Solution](#)

## Subjective

1. A bob of mass  $m$  hangs from the ceiling of a smooth trolley car which is moving with a constant acceleration  $a$ . If young's modulus, radius and length of the string are  $Y$ ,  $r$  and  $l$ , respectively, find the (a) stress in the string

and (b) extension of the string when it makes a constant angle relative to vertical.



[Watch Video Solution](#)

2. A smooth uniform, string of natural length  $l$ , cross-sectional area  $A$  and Young's modulus  $Y$

is pulled along its length by a force  $F$  on a horizontal surface. Find the elastic potential energy stored in the string.



[Watch Video Solution](#)

**3.** A narrow capillary tube is dipped  $10\text{cm}$  below water surface and a liquid bubble of radius  $2\text{mm}$  formed at the lower end by blowing air through the tube.

a. Calculate the excess pressure due to surface tension.

b. What is the pressure required in the tube in order to blow a hemispherical bubble at its end in water?


The surface tension of water at temperature of the experiment is  $7.30 \times 10^{-2} N/m$ . 1 atmospheric pressure =  $10^5 Pa$ , density of water =  $1000 kg/m^3$



**Watch Video Solution**

4. A thin plate of large area is placed midway in a gap of height  $h$  filled with oil of viscosity

and the plate is pulled at constant velocity  $v$  by applying the same drag force on the plate. If a lighter oil of viscosity  $\eta$  is then substituted in the gap. it is found that for the velocity  $v$ , and the same drag force as previous case the plate is located unsymmetrically in the gap but parallel to the walls. Find  $\eta$  in terms of distance from nearer wall to the plane  $y$ .




**Watch Video Solution**

5. When a vertical capillary of length with the sealed upper end was brought in contact with the surface of a liquid, the level of this liquid rose to the height  $h$ . The liquid density is  $\rho$ , the inside diameter the capillary is  $d$ , the contact angle is  $\theta$ , the atmospheric pressure is  $\rho_0$ . Find the surface tension of the liquid.



(Temperature this process remains constant.)



[Watch Video Solution](#)

6. A rubber cord has a cross-sectional area  $1\text{mm}^2$  and total un stretched length  $10.0\text{cm}$ . It is stretched to  $12.0\text{cm}$  and then released to project a missile of mass  $5.0\text{g}$ . Taking young's modulus  $Y$  for rubber as  $5.0 \times 10^8\text{N/m}^2$ . Calculate the velocity of projection .




**Watch Video Solution**

7. Two wires of diameter  $0.25\text{cm}$ , one made of steel and other made of brass, are loaded as

shown in the figure. The unloaded length of the steel wire is  $1.5m$  and that of brass is  $1.0m$ . Young's modulus of steel is  $2.0 \times 10^{11} Pa$  and that of brass is  $1.0 \times 10^{11} Pa$ . Compute the ratio of elongations of steel and brass

wires.  $\frac{\Delta l_{\text{steel}}}{\Delta l_{\text{brass}}} = ?$





Watch Video Solution


8. A steel rod of length  $l_1 = 30\text{cm}$  and two identical brass rod of length  $l_2 = 20\text{cm}$  each support a light horizontal platform as shown in Fig. Cross-sectional area of each of the three rods is  $A = 1\text{cm}^2$ . A vertically downward force  $F = 5000\text{N}$  is applied on the platform.

Young's modulus of elasticity for steel

$$Y_s = 2 \times 10^{11} \text{Nm}^{-2} \quad \text{and} \quad \text{brass}$$

$$Y_b = 1 \times 10^{11} \text{Nm}^{-2}. \text{ Find stress (in MPa)}$$

developed in a. Steel rod b. Brass rod



[Watch Video Solution](#)

9. The two wires shown in figure are made of the same material which has a breaking stress of  $8 \times 10^8 \text{ Nm}^{-2}$ . The area of cross section of

the upper wire is  $0.006\text{cm}^2$  and that of the lower wire is  $0.003\text{cm}^2$ . The mass  $m_1 = 10\text{kg}$ ,  $m_2 = 20\text{kg}$  and the hanger is light. a. Find the maximum load that can be put on the hanger without breaking a wire. Which wire will break first if the load is increased? b. Repeat the above part  $m_1 = 10\text{kg}$  and  $m_2 = 36\text{kg}$ .



[Watch Video Solution](#)

**10.** A copper wire of negligible mass,  $1m$  length and cross-sectional area  $10^{-6}m^2$  is kept on a smooth horizontal table with one end fixed. A ball of mass  $1kg$  is attached to the other end. The wire and the ball are rotating with an angular velocity of  $20rad/s$ . If the elongation in the wire is  $10^{-3}m$ .

a. Find the Young's modulus of the wire (in terms of  $\times 10^{11}N/m^2$ ).

b. If for the same wire as stated above, the angular velocity is increased to  $100rad/s$  and



the wire breaks down, find the breaking stress (in terms of  $\times 10^{10} N/m^2$ ).




**Watch Video Solution**

**11.** An amusement park ride consists of airplane shaped cars attached to steel rods. Each rod has a length of  $20.0m$  and a cross-sectional area of  $8.00cm^2$ . Young's modulus for steel is  $2 \times 10^{11} N/m^2$ .

a. How much is the rod stretched (in  $mm$ ) when the ride is at rest ? (Assume that each

car plus two people seated in it has a total weight of  $2000N$ .)

b. When operating, the ride has a maximum angular speed of  $\sqrt{\frac{19}{5}} \text{ rad/s}$ . How much is the rod stretched (in mm) then?



**Watch Video Solution**


**12.** A capillary tube of length (i)  $l = 60\text{cm}$ , (ii)  $l = 50\text{cm}$  and radius  $r = 1/4\text{mm}$  is immersed vertically into water. Find the capillary rise in both cases. Angle of contact  $= 0^\circ$ . Take coefficient of surface tension as  $72\text{dyne/cm}$ ,  $g = 1000\text{cm s}^{-12}$ .



**Watch Video Solution**

**13.** An annular disc of radius  $r_1 = 10\text{cm}$  and  $r_2 = 5\text{cm}$  is placed on a water surface. Find

the surface tension force on the disc if we want to pull it from water surface. Take coefficient of surface tension as  $\sigma = 7 \times 10^{-3} \text{ N/m}$ ,  $g = 10 \text{ m/s}^2$ .



[Watch Video Solution](#)

**14.** Two soap bubbles of radii  $a$  and  $b$  coalesce to form a single bubble of radius  $c$ . If the


external pressure is  $P$ , find the surface tension of the soap solution.



[Watch Video Solution](#)

**15.** Calculate the pressure inside a small air bubble of radius  $0.01\text{mm}$  situated at a depth of  $h = 20\text{m}$  below the free surface of liquid of density  $\rho_1 = 10^3\text{kg}/\text{m}^3$ ,  $\rho_2 = 800\text{kg}/\text{m}^3$  and surface tension  $T_2 = 7.5 \times 10^{-2}\text{N}/\text{m}$ . The thickness of the first liquid is  $h_1 = 15\text{m}$ .

and  $h_2 = 25\text{m}$ .



[Watch Video Solution](#)

**16.** A mercury drop of radius  $R$  is sprayed into  $n$  droplets of equal size. Calculate the energy expended if surface tension of mercury is  $T$ .



Watch Video Solution

## Single Correct

1. The ratio of diameters of two wires of same material is  $n : 1$ . The length of each wire is  $4m$ . On applying the same load, the increases in the length of the thin wire will be ( $n > 1$ )

A.  $n^2$  times

B.  $n$  times

C.  $2n$  times

D.  $(2n + 1)$  times

**Answer: A**



**Watch Video Solution**

2. A nylon rope  $2\text{cm}$  in diameter has a breaking strength of  $1.5 \times 10^5 \text{N}$ . The breaking strength of a similar rope  $1\text{cm}$  in diameter is

A.  $0.375 \times 10^5 \text{N}$



B.  $2 \times 10^5 N$

C.  $6 \times 10^5 N$

D.  $9 \times 10^4 N$

**Answer: A**



**Watch Video Solution**

**3.** The dimensions of four wires of the same material are given below. In which wire the increase in the length will be maximum?

A. Length  $100\text{cm}$ , diameter  $1\text{mm}$

B. Length  $200\text{cm}$ , diameter  $2\text{mm}$

C. Length  $300\text{cm}$ , diameter  $3\text{mm}$

D. Length  $50\text{cm}$ , diameter  $0.5\text{mm}$

**Answer: D**



**Watch Video Solution**

4. Two wires of the same material and length but diameters in the ratio  $1:2$  are stretched by the same force. The potential energy per

unit volume for the two wires when stretched will be of the ratio.

A. 16:1

B. 4:1

C. 2:1

D. 1:1

**Answer: A**



**Watch Video Solution**

5. Two wires of the same material and length are stretched by the same force. Their masses are in the ratio 3:2. Their elongations are in the ratio

A. 3:2

B. 9:4

C. 2:3

D. 4:9

**Answer: C**



Watch Video Solution

6. Two wires of the same length and same material but radii in the ratio of 1:2 are stretched by unequal forces to produce equal elongation. The ratio of the two forces is

A. 1:1

B. 1:2

C. 1:3

D. 1:4

**Answer: D**



**Watch Video Solution**

7. When a weight of  $5kg$  is suspended from a copper wire of length  $30m$  and diameter  $0.5mm$ , the length of the wire increases by  $2.4cm$ . If the diameter is doubled, the extension produced is

A.  $1.2cm$

B.  $0.6$

C.  $0.3\text{cm}$

D.  $0.15\text{cm}$

**Answer: B**



**Watch Video Solution**

8. The length of a wire is increased by  $1\text{mm}$  on the application, of a given load. In a wire of the same material, but of length and radius twice that of the first, on application of the same load, extension is

A.  $0.25\text{mm}$

B.  $0.5\text{mm}$

C.  $2\text{mm}$

D.  $4\text{mm}$

**Answer: B**



**Watch Video Solution**

9. An cube is shifted to a depth of  $100\text{m}$  is a lake. The change in volume is  $0.1\%$ . The bulk modulus of the material is



A.  $10Pa$

B.  $10^4 Pa$

C.  $10^7 Pa$

D.  $10^9 Pa$

**Answer: D**



**Watch Video Solution**

**10.** If 'S' is stress and 'Y' is young's modulus of material of a wire, the energy stored in the wire per unit volume is

A.  $\frac{SY}{2}$

B.  $\frac{S^2}{2Y}$

C.  $\frac{S}{2y}$

D.  $\frac{2S}{Y}$

**Answer: B**



**Watch Video Solution**

**11.** What amount of work is done in increasing the length of a wire through unity?

A.  $\frac{YL}{2A}$

B.  $\frac{YL^2}{2A}$

C.  $\frac{YA}{2L}$

D.  $\frac{YL}{A}$

**Answer: C**



**Watch Video Solution**

**12.** Young's modulus of rubber is  $10^4 N/m^2$  and area of cross section is  $2cm^{-2}$ . If force of

$2 \times 10^5 \text{ dyn}$  is applied along its length, then

its initial  $l$  becomes

A.  $3l$

B.  $4l$

C.  $2l$

D. none of these

**Answer: C**



**Watch Video Solution**

**13.** When a certain weight is suspended from a long uniform wire, its length increases by  $1\text{cm}$ . If the same weight is suspended from another wire of the same material and length but having a diameter half of the first one, the increase in length will be

A.  $0.5\text{cm}$

B.  $2\text{cm}$

C.  $4\text{cm}$

D.  $8\text{cm}$

**Answer: C**



**Watch Video Solution**

**14.** Two wires of the same material have lengths in the ratio  $1:2$  and their radii are in the ratio  $1:\sqrt{2}$ . If they are stretched by applying equal forces, the increase in their lengths will be in the ratio

A.  $\sqrt{2}:2$

B.  $2:\sqrt{2}$

C. 1:1

D. 1:2

**Answer: C**



**Watch Video Solution**

**15.** A piece of copper wire has twice the radius of a piece of steel wire. Young's modulus for steel is twice that of the copper. One end of the copper wire is joined to one end of the steel wire so that both can be subjected to the

same longitudinal force. By what fraction of its length will the steel have stretched when the length of the copper has increased by 1 % ?

A. 1 %

B. 2 %

C. 2.5 %

D. 3 %

**Answer: B**



**Watch Video Solution**



16. The breaking stress for a substance is  $10^6 \text{ N/m}^2$ . What length of the wire of this substance should be suspended vertically so that the wire breaks under its own weight? (Given: density of material of the wire  $= 4 \times 10^3 \text{ kg/m}^3$  and  $g = 10 \text{ ms}^{-2}$ )

A.  $10\text{m}$

B.  $15\text{m}$

C.  $25\text{m}$

D.  $34\text{m}$

**Answer: C**



**Watch Video Solution**

17. Water rises to a height of  $2\text{cm}$  in a capillary tube. If the tube is tilted  $60^\circ$  from the vertical, water will rise in the tube to a length of

A.  $4.0\text{cm}$

B.  $2.0\text{cm}$

C.  $1.0\text{cm}$

D. water will not rise at all

**Answer: A**



**Watch Video Solution**

**18.** A spherical liquid drop of radius  $R$  is divided into eight the surface tension is  $T$ , then the work equal droplets. If the surface tension is  $T$ , then the work done in this process will be

A.  $2\pi R^2 T$

B.  $3\pi R^2 T$

C.  $4\pi R^2 T$

D.  $2\pi RT^2$

**Answer: C**



**Watch Video Solution**

**19.** Air is pushed into a soap bubble of radius  $r$  to double its radius. If the surface tension of the soap solution is  $S$ , the work done in the process is

A.  $2\pi D^2 T$

B.  $4\pi D^2 T$

C.  $6\pi D^2 T$

D.  $8\pi D^2 T$

**Answer: C**



**Watch Video Solution**

**20.** A water drop is divided into eight equal droplets. The pressure difference between inner and outer sides of big drop

A. will be the same as for smaller droplet

B. will be half of that for smaller droplet

C. will be one-fourth of that for smaller  
droplet

D. will be twice of that for smaller droplet

**Answer: B**



**Watch Video Solution**

21. A vessel whose , bottom has round holes with diameter  $0.1\text{mm}$ , is filled with water. The maximum height up to which water can be filled without leakage is

A.  $100\text{cm}$

B.  $75\text{cm}$

C.  $50\text{cm}$

D.  $30\text{cm}$

**Answer: D**



Watch Video Solution

22. Water rises to a height of  $10\text{cm}$  in a capillary tube and mercury falls to a depth of  $3.42\text{cm}$  in the same capillary tube. If the density of mercury is  $13.6\text{g}/\text{c. c.}$  and the angles of contact for mercury and water are  $135^\circ$  and  $0^\circ$ , respectively, the ratio of surface tension for water and mercury is

A.  $1 : 0.15$

B.  $1 : 3$



C. 1 : 6.5

D. 1.5 : 1

**Answer: C**



**Watch Video Solution**

**23.** The velocity of small ball of mass  $M$  and density  $d_1$  when dropped in a container filled with glycerin becomes constant after sometime. If the density glycerin of is  $d_2$ , the viscous force acting on ball is

A.  $\frac{Md_1g}{d_2}$

B.  $Mg\left(1 - \frac{d_2}{d_1}\right)$

C.  $\frac{M(d_1 + d_2)}{g}$

D.  $Md_1d_2$

**Answer: B**



**Watch Video Solution**

**24.** Two soap bubbles, one of radius  $50mm$  and the other of radius  $80mm$ , are brought in contact so that they have a common interface.

The radius of the curvature of the common interface is

A.  $0.003m$

B.  $0.133m$

C.  $1.2m$

D.  $8.9m$

**Answer: B**



**Watch Video Solution**

25. A glass rod of radius  $r_1$  is inserted symmetrically into a vertical capillary tube of radius  $r_2$  such that their lower ends are at the same level. The arrangement is now dipped in water. The height to which water will rise into the tube will be ( $\sigma =$  surface tension of water,  $\rho =$  density of water)

A. 
$$\frac{2\sigma}{(r_2 - r_1)\rho g}$$

B. 
$$\frac{\sigma}{(r_2 - r_1)\rho g}$$

C. 
$$\frac{2\sigma}{(r_2 + r_1)\rho g}$$

$$D. \frac{2\sigma}{(r_2^2 + r_1^2)\rho g}$$

**Answer: A**



**Watch Video Solution**

**26.** A large number of droplets, each of radius  $a$ , coalesce to form a bigger drop of radius  $b$ . Assume that the energy released in the process is converted into the kinetic energy of the drop. The velocity of the drop is  $\sigma =$  surface tension,  $\rho =$  density)

A.  $\left[ \frac{\sigma}{\rho} \left( \frac{1}{a} - \frac{1}{b} \right) \right]^{\frac{1}{2}}$

B.  $\left[ \frac{2\sigma}{\rho} \left( \frac{1}{a} - \frac{1}{b} \right) \right]^{\frac{1}{2}}$

C.  $\left[ \frac{3\sigma}{\rho} \left( \frac{1}{a} - \frac{1}{b} \right) \right]^{\frac{1}{2}}$

D.  $\left[ \frac{6\sigma}{\rho} \left( \frac{1}{a} - \frac{1}{b} \right) \right]^{\frac{1}{2}}$

**Answer: D**



**Watch Video Solution**

**27.** A thick rope of density  $\rho$  and length  $L$  is hung from a rigid support. The increase in

length of the rope due to its own weight is ( $Y$   
is the Young's modulus)

A.  $\frac{0.1}{4Y} \rho L^2 g$

B.  $\frac{1}{2Y} \rho L^2 g$

C.  $\frac{\rho L^2 g}{Y}$

D.  $\frac{\rho L g}{Y}$

**Answer: B**



**Watch Video Solution**

28. When the load on a wire is slowly increased from 3 to 5 *kgwt*, the elongation increases from 0.61 to 1.02 *mm*. The work done during the extension of wire is

A. 0.16 *J*

B. 0.016 *J*

C. 1.6 *J*

D. 16 *J*

**Answer: B**



Watch Video Solution



29. Two identical wires of iron and copper with their Young's modulus in the ratio 3:1 are suspended at same level. They are to be loaded so as to have the same extension and hence level. Ratio of the weight is

A. 1:3

B. 2:1

C. 3:1

D. 4:1

**Answer: C**



**Watch Video Solution**

**30.** A wire of cross section  $A$  is stretched horizontally between two clamps located  $2lm$  apart. A weight  $Wkg$  is suspended from the mid-point of the wire. If the mid-point sags vertically through a distance  $x < 1$  the strain produced is

A.  $\frac{2x^2}{l^2}$

B.  $\frac{x^2}{l^2}$

C.  $\frac{x^2}{2l^2}$

D. none of these

**Answer: C**



**Watch Video Solution**

**31.** A long wire hangs vertically with its upper end clamped. A torque of  $8Nm$  applied to the free end twists it through  $45^\circ$ . The potential energy of the twisted wire is

A.  $\pi J$

B.  $\frac{\pi}{2} J$

C.  $\frac{\pi}{4} J$

D.  $\frac{\pi}{8} J$

**Answer: A**



**Watch Video Solution**

**32.** The bulk modulus of water is  $2.0 \times 10^9 N/m^2$ . The pressure required to increase the density of water by 0.1 % is

A.  $2 \times 10^9 N / m^2$

B.  $2 \times 10^8 N / m^2$

C.  $2 \times 10^4 N / m^2$

D.  $2 \times 10^4 N / m^2$

**Answer: C**



**Watch Video Solution**

**33.** Two rods of different materials having coefficients of thermal expansion  $\alpha_1$ ,  $\alpha_2$  and Young's moduli  $Y_1$ ,  $Y_2$  respectively are fixed

between two rigid massive walls. The rods are heated such that they undergo the same increase in temperature. There is no bending of the rods. If  $\alpha_1 : \alpha_2 = 2 : 3$ , the thermal stresses developed in the two rods are equal provided  $Y_1 : Y_2$  is equal to

A. 2 : 3

B. 1 : 1

C. 3 : 2

D. 4 : 9

**Answer: C**



Watch Video Solution

34. One end of uniform wire of length  $L$  and of weight  $W$  is attached rigidly to a point in the roof and a weight  $W_1$  is suspended from its lower end. If  $s$  is the area of cross section of the wire, the stress in the wire at a height ( $3L/4$ ) from its lower end is

A.  $\frac{W_1}{s}$

B.  $\left[ W_1 + \frac{W}{4} \right] s$

C.  $\left[ W_1 + \frac{3W}{4} \right] / s$

D.  $\frac{W_1 + W}{s}$

**Answer: C**



**Watch Video Solution**

**35.** A wire is stretched  $1\text{mm}$  by a force of  $1\text{kN}$ .

How far would a wire of the same material and length but of four times that diameter be stretched by the same force?

A.  $\frac{1}{2}\text{mm}$



B.  $\frac{1}{4}mm$

C.  $\frac{1}{8}mm$

D.  $\frac{1}{16}mm$

**Answer: D**



**Watch Video Solution**

**36.** Young's modulus of brass and steel are  $10 \times 10^{10} N/m$  and  $2 \times 10^{11} N/m^2$ , respectively. A brass wire and a steel wire of the same length are extended by  $1mm$  under

the same force. The radii of the brass and steel wires are  $R_B$  and  $R_S$ , respectively. Then

A.  $R_s = \sqrt{2}R_B$

B.  $R_S = \frac{R_B}{\sqrt{2}}$

C.  $R_S = 4R_B$

D.  $R_S = (R_B) / 4$

**Answer: B**



**Watch Video Solution**

37. The length of a steel wire is  $l_1$  when the stretching force is  $T_1$  and  $l_2$  when the stretching force is  $T_2$ . The natural length of the wire is

A.  $\frac{T_2}{R} - 1(l_1 + l_2)$

B.  $T_1l_1 + T_2l_2$

C.  $\frac{l_1T_2 - l_2T_1}{T_2 - T_1}$

D.  $\frac{l_1T_2 + l_2T_1}{T_2 + T_1}$

**Answer: C**




Watch Video Solution

**38.** Two blocks of masses  $1\text{kg}$  and  $2\text{kg}$  are connected by a metal wire going over a smooth pulley as shown in figure.

The breaking stress of the metal is  $(40/3\pi) \times 10^6 \text{N/m}^2$ . If  $g = 10\text{ms}^{-2}$ , then what should be the minimum radius of the

wire used if it is not to break?



A.  $0.5\text{mm}$

B.  $1\text{mm}$

C.  $1.5\text{mm}$

D.  $2\text{mm}$

**Answer: B**



**Watch Video Solution**

**39.** A long elastic spring is stretched by  $2\text{cm}$  and its potential energy is  $U$ . If the spring is stretched by  $10\text{cm}$ , the  $PE$ ..... will be

A.  $5U$

B.  $25U$

C.  $U/5$

D.  $U/20$

**Answer: B**



**Watch Video Solution**

**40.** A copper bar of length  $L$  and area of cross section  $A$  is placed in a chamber at atmospheric pressure. If the chamber is

evacuated, the percentage change in its volume will be (compressibility of copper is  $8 \times 10^{-12} m^2 / N$  and  $1 atm = 10^5 N / m$ )

A.  $8 \times 10^{-7}$

B.  $8 \times 10^{-5}$

C.  $1.25 \times 10^{-4}$

D.  $1.25 \times 10^{-5}$

**Answer: B**



**Watch Video Solution**



41. A small but heavy block of mass  $10\text{kg}$  is attached to a wire  $0.3\text{m}$  long. Its breaking stress is  $4.8 \times 10^7 \text{N}/\text{m}^2$ . The area of the cross section of the wire is  $10^{-6}\text{m}^2$ . The maximum angular velocity with which the block can be rotated in the horizontal circle is

A.  $4\text{rad}/\text{s}$

B.  $8\text{rad}/\text{s}$

C.  $10\text{rad}/\text{s}$

D.  $32\text{rad}/\text{s}$

**Answer: A**



**Watch Video Solution**

**42.** A ball falling in a lake of depth  $200m$  shows a decrease of  $0.1\%$  in its volume at the bottom. The bulk modulus of the elasticity of the material of the ball is (take  $g = 10ms^{-12}$ )

A.  $10^9 N/m^2$

B.  $2 \times 10^9 N/m^2$

C.  $3 \times 10^9 N/m^2$

$$D. 4 \times 10^9 N/m^2$$

**Answer: B**



**Watch Video Solution**

**43.** A massive stone pillar  $20m$  high and of uniform cross section rests on a rigid base and supports a vertical load of  $5.0 \times 10^5 N$  at its upper end. If the compressive stress in the pillar is not exceed  $16 \times 10^6 N/m$ , what is the minimum cross-sectional area of the pillar?

(Density of the stone =  $2.5 \times 10^3 \text{ kg/m}^3$  take

$g = 10 \text{ N/kg}$ )

A.  $0.15 \text{ m}^2$

B.  $0.25 \text{ m}^2$

C.  $0.35 \text{ m}^2$

D.  $0.45 \text{ m}^2$

**Answer: D**



**Watch Video Solution**

44. If the work done in stretching a wire by  $1\text{mm}$  is  $2J$ , then work necessary for stretching another wire of same material but with double radius of cross -section and half of the length by  $1\text{mm}$  is

A.  $\frac{1}{4}J$

B.  $4J$

C.  $8J$

D.  $16J$

**Answer: B**



Watch Video Solution

**45.** Two wires of the same material and same mass are stretched by the same force. Their lengths are in the ratio

A. 3 : 2

B. 2 : 3

C. 4 : 9

D. 9 : 4

**Answer: C**



Watch Video Solution

**46.** A wire is suspended vertically from a rigid support. When loaded with a steel weight in air, the wire extends by  $16\text{cm}$ . When the weight is completely immersed in Water, the extension is reduced to  $14\text{cm}$ . The relative density of the material of the weight is

A.  $2\text{g} / \text{cm}^3$

B.  $6\text{g} / \text{cm}^3$

C.  $8\text{g} / \text{cm}^3$

D.  $16g/cm^3$

**Answer: C**



**Watch Video Solution**

**47.** Two bars  $A$  and  $B$  of circular cross section, same volume and made of the same material, are subjected to tension. If the diameter of  $A$  is half that of  $B$  and if the force applied to both the rod is the same and it is in the elastic



limit, the ratio of extension of  $A$  to that of  $B$  will be

A. 16

B. 8

C. 4

D. 2

**Answer: A**



**Watch Video Solution**

**48.** A uniform cylindrical wire is subjected to a longitudinal tensile stress of  $5 \times 10^7 \text{ N/m}^2$ . Young's modulus of the material of the wire is  $2 \times 10^{11} \text{ N/m}^2$ . The volume change in the wire is  $0.02\%$ . The fractional change in the radius is

A.  $0.25 \times 10^{-4}$

B.  $0.5 \times 10^{-4}$

C.  $1.0 \times 10^{-4}$

D.  $1.5 \times 10^{-4}$

**Answer: A**



**Watch Video Solution**

**49.** A material has normal density  $\rho$  and bulk modulus  $K$ . The increase in the density of the material when it is subjected to an external pressure  $P$  from all sides is

A.  $P / \rho K$

B.  $K / \rho P$

C.  $\rho P / K$

$$D. \rho K / P$$

**Answer: C**



**Watch Video Solution**

**50.** A rubber rope of length  $8m$  is hung from the ceiling of a room. What is the increase in length of rope due to its own weight? (Given: Young's modulus of elasticity of rubber  $= 5 \times 10^6 N/m$  and density of rubber  $= 1.5 \times 10^3 kg/m^3$ . Take  $g = 10ms^{-12}$ )

A.  $1.5\text{mm}$

B.  $6\text{mm}$

C.  $24\text{mm}$

D.  $96\text{mm}$

**Answer: D**



**Watch Video Solution**

**51.** A straw  $6\text{cm}$  long floats on water. The water film on one side has surface tension of  $50\text{dyn}/\text{cm}$ . On the other side, camphor

reduces the surface tension to  $40\text{dyn}/\text{cm}$ .

The resultant force acting on the straw is

A.  $(50 \times 6 - 40 \times 6)\text{dyn}$

B.  $10\text{dyn}$

C.  $\left(\frac{50}{6} - \frac{40}{6}\right)\text{dyn}$

D.  $90\text{dyn}$

**Answer: A**



**Watch Video Solution**

52. Two glass plates are separated by water. If surface tension of water is  $75 \text{ dyn/cm}$  and the area of each plate wetted by water is  $8 \text{ cm}^2$  and the distance between the plates is  $0.12 \text{ mm}$ , then the force applied to separate the two plates is

A.  $10^2 \text{ dyn}$

B.  $10^4 \text{ dyn}$

C.  $10^5 \text{ dyn}$

D.  $10^6 \text{ dyn}$

**Answer: C**



**Watch Video Solution**

**53.** A ring is cut from a platinum tube  $8.5\text{cm}$  internal and  $8.7\text{cm}$  external diameter. It is supported horizontally from the pan of a balance, so that it comes in contact with the water in a glass vessel. If an extra  $3.103\text{gf}$  is required to pull it away from water, the surface tension of water is



A.  $72 \text{ dyn} / \text{ cm}$

B.  $70.80 \text{ dyn} / \text{ cm}$

C.  $63.35 \text{ dyn} / \text{ cm}$

D.  $60 \text{ dyn} / \text{ cm}$


**Answer: A**



**Watch Video Solution**

**54.** A soap film of surface tension  $3 \times 10^{-2}$  formed in a rectangular frame can support a straw as shown in Fig. If  $g = 10 \text{ m s}^{-2}$ , the

mass of the straw is



A.  $0.006g$

B.  $0.06g$

C.  $0.6g$

D.  $6g$

**Answer: C**



Watch Video Solution

55. The lower end of a capillary tube is at a depth of  $12\text{cm}$  and water rises  $3\text{cm}$  in it. The mouth pressure required to blow an air bubble at the lower end will be  $x\text{cm}$  of water column, where  $x$  is

A. 12

B. 15

C. 3

D. 9

**Answer: B**




**Watch Video Solution**

**56.** A light wire AB of length 10 cm can slide on a vertical frame as shown in figure. There is a film of soap solution trapped between the frame and the wire. Find the load  $W$  that should be suspended from the wire to keep it in equilibrium. Neglect friction. Surface

tension of soat solution =  $25 \text{ dyn cm}^{-1}$ . Take

$g = 10 \text{ ms}^{-2}$



A.  $0.2g$

B.  $0.3g$

C.  $0.4g$

D.  $0.5g$

**Answer: D**



Watch Video Solution

57. The angle of contact between glass and water is  $0^\circ$  and water (surface tension  $70 \text{ dyn/cm}$ ) rises in a glass capillary up to  $6 \text{ cm}$ . Another liquid of surface tension  $140 \text{ dyn/cm}$ , angle of contact  $60^\circ$  and relative density 2 will rise in the same capillary up to

A.  $12 \text{ cm}$

B.  $24 \text{ cm}$

C.  $3 \text{ cm}$

D.  $6\text{cm}$

**Answer: C**



**Watch Video Solution**

**58.** A hollow sphere has a small hole in it. On lowering the sphere in a tank of water, it is observed that water enters into the hollow sphere at a depth of  $40\text{cm}$  below the surface. Surface tension of water is  $7 \times 10^{-2}\text{N/m}$ . The diameter of the hole is

A.  $\frac{1}{28}mm$

B.  $\frac{1}{21}mm$

C.  $\frac{1}{14}mm$

D.  $\frac{1}{7}mm$

**Answer: C**



**Watch Video Solution**

**59.** Work  $W$  is required to form a bubble of volume  $V$  from a given solution. What amount



of work is required to be done to form a bubble of volume  $2V$  ?

A.  $W$

B.  $2W$

C.  $2^{\frac{1}{3}}W$

D.  $4^{\frac{1}{3}}W$

**Answer: D**



**Watch Video Solution**

60. The surface energy of a liquid drop is  $E$ . It is sprayed into 1000 equal droplets. Then its surface energy becomes

A.  $1000E$

B.  $100E$

C.  $10E$

D.  $E$

**Answer: C**



**Watch Video Solution**

61. A cube with a mass  $= 20g$  wettable water floats on the surface of water. Each face of the cube is  $a = 3cm$  long. Surface tension of water is  $70dyn/cm$ . The distance of the lower face of the cube from the surface of water is ( $g = 980cm/s^{-2}$ )

A.  $2.3cm$

B.  $4.6cm$

C.  $9.7cm$

D.  $12.7cm$


**Answer: A**



**Watch Video Solution**

**62.** A liquid is contained in a vertical tube of semicircular cross section figure. The contact angle is zero. The force of surface tension on the curved part and on the flat part are in ratio





A.  $2 : \pi$

B.  $1 : \pi$

C.  $3 : \pi$

D.  $2.7 : \pi$

**Answer: A**



**Watch Video Solution**

**63.** Two vertical parallel glass plates are partially submerged in water. The distance between the plates is  $d$  and the length is  $l$ .

Assume that the water between the plates does not reach the upper edges of the plates and the wetting is complete. The water will rise to height ( $\rho =$  density of water and  $\alpha =$  surface tension of water)

A.  $\frac{2\sigma}{\rho g d}$

B.  $\frac{\sigma}{2\rho g d}$

C.  $\frac{4\sigma}{\rho g d}$

D.  $\frac{5\sigma}{\rho g d}$

**Answer: A**



64. A number of droplets, each of radius  $r$ , combine to form a drop of radius  $R$ . If  $T$  is the surface tension, the rise in temperature will be

A.  $\frac{2T}{r}$

B.  $\frac{3T}{R}$

C.  $2T \left[ \frac{1}{r} - \frac{1}{R} \right]$

D.  $3T \left[ \frac{1}{r} - \frac{1}{R} \right]$

**Answer: D**





65. A drop of liquid of density  $\rho$  is floating half-immersed in a liquid of density  $d$ . If  $\rho$  is the surface tension the diameter of the drop of the liquid is

A.  $\sqrt{\frac{\sigma}{g(2\rho - d)}}$

B.  $\sqrt{\frac{2\sigma}{g(2\rho - d)}}$

C.  $\sqrt{\frac{6\sigma}{g(2\rho - d)}}$

D.  $\sqrt{\frac{12\sigma}{g(2\rho - d)}}$

**Answer: D**



**Watch Video Solution**

**66.** A drop of liquid of density  $\rho$  is floating half-immersed in a liquid of density  $d$ . If  $\rho$  is the surface tension the diameter of the drop of the liquid is

A.  $\frac{TA^2}{V}$

B.  $\frac{2TA^2}{V}$

C.  $\frac{4TA^2}{V}$

D.  $\frac{TA^2}{2V}$

**Answer: B**



**Watch Video Solution**

**67.** Two soap bubbles of radii  $a$  and  $b$  coalesce to form a single bubble of radius  $c$ . If the external pressure is  $P$ , find the surface tension of the soap solution.

A.  $\frac{P(c^3 + a^3 + b^3)}{4(a^2 + b^2 - c^2)}$

B.  $\frac{P(c^3 - a^3 - b^3)}{4(a^2 + b^2 - c^2)}$

C.  $Pc^3 - 4a^2 - 4b^2$

D.  $Pc^3 - 2a^2 - 3b^2$

**Answer: B**



**Watch Video Solution**

**68.** A thin square plate of side  $5\text{cm}$  is suspended vertically a balance so that lower side just dips into water with side to surface. When the plate is clean ( $\theta = 0^\circ$ ), it appears

to weigh  $0.044\text{N}$ . But when the plate is greasy ( $\theta = 180^\circ$ ) it appears to weigh  $0.03\text{N}$ . The surface tension of water is

- A.  $3.5 \times 10^{-2}\text{N/m}$
- B.  $7.0 \times 10^{-2}\text{N/m}$
- C.  $14.0 \times 10^{-2}\text{N/m}$
- D.  $1.08\text{N/m}$

**Answer: B**



**Watch Video Solution**

69. A wire forming a loop is dipped into soap solution and taken out so that a film of soap solution is formed. A loop of 6.28 cm long thread is gently put on the film and the film is pricked with a needle inside the loop. The thread loop takes the shape of a circle. Find the tension in the thread. Surface tension of soap solution =  $0.030 \text{ Nm}^{-1}$ .

A.  $1 \times 10^4 \text{ n}$

B.  $2 \times 10^{-4} \text{ N}$

C.  $3 \times 10^{-4} \text{ N}$

$$D. 4 \times 10^{-4} N$$

**Answer: C**



**Watch Video Solution**

**70.** A 20 cm long capillary tube is dipped in water. The water rises up to 8 cm. If the entire arrangement is put in a freely falling elevator, the length of water column in the capillary tube will be

A.  $20\text{cm}$

B.  $4\text{cm}$

C.  $10\text{cm}$

D.  $8\text{cm}$

**Answer: A**



**Watch Video Solution**

71. A marble of mass  $x$  and diameter  $2r$  is gently released in a tall cylinder containing honey. If the marble displaces mass  $y$  ( $< x$ ) of



the liquid, then the terminal velocity is proportional to

A.  $x + y$

B.  $x - y$

C.  $\frac{x + y}{r}$

D.  $\frac{x - y}{r}$

**Answer: D**



**Watch Video Solution**

72. A small metal ball of diameter  $4\text{mm}$  and density  $10.5\text{g}/\text{cm}^3$  is dropped in glycerine of density  $1.5\text{g}/\text{cm}^3$ . The ball attains a terminal velocity of  $8/\text{cm}\text{s}^{-1}$ . The coefficient of viscosity of glycerine is

A. 4.9 poise

B. 9.8 poise

C. 98 poise

D. 980 poise

**Answer: B**



Watch Video Solution

**73.** A capillary tube is attached horizontally to a constant pressure head arrangement. If the radius of the capillary tube is increased by 10 % , then the rate of flow of the liquid shall change nearly by

A. + 10 %

B. 46 %

C. - 10 %

D.  $-40\%$

**Answer: B**



**Watch Video Solution**

**74.** A sphere of brass released in a long liquid column attains a terminal speed  $v_0$ . If the terminal speed is attained by a sphere of marble of the same radius and released in the same liquid is  $nv_0$ , then the value of  $n$  will be

(Given: The specific gravities of brass, marble and liquid are 8.5, 2.5 and 0.8, respectively)

A.  $\frac{5}{17}$

B.  $\frac{17}{77}$

C.  $\frac{1}{31}$

D.  $\frac{17}{5}$

**Answer: B**



**Watch Video Solution**

75. Between a plate of area  $100\text{cm}^2$  and another plate of area  $100\text{m}^2$  there is a  $1\text{mm}$ , thick layer of water, if the coefficient of viscosity of water is  $0.01$  poise, then the force required to move the smaller plate with a velocity  $10\text{cm s}^{-1}$  with reference to large plate is

A.  $100\text{dyn}$

B.  $10^4\text{dyn}$

C.  $10^6\text{dyn}$

D.  $10^9 \text{ dyn}$

**Answer: A**



**Watch Video Solution**

**76.** A river  $10\text{m}$  deep is flowing at  $5\text{ms}^{-1}$ . The shearing stress between horizontal layers of the rivers is ( $\eta = 10^{-3}$   $SI$  units)

A.  $10^{-3} \text{ N/m}^2$

B.  $0.8 \times 10^{-3} \text{ N/m}^2$

C.  $0.5 \times 10^{-3} N / m^2$

D.  $1 N / m^2$

**Answer: C**



**Watch Video Solution**

**77.** A ball rises to the surface of a liquid with constant velocity. The density of the liquid is four times the density of the material of the ball. The frictional force of the liquid on the



rising ball is greater than the weight of the ball by a factor of

A. 2

B. 3

C. 4

D. 6

**Answer: B**



**Watch Video Solution**

78. A spherical ball falls through viscous medium with terminal velocity  $v$ . If this ball is replaced by another ball of the same mass but half the radius, then the terminal velocity will be (neglect the effect of buoyancy.)

A.  $v$

B.  $2v$

C.  $4v$


D.  $8v$




**Answer: B**



Watch Video Solution

79. Each of the pictures shows four objects tied together with rubber bands being pulled to the right across a horizontal frictionless surface by a horizontal force  $F$ . All the objects have the same mass, all the rubber bands obey Hooke's law and have the same equilibrium length and the same force constant. Which of these pictures is drawn most correctly?





**Answer: B**



**Watch Video Solution**

**80.** A glass rod of radius  $1\text{mm}$  is inserted symmetrically into a glass capillary tube with inside radius  $2\text{mm}$ . Then the whole arrangement is brought in contact of the

surface of water. Surface tension of water is  $7 \times 10^{-2} \text{ N/m}$ . To what height will the water rise in the capillary? ( $\theta = 0^\circ$ )

A.  $1.4 \text{ cm}$

B.  $4.2 \text{ cm}$

C.  $2.1 \text{ cm}$


D.  $6.8 \text{ cm}$

**Answer: A**



**Watch Video Solution**

81. Two soap bubbles  $A$  and  $B$  of different diameters are blown at the two ends of a bent tube. By opening the stopcock  $S$ , the two bubbles are put in communication. What will happen?




- A. There will be change in the size of the bubbles
- B. The bubbles will become of equal size
- C. The bubbles will become of equal size
- D. The bubbles will become of equal size

**Answer: C**



**Watch Video Solution**

82. In the figure shown, forces of equal magnitude are applied to the two ends of a uniform rod. Consider  $A$  as the cross-sectional area of the rod. For this situation, mark out the incorrect statements.



A. The rod is in compressive stress.

B. The numerical value of stress developed

in the rod is equal to  $F / A$ .



C. The stress is defined as internal force developed at any cross section per unit area.

D. none of these

**Answer: D**



**Watch Video Solution**

**83.** The space between two large horizontal metal plates,  $6\text{cm}$  apart, is filled with a liquid of viscosity  $0.8\text{N} / \text{m}^2$ . A thin plate of surface

area  $0.01m^2$  is moved parallel to the length of the plate such that the plate is at a distance of  $2m$  from one of the plates and  $4cm$  from the other. If the plate moves with a constant speed of  $1ms^{-1}$ , then

A. the layer of the fluid, which is having the maximum velocity, is lying mid-way between the plates

B. the layers of the fluid, which is in contact with the moving plate, is having the maximum velocity

C. the layer of the fluid, which is in contact with the moving plate and is on the side of farther plate, is moving with the maximum velocity

D. the layer of the fluid, which is in contact with the moving plant and is on the side of nearer plate, is moving with the maximum velocity

**Answer: B**



**Watch Video Solution**

**84.** The pressure that has to be applied to the ends of a steel wire of length 10cm to keep its length constant when its temperature is raised by  $100^{\circ}C$  is : (For steel Young's modulus is  $2 \times 10^{11} Nm^{-2}$  and coefficient of thermal expansion is  $1.1 \times 10^{-5} K^{-1}$ )

A.  $22 \times 10^7 atm$

B.  $2.2 \times 10^3 atm$

C. zero

$$D. 4.3 \times 10^3 \text{ atm}$$

**Answer: B**



**Watch Video Solution**

**85.** Maximum excess pressure inside a thin-walled steel tube of radius  $r$  and thickness  $\Delta r$  ( $\Delta r \ll r$ ), so that the tube would not rupture would be (breaking stress of steel is

$\sigma_{\max}$

$$A. \sigma_{\max} \times \frac{r}{\Delta r}$$

B.  $\sigma_{\max} \times \frac{\Delta r}{r}$

C.  $\sigma_{\max}$

D.  $\sigma_{\max} \times \frac{\Delta 2r}{r}$


**Answer: B**



**Watch Video Solution**

**86.** Two equal and opposite point forces applied at mid-points of the ends of a rod of square cross shown. Consider the dotted section  $ABCD$ . If the rod is cut across this

Gloss section, the force exerted by the right part of the rod on left part across this cross section is



A. acting at point passing through cross section acting at point passing through cross section  $ABCD$

B. acting at a point but not passing through the centre of cross section

*ABCD*

C. uniformly distributed across the cross section *ABCD*

D. non-uniformly distributed across the cross section *ABCD*

**Answer: C**



**Watch Video Solution**



**87.** A  $5\text{kg}$  rod of square cross section  $5\text{cm}$  on a side and  $1\text{m}$  long is pulled along a smooth horizontal surface by a force applied at one end. The rod has a constant acceleration of  $2\text{ms}^{-2}$ . Determine the elongation in the rod. (Young's modulus of the material of the rod is  $5 \times 10^3 \text{N/m}^2$ ).

- A. Zero, as for elongation to be there, equal and opposite forces must act on the rod
- B. Non-zero but cannot be determined from the given situation

C.  $0.4\mu m$

D.  $16\mu m$

**Answer: C**



**Watch Video Solution**

**88.** If two soap bubbles of different radii are connected by a tube

A. air follows from the larger bubble into smaller bubble till both bubbles acquire-

same size

B. air follows from the smaller bubble into larger bubble and the larger bubble grows in size with decrease in size of the smaller bubble

C. air does not flow but the sizes of the bubbles changes

D. sizes of the bubbles remain unchanged

**Answer: B**



**Watch Video Solution**

**89.** A paper disc of radius  $R$  from which a hole of radius  $r$  is cut out is floating in a liquid of the surface tension  $S$ . The force on the disc due to the surface tension is

A.  $S \times 2\pi R$

B.  $S \times 2\pi r$

C.  $S \times 2\pi(R - r)$

D.  $S \times 2\pi(R + r)$

**Answer: D**



Watch Video Solution

90. A glass rod of radius  $1\text{mm}$  is inserted symmetrically into a glass capillary tube with inside radius  $2\text{mm}$ . Then the whole arrangement is brought in contact with the surface of water. Surface tension of water is  $7 \times 10^{-2}\text{N/m}$ . To what height will the water rise in the capillary? ( $\theta = 0^\circ$ )

A.  $1.44\text{cm}$

B.  $6\text{cm}$

C. 4.86

D. none of these

**Answer: A**



**Watch Video Solution**

**91.** The elastic limit of an elevator cable is  $2 \times 10^9 \text{ N/m}^2$ . The maximum upward acceleration that an elevator of mass  $2 \times 10^3 \text{ kg}$  can have when supported by a cable whose cross sectional area is  $10^{-4} \text{ m}^2$ ,

provided the stress in cable would not exceed half to the elastic limit would be

A.  $10ms^{-2}$

B.  $50ms^{-2}$

C.  $40ms^{-2}$

D. Not possible to move up

**Answer: C**



**Watch Video Solution**

92. A wire can sustain the weight of 20 kg before breaking. If the wire is cut into two equal parts each part can sustain a weight of

A.  $10\text{kg}$

B.  $20\text{kg}$

C.  $40\text{kg}$

D.  $35\text{kg}$

**Answer: B**



**Watch Video Solution**



**93.** A wire of length  $L$  and radius  $r$  is fixed at one end. When a stretching force  $F$  is applied at free end, the elongation in the wire is  $l$ . When another wire of same material but of length  $2L$  and radius  $2r$ , also fixed at one end is stretched by a force  $2F$  applied at free end, then elongation in the second wire will be

A.  $\frac{l}{2}$

B.  $l$

C.  $2l$

D.  $\frac{l}{4}$

**Answer: B**



**Watch Video Solution**

**94.** On applying a stress of  $xN/m^2$ , the length of wire of some material gets doubled. Value of Young's modulus for the material of the wire in  $N/m^2$ , is (assume Hooke's law to be valid and go for approx. results)

A.  $x$

B.  $2x$

C.  $\frac{x}{2}$

D. Insufficient information

**Answer: A**



**Watch Video Solution**

**95.** A Copper wire and steel of the same diameter and length are connected end to end and a force is applied, which stretches their

combined length by 1 cm. The two wires will have

- A. same stress and same strain
- B. same stress and different strains
- C. different stresses and same strain
- D. different stresses and different strains

**Answer: B**



**Watch Video Solution**

**96.** A steel wire of length  $4.7m$  and cross-sectional area  $3 \times 10^{-6}m^2$  stretches by the same amount as a copper wire of length  $3.5m$  and cross-sectional area of  $4 \times 10^{-6}m^2$  under a given load. The ratio of Young's modulus of steel to that of copper is

A. 1.8

B. 3.6

C. 0.6

D. 8.7

**Answer: A**



**Watch Video Solution**

**97.** The edges of an aluminum cube are  $10\text{cm}$  long. One face of the cube is firmly fixed to a vertical wall. A mass of  $100\text{kg}$  is then attached to the opposite face of the cube. Shear modulus of aluminum is  $25 \times 10^9 \text{ Pa}$ , the vertical deflection in the face to which mass is attached is

A.  $4 \times 10^{-4}m$

B.  $4 \times 10^{-7}m$

C.  $25 \times 10^{-6}m$

D.  $6 \times 10^{-7}m$

**Answer: B**



**Watch Video Solution**

**98.** A solid sphere of radius  $R$  made of a material of bulk modulus  $K$  is surrounded by a liquid in a cylindrical container. A massless

piston of area  $A$  floats on the surface of the liquid. When a mass  $M$  is placed on the piston to compress the liquid the fractional change in the radius of the sphere,  $\delta R / R$ , is .....

A.  $\frac{Mg}{AK}$

B.  $\frac{Mg}{3AK}$

C.  $\frac{3Mg}{AK}$

D.  $\frac{Mg}{2AK}$

**Answer: B**



**Watch Video Solution**



**99.** A film of water is formed between two straight parallel wires each  $10\text{cm}$  long and at a separation of  $0.5\text{cm}$ . Calculate the work required to increase  $1\text{mm}$  distance between the wires. Surface tension of water  $= 72 \times 10^{-3}\text{N/m}$ .

A.  $1.44 \times 10^{-5}\text{J}$

B.  $1.72 \times 10^{-5}\text{J}$

C.  $1.44 \times 10^{-4}\text{J}$

D.  $1.72 \times 10^{-4}\text{J}$

**Answer: A**



**Watch Video Solution**

**100.** The length of a needle floating on water is  $2.5\text{cm}$ . The minimum force in addition to its weight needed to lift the needle above the surface of water will be (surface tension of water is  $0.072\text{N}/\text{m}$ )

A.  $3.6 \times 10^{-3}\text{N}$

B.  $10^{-2}\text{N}$

C.  $9 \times 10^{-4} N$

D.  $6 \times 10^{-4} N$

**Answer: A**



**Watch Video Solution**

**101.** A steel wire is stretched by  $1kgwt$ . If the radius of the wire is doubled, its Young's modulus will

A. remain unchanged

B. become half

C. become double

D. become four times

**Answer: A**



**Watch Video Solution**

**102.** Two long metallic strips are joined together by two rivets each of radius  $2mm$ . Each rivet can withstand a maximum shearing stress of  $1.5 \times 10^9 N/m^2$ . Assuming that each

rivet shares the stretching load equally, the maximum tensile force the strip can exert without rupture is

A.  $1.88 \times 10^4 N$

B.  $3.8 \times 10^4 N$

C.  $6 \times 10^7 N$

D.  $3 \times 10^4 N$

**Answer: B**



**Watch Video Solution**

**103.** A solid sphere falls with a terminal velocity of  $20\text{ms}^{-1}$  in air. If it is allowed to fall in vacuum

A. terminal velocity will be  $20\text{ms}^{-1}$

B. terminal velocity will be less than  $20\text{ms}^{-1}$

C. terminal velocity will be greater than  $20\text{ms}^{-1}$

D. no terminal velocity will be attained

**Answer: D**



Watch Video Solution

**104.** The density of water at the surface of ocean is  $\rho$ . If the bulk modulus of water is  $B$ , then the density of ocean water at depth, when the pressure is  $\alpha p_0$  and  $p_0$  is the atmospheric pressure is

A. 
$$\frac{pB}{B - (\alpha - 1)p_0}$$

B. 
$$\frac{pB}{B + (\alpha - 1)p_0}$$

C. 
$$\frac{pB}{B - \alpha p_0}$$

D.  $\frac{pB}{B + \alpha p_0}$

**Answer: A**



**Watch Video Solution**

**105.** Water rises to a height  $h$  in a capillary tube of cross-sectional area  $A$ . the height to which water will rise in a capillary tube of cross-sectional area  $4A$  will be

A.  $h$



B.  $h / 2$

C.  $h / 4$

D.  $4h$

**Answer: B**



**Watch Video Solution**

**106.** Neglecting the density of air, the terminal velocity obtained by a raindrop of radius  $0.3\text{mm}$  falling through the air of viscosity  $1.8 \times 10^{-5} \text{N/m}^2$  will be

A.  $10.9m / s$

B.  $8.3m / s$

C.  $9.2m / s$

D.  $7.6m / s$

**Answer: A**



**Watch Video Solution**

**107.** A composite rod consists of a steel rod of length  $25cm$  and area  $2A$  and a copper rod of length  $50cm$  and area  $A$ . The composite rod is

subjected to an axial load  $F$ . If the Young's moduli of steel and copper are in the ratio 2:1 then

A. the extension produced in copper rod will be more

B. the extension in copper and steel parts will be in the ratio 1:2

C. the stress applied to copper rod will be more

D. no extension will be produced in the  
steel rod

**Answer: A::B::C**



**Watch Video Solution**

**108.** Four rods  $A, B, C$  and 1) of the same length and material but of different radii  $r, r\sqrt{2}, r\sqrt{3}$  and  $2r$ , respectively, are held between two rigid walls. The temperature of

all rods is increased through the same range.

If the rods do not bend, then

A. the stress in the rods  $A$ ,  $B$ ,  $C$  and  $D$  is

in the ratio  $1:2:3:4$

B. the forces on them exerted by the wall

are in the ratio  $1:2:3:4$

C. the energy stored in the rods due to

elasticity is in the ratio  $1:2:3:4$

D. it is independent of area like surface

tension while friction depends

**Answer: B::C**



**Watch Video Solution**

**109.** Viscous force is somewhat like friction as it opposes the motion and is non-conservative but not exactly so because

A. it is velocity dependent while friction is not

B. it is velocity independent while friction is

C. it is temperature dependent while friction is not

D. it is independent of area is like surface tension while friction is dependent

**Answer: A:C**



**Watch Video Solution**

**110.** Excess pressure can be  $(2T / R)$  for

A. spherical drop

B. spherical meniscus

C. cylindrical bubble in air

D. spherical bubble in water

**Answer: A::B::C::D**



**Watch Video Solution**

**111.** If a liquid rises to the same height in two capillaries of the same material at the same temperature, then



- A. the weight of liquid in both capillaries must be equal
- B. the radius of meniscus must be equal
- C. the capillaries must be cylindrical and vertical
- D. the hydrostatic pressure at the base of capillaries must be same


**Answer: A::B**



**Watch Video Solution**

**112.** The wires  $A$  and  $B$  shown in Fig. are made of the same material and have radii  $r_A$  and  $r_B$ , respectively. The block between them has a mass  $m$ . When the force  $F$  is  $mg/3$ , one of

the wires breaks. Then



A.  $A$  will break before  $B$  if  $r_A = r_B$

B.  $A$  will break before  $B$  if  $r_A < 2r_B$

C. either  $A$  or  $B$  may break if  $r_A = 2r_B$

D. the lengths of  $A$  and  $B$  must be known  
to predict which wire will break.

**Answer: A::B::C**



**Watch Video Solution**

**Multiple Correct**

1. If  $n$  drops of a liquid, form a single drop, then

A. some energy will be released in the process

B. some energy will be absorbed in the process

C. the energy released or absorbed will be

$$E\left(n - n^{\frac{2}{3}}\right)$$

D. the energy released or absorbed will be

$$nE\left(2^{\frac{2}{3}} - 1\right)$$

**Answer: A::C**



**Watch Video Solution**

2. When a capillary tube is dipped in a liquid, the liquid rises to a height  $h$  in the tube. The free liquid surface inside, the tube is hemispherical in shape. The tube is now pushed down so that the height of the tube outside the liquid is less than  $h$ . Then

A. the liquid will come out of the tube like  
in a small fountain

B. the liquid will ooze out of the tube  
slowly

C. the liquid will fill the tube but not come  
out of its upper end

D. the free liquid surface inside the tube  
will not be hemispherical


**Answer: C::D**




**Watch Video Solution**



3. A vertical glass capillary tube, open at both ends, contains some water. Which of the following shapes may not be taken by the water -in the tube?




**b.**




B.

**c.**



C.

**d.**



D.

**Answer: A::B::C**



**Watch Video Solution**

4. A uniform plank is resting over a smooth horizontal floor and is pulled by applying a horizontal force at its one end. Which of the following statements are not correct?

A. Stress developed in plank material is maximum at the end at which force is

applied and decrease linearly to zero at the other end.

B. A uniform tensile stress is developed in the plank material.

C. Since plank is pulled at one end only, plank starts to accelerate along direction of the force. Hence, no stress developed in the plank material.

D. none of these


**Answer: B::C**



Watch Video Solution

5. A rod is made of uniform material and has non-uniform cross section. It is fixed at both the ends as shown and heated at mid-section.

Which of the following are not correct?




- A. Force of compression in the rod will be maximum at mid section
- B. compressive stress in the rod will be maximum at left end
- C. since rod is fixed at both the ends, its length will remain unchanged. Hence, no strain will be induced in it.
- D. none of these

**Answer: A::C**



**Watch Video Solution**

6. Figure shows the stress-strain graphs for materials  $A$  and  $B$ . From the graph it follows that



A. to material  $A$  has a higher Young's modulus

B. material  $B$  is more ductile

C. material  $A$  can withstand greater stress

D. material  $B$  can withstand greater stress

**Answer: A::D**



**Watch Video Solution**

7. Two wires  $A$  and  $B$  have the same cross section and are made of the same material, but the length of wire  $A$  is twice that of  $B$ . Then, for a given load



- A. the extension of  $A$  will be twice that of  $B$
- B. the extensions of  $A$  and  $B$  will be equal
- C. the strain in  $A$  will be half that in  $B$
- D. the strains in  $A$  and  $B$  will be equal

**Answer: A::D**



**Watch Video Solution**

8. Two wires  $A$  and  $B$  have equal lengths and are made of the same material, but diameter of wire  $A$  is twice that of wire  $B$ . Then, for a given load,

A. The extension of  $B$  will be four times that of  $A$

B. the extension of  $A$  and  $B$  will be equal

C. the strain in  $B$  is four times that in  $A$

D. the strains in  $A$  and  $B$  will be equal

**Answer: A::C**



**Watch Video Solution**

**9.** Choose the correct statements from the following:

A. Steel is more elastic than rubber.

B. The stretching of a coil spring is determined by the Young's modulus of the wire of the spring.

C. The frequency of a tuning fork is determined by the shear modulus of the material of the fork.

D. When a material is subjected to a tensile (stretching) stress the restoring forces are caused by interatomic attraction.

**Answer: A**



**Watch Video Solution**

**10.** Which of the following are correct?

A. For a small deformation of a material,  
the ratio (stress/ strain) remains same.

B. For a large deformation of a material,  
the ratio (stress/ strain) decreases.

C. Two wires made of different materials,  
having the same diameter and length  
are connected end to end. A force is  
applied. This stretches their combined

length by  $2mm$ . Now, the strain is same in both the wire but stress is different.

D. None of these is correct.

**Answer: A::B**



**Watch Video Solution**

**11.** A light rod of length  $2m$  is suspended from the ceiling horizontally by means of two vertical wires of equal length tied to its ends. One of the wires is made of steel and is of

cross section  $0.1\text{cm}^2$ . The other wire is a brass of cross section  $0.2\text{cm}^2$ . A weight is suspended from a certain point of the rod such that equal stress are produced in both the wires. Which of the following are correct?

- A. The ratio of tension in the steel and brass wires is 0.5
- B. The load is suspended at a distance of  $400/3$  cm from the steel wire.
- C. Both (a) and (b) are correct
- D. Neither (a) nor b) is correct.

**Answer: A::B::C**



**Watch Video Solution**

**12. Which of the following are correct?**

A. The product of bulk' modulus of elasticity and compressibility is 1

B. A rope  $1\text{cm}$  in diameter breaks if the tension in it exceeds  $500\text{N}$ . The



maximum tension that may be given to a similar rope of diameter  $2\text{cm}$  is  $2000\text{N}$ .

C. Both (a) and (b) are correct.

D. Neither (a) nor (b) is correct.

**Answer: A::B::C**



**Watch Video Solution**

**13.** Which of the following are correct?

A. The shear modulus of a liquid is infinite.

B. Bulk modulus of a perfectly rigid body is infinite.

C. According to Hooke's law, the ratio of the stress and strain remains constant.

D. None of the above


**Answer: B::C**



**Watch Video Solution**

**14.** A heavy block of mass  $150kg$  hangs with the help of three vertical wires of equal length and equal cross-sectional area as shown in Fig. Wire is attached to the mid-point (centre of mass) of block. Take  $Y_2 = 2Y_1$ . For this arrangement mark out the correct

statement(s).



A. The wire I and III should have same Young's modulus.

B. Tension in I and III would be always equal.

C. Tension in I and III would be different.

D. Tension in II is  $75g$

**Answer: A::B::D**



**Watch Video Solution**

**15.** A body of mass  $M$  is attached to the lower end of a metal wire, whose upper end is fixed .  
The elongation of the wire is  $l$ .

A. Loss in gravitational potential energy of

$$M \text{ is } Mgl$$

B. Elastic potential energy stored in the

$$\text{wire is } \frac{Mgl}{2}$$

C. Elastic potential energy stored in the

$$\text{wire is } Mgl$$

D. Elastic potential energy stored in the

$$\text{wire is } \frac{Mgl}{3}$$

**Answer: A::B**



**Watch Video Solution**

**16.** A metal wire of length  $L$ , area of cross-section  $A$  and young's modulus  $Y$  is stretched by a variable force  $F$  such that  $F$  is always slightly greater than the elastic forces of resistance in the wire. When the elongation of the wire is  $l$

A. the work done by  $F$  is  $\frac{YAl^2}{2L}$

B. the work done by  $F$  is  $\frac{YAl^2}{L}$

C. the elastic potential energy stored in

wire is  $\frac{YAl^2}{2L}$

D. no energy is lost during elongation

**Answer: A::C::D**




**Watch Video Solution**

**17.** A tank of large base area is filled with water up to a height of  $5m$ . A hole of  $2cm^2$  cross section in the bottom allows the water to drain out in continuous streams. For this



situation, mark out the correct statement(s)

(take  $\rho_{\text{water}} = 1000\text{kg}/\text{m}^3$ ,  $g = 10\text{ms}^{-2}$ )



A. The cross-sectional area of the emerging stream of water decreases as it falls down.

B. The cross-sectional area of the emerging stream of water increases as it falls down.

C. At a distance of  $5m$  below the bottom of the tank, the cross-sectional area of the stream is  $1.414cm^2$ .

D. At a distance of  $5m$  below the bottom of the tank, the cross-sectional area of the stream is  $2.86cm^2$ .

**Answer: A::C**



Watch Video Solution

## Assertion- Reasoning

1. Statement I: Surface tension has the same units as force gradient.

Statement II: Surface tension is the force gradient along the surface of liquid.

A. Statement I is true, statement II is true and Statement II is a correct explanation

for Statement I.

B. Statement I is true, Statement II is true  
and Statement II is NOT the correct  
explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: A**



**Watch Video Solution**

2. Statement I: Small liquid drops assume spherical shape.

Statement II: Due to surface tension liquid drops tend to have minimum surface area.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true and Statement II is NOT the correct explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: A**



**Watch Video Solution**

**3.** Statement I: A small drop of mercury is spherical & bigger drops are oval in shape.

Statement II: Surface tension of liquid decreases with increase in temperature.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true and Statement II is NOT the correct explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: B**



**Watch Video Solution**

4. Statement I: Droplets of liquid are usually more spherical in shape than large drops of the same liquid.

Statement II: Force of surface tension predominates force of gravity in case of small drops.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.



B. Statement I is true, Statement II is true  
and Statement II is NOT the correct  
explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: A**



**Watch Video Solution**

5. Statement I: Finer the capillary, greater is the height to which the liquid rises in the tube

Statement II: This is in accordance with the ascent formula.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true and Statement II is NOT the correct explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: A**



**Watch Video Solution**

**6.** Statement I: A raindrop after falling through some height attains a constant velocity.

Statement II: At constant velocity, the viscous drag is just equal to its weight.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true and Statement II is NOT the correct explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: A**



**Watch Video Solution**

7. Statement I: A needle placed carefully on the surface of water may float, whereas the ball of the same material will always sink.

Statement II: The buoyancy of an object depends both on the material and shape of the object.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true  
and Statement II is NOT the correct  
explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: C**



**Watch Video Solution**

**8.** Statement I: Dust particles generally settle down in a closed room.

Statement II: The terminal velocity is inversely proportional to the square of their radii.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true and Statement II is NOT the correct explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: C**



**Watch Video Solution**

**9.** Statement I: Smaller drops of liquid resist deforming forces better than the larger drops.

Statement II: Excess pressure inside a drop is directly proportional to its surface area.



A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true and Statement II is NOT the correct explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: C**



**Watch Video Solution**

**10.** Statement I: An object from a greater height reaches a steady terminal velocity.

Statement II: The viscous forces on a body depends upon its velocity. The greater the velocity the greater is the viscous force.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true  
and Statement II is NOT the correct  
explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: B**



**Watch Video Solution**

**11.** Statement I: Spraying of water causes cooling.

Statement II: For an isolated system, surface energy increase on the expense of internal energy.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true and Statement II is NOT the correct

explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: A**



**Watch Video Solution**

**12.** Statement I: While blowing a soap bubble. to increase the size of soap bubble, we have to increase the air pressure within the soap bubble.

Statement II: To increase the size of soap bubble more air has to be pushed into the bubble.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true and Statement II is NOT the correct explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: D**



**Watch Video Solution**

**13.** Statement I: More is the cohesive force, more is the surface tension.

Statement II: More cohesive force leads to more shrinking of liquid surface.

A. Statement I is true, statement II is true and Statement II is a correct explanation for Statement I.

B. Statement I is true, Statement II is true and Statement II is NOT the correct explanation for Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

**Answer: A**



**Watch Video Solution**




## Linked Comprehension

1. A light rod of length  $L = 2m$  is suspended horizontally from the ceiling by two wires  $A$  and  $B$  of equal lengths. The wire  $A$  is made of steel with the area of cross section  $A_s = 1 \times 10^{-5} m^2$ , while the wire  $B$  is made of brass of cross sectional area  $A_b = 2 \times 10^{-5} m^2$ . A weight  $W$  is suspended at a distance  $x$  from the wire  $A$  as shown in figure.

Take, Young's modulus of steel and brass as

$$Y_s = 2 \times 10^{11} \text{ Nm}^{-2} \quad \text{and}$$

$$Y_b = 1 \times 10^{11} \text{ Nm}^{-2}.$$



Determine the value of  $x$  so that equal stresses are produced in each wire.

A.  $1.33m$

B.  $2.5m$

C.  $3.6m$

D.  $2.1m$

**Answer: A**



**Watch Video Solution**


2. A light rod of length  $L = 2m$  is suspended horizontally from the ceiling by two wires  $A$  and  $B$  of equal lengths. The wire  $A$  is made of steel with the area of cross section

$A_s = 1 \times 10^{-5} m^2$ , while the wire  $B$  is made of brass of cross sectional area  $A_b = 2 \times 10^{-5} m^2$ . A weight  $W$  is suspended at a distance  $x$  from the wire  $A$  as shown in figure.

Take, Young's modulus of steel and brass as

$$Y_s = 2 \times 10^{11} Nm^{-2} \quad \text{and}$$

$$Y_b = 1 \times 10^{11} Nm^{-2}.$$



Determine the value of  $x$  so that equal strains are produced in each wire

A.  $1m$

B.  $2m$

C.  $3m$

D.  $2.2m$

**Answer: A**



**Watch Video Solution**

**3.** A lead sphere of  $1.0mm$  diameter and relative density  $11.20$  attains a terminal velocity of  $0.7cms^{-1}$  in a liquid of relative density  $1.26$ .

Determine the coefficient of dynamic viscosity of the liquid.

A.  $0.45N / m^2$

B.  $0.85N / m^2$

C.  $0.56N / m^2$

D.  $0.77N / m^2$

**Answer: D**



**Watch Video Solution**

4. A lead sphere of  $1.0mm$  diameter and relative density  $11.20$  attains a terminal velocity of  $0.7cms^{-1}$  in a liquid of relative

density 1.26.

What is the value of the Reynolds number?

A. 0.01

B. 0.03

C. 0.15

D. 0.26

**Answer: A**



**Watch Video Solution**



5. A long capillary tube of radius  $0.2\text{mm}$  is placed vertically inside a beaker of water.

If the surface tension of water is  $7.2 \times 10^{-2}\text{N/m}$  the angle of contact between glass and water is zero, then determine the height of the water column in the tube.

A.  $3\text{cm}$

B.  $9\text{cm}$

C.  $7\text{cm}$

D.  $5\text{cm}$

**Answer: C**



**Watch Video Solution**

6. A long capillary tube of radius  $0.2\text{mm}$  is placed vertically inside a beaker of water.

If the tube is now pushed into water so that only  $5.0\text{cm}$  of its length is above the surface, then determine the angle of contact between the liquid and glass surface.

A.  $\cos^{-1}\left(\frac{4}{5}\right)$

B.  $\cos^{-1}\left(\frac{5}{7}\right)$

C.  $\cos^{-1}\left(\frac{3}{5}\right)$

D.  $\cos^{-1}\left(\frac{5}{4}\right)$

**Answer: B**



**Watch Video Solution**

7. An oil of relative density 0.9 and viscosity  $0.12 \text{ kg/ms}$  flows through a  $2.5 \text{ cm}$  diameter pipe with a pressure drop of  $38.4 \text{ kN/m}^2$  in a

length of  $30m$ . Determine

Determine the discharge

A.  $2.16 \times 10^{-4} m^3 / s$

B.  $2.9 \times 10^{-3} m^3 / s$

C.  $1 \times 10^{-4} m^3 / s$

D.  $2 \times 10^{-4} m^3 / s$

**Answer: C**



**Watch Video Solution**

8. An oil of relative density 0.9 and viscosity  $0.12 \text{ kg/ms}$  flows through a  $2.5 \text{ cm}$  diameter pipe with a pressure drop of  $38.4 \text{ kN/m}^2$  in a length of  $30 \text{ m}$ . Determine

Determine the shear stress at the pipe wall

A.  $8 \times 10^{-6} \text{ N/m}^2$

B.  $3.9 \times 10^{-6} \text{ N/m}^2$

C.  $2.3 \times 10^{-6} \text{ N/m}^2$

D.  $10.6 \times 10^{-6} \text{ N/m}^2$

**Answer: A**



Watch Video Solution

9. An oil of relative density 0.9 and viscosity  $0.12 \text{ kg/ms}$  flows through a  $2.5 \text{ cm}$  diameter pipe with a pressure drop of  $38.4 \text{ kN/m}^2$  in a length of  $30 \text{ m}$ . Determine  
Determine the power required to maintain the flow

A.  $2.2 \text{ W}$

B.  $3.84 \text{ W}$

C.  $5.6 \text{ W}$

D. 9.3W


**Answer: B**



**Watch Video Solution**

**10.** A steel bolt of cross-sectional area  $A_b = 5 \times 10^{-5} m^2$  is passed through a cylindrical tube made of aluminium. Cross-sectional area of the tube material is  $A_t = 10^{-4} m^2$  and its length is  $l = 50 cm$ . The bolt is just taut so that there is no stress in

the bolt and temperature of the assembly increases through  $\Delta \theta = 10^\circ C$ . Given, coefficient of linear thermal expansion of steel,  $\alpha_b = 10^{-5} / ^\circ C$ .



Young's modulus of steel

$$Y_b = 2 \times 10^{11} N/m^2$$

Young's modulus of Al,  $Y_t = 10^{11} N/m^2$ ,

coefficient of linear thermal expansion of

$$Al \alpha_t = 2 \times 10^{-5} / ^\circ C$$

The compressive strain in tube is



A.  $10^{-4}$

B.  $5 \times 10^{-5}$

C.  $2 \times 10^{-3}$

D.  $10^{-6}$


**Answer: B**



**Watch Video Solution**

**11.** A steel bolt of cross-sectional area  $A_b = 5 \times 10^{-5} m^2$  is passed through a cylindrical tube made of aluminium. Cross-

sectional area of the tube material is  $A_t = 10^{-4} m^2$  and its length is  $l = 50 cm$ . The bolt is just taut so that there is no stress in the bolt and temperature of the assembly increases through  $\Delta \theta = 10^\circ C$ . Given, coefficient of linear thermal expansion of steel,  $\alpha_b = 10^{-5} / ^\circ C$ .



Young's modulus of steel

$$Y_b = 2 \times 10^{11} N / m^2$$

Young's modulus of Al,  $Y_t = 10^{11} N / m^2$ ,

coefficient of linear thermal expansion of  $Al$

$$\alpha(t) = 2 \times 10^{-5} / ^\circ C$$

The compressive stress in tube is

A.  $5 \times 10^6 N / m^2$

B.  $10^5 N / m^2$

C.  $10^8 N / m^2$


D.  $10^3 N / m^2$

**Answer: A**



**Watch Video Solution**

12. A steel bolt of cross-sectional area  $A_b = 5 \times 10^{-5} m^2$  is passed through a cylindrical tube made of aluminium. Cross-sectional area of the tube material is  $A_t = 10^{-4} m^2$  and its length is  $l = 50 cm$ . The bolt is just taut so that there is no stress in the bolt and temperature of the assembly increases through  $\Delta \theta = 10^\circ C$ . Given, coefficient of linear thermal expansion of steel,  $\alpha_b = 10^{-5} / ^\circ C$ .



Young's modulus of steel

$$Y_b = 2 \times 10^{11} N/m^2$$

Young's modulus of Al,  $Y_t = 10^{11} N/m^2$ ,

coefficient of linear thermal expansion of

$$Al\alpha_t = 2 \times 10^{-5} / ^\circ C$$

The tensile stress in bolt is

A.  $10^4 N/m^2$

B.  $10^7 N/m^2$

C.  $2 \times 10^8 N/m^2$

D.  $10^{10} N/m^2$

**Answer: B**



Watch Video Solution

13. On gradual loading , stress - strain relationship for a metal wire is as follows .


Within proportionality limit , stress  $\propto$  strain

or,  $\frac{\text{Stress}}{\text{strain}}$  = a constant for the material of

wire.

Two wires of same material have length and radius  $(L, r)$  and  $(2L, \frac{r}{2})$ . The ratio of their

young's moduli is



A. 1 : 2

B. 2 : 3

C. 2 : 1

D. 1 : 1

**Answer: D**



**Watch Video Solution**

**14.** On gradual loading , stress - strain relationship for a metal wire is as follows .

Within proportionality limit , stress


proportional to strain or,  $\frac{\text{Stress}}{\text{strain}} = \text{a constant}$

for the material of wire.

Just on crossing the yield region, the material



will have



- A. reduced stress
- B. increased stress
- C. breaking stress
- D. constant stress

**Answer: A::C**



**Watch Video Solution**

**15.** According to Hooke's law, within the elastic limit  $\text{stress/strain} = \text{constant}$ . This constant depends on the type of strain or the type of force acting. Tensile stress might result in compressional or elongative strain, however, a tangential stress can only cause a shearing strain. After crossing the elastic limit, the material undergoes elongation and beyond a

stage beaks. All modulus of elasticity are basically constants for the materials under stress.

If stress/strain is  $x$  in elastic region and  $y$  in the region of yield, then

A.  $x = y$

B.  $x > y$

C.  $x < y$

D.  $x = 2y$

**Answer: B**



**16.** Molecular forces exist between the molecules of a liquid in a container. The molecules on the surface have unequal force leading to a tension on the surface. If this is not compensated by a force, the equilibrium of the liquid will be a difficult task. This leads to an excess pressure on the surface. The nature of the meniscus can inform us of the direction of the excess pressure. The angle of contact of the liquid decided by the forces

between the molecules, air and container can make the angle of contact.

The direction of the excess pressure in the meniscus of a liquid of angle of contact  $2\pi / 3$  is

- A. upward
- B. downward
- C. horizontal
- D. cannot be determined

**Answer: A**



17. Molecular forces exist between the molecules of a liquid in a container. The molecules on the surface have unequal force leading to a tension on the surface. If this is not compensated by a force, the equilibrium of the liquid will be a difficult task. This leads to an excess pressure on the surface. The nature of the meniscus can inform us of the direction of the excess pressure. The angle of contact of the liquid decided by the forces

between the molecules, air and container can make the angle of contact.

If the excess pressure in a soap bubble is  $p$ , the excess pressure in an air bubble is

A.  $\frac{p}{2}$

B.  $p$

C.  $2p$

D.  $4p$

**Answer: A**



**Watch Video Solution**

**18.** Molecular forces exist between the molecules of a liquid in a container. The molecules on the surface have unequal force leading to a tension on the surface. If this is not compensated by a force, the equilibrium of the liquid will be a difficult task. This leads to an excess pressure on the surface. The nature of the meniscus can inform us of the direction of the excess pressure. The angle of contact of the liquid decided by the forces between the molecules, air and container can



make the angle of contact.

In a meniscus of radius  $r$ , with excess pressure  $p$  in atmospheric pressure  $p_0$ , the force experienced is

A.  $(p - p_0)pr^2$

B.  $(p - p_0)2\pi r$

C.  $p\pi r^2$

D.  $p_0 2\pi r$

**Answer: C**




**Watch Video Solution**

**19.** Materials get deformed when force is applied. Some of them regain their status when the applied force is removed. They are termed as elastic. Those of which not regaining are called plastic. There may be delay in the regaining in some materials. They are said to have got elastic aftereffect, since they have gone beyond the elastic limit. Repeated application and removal of force leads to fatigueness in the material. Fatigued materials may break at any point time and so

are avoided.

The stress strain graph for two materials  $A$  and  $B$  is shown in the following figure:



If the intensity of  $A$  and  $B$  is  $E_A$  and  $E_B$ , respectively

A.  $E_A = E_B$

B.  $E_A > E_B$

C.  $E_A < E_B$

D.  $E_A < < E_B$

**Answer: B**




**Watch Video Solution**

20. Materials get deformed when force is applied. Some of them regain their status when the applied force is removed. They are termed as elastic. Those of which not regaining are called plastic. There may be

delay in the regaining in some materials. They are said to have got elastic aftereffect, since they have gone beyond the elastic limit. Repeated application and removal of force leads to fatigueness in the material. Fatigued materials may break at any point time and so are avoided.

The stress strain graph for two materials  $A$  and  $B$  is shown in the following figure:



The strength of the material  $A$  and  $B$  is  $S_A$  and  $S_B$ , respectively, while the longevity of plastic behaviour is  $L_A$  and  $L_B$ . Then

A.  $S_A > S_B, L_A < L_B$

B.  $S_A = S_B, L_A = L_B$

C.  $S_A > S_B, L_A > L_B$

$$D. S_A < S_B, L_A < L_B$$

**Answer: C**




**Watch Video Solution**

**21.** Materials get deformed when force is applied. Some of them regain their status when the applied force is removed. They are termed as elastic. Those of which not regaining are called plastic. There may be delay in the regaining in some materials. They

are said to have got elastic aftereffect, since they have gone beyond the elastic limit. Repeated application and removal of force leads to fatigueness in the material. Fatigued materials may break at any point time and so are avoided.

The stress strain graph for two materials A and B is shown in the following figure:





The time in which the two materials regain their original status is  $t_A$  and  $t_B$  related as  $t_B = 2t_A$ . Then the material under elastic aftereffect (relatively) is

A.  $B$

B.  $A$

C. both  $A$  and  $B$


D. neither  $A$  nor  $B$

**Answer: A**



**Watch Video Solution**

22. Figure shows a capillary tube of radius  $r$  dipped into water. If the atmosphere pressure is  $P_0$ , the pressure at point A is



A.  $P_0$

B.  $P_0 + \frac{2s}{r}$

C.  $P_0 - \frac{2s}{r}$


D.  $P_0 - \frac{4s}{r}$

**Answer: C**



**Watch Video Solution**

**23.** Figure shows a capillary tube of radius  $r$  dipped into The atmospheric pressure is  $P_0$  and the capillary rise of water is  $h$ .  $s$  is the surface tension for water-glass.



Initially,  $h = 10\text{cm}$ . If the capillary tube is now incline at  $45^\circ$ , the length of water rising in the tube will be

A.  $10\text{cm}$

B.  $10\sqrt{2}\text{cm}$

C.  $\frac{10}{\sqrt{2}}\text{cm}$

D. none of these

**Answer: B**



**Watch Video Solution**

**24.** Figure shows a capillary tube of radius  $r$  dipped into The atmospheric pressure is  $P_0$  and the capillary rise of water is  $h$ .  $s$  is the

surface tension for water-glass.



Initially,  $h = 10\text{cm}$ . If the capillary tube is now incline at  $45^\circ$ , the length of water rising in the tube will be

A.

B.

C.

D.

**Answer: C**



25. In the figure shown,  $A$  and  $B$  are two short steel rods each of cross-sectional area  $5\text{cm}^2$ . The lower ends of  $A$  and  $B$  are welded to a fixed plate  $CD$ . The upper end of  $A$  is welded to the  $L$ -shaped piece  $EFG$ , which can slide without friction on upper end of  $B$ . A horizontal pull of  $1200\text{N}$  is exerted at  $G$  as shown. Neglect the weight of  $EFG$ .



Mark out the correct statement(s).

A. Shearing stress in  $A$  is zero.

B. Shearing stress in  $B$  is zero

C. Shearing stress in both  $A$  and  $B$  is zero

D. Shearing stress in both  $A$  and  $B$  is non-zero

**Answer: B**



**Watch Video Solution**



**26.** In the figure shown,  $A$  and  $B$  are two short steel rods each of cross-sectional area  $5\text{cm}^2$ . The lower ends of  $A$  and  $B$  are welded to a fixed plate  $CD$ . The upper end of  $A$  is welded to the  $L$ -shaped piece  $EFG$ , which can slide without friction on upper end of  $B$ . A horizontal pull of  $1200\text{N}$  is exerted at  $G$  as shown. Neglect the weight of  $EFG$ .



Longitudinal stress in  $A$  is

A. tensile in nature and having magnitude

$$180N / m^2$$

B. tensile in nature and having magnitude

$$240N / m^2$$

C. compressive in nature and having

$$\text{magnitude } 180N / m^2$$

D. compressive in nature and having

$$\text{magnitude } 240N / m^2$$

**Answer: A**



**Watch Video Solution**

27. In the figure shown,  $A$  and  $B$  are two short steel rods each of cross-sectional area  $5\text{cm}^2$ . The lower ends of  $A$  and  $B$  are welded to a fixed plate  $CD$ . The upper end of  $A$  is welded to the  $L$ -shaped piece  $EFG$ , which can slide without friction on upper end of  $B$ . A horizontal pull of  $1200\text{N}$  is exerted at  $G$  as shown. Neglect the weight of  $EFG$ .



Longitudinal stress in  $B$  is

A. tensile in nature and having magnitude

$$180N / cm^2$$

B. tensile in nature and having magnitude

$$240N / cm^2$$

C. compressive in nature and having

$$\text{magnitude } 180N / cm^2$$

D. compressive in nature and having

$$\text{magnitude } 240N / cm^2$$

**Answer: C**




**Watch Video Solution**

28. Two opposite forces  $F_1 = 120N$  and  $F_2 = 80N$  act on an elastic plank of modulus of elasticity  $Y = 2x10^{11}N/m^2$  and length  $l = 1m$  placed over a smooth horizontal surface. The cross-sectional area of the plank is  $S = 0.5m^2$ . The change in length of the plank is  $x \times 10^{-11}m$ . Find the value of  $x$ .



[Watch Video Solution](#)

29. A ring of radius  $r$  made of wire of density  $\rho$  is rotated about a stationary vertical axis passing through its centre and perpendicular to the plane of the ring as shown in the figure. Determine the angular velocity (in rad/s) of ring at which the ring breaks. The wire breaks at tensile stress  $\sigma$ . Ignore gravity. Take  $\sigma / \rho = 4$  and  $r = 1m$ .



**30.** A cube of side  $a$  and mass  $m$  just floats on the surface of water as shown in the figure. The surface tension and density of water are  $T$  and  $\rho_w$  respectively. If angle of contact between cube and water surface is zero, find the distance  $h$  (in metres) between the lower face of cube and surface of the water.

(Take  $m = 1\text{kg}$ ,  $g = 10\text{ms}^{-2}$ ,  $aT = \frac{10}{4}$  unit and  $\rho_w a^2 g = 10$  unit)





Watch Video Solution

## Integer

1. A thin plate  $AB$  of large area  $A$  is placed symmetrically in a small gap of height  $h$  filled with water of viscosity  $\eta_0$  and the plate has a constant velocity  $v$  by applying a force  $F$  as shown in the figure. If the gap is filled with some other liquid of viscosity  $0.75\eta_0$  at what minimum distance (in cm) from top wall should the plate be placed in the gap, so that



the plate can again be pulled at the same constant velocity  $V$ . by applying the same force  $F$ ? (Take  $h = 20\text{cm}$ )



[Watch Video Solution](#)

2. The diameter of a gas bubble formed at the bottom of a pond is  $d = 4\text{cm}$ . When the bubble rises to the surface, its diameter tension of water  $= T = 0.07\text{Nm}^{-1}$



[Watch Video Solution](#)

3.  $n$  drops of water, each of radius  $2\text{mm}$ , fall through air at a terminal velocity of  $8\text{cm}^{-1}$ . If they coalesce to form a single drop, then the terminal velocity of the combined drop is  $32\text{cm}^{-1}$ . The value of  $n$  is



[Watch Video Solution](#)

4. A substance breaks down under a stress of  $10^5\text{Pa}$ . If the density of the wire is  $2 \times 10^3\text{kg}/\text{m}^3$ , find the minimum length of

the wire which will break under its own weight

$$(g = 10ms^{-2}).$$



**Watch Video Solution**

5. A wire of length  $L$  and cross sectional area  $A$  is made of a material of Young's modulus  $Y$ . If the wire is stretched by an amount  $x$ , the work done is.....



**Watch Video Solution**


6. A solid sphere of radius  $R$  made of a material of bulk modulus  $K$  is surrounded by a liquid in a cylindrical container. A massless piston of area  $A$  floats on the surface of the liquid. When a mass  $M$  is placed on the piston to compress the liquid the fractional change in the radius of the sphere,  $\delta R / R$ , is .....



[Watch Video Solution](#)

**Fill In The Blanks**

1. A uniform rod of length  $L$  and density  $\rho$  is being pulled along a smooth floor with a horizontal acceleration  $\alpha$  (see Fig.) The magnitude of the stress at the transverse cross-section through the mid-point of the rod is.....



[Watch Video Solution](#)

1. The following four wires are made of the same material. Which of these will have the largest extension when the same tension is applied?

A. Length =  $50\text{cm}$ , diameter =  $0.5\text{mm}$

B. Length  $100\text{cm}$ , diameter =  $1$

C. Length =  $200\text{cm}$ , diameter =  $2\text{mm}$

D. Length =  $300\text{cm}$ , diameter =  $3\text{mm}$

**Answer: A**



**Watch Video Solution**

2. Two rods of different materials having coefficients of thermal expansion  $\alpha_1$ ,  $\alpha_2$  and Young's moduli  $Y_1$ ,  $Y_2$  respectively are fixed between two rigid massive walls. The rods are heated such that they undergo the same increase in temperature. There is no bending of the rods. If  $\alpha_1 : \alpha_2 = 2 : 3$ , the thermal

stresses developed in the two rods are equal  
provided  $Y_1 : Y_2$  is equal to

A. 2 : 3

B. 1 : 1

C. 3 : 2

D. 4 : 9


**Answer: C**



**Watch Video Solution**



3. The adjacent graph shows the estension ( $\Delta l$ ) of a wire of length 1m suspended from the top of a roof at one end and with a load  $W$  connected to the other end. If the cross-sectional area of the wire is  $10^{-6}m^2$ , calculate the Young's modulus of the material of the wire.



A.  $2 \times 10^{11} N/m$

B.  $2 \times 10^{11} N/m$

C.  $3 \times 10^{-12} N/m$

D.  $2 \times 10^{-3} N/m$

**Answer: A**



**Watch Video Solution**

4. When temperature of a gas is  $20^\circ C$  and pressure is changed from  $p_1 = 1.01 \times 10^5 Pa$

to  $p_2 = 1.165 \times 10^5 Pa$ , the volume changes by 10 %. The bulk modulus is

A.  $1.55 \times 10^5 Pa$

B.  $0.115 \times 10^5 Pa$

C.  $1.4 \times 10^5 Pa$


D.  $1.01 \times 10^5 Pa$

**Answer: A**



**Watch Video Solution**

5. A glass tube of uniform internal radius( $r$ ) has a valve separating the two identical ends. Initially, the valve is in a tightly closed position. Initially, the valve is in a tightly closed position. End 1 has a hemispherical soap bubble of radius  $r$ . End 2 has sub-hemispherical soap bubble as shown in figure. Just after opening the valve,



A. Air from end 1 flows towards end 2.

There is no change in the volume of the soap bubble.

B. Air from end 1 flows towards end 2.

Volume of the soap bubble at end 1 decreases.

C. No change occurs.

D. Air from end 2 flows towards end 1.

Volume of the soap bubble at end 1 increases.

**Answer: B**



**Watch Video Solution**

6. One end of a horizontal thick copper wire of length  $2L$  and radius  $2R$  is welded to an end of another horizontal thin copper wire of length  $L$  and radius  $R$ . When the arrangement is stretched by applying forces at two ends, the ratio of the elongation in the thin wire to that in the thick wire is

A. 0.25

B. 0.50

C. 2.00

D. 4.00

**Answer: C**



**Watch Video Solution**

**LC\_TYPE**

1. When liquid medicine of density  $\rho$  is to be put in the eye, it is done with the help of a dropper. As the bulb on the top of the dropper is pressed, a drop forms at the opening of the dropper. We wish to estimate the size of the drop. We first assume that the drop formed at the opening is spherical because that requires a minimum increase in its surface energy. To determine the size, we calculate the net vertical force due to the surface tension  $T$  when the radius of the drop is  $R$ . When this force becomes smaller than the weight of the



drop, the drop gets detached from the dropper.

If the radius of the opening of the dropper is  $r$ , the vertical force due to the surface tension on the drop of radius  $R$  (assuming  $r \ll R$ ) is

A.  $2\pi rT$

B.  $2\pi RT$

C.  $2\pi r^2T / R$

D.  $2\pi R^2T / r$

**Answer: C**



2. When liquid medicine of density  $\rho$  is to put in the eye, it is done with the help of a dropper. As the bulb on the top of the dropper is pressed, a drop forms at the opening of the dropper. We wish to estimate the size of the drop. We first assume that the drop formed at the opening is spherical because that requires a minimum increase in its surface energy. To determine the size, we calculate the net vertical force due to the surface tension  $T$

when the radius of the drop is  $R$ . When this force becomes smaller than the weight of the drop, the drop gets detached from the dropper.

If  $r = 5 \times 10^{-4} m$ ,  $\rho = 10^3 kgm^{-3}$ ,  
 $g = 10ms^{-2}$ ,  $T = 0.11Nm^{-1}$ , the radius of the drop when it detaches from the dropper is approximately

A.  $1.4 \times 10^{-3} m$

B.  $3.3 \times 10^{-3} m$

C.  $2.0 \times 10^{-3} m$

$$D. 4.1 \times 10^{-3}m$$

**Answer: A**



**Watch Video Solution**

**3.** When liquid medicine of density  $\rho$  is to put in the eye, it is done with the help of a dropper. As the bulb on the top of the dropper is pressed, a drop forms at the opening of the dropper. We wish to estimate the size of the drop. We first assume that the drop formed at

the opening is spherical because that requires a minimum increase in its surface energy. To determine the size, we calculate the net vertical force due to the surface tension  $T$  when the radius of the drop is  $R$ . When this force becomes smaller than the weight of the drop, the drop gets detached from the dropper.

After the drop detaches, its surface energy is

A.  $1.4 \times 10^{-6} J$

B.  $2.7 \times 10^{-6} J$

C.  $5.4 \times 10^{-6} J$

$$D. 8.1 \times 10^{-6} J$$

**Answer: B**



**Watch Video Solution**

**INTEGER\_TYPE**

1. Two soap bubbles A and B are kept in a closed chamber where the air is maintained at pressure  $8N/m^2$ . The radii of bubbles A and B are 2cm and 4cm, respectively. Surface tension

of the soap-water used to make bubbles is  $0.04\text{N}/\text{m}$ . Find the ratio  $n_B/n_A$ , where  $n_A$  and  $n_B$  are the number of moles of air in bubbles A and B, respectively. [Neglect the effect of gravity.]



[Watch Video Solution](#)

2. A  $0.1\text{kg}$  mass is suspended from a wire of negligible mass. The length of the wire is  $1\text{m}$  and its cross sectional area is  $4.9 \times 10^{-7}\text{m}^2$ . If the mass is pulled a little in the vertically

downward direction and released, it performs simple harmonic motion of angular frequency  $140\text{rads}^{-1}$ . If the Young's modulus of the material of the wire is  $n \times 10^9 \text{Nm}^{-2}$ , the value of n is



**Watch Video Solution**