

PHYSICS

BOOKS - CENGAGE PHYSICS (HINGLISH)

THERMODYNAMICS

1

1. One mole of an ideal gas is warmed slowly so that it goes from the PV state $(P_f V_i)$ to

$(3P_i, 3V_i)$ in such a way that the pressure of the gas is directly proportional to the volume.

(a) How much work is done on the gas in the process?

(b) How is the temperature of the gas related to its volume during this process?



Watch Video Solution

2. We consider a thermodynamic system. If ΔU represents the increase in its internal

energy and W the work done by the system,
which of the following statements is true?

A. $\Delta U = -W$ in an adiabatic process

B. $\Delta U = W$ in an isothermal process

C. $\Delta U = -W$ in an isothermal process

D. $\Delta U = W$ in an adiabatic process

Answer:



Watch Video Solution

3. Two moles of a monatomic gas in state A having critical pressure P_0 and temperature $3T_0$ is taken to a state B having pressure $3P_0$ and temperature $T_0/3$ by the process of equation $P^2T = \text{constant}$. Then state B is taken to state C keeping the volume constant and it comes back to initial state A keeping temperature constant.

a. Plot a P and T graph. (P on the y-axis and T on the x-axis).

Find the net work done and heat supplied to the gas during the complete cycle.



Watch Video Solution

2

1. An ideal gas is taken through a quasi-static process described by $P = \alpha V^2$, with $\alpha = 5.00 \text{ atm} / \text{m}^6$. The gas is expanded to twice its original volume of 1.00 m^3 . How much work is done by the gas in expanding gas in this process?



Watch Video Solution

2. When heat is given to a gas in an isobaric process, then


- A. the work is done by the gas
- B. internal energy of the gas increases
- C. both (a) and (b)
- D. none from (a) and (b)

Answer:



Watch Video Solution

3. For one complete cycle of a thermodynamic process gas as shown in the P-V diagram, which of following correct?



A. $\Delta E_{\text{int}} = 0, Q < 0$

B. $\Delta E_{\text{int}} = 0, Q > 0$

C. $\Delta E_{\text{int}} > 0, Q < 0$

$$D. \Delta E_{\text{int}} < 0, Q > 0$$

Answer: A




Watch Video Solution

3

1. The cyclic process for 1 mole of an ideal gas is shown in the V-T diagram. The work done in

AB, BC and CA respectively is



A. $0, RT_2 \ln \left(\frac{V_1}{V_2} \right), R(T_1 - T_2)$

B. $R(T_1 - T_2), 0, RT_1 \ln \frac{V_1}{V_2}$

C. $0, RT_2 \ln \left(\frac{V_2}{V_1} \right), R(T_1 - T_2)$

D. $0, RT_2 \ln \left(\frac{V_2}{V_1} \right), R(T_2 - T_1)$

Answer: D



Watch Video Solution

2. Which of the following is correct in terms of increasing work done for the same initial and final state?

A. Adiabatic $<$ Isothermal $<$ Isobaric

B. Isobaric $<$ Adiabatic $<$ Isothermal


C. Adiabatic $<$ Isobaric $<$ Isothermal

D. None of these

Answer: d



Watch Video Solution



3.

A sample of an ideal gas is taken through the cyclic process $abca$ in the given figure. It absorbs 50 J of heat during the path ab , no

heat during bc and rejects 70J of heat during ca, 40 J of work is done on the gas during the part bc.

(a) Find the internal energy of the gas at b and c if it is 1500 J at a. (b) Calculate the work done by the gas during the part ca.

A. 1590J

B. 1620J

C. 1540J

D. 1570J

Answer: A




Watch Video Solution

4

1. A thermodynamic system undergoes cyclic process $ABCD$ as shown in figure. The work

done by the system is



- A. P_0V_0
- B. $2P_0V_0$
- C. $\frac{P_0V_0}{2}$
- D. Zero

Answer: D



Watch Video Solution

2. When an ideal diatomic gas is heated at constant pressure, the fraction of the heat energy supplied, which increases the internal energy of the gas, is

A. $\frac{2}{5}$

B. $\frac{3}{5}$

C. $\frac{3}{7}$

D. $\frac{5}{7}$

Answer: d



Watch Video Solution

3. A thermodynamic process of one mole ideal monoatomic gas is shown in figure. The efficiency of cyclic process $ABCA$ will be

A. 25 %

B. 12.5 %

C. 50 %

D. $\frac{100}{13}$ %

Answer: D




Watch Video Solution

5

1. Consider a process shown in the figure.
During this process the work done by the

system



- A. Continuously increases
- B. Continuously decreases
- C. First increases, then decreases
- D. First decreases, then increases

Answer: A



Watch Video Solution

2. The molar heat capacity in a process of a diatomic gas if it does a work of $\frac{Q}{4}$ when a heat of Q is supplied to it is

A. $\frac{2}{5}R$

B. $\frac{5}{2}R$

C. $\frac{10}{3}R$

D. $\frac{6}{7}R$


Answer: a



Watch Video Solution

3. An ideal gas is taken from state 1 to state 2 through optional path A , B , C and D as shown in the PV diagram. Let Q , W and U represent the heat supplied, work done and change in internal energy of the gas respectively.

Then,



A. $Q_A - Q_D = W_A - W_D$

B. $Q_B - W_B > Q_C - W_C$

C. $W_A < W_B < W_C < W_D$

D. $Q_A < Q_B < Q_C < Q_D$

Answer: A




Watch Video Solution

6

1. Six moles of an ideal gas performs a cycle shown in figure. If the temperature are $T_D = 600K$, $T_B = 800K$, $T_C = 2200K$ and

$T_D = 1200K$, the work done per cycle is



A. 20 kJ

B. 30 kJ

C. 40 kJ

D. 60 kJ

Answer: C



Watch Video Solution

2. An insulator container contains 4 moles of an ideal diatomic gas at temperature T . Heat Q is supplied to this gas, due to which 2 moles of the gas are dissociated into atoms but temperature of the gas remains constant.

Then

A. $Q = 2RT$

B. $Q = RT$

C. $Q = 3RT$

D. $Q = 4RT$


Answer: C



Watch Video Solution

3. Following figure shows on adiabatic cylindrical container of volume V_0 divided by an adiabatic smooth piston (area of cross-section = A) in two equal parts. An ideal gas

$(C_p/C_v = \lambda)$ is at pressure P_1 and temperature T_1 in left part and gas at pressure P_2 and temperature T_2 in right part. The piston is slowly displaced and released at a position where it can stay in equilibrium. The final pressure of the two parts will be (Suppose x = displacement of the piston)



A. P_2

B. P_1

C.
$$\frac{P_1 \left(\frac{V_0}{2} \right)^\gamma}{\left(\frac{V_0}{2} + Ax \right)^\gamma}$$


D.
$$\frac{P_2 \left(\frac{V_0}{2} \right)^\gamma}{\left(\frac{V_0}{2} + Ax \right)^\gamma}$$

Answer: C



Watch Video Solution

1. $P - V$ diagram of an ideal gas is as shown in figure. Work done by the gas in process $ABCD$ is



A. $4P_0V_0$

B. $2P_0V_0$

C. $3P_0V_0$

D. P_0V_0

Answer: C



Watch Video Solution

2. Which one of the following gases possesses the largest internal energy

A. 2 moles of helium occupying 1 m^3 at 300

K

B. 56 kg of nitrogen at $10^7 Nm^{-2}$ and 300

K

C. 8 grams of oxygen at 8 atm and 300 K

D. 6×10^{26} molecules of argon occupying


$40m^3$ at 900 K

Answer: c



Watch Video Solution

3. In the following P-V diagram two adiabatics cut two isothermals at temperature T_1 and T_2 (fig). The value of $\frac{V_a}{V_d}$ will be



- A. $\frac{V_b}{V_c}$
 B. $\frac{V_c}{V_b}$

C. $\frac{V_d}{V_a}$

D. $V_b V_c$

Answer: A



Watch Video Solution

8

1. A gas expands with temperature according to the relation $V = kT^{2/3}$. What is the work

done when the temperature changes by $30^{\circ}C$

?

A. $10R$

B. $20R$

C. $30R$


D. $40R$

Answer: B



Watch Video Solution

2. In the figure given two processes A and B are shown by which a thermodynamic system goes from initial to final state F . if ΔQ_A and ΔQ_B are respectively the heats supplied to the systems then



A. $\Delta Q_A = \Delta Q_B$

B. $\Delta Q_A \geq \Delta Q_B$

C. $\Delta Q_A < \Delta Q_B$

D. $\Delta Q_A > \Delta_B$


Answer: b



Watch Video Solution

3. An ideal gas undergoes a cyclic process abcda which is shown by pressure-density

curve,



A. Work done by the gas in the process 'bc'

is zero

B. Work done by the gas in the process 'cd'

is negative

C. Internal energy of the gas at point 'a' is greater than at state 'c'

D. Net work done by the gas in the cycle is neagative.


Answer: A,B,D



Watch Video Solution

1. One mole of an ideal gas is kept enclosed under a light piston (area = $10^{-2}m^2$) connected by a compressed spring (spring constant $100N/m$). The volume of gas is $0.83m^3$ and its temperature is $100K$. The gas is heated so that it compresses the spring further by $0.1m$. The work done by the gas in the process is $N \times 10^{-1}J$. Find N . (Take $R = 8.3J/K - \text{mole}$) and suppose there is

no atmosphere).



A. $3J$

B. $6J$

C. $9J$

D. $1.5J$


Answer: D



Watch Video Solution

2. When a system is taken from state f along path iaf , $Q = 50J$ and $W = 20J$. Along path ibf , $Q = 35J$. If $W = -13J$ for the curved

return path fI, Q for this path is



A. 33 J

B. 23 J

C. $-7J$

D. $-43J$

Answer: d



Watch Video Solution

3. An ideal gas can be expanded from an initial state to a certain volume through two different processes $PV^2 = \text{constant}$ and (ii) $P = KV^2$ where K is a positive constant.

Then

A. Final temperature in (i) will be greater than in (ii)

B. Final temperature in (ii) will be greater than in (i)

C. Total heat given to the gas in (i) case is greater than in (ii)


D. Total heat is given to the gas in (ii) case is greater than in (i)

Answer: B,D



Watch Video Solution

1. n moles of a gas fill in a container temperature T is in thermodynamic equilibrium initially. If the gas is compressed slowly and isothermally so half its initial volume the work done by the atmosphere on the piston is :



A. $\frac{nRT}{2}$

B. $-\frac{nRT}{2}$

C. $nRT \left(\ln 2 - \frac{1}{2} \right)$

D. $-nRT \ln 2$


Answer: A



Watch Video Solution

2. The $P - V$ diagram of a system undergoing thermodynamic transformation is shown in figure. The work done by the system in going

from $A \rightarrow B \rightarrow C$ is $30J$ and $40J$ heat is given to the system. The change in internal energy between A and C is




- A. 10 J
- B. 70 J
- C. 84 J
- D. 134 J

Answer: a



Watch Video Solution

3. A monatomic ideal gas of 2 mol is taken through a cyclic process starting from A as shown in figure. The volume ratios are $V_B/V_A = 2$ and $V_D/V_A = 4$. If the temperature T_A at A is 27°C , and gas constant is R . Calculate.



The temperature of the gas at point B

A. 600K

B. 450K

C. 400K

D. 900K


Answer: A



Watch Video Solution

11

1. Find the work done by the gas in the process *ABC*.



A. $\frac{3}{2}P_0V_0$

B. $\frac{5}{2}P_0V_0$

C. $\frac{7}{2}P_0V_0$

D. $4P_0V_0$


Answer: C



Watch Video Solution

2. The $P - V$ diagram of 2 gm of helium gas for a certain process $A \rightarrow B$ is shown in the figure. What is the heat given to the gas

during the process $A \rightarrow B$?



A. $4P_0V_0$

B. $6P_0V_0$


C. $4.5P_0V_0$

D. $2P_0V_0$

Answer: c



Watch Video Solution



3.

A monoatomic ideal gas of two moles is taken through a cyclic process starting from A as

shown

$$\frac{V_B}{V_A} = 2 \text{ and } \frac{V_D}{V_A} = 4$$

Temperature T_A at A is $27^\circ C$.

Q. Work done during complete cyclic process

A. 1200R

B. 1500R

C. 1400R

D. 1000R

Answer: B,D



Watch Video Solution


1. In an isothermal reversible expansion, if the volume of 96 gm of oxygen at 27°C is increased from 70 litres to 140 litres, then the work done by the gas will be



[Watch Video Solution](#)

2. Volume versus temperature graph of two moles of helium gas is as shown in figure. The

ratio of heat absorbed and the work done by the gas in process 1 – 2 is



- A. 3
- B. $\frac{5}{2}$
- C. $\frac{5}{3}$
- D. $\frac{7}{2}$

Answer: B



Watch Video Solution


3. A monatomic ideal gas of two moles is taken through a cyclic process starting from A as shown

$$V_B/V_A = 2 \text{ and } V_D/V_A = 4$$

Temperature T_A is 27°C

The work done during process $B \rightarrow C$

(approx) is



A. 1000R

B. 800R

C. 832R

D. 945R

Answer: C



[Watch Video Solution](#)

13

1. 540 calories of heat convert 1 cubic centimeter of water at $100^{\circ}C$ into 1671 cubic centimeter of steam at $100^{\circ}C$ at a pressure of one atmosphere. Then the work done against the atmospheric pressure is nearly



[Watch Video Solution](#)

2. Heat is supplied to a diatomic gas at constant pressure.

The ratio of $\Delta Q : \Delta U : \Delta W$ is

A. 5 : 3 : 2

B. 5 : 2 : 3

C. 7 : 5 : 2

D. 7 : 2 : 5


Answer: C

3. A monatomic ideal gas of two metal is taken through a cyclic process straining from A as

shown $V_B/V_A = 2$ and $V_D/V_A = 4$

Temperature T_A is $27^\circ C$

The work done during the process $C \rightarrow D$ is



A. 900R (absorbed)

B. 900R(released)

C. 1200R(absorbed)

D. 1200R(released)

Answer: B,D




Watch Video Solution

14

1. A system changes from the state (P_1, V_1) to (P_2, V_2) as shown in the diagram. The

workdone by the system is



[Watch Video Solution](#)

2. N moles of an ideal diatomic gas are in a cylinder at temperature T . Suppose on supplying heat to the gas, its temperature

remain constant but n moles get dissociated into atoms. Heat supplied to the gas is

A. zero

B. $\frac{1}{2}nRT$

C. $\frac{3}{2}nRT$

D. $\frac{3}{2}(N - n)RT$

Answer: B




Watch Video Solution

3. A monatomic ideal gas of two metal is taken through a cyclic process straining from A as shown $V_B/V_A = 2$ and $V_D/V_A = 4$

Temperature T_A is $27^\circ C$

The work done during the process $D \rightarrow A$ is



A. 900R (absorbed)

B. 900R(released)

C. 1200R(absorbed)


D. 1200R(released)

Answer: D



Watch Video Solution

1. A sample of an ideal monoatomic gas is taken round the cycle ABCA as shown in the figure the work done during the cycle is



[Watch Video Solution](#)

2. In thermodynamic process, pressure of a fixed mass of a gas is changes in such a manner that the gas molecules gives out 20 J of heat and 10 J of work is done in the gas. If the initial internal energy of the gas was 40 J, then the final internal energy will be




[Watch Video Solution](#)

3. A monatomic ideal gas of two metal is taken through a cyclic process straining from A as shown

$$V_B/V_A = 2 \text{ and } V_D/V_A = 4$$

Temperature T_A is 27°C

The work done during the entire process is



A. 400R

B. 600R

C. 450R

D. 800R

Answer: B



Watch Video Solution

16

1. A perfect gas goes from a state A to another state B by absorbing 8×10^5 J of heat and doing 6.5×10^5 J of external work. It is now transferred between the same two states in

another process in which it absorbs 105 J of heat. In the second process



[Watch Video Solution](#)

17

1. The specific heat of a gas in an isothermal process is



[Watch Video Solution](#)

1. A thermally insulated container is divided into two parts by a screen. In one part the pressure and temperature are P and T for an ideal gas filled. In the second part it is vacuum. If now a small hole is created in the screen, then the temperature of the gas will



[Watch Video Solution](#)

1. When an ideal gas in a cylinder was compressed isothermally by a piston, the work done on the gas found to be 1.5×10^4 cal. During this process about



[Watch Video Solution](#)

20

1. In adiabatic expansion ΔU is ...
(positive/negative/zero)



[Watch Video Solution](#)

21

1. A gas at NTP is suddenly compressed to one-fourth of its original volume. If λ is supposed to be $\frac{3}{2}$, then the final pressure is



[Watch Video Solution](#)

22

1. An ideal gas at $27^{\circ}C$ is compressed adiabatically to $8/27$ of its original volume. If $\gamma = 5/3$, then the rise in temperature is



[Watch Video Solution](#)

23

1. Two identical samples of gases are allowed to expand to the same final volume (i) isothermally (ii) adiabatically. Work done is



[Watch Video Solution](#)

1. During the adiabatic expansion of 2 moles of a gas, the internal energy of the gas is found to decrease by 2 joules , the work done during the process on the gas will be equal to



[Watch Video Solution](#)