

PHYSICS


BOOKS - CENGAGE PHYSICS (HINGLISH)

TRANSMISSION OF HEAT

Single Correct Answer Type

1. Two bars of thermal conductivities K and $3K$ and lengths 1cm and 2cm respectively

have equal cross-sectional area, they are joined length wise as shown in the figure. If the temperature at the ends of this composite bar is 0°C and 100°C respectively (see figure), then the temperature ϕ of the interface is



A. 50°C

B. $\left(\frac{100}{3}\right)^\circ\text{C}$

C. 60°C

D. $\left(\frac{200}{3}\right)^{\circ} \text{C}$

Answer: C



Watch Video Solution

2. A wall has two layers A and B each made of different materials. Both the layers have the same thickness. The thermal conductivity of materials A is twice of B. Under thermal equilibrium temperature difference across the

layer B is 36°C . The temperature difference across layer A is

A. 6°C

B. 12°C

C. 18°C

D. 24°C


Answer: B



Watch Video Solution

3. A cylinder of radius R made of a material of thermal conductivity K_1 is surrounded by cylindrical shell of inner radius R and outer radius $2R$ made of a material of thermal conductivity K_2 . The two ends of the combined system are maintained at two different temperatures. There is no loss of heat across the cylindrical surface and system is in steady state. What is the effective thermal

conductivity of the system



A. $K_1 + K_2$

B. $\frac{K_1 K_2}{K_1 + K_2}$

C. $\frac{K_1 + 3K_2}{4}$

D. $\frac{3K_1 + K_2}{4}$

Answer: C




Watch Video Solution

4. Three rods made of the same material and having the same cross-section have been joined as shown in the figure. Each rod is of the same length. The left and right ends are kept at $0^{\circ}C$ and $90^{\circ}C$, respectively. The temperature of junction of the three rods will be

(a) $45^{\circ}C$ (b) $60^{\circ}C$

(c) $30^{\circ}C$ (d) $20^{\circ}C$.



A. $45^{\circ}C$

B. $60^{\circ}C$

C. $30^{\circ}C$

D. $20^{\circ}C$

Answer: B



Watch Video Solution

5. Ice has formed on a shallow pond, and a steady state has been reached, with the air above the ice at $-5.0^{\circ}C$ and the bottom of the pond at $40^{\circ}C$. If the total depth of ice + water is $1.4m$, (Assume that the thermal conductivities of ice and water are 0.40 and $0.12cal / mC^{\circ} s$, respectively.)

The thickness of ice layer is

A. $1.1 m$

B. 0.4 m

C. 2.1 m

D. 3.6 m

Answer: A



Watch Video Solution

6. Two identical conducting rods are first connected independently to two vessels, one containing water at $100^{\circ}C$ and the other containing ice at $0^{\circ}C$. In the second case, the

rods are joined end to end and connected to the same vessels. Let q_1 and q_2 gram per second be the rate of melting of ice in the two cases respectively. The ratio $\frac{q_1}{q_2}$ is

A. $\frac{1}{2}$

B. $\frac{2}{1}$

C. $\frac{4}{1}$

D. $\frac{1}{4}$

Answer: C



Watch Video Solution

7. A solid cube and a solid sphere of the same material have equal surface area. Both are at the same temperature $120^{\circ} C$, then

A. Both the cube and the sphere cool down at the same rate

B. The cube cools down faster than sphere

C. The sphere cools down faster than the cube

D. Whichever is having more mass will cool
down faster

Answer: B



Watch Video Solution

8. The two opposite faces of a cubical piece of iron (thermal conductivity = 0.2 CGS units) are at $100^{\circ}C$ and $0^{\circ}C$ in ice. If the area of a surface is $4cm^2$, then the mass of ice melted in 10 minutes will b

A. 30 gm

B. 300 gm

C. 5 gm

D. 50 gm





Answer: B



Watch Video Solution

9. A conducting cylindrical rod of uniform cross-sectional area is kept between two large chambers which are at temperatures 100°C

and 0°C , respectively. The conductivity of the rod increases with x , where x is distance from 100°C end. The temperature profile of the rod in steady-state will be as



Answer: B




View Text Solution

Multiple Correct Answer Type

1. Three different arrangements of materials 1 and 2, 3 to form a wall. Thermal conductivities are $k_1 > k_2 > k_3$. The left side of the wall is $20^\circ C$ higher than the right side. Temperature difference ΔT across the material 1 has

following relation in three cases



A. In steady state, the rate of energy conduction through the wall III is greatest

B. In steady state, the rate of energy conduction through all the walls (I), (II) and (III) are same

C. In steady state, temperature difference across material I is greatest in wall *II*.

D. In steady state, temperature difference across material I is same in all the walls

Answer: B::D



Watch Video Solution

2. The ends of a rod of uniform thermal conductivity are maintained at different

(constant) temperatures. After the steady state is achieved:

A. heat flows in the rod from high temperature to low temperature even if the rod has the non-uniform cross sectional area

B. temperature gradient along length is same even if the rod has the non-uniform cross sectional area

C. heat current is same even if the rod has non-uniform cross sectional area

D. of the rod has uniform cross-sectional area the temperature is same all the point at rod.

Answer: A::C




Watch Video Solution

Comprehension Type

1. Liquid water coats an active (growing) icicle and extends up a short, narrow tube along the central axis. Because the water-ice interface must have a temperature of 0°C , the water in the tube cannot lose energy through the sides of the icicle or down through the tip because there is no temperature change in those directions. It can lose energy and freeze only by sending energy up (through distance L) to the top of the icicle, where the temperature T_r can be below 0°C . Take $L=0.10\text{m}$ and $T_r=-5^{\circ}\text{C}$. Assume that the central tube and the

upward conduction path both have cross-sectional area $A=0.5\text{m}^2$. The thermal conductivity of ice is $0.40\text{ W}/\text{m}\cdot\text{K}$, latent heat of fusion is $L_F = 4.0 \times 10^5\text{ J}/\text{K}$ and the density of liquid water is $1000\text{ kg}/\text{m}^3$.



The rate at which the energy conducted upward is

A. 20 W

B. 15 W

C. 5 W

D. 10 W


Answer: D



View Text Solution

2. Liquid water coats an active (growing) icicle and extends up a short, narrow tube along the central axis. Because the water-ice interface must have a temperature of 0°C , the water in the tube cannot lose energy through the sides of the icicle or down through the tip because there is no temperature change in those directions. It can lose energy and freeze only by sending energy up (through distance L) to the top of the icicle, where the temperature T_r can be below 0°C . Take $L=0.10\text{m}$ and $T_r=-5^\circ\text{C}$. Assume that the central tube and the

upward conduction path both have cross-sectional area $A=0.5\text{m}^2$. The thermal conductivity of ice is $0.40\text{ W}/\text{m}\cdot\text{K}$, latent heat of fusion is $L_F = 4.0 \times 10^5\text{ J}/\text{K}$ and the density of liquid water is $1000\text{ kg}/\text{m}^3$.



The rate at which mass converted from liquid to ice at the top of the central tube is

A. $7.5 \times 10^{-3} \text{ kg/s}$

B. $2.5 \times 10^{-5} \text{ kg/s}$

C. $5 \times 10^{-3} \text{ kg/s}$

D. $5 \times 10^{-5} \text{ kg/s}$


Answer: B



View Text Solution

3. Liquid water coats an active (growing) icicle and extends up a short, narrow tube along the central axis. Because the water-ice interface must have a temperature of 0°C , the water in the tube cannot lose energy through the sides of the icicle or down through the tip because there is no temperature change in those directions. It can lose energy and freeze only by sending energy up (through distance L) to the top of the icicle, where the temperature T_r can be below 0°C . Take $L=0.10\text{m}$ and $T_r=-5^\circ\text{C}$. Assume that the central tube and the

upward conduction path both have cross-sectional area $A=0.5\text{m}^2$. The thermal conductivity of ice is $0.40\text{ W/m}\cdot\text{K}$, latent heat of fusion is $L_F = 4.0 \times 10^5\text{ J/K}$ and the density of liquid water is 1000 kg/m^3 .



At what rate does the top of the tube move downward because of water freezing there?

A. $5 \times 10^{-8} m / s$

B. $2.5 \times 10^{-3} m / s$

C. $2.5 \times 10^{-8} m / s$

D. $4.5 \times 10^{-2} m / s$

Answer: C



View Text Solution

1. The thermal conductivity of a material in CGS system is 0.4. In steady state, the rate of flow of heat 10 cal/sec-cm, then the thermal gradient will be



[Watch Video Solution](#)

2. Two vessels of different materials are similar in size in every respect. The same quantity of ice filled in them gets melted in 20 minutes

and 30 minutes. The ratio of their thermal conductivities will be



[Watch Video Solution](#)

3. Two walls of thickness d and d and thermal conductivities k and k are in contact. In the steady state, if the temperature at the outer faces are T_1 and T_2 , the temperature at the common wall is



[Watch Video Solution](#)

4. Two identical plates of different metals are joined to form a single plate whose thickness is double the thickness of each plate. If the coefficients of conductivity of each plate are 2 and 3 respectively, Then the conductivity of composite plate will be



[Watch Video Solution](#)

5. The ratio of the diameters of two metallic rods of the same material is 2:1 and their lengths are in the ratio 1:4. If the

temperature difference between their ends are equal, the rate of flow of heat in them will be in the ratio



[Watch Video Solution](#)

6. Two cylinders P and Q have the same length and diameter and are made of different materials having thermal conductivities in the ratio $2 : 3$. These two cylinders are combined to make a cylinder. One end of P is kept at

$100^{\circ}C$ and another end of Q at $0^{\circ}C$. The temperature at the interface of P and Q is



[Watch Video Solution](#)

Single Corrcr Answer

1. Half part of ice block is covered with black cloth and rest half is covered with white cloth and then it is kept in sunlight. After some time clothes are removed to see the melted ice.

Which of the following statements is correct

A. Ice covered with white cloth will melt more.

B. Ice covered with black cloth will melt more.

C. Equal ice will melt under both clothes.

D. It will depend on the temperature of surroundings of ice.

Answer: (b)



Watch Video Solution

2. Two thermometers A and B are exposed in sunlight. The bulb of A is painted black, But that of B is not painted. The correct statement regarding this case is

A. Temperature of A will rise faster than b
but the final temperature will be the
same in both

B. Both A and B show equal rise in
beginning

C. Temperature of A will remain more than

B

D. Temperature of b will rise faster

Answer: (a)



Watch Video Solution

3. If 'p' calorie of heat energy is incident on a body and absorbs 'q' calories of heat absorbed then its coefficient of absorption is .

A. p/q

B. q/p

C. p^2/q^2

D. q^2/p^2

Answer: (b)



Watch Video Solution

4. The total energy radiated from a black body source at constant temperature is collected for one minute and is used to heat a quantity

of water. The temperature of water is found to increase from $20^{\circ} C$ to $20.5^{\circ} C$. If the absolute temperature of the blackbody is doubled and the experiment is repeated with the same quantity of water of $20^{\circ} C$, the temperature of water will be:

A. $21^{\circ} C$

B. $22^{\circ} C$

C. $24^{\circ} C$

D. $28^{\circ} C$

Answer: (d)



Watch Video Solution

5. The surface temperature of the sun is T_0 and it is at average distance d from a planet. The radius of the sun is R . The temperature at which planet radiates the energy is

A. $T_0 \sqrt{\frac{R}{2d}}$

B. $T_0 \sqrt{\frac{2R}{d}}$

C. $T_0 \sqrt{\frac{R}{d}}$

D. $T_0 \left(\frac{R}{d}\right)^{1/4}$

Answer: (a)



Watch Video Solution

6. In the previous problem, if θ is the angle subtended by the Sun at the planet, then temperature at which planet radiates energy is proportional to

A. θ

B. θ^2

C. $\theta^{1/2}$

D. $\theta^{-1/2}$

Answer: C



View Text Solution

7. The power radiated by a black body is P and it radiates maximum energy around the wavelength λ_0 . If the temperature of the black body is now changed so that it radiates maximum energy around a wavelength $3\lambda_0/4$

the power radiated by it will increase by a factor of .

A. $\frac{4}{3}$

B. $\frac{16}{9}$

C. $\frac{64}{27}$

D. $\frac{256}{81}$

Answer: (d)



Watch Video Solution

Multiple Correct Answer

1. A heated body emits radiation which has maximum intensity at frequency ν_m . If the temperature of the body is doubled:

A. the maximum intensity radiation will be

at frequency $2\nu_m$

B. the maximum intensity radiation will be

at frequency $(1/2)\nu_m$

C. the total emitted energy will increase by

a factor 16

D. the total emitted energy will increase by
a factor 2

Answer: (A, C)



Watch Video Solution

2. The solar constant for a planet is S . The surface temperature of the sun is T_K . The sun subtends an angle θ at the planet:

A. $S \propto T^4$

B. $S \propto T^\theta$

C. $S \propto \theta^2$

D. $S \propto \theta$

Answer: (A, C)



Watch Video Solution

Comprehension

1. A body is kept inside a container the temperature of the body is T_1 and the

temperature of the container is T_2 the rate at which body absorbs the energy is α . The emissivity of the body is e . The radiation striking the body is either absorbed or reflected.. After a long time the temperature of the body will be

A. T_1

B. T_2

C. $T_1 + \frac{(T_2 - T_1)}{2}$

D. None the these

Answer: (b)



Watch Video Solution

2. A body is kept inside a container the temperature of the body is T_1 and the temperature of the container is T_2 the rate at which body absorbs the energy is α The emissivity of the body is The radiation striking the body is either absorbed or reflected

At what rate of body will absorb the radiant energy .

A. if t is the time , rate is $(T_1 - T_2) / t$

B. e

C. Both of the above

D. None the above

Answer: (b)



Watch Video Solution

3. A body is kept inside a container the temperature of the body is T_1 and the temperature of the container is T_2 the rate at which body absorbs the energy is α The

emissivity of the body is The radiation striking the body is either absorbed or reflected

At what rate of body will absorb the radiant energy .

A. α , but $\alpha \neq e$

B. $(T_1 - T_2) / t$, where t is the time

C. e, but $e = \alpha$

D. None of above

Answer: C



Watch Video Solution

4. A body is kept inside a container the temperature of the body is T_1 and the temperature of the container is T_2 the rate at which body absorbs the energy is α The emissivity of the body is The radiation striking the body is either absorbed or reflected

A good absorber is .

- A. good reflector
- B. poor reflector
- C. average reflector

D. Assessment not possible

Answer: (b)



Watch Video Solution

Fill In The Blacks Type

1. If wavelengths of maximum intensity of radiations emitted by the sun and the moon are $0.5 \times 10^{-6}m$ and $10^{-4}m$ respectively, the ratio of their temperature is



[Watch Video Solution](#)

2. A black body emits radiations of maximum intensity at a wavelength of $\text{\AA} 5000$, when the temperature of the body is $1227^\circ C$. If the temperature of the body is increased by $1000^\circ C$, the maximum intensity of emitted radiation would be observed at



[Watch Video Solution](#)

3. The intensity of radiation emitted by the sun has its maximum value at a wavelength of 510 nm and that emitted by the North star has the maximum value at 350 nm. If these stars behave like black bodies, then the ratio of the surface temperatures of the sun and the north star is



Watch Video Solution

4. A black body radiates energy at the rate of E W/m at a high temperature T_K . When the temperature is reduced to $\frac{T}{2}K$, the radiant energy will be



[Watch Video Solution](#)

5. Energy is being emitted from the surface of a black body at $127^\circ C$ temperature at the rate of $1.0 \times 10^6 J/sec - m^2$. Temperature of

the black at which the rate of energy emission is $16.0 \times 10^6 \text{ J/sec} - \text{m}^2$ will be



Watch Video Solution

6. The rectangular surface of area $8\text{cm} \times 4\text{cm}$ of a black body at temperature 127°C emits energy E per second. If length and breadth are reduced to half of the initial value and the temperature is raised to 327°C , the emission energy becomes

A. $\frac{3}{8}E$

B. $\frac{81}{16}E$

C. $\frac{9}{16}E$

D. $\frac{81}{64}E$

Answer: D



Watch Video Solution

Single Correct

1. Consider two hot bodies B_1 and B_2 which have temperature 100°C and 80°C

respectively at $t = 0$. The temperature of surroundings is 40°C . The ratio of the respective rates of cooling R_1 and R_2 of these two bodies at $t = 0$ will be

A. $R_1 : R_2 = 3 : 2$

B. $R_1 : R_2 = 5 : 4$

C. $R_1 : R_2 = 2 : 3$

D. $R_1 : R_2 = 4 : 5$

Answer:



Watch Video Solution

2. Equal masses of two liquids are filled in two similar calorimeters. The rate of cooling will

- A. Depend on the nature of the liquids
- B. Depend on the specific heats of liquids
- C. Be same for both the liquids
- D. Depend on the mass of the liquids

Answer:



Watch Video Solution

3. A body takes 4 minutes to cool from $100^{\circ}C$ to $70^{\circ}C$. To cool from $70^{\circ}C$ to $40^{\circ}C$ it will take (room temperature is $15^{\circ}C$)

A. 7 min

B. 6 min

C. 5 min

D. 4 min

Answer:



Watch Video Solution

4. A body cools from $60^{\circ}C$ to $50^{\circ}C$ in 10 minutes . If the room temperature is $25^{\circ}C$ and assuming Newton's law of cooling to hold good, the temperature of the body at the end of the next 10 minutes will be

A. $38.5^{\circ}C$

B. $40^{\circ}C$

C. $42.85^{\circ}C$

D. $45^{\circ}C$

Answer:



Watch Video Solution

5. The temperature of a liquid drops from 365 K to 361 K in 2 minutes . Find the time during which temperature of the liquid drops from 344 K to 342 K. Temperature of room is 293 K

A. 84 sec

B. 72 sec

C. 66 sec

D. 60 sec

Answer:



Watch Video Solution

6. Hot water kept in a beaker placed in a room cools from $70^{\circ}C$ to $60^{\circ}C$ in 4 minutes . The time taken by it to cool from $69^{\circ}C$ to $59^{\circ}C$ will be

- A. The same 4 minutes
- B. More than 4 minutes
- C. Less than 4 minutes

D. We cannot say definitely

Answer:



Watch Video Solution

7. In a room where the temperature is $30^{\circ}C$, a body cools from $61^{\circ}C$ to $59^{\circ}C$ in 4 minutes.

The time (in min.) taken by the body to cool from $51^{\circ}C$ to $49^{\circ}C$ will be

A. 4 min

B. 6 min

C. 5 min

D. 8 min

Answer:



Watch Video Solution

8. A calorimeter of mass 0.2kg and specific heat $900\text{J}/\text{kg} - \text{K}$. Containing 0.5kg of a liquid of specific heat $2400\text{J}/\text{kg} - \text{K}$. Its

temperature falls from $60^{\circ}C$ to $55^{\circ}C$ in one minute. Find the rate of cooling.

A. $5J/s$

B. $15J/s$

C. $100J/s$

D. $115J/s$

Answer:



Watch Video Solution

9. A cane is taken out from a refrigerator at 0°C . The atmospheric temperature is 25°C . If t_1 is the time taken to heat from 0°C to 5°C and t_2 is the time taken from 10°C to 15°C , then the wrong statements are

(1) $t_1 > t_2$

(2) $t_1 = t_2$

(3) There is no relation

(4) $t_1 < t_2$

A. $t_1 > t_2$

B. $t_1 < t_2$

C. $t_1 = t_2$

D. There is no relation

Answer:



Watch Video Solution