doubtnut
India's Number 1 Education App

# BIOLOGY

## BOOKS - PRADEEP BIOLOGY (HINGLISH)

## MORPHOLOGY OF FLOWERING PLANTS

### Curiosity Questions

**1.** Why do plants belonging to pea family bear nodulated roots ? What is inside the nodules ?

Watch Video Solution

**2.** Why the plants growing in saline swamps near the sea shores develop pneumatophores ?

Watch Video Solution

**3.** Why are canopy of Banyan tree extend over a large area ?

Watch Video Solution

**4.** Why the stems of flowering plants attain diverse forms ?

Watch Video Solution

**5.** Why do some modified stems store reserve food materials ?

Watch Video Solution

**6.** What is maize grain usually called as a fruit and not a seed ?

Watch Video Solution

## Notable Questions

**1.** Underground storage organs (such as potato tubers, roots of sweet potato, beet, carrot, etc.) generally store

carbohydrates in the form of starch grains. How do they store starch when they do not get light for photosynthesis ?

Watch Video Solution

**2.** Tubers of Sweet potato (Ipomoea batata) take part in vegetative propagation and grow to form new plants. How can we ascertain that they are not modified stems.

Watch Video Solution

**3.** What are angiosperms ?

Watch Video Solution

**4.** What is Floral Formula ?

Watch Video Solution

**5.** What is a floral diagram ? What special features does a floral diagram inform ?

Watch Video Solution

## Ncert Exercises With Answers

**1.** What is meant by modification of root? What type of modification of root is found in the

(a) Banyan tree

(b) Turnip

(c) Mangrove trees

Watch Video Solution

**2.** Justify the following statements on the basis of external features

(i) Underground parts of a plant are not always roots

(ii) Flower is a modified shoot

Watch Video Solution

**3.** How is pinnately compound leaf different from palmately compound leaf?

Watch Video Solution

**4.** Explain with suitable examples the different types of phyllotaxy?

Watch Video Solution

**5.** Define the following terms:

(a) Aestivation

(b) Placentation

(c) Actinomorphic

(d) Zygomorphic

(e) Superior ovary

(f) Perigynous flower

(g) Epipetalous Stamen

Watch Video Solution

**6.** Differentiate between

(a) Racemose and cymose inflorescence

(b) Fibrous roots and adventitious roots

(c) Apocarpous and syncarpous ovary

Watch Video Solution

**7.** Draw the labelled diagram of the following : (i) gram seed (ii) V.S. of maize seed.

Watch Video Solution

**8.** Describe the modification of stem with suitable examples

Watch Video Solution

**9.** Take one flower each of families Fabaceae and Solanaceae and write its semitechnical description. Also draw their floral diagrams after studying them.

Watch Video Solution

**10.** Describe the various types of placentations found in flowering plants.

Watch Video Solution

**11.** What is a flower ? Describe the parts of a typical angiosperm flower.

Watch Video Solution

**12.** How do the various leaf modifications help plants?

Watch Video Solution

**13.** Define the term inflorescence. Explain the basis for the different types of inflorescence in flowering plants.

Watch Video Solution

**14.** Write the floral formula of an actinomorphic bisexual, hypogynous flower with five united sepals, five free petals. Five free stamens and two united carpals with superior ovary and axile placentation.

Watch Video Solution

**15.** Describe the arrangement of floral members in relation to their insertion on thalamus?

Watch Video Solution

## Additional Questions Very Short Answer Questions

**1.** What are angiospermic plants ?

Watch Video Solution

**2.** What are Pneumatophores ?

Watch Video Solution

**3.** Name the roots that arise from any other plant part except the radicle.

Watch Video Solution

**4.** Name the plant with lightest seeds.

Watch Video Solution

**5.** Give examples of angiosperms in which stems are not developed.

Watch Video Solution

**6.** Give examples of angiosperms in which roots are not developed.

Watch Video Solution

**7.** Give examples of angiosperms in which plant is devoid of green leaves.

Watch Video Solution

**8.** Name the different kinds of compound leaves.

Watch Video Solution

**9.** What is the function of pulvinus? Give one example of the plant having it.

Watch Video Solution

**10.** What term is given to the arrangement of leaves on the stem ?

Watch Video Solution

**11.** What are adventitious roots ? Give one example.

Watch Video Solution

**12.** Name any one plant in which adventitious roots develop chlorophyll and photosynthesise.

Watch Video Solution

**13.** What takes over the function of photosynthesis in Opuntia ?

Watch Video Solution

**14.** What is a bulbil ?

Watch Video Solution

**15.** Why are potato and sweet potato called tubers ?

Watch Video Solution

**16.** Which same plant part has transformed Into the following different modlftcatlons (i) tendril of pumpkin (ii) thorn of Citrus.

Watch Video Solution

**17.** Name an insectivorous angiosperm without roots,

Watch Video Solution

**18.** How annuals differ from biennial?

Watch Video Solution

**19.** Name a free floating aquatic plant with photosynthetic roots.

Watch Video Solution

**20.** Name two plants that produce rhizomes.

Watch Video Solution

**21.** Cuscuta develops roots to penetrate the host tissue and obtain nutrition. What are these roots called?

Watch Video Solution

**22.** What is heterophylly ?

Watch Video Solution

**23.** Define venation.

Watch Video Solution

**24.** Define phyllotaxy.

Watch Video Solution

**25.** Name a plant in which leaf is bipinnately compound.

Watch Video Solution

**26.** What is phyllode ? Give one example of it.

Watch Video Solution

**27.** In which type of leaf the stomata remain distributed only on lower surface ?

Watch Video Solution

**28.** What type of fruit tomato is ?

Watch Video Solution

**29.** Give the names of the plants in which following types of inflorescences are found : Catkin, Capitulum and Spikelet.

Watch Video Solution

**30.** Name as cultivated plant in which neither fruits nor seeds are formed.

Watch Video Solution

**31.** Give three examples of false fruit.

Watch Video Solution

**32.** In swampy areas like the Sundarbans in West Bengal, plants bear special kind of roots called .....

Watch Video Solution

**33.** Reticulate and parallel venation are the characteristic of ..... and ..... Respectively.

Watch Video Solution

**34.** Which parts of ginger and onion are edible ?

Watch Video Solution

**35.** In epigynous flower , ovary is situated below the ..... .

Watch Video Solution

**36.** Give two examples of roots that develop from different parts of the angiospermic plant other than the radicle.

Watch Video Solution

## Additional Questions Short Answer Questions

**1.** Distinguish between alternate and whorled phyllotaxy.

Watch Video Solution

**2.** Pick out the plant with whorled arrangement of leaves from the list given below anti say why It Is categorised so? Neem, Nerlum , Nepenthes.

Watch Video Solution

**3.** Pick out the plant with unipinnate type of compound leaves from the list given below and say why It Is

categorised so ? Trifolium, Neem, Moringa.

Watch Video Solution

**4.** Differentiate between root climbers and stem climbers.

Watch Video Solution

**5.** What are false fruits ? Give two examples .

Watch Video Solution

**6.** Mention two types of venation with examples.

Watch Video Solution

**7.** Mention the function of (i) Underground stem of ginger (ii) Axillary bud of citrus.

Watch Video Solution

**8.** Mention the function performed by

(i) Scale leaf of onion (ii) axillary bud of cucumber (iii) Tap root of beet.

Watch Video Solution

**9.** Write the floral formula of family-Solanaceae.

Watch Video Solution

**10.** Write short notes on tetradynamous condition of stamens.

Watch Video Solution

**11.** What is imbricate aestivation ?

Watch Video Solution

**12.** What type of ovary is found in family-Solanaceae ?

Watch Video Solution

**13.** Write the botanical names of (i) Tomato, (ii) Chillies and (iii) Tobacco.

Watch Video Solution

**14.** A typical angiosperm flower consists of four floral parts. Give the names of the floral parts and their arrangements sequentially.

Watch Video Solution

**15.** You have heard about several insectivorous plants that feed on insects. Nepenthes or the pitcher plant is one such example, which usually grows in shallow water or in

marsh lands. What part of the plant is modified into a pitcher ? How does this modification help the plant for food even through it can photosynthesize like any other green plant ?

Watch Video Solution

**16.** How can you differentiated between free central and axile placentation ?

Watch Video Solution

**17.** Tendrils are found in the following plants . Identify whether they are stem tendrils of leaf tendrils.

(a) Cucumber (b) Peas

(c) Pumpkins (d) Grapevine

(e) Watermelon

Watch Video Solution

**18.** What do you understand by special types of inflorescence ? Describe them with diagrams and examples.

Watch Video Solution

**19.** Give any four differences between rhizome and bulb.

Watch Video Solution

**20.** Differentiate between stolon and sucker.

Watch Video Solution

**21.** Differentiate between runner and stolon.

Watch Video Solution

**22.** What is aestivation? Describe its various types found in petals.

Watch Video Solution

**23.** Describe the various types of placentations found in flowering plants.

Watch Video Solution

**24.** How would you differentiate leaflets of a compound leaf from simple leaves of a branch ? Give an example of a plant with bipinnate leaves.

Watch Video Solution

**25.** Why is the leaf of Bombax categorised as palmately compound multifoliate leaf?

Watch Video Solution

**26.** Describe the corolla of family-Fabaceae.

Watch Video Solution

**27.** (A) The famous Bodhi tree (*Ficus religiosa*) at Gaya is about ..... years old. (B) Flower is a modified ..... and produces fruits and seeds. (C) In bamboos, the stem is jointed with solid ..... and internodes and is known as ..... (D) The floral envelop including both calyx and corolla is called ..... . (E) A seed is a ripened ..... which contains the ..... or the miniature of plant body.

Watch Video Solution

**28.** Give one example each of the following :

(A) A plant having hypanthodium inflorescence

(B) A palmately compound leaf having 5 or more terminal leaflets

(C) A plant with decompound leaves

(D) A cladode

(E) A plant which reproduces vegetatively by root tubers

(F) A plant in which the floral buds of inflorescence transform into bulbils

(G) A plant in which the free ends of adnate stipules are modified into tendrils

(H) An endospermic dicotyledonous seed

(I) A plant which completes its life-cycle in two years

(J) A plant in which the main plant body is a thalloid root which carries out the process of photosynthesis.

Watch Video Solution

**29.** Make corrections wherever you find mistake in spellings/words in the following paragraph/sentences.

(A) The endospermic seeds of castor and custard apple contain an outer hard shell called tegmen. There is an outgrowth at the micropyle called aril, which is spongy and secretes moisture during seed germination.

B) In dicotyledonous seeds, the plumule is covered by a sheath called coleorhiza and radicle is protected by coleoptile.

(C) The fruits developed from the simple ovary are called simple type. If there is cluster of simple fruits which is borne on the same thalamus and developed from

bicarpellary, syncarpous ovary, they are aggregate. When the fruit develops from inflorescence, it is composite.

Watch Video Solution

**30.** Differentiate between

(a) Bract and Bracteole

(b) Pulvinus and petiole

(c) Pedicel and peduncle

(d) Spike and spadix

(e) Stamen and staminode

(f) Pollen and pollinium

Watch Video Solution

**31.** Given below are a few formulae of some well known plants. Draw Doral diagrams from these formulae.

(i) $\oplus \male\!\!\female K_{(5)}, C_{(5)}, A_{(5)}, G_{(2)}$ (ii) $\Phi \male\!\!\female K_{(5)}C_{1+2+(2)} A_{(9)+1} G_1$ (iii) $\oplus \male\!\!\female K_5C_5A_{5+5}G_{(5)}$

View Text Solution

## Additional Questions Long Answer Questions

**1.** Describe the various roles of roots and stem of an angiospermic plant.

Watch Video Solution

**2.** What are underground stems ? Describe various forms of underground stem. Draw diagram and give one example of each form.

Watch Video Solution

**3.** What are edible fruits ? Describe chief edible fruits you have studied giving suitable examples.

Watch Video Solution

**4.** What are fleshy fruits ? Describe the various types of fleshy fruits with suitable examples.

Watch Video Solution

**5.** Describe with the help of diagram the structure of Maize grain. How does it differ from Bean seed ?

Watch Video Solution

**6.** Give a scientific description of family Liliaceae. Mention the economic importance.

Watch Video Solution

**7.** Give an account of the distinguishing features of family Solanaceae. Give botanical names of four plants of economic value belonging to this family.

Watch Video Solution

**8.** Write an explanatory note on defense mechanisms in plants.

Watch Video Solution

**9.** Define a flower. Name the various parts of a typical flower and describe only non-essential parts.

Watch Video Solution

**10.** Match the items in column (A) with column (B). Each point in column (A) has minimum one match in coiumn (B)

and maximum three matches

| Column (A) | Column (B) |
| --- | --- |
| 1. Perianth | (*a*) Pistil |
| 2. Cladode | (*b*) *Ruscus* |
| 3. *Ficus bengalensis* | (*c*) Tepal |
| 4. Gynoecium | (*d*) Prop roots |
| | (*e*) Carpels |
| | (*f*) Syconus |
| | (*g*) *Asparagus* |
| | (*h*) Hypanthodium |

Watch Video Solution

**11.** Longitudinal section of Maize grain is shown in the diagram. Label the parts 1 to 9, and also write the function

of parts three and four.




View Text Solution

**12.** (a) Write the type of fruit against the following and furnish the other information asked :

Various types of fruits : Drupe, Achene, Capsella, Hesperidium, Cremocarp, Cypsella, Schizocarp

(i) Cococut - Edible part

(ii) Orange - Type of placentation

(iii) Coriander - Type of inflorescence

(b) Write the type of inflorescence against the foil owing and furnish the other information asked :

Various types of inflorescence : Racemose, Umbel, Cyathium, Capitulum, Cymose

(i) Euphorbia - Type of fruit (ii) Marigold - Type of fruit

Watch Video Solution

**13.** The arrangement of ovules within the ovary is known as placentation. What does the term placenta refer to ?

Name and draw various types of placentations in the flower as seen in T.S. or V.S.

Watch Video Solution

**14.** Describe various stem modifications associated with food storage climbing and protection.

Watch Video Solution

**15.** Name the body part modified for food storage in the following

$(a)$ Carrot.....
$(b)$ Colocasia.....
$(c)$ Sweet potato.....
$(d)$ Asparagus.....
$(e)$ Radish..... .
$(f)$ Potato.....
$(g)$ Dahlia.....
$(h)$ Tumeric.....
$(g)$ Gladiolus.....
$(j)$ Ginger.....
$(k)$ Portulaca.....

Watch Video Solution

## Analytical Questions With Answers

**1.** What are phylloclades? What are their functions ?

Watch Video Solution

**2.** What is phyllotaxy ? Give its types with one example.

Watch Video Solution

**3.** How would you differentiate between aggregate fruits and composite fruits ? Give one example of each.

Watch Video Solution

**4.** Whether mango is a true fruit or false fruit? Justify. Give its structure. Which branch of horticulture deals with the study and cultivation of fruits ?

Watch Video Solution

**5.** Give any two reasons to justify that Onion bulb is a modified stem.

Watch Video Solution

**6.** Mango and coconut are'drupe' type of fruits. In mango fleshy mesocarp is edible. What is the edible part of coconut? What does milk of tender coconut represent?

Watch Video Solution

**7.** How does the niustard inflorescence differ from the banana inflorescence in arrangement '? Give the technical term for each.

Watch Video Solution

**8.** Why are flowers of Cucumber referred to as epigynous ?

Watch Video Solution

**9.** How do you distinquish between hypogeal germination and epigeal germination ? What is the role of cotyledon (s) and the endosperm in the germination of seeds ?

Watch Video Solution

**10.** In the given examples, which body part is modified for food storage :

(a) Potato (b) Radish (c) Portulaca (d) Dahlia

Watch Video Solution

**11.** hree students described the floral characters of a plant belonging to family Liliaceae. One of them described correctly and the others two were wrong. Who was correct and which characters were described wrong by the other two ?

Student A : Tetramerous, actinomorphic, polyphyllus, unilocular ovary, axile placentation.

Student B : Trimerous, actinomorphic, polyandrous, superior ovary, axile placentation

Student C : Trimerous, zygomorphic, gamophyllus, inferior ovary, marginal placentation.

Watch Video Solution

12.

Study the above figure and answer the following questions :

(a) Write the names of A, B, C and D (b) What is the type of fruit ?

(c) Write the type of ovary from which the fruit is formed.

(d) Which part is edible?

View Text Solution

13.

Observe the above figure and answer the following questions -

(a) Which one of the above represents the characteristic type of corolla of family Fabaceae ? Name the type of

corolla.

(b) Which of the above figures shows tubular type of corolla.

View Text Solution

**14.** (a) Name the layer which separates endosperm from embryo in a monocotyledonous seed.

(b) Where lies the cotyledon of a maize grain (monocotyledonous seed) ? What is it called ? Give its functions.

Watch Video Solution

**15.** Select a plant which grows in swampy areas, giving reasons.

Plant A : Develops haustoria for absorption of food.

Plant B : Develops prop roots for support.

Plant C : Develops pneumatophores to get oxygen for respiration.

Watch Video Solution

**16.** Write the symbols for the following floral characters -

(a) Calyx

(b) Actinomorphic flower

(c) Bracteate

( d) Inferior ovary

( e) bisexual flower

Watch Video Solution

**17.** Identify the family which shows the following diagnostic features. Name two plants of the family which are used as source of food.

**Features** : Flowers pentamerous, Gynoecium - bicarpellary, syncarpous, ovary placed obliquely, placentation axile, placenta swollen.

Watch Video Solution

**18.** Describe the Rower on the basis of the following floral formula

View Text Solution

**19.** Give the technical terms for the following:

(a) A sterile stamen (b) Stamens are united in one bunch

(c) Stamens are attached to the petals

Watch Video Solution

**20.** Complete the following table by givlng the correct options :

| Family | Inflorescence | Flower | Stamens | Gynoecium |
|---|---|---|---|---|
| Fabaceae | — | — | 10 | — |
| Solanaceae | Solitary, axillary or cymose | — | — | — |
| Liliaceae | — | Actinomorphic | — | Tricarpellary |

Watch Video Solution

## Practice Questions Multiple Choice Questions

**1.** Edible part of mango is

A. Receptacle

B. Epicarp

C. Mesocarp

D. Endocarp

**Answer: C**

Watch Video Solution

**2.** Thorn is a modified branch because

A. it is hard, straight and pointed

B. it is a part of the plant

C. it arises in the axil of a leaf

D. it is a defensive organ

**Answer: C**

Watch Video Solution

**3.** Flower in which only one set of essential organ is present are said to be

A. Bisexual

B. Monoecious

C. Dioecious

D. Unisexual

**Answer: D**

Watch Video Solution

**4.** The example for trimerous, unisexual flower is

A. Pea

B. Tamarind

C. Hibiscus

D. Cocos nucifera

**Answer: D**

Watch Video Solution

**5.** The condition where filaments and anthers are fused throughout the entire length is

A. Synandrous

B. Gynandrous

C. Protandrous

D. Syngenesis

**Answer: A**

Watch Video Solution

**6.** A flower which can be divided into equal vertical halves by more than one plane of division is

A. Actinomorphic

B. Zygomorphic

C. Heteromorphic

D. Cyclic

**Answer: A**

Watch Video Solution

**7.** Which of the following pairs is not correct ?

A. Corymb-Candituft

B. Capitulum-Sunflower

C. Catkin-Mulberry

D. Raceme-Wheat

**Answer: D**

Watch Video Solution

**8.** Which of the following represents the edible part of the fruit of litchi ?

A. Mesocarp

B. Endocarp

C. Pericarp

D. Juicy aril

**Answer: D**

Watch Video Solution

**9.** How many female flowers occur in a cyathium ?

A. One

B. Two

C. Three

D. Four

**Answer: A**

Watch Video Solution

**10.** The receptacle is flattened at the top and bears numerous sessile flowers in centripetal manner in

A. Cyathium

B. Catkin

C. Umbel

D. Capitulum

**Answer: D**

Watch Video Solution

**11.** Maize grain is

A. Seed

B. Embryo

C. Ovule

D. Fruit

**Answer: D**

Watch Video Solution

**12.** Rhizome which grows vertically upwards are

A. Corms

B. Stolon

C. Bulbils

D. Root stock

**Answer: D**

Watch Video Solution

**13.** Fibrous root system is mostly found in

A. Dicot plants

B. Monocot plants

C. Pteridophytes

D. Bryophytes

**Answer: B**

Watch Video Solution

**14.** The leaves of Utricularia plant are modified into

A. Hooks

B. Tendrils

C. Bladders

D. Pitchers

**Answer: C**

Watch Video Solution

**15.** If position of ovary is below sepals, petals, stamens, the flower is called

A. Epigynous

B. Perigynous

C. Mesogynous

D. Metagynous

**Answer: A**

Watch Video Solution

**16.** Trimerous flower, superior ovary and axile placentation is characteristic of

A. Cucurbitaceae

B. Solanaceae

C. Compositae

D. Liliaceae

**Answer: D**

Watch Video Solution

**17.** What type of placentation is seen in sweet pea

A. Marginal

B. Basal

C. Axile

D. Free central

**Answer: A**

Watch Video Solution

**18.** Long filamentous threads protruding at the end of a young cob of maize are-

A. Hair

B. Anthers

C. Styles

D. Ovaries

**Answer: C**

Watch Video Solution

**19.** In a cereal grain the single cotyledon of embryo is represented by

A. Coleoptile

B. Coleorhiza

C. Scutellum

D. Prophyll

**Answer: C**

Watch Video Solution

**20.** Pineapple (ananas) fruit develops from

A. A multilocular monocarpellary flower

B. A unilocular poly carpellary flower

C. A multipistillate syncarpous flower

D. A cluster of completely borne flowers on a common axis

**Answer: D**

Watch Video Solution

**21.** In the members of family malvaceae, anthers are described as

A. Diadelphous and dithecous

B. Diadelphous and monothecous

C. Monadelphous and dithecous

D. Monadelphous and monothecous

**Answer: D**

Watch Video Solution

**22.** Corm is a modification of

A. Root

B. Leaf

C. Stem

D. Bud

**Answer: C**

Watch Video Solution

**23.** The leaves are modified into tendrils, hooks, pitcher and bladder in the following plants respectively

A. Sweet pea, Cat's nail, Nepenthes, Utricularia

B. Sweet pea, Cat's nail, Utricularia, Nepenthes

C. Nepenthes, Cat's nail, sweet pea, Utricularia

D. Nepenthes, sweet pea, Cat's nail, Utricularia

**Answer: A**

Watch Video Solution

**24.** Which of the following is not a characteristic feature of Fabaceae

A. Descending imbricate aestivation, ten stamens diadelphous, ovary superior

B. Sepals five, gamosepalous, imbricate aestivation, placentation marginal

C. Monocarpellary, ovary superior, style long, slightly bent at the apex

D. Corolla five petals, polypetalous, anterior one large and outermost

**Answer: D**

Watch Video Solution

**25.** Pattern of arrangement of leaves on stem is known as

A. Phyllode

B. Pbyllotaxy

C. Phylloclade

D. Heterophylly

**Answer: B**

Watch Video Solution

**26.** A flower which can be divided in the halves in only one vertical plane is called

A. Actinomorphic

B. Heteromorphic

C. Isomerous

D. Zygomorphic

**Answer: D**

Watch Video Solution

**27.** An example of composite sorosis type of fruit is

A. Fig

B. Custard apple

C. Pineapple

D. Raspberry

**Answer: C**

Watch Video Solution

**28.** The plwnule is covered by protective sheath called:

A. Coleorhiza

B. Scutellum

C. Coleoptile

D. Tegmen

**Answer: C**

Watch Video Solution

**29.** Tricarpellary, syncarpous superior ovary is seen in :

A. Allium

B. Oenothera

C. Solanum

D. Dolichus

**Answer: A**

Watch Video Solution

**30.** Tendrils in Passiflora develop from

A. Apical roots

B. Apices of roots

C. Axillary buds

D. Terminal buds

**Answer: C**

Watch Video Solution

**31.** number of female flowers in a cyathium I

A. 1

B. 2

C. 3

D. Several

**Answer: A**

Watch Video Solution

**32.** Ginger is a stem which can be differentiated from root because it

A. It lacks chlorophyll

B. It stores food

C. It has nodes and internodes

D. It has xylem and vessels

**Answer: C**

Watch Video Solution

**33.** Gynoecium with fused carpels

A. Syncarpous

B. Apocarpous

C. Syngenecium

D. None of these

**Answer: A**

Watch Video Solution

**34.** Beauty of Bougainvillea flower is because of

A. Petals

B. Calyx

C. Bracts

D. Sepals

**Answer: A**

Watch Video Solution

**35.** An orthotropous ovule is one in which micropyle and chalaza are

A. At right angles to funicle

B. Parallel to the funicle

C. In straight line of funicle

D. Parallel along with ovule

**Answer: C**

Watch Video Solution

**36.** Match the following with correct combination

| | Column I | Column II |
|---|---|---|
| A | Cuscuta | 1 Saprophyte |
| B | Eichornia | 2Pneumatophore |
| C | Monotropa | 3 Insectivorous plant |
| D | Rhizophora | 4 Parasite |
| E | Utricularia | 5 Root pocket |

A. A-4, B-3, C-1, D-5, E-2

B. A-4, B-5, C-1, D-2, E-3

C. A-2, B-3, C-1, D-5, E-4

D. A-3, B-1, C-5, D-4, E-2

**Answer: B**

Watch Video Solution

**37.** The fruit is chambered, developed from inferior ovary and has seeds with succulent testa in

A. Guava

B. Cucumber

C. Pomegranate

D. Orange

**Answer: C**

Watch Video Solution

**38.** Dry indehiscent single-seeded fruit formed from biscarpellary syncarpous inferior ovary is

A. Berry

B. Cremocarp

C. Caryopsis

D. Cypsella

**Answer: D**

**39.** The fleshy receptacle of syconus of fig encloses a number of

A. Berries

B. Mericarps

C. Achenes

D. Samaras

**Answer: C**

Watch Video Solution

**40.** Replum is present in the ovary of flower of

A. Sunflower

B. Pea

C. Lemon

D. Mustard

**Answer: D**

Watch Video Solution

**41.** Zygomorphic condition can be represented as

A. $\oplus$

B. $\%$

C. P

D. G

**Answer: B**

Watch Video Solution

**42.** Aril is the edible part of

A. Apple

B. Litchi

C. Banana

D. Mango

**Answer: B**

Watch Video Solution

**43.** In which plant the fruit is a drupe, seed coat is thin, embryo in inconspicuous and endosperm is edible

A. Ground nut

B. Wheat

C. Apple

D. Coconut

**Answer: D**

Watch Video Solution

**44.** A fibrous root system is excellent for

A. Food storage

B. Nitrogen fixation

C. Absorbing water from deeper layers of soil

D. Providing good anchorage for the plant

**Answer: D**

Watch Video Solution

**45.** Cotyledons and testa respectively are edible parts in

A. Walnut and tamarind

B. French bean and coconut

C. Cashew nut and litchi

D. Ground nut and pomegranate

**Answer: D**

Watch Video Solution

**46.** An example of a seed with endosperm perisperm and caruncle is

Or

Which one of the following is an endosperm seed

Or

In which of the following plants, cotyledons form the first pair of leaves.

A. Coffee

B. Lily

C. Castor

D. Cotton

**Answer: C**

Watch Video Solution

Watch Video Solution

**47.** Rearrange the following zones as seen in the root in vertical section and choose the correct option.

A. Root hair zone

B. Zone of meristems

C. Root cap zone

D. Zone of maturation

E. Zone of elongation

A. C, B, E, A, D

B. A, B, C, D, E

C. D, E, A, C, B

D. E, D, C, B, A

**Answer: A**

Watch Video Solution

**48.** In an inflorescence where flowers are borne laterally in an acropetal succession,the position of the youngest floral bud shall be

A. Proximal

B. Distal

C. Intercalary

D. Any where

**Answer: B**

Watch Video Solution

Watch Video Solution

**49.** The mature seeds of plants such as gram and peas, possess no endosperm, because

A. These plants are not angiosperms

B. There is no double fertilization in them

C. Endosperm is not formed in them

D. Endosperm gets used up by the developing embryo during seed development

**Answer: D**

Watch Video Solution

**50.** Roots developed from parts of the plant other than radicle are called

A. Taproots

B. Fibrous roots

C. Adventitious roots

D. Nodular roots

**Answer: C**

Watch Video Solution

**51.** Venation is a term used to describe the pattern of arrangement of

A. Floral organs

B. Flower in inflorescence

C. Veins and veiitlets in a lamina

D. All of them

**Answer: C**

Watch Video Solution

**52.** Endosperm, a product of double fertilization in angiosperms is absent in the seeds of

A. Gram

B. Orchids

C. Maize

D. Castor

**Answer: B**

Watch Video Solution

**53.** Many pulses of daily use belong to one of the families below (tick the correct answer)

A. Solanaceae

B. Fabaceae

C. Liliaceae

D. Poaceae

**Answer: B**

Watch Video Solution

**54.** The placenta is attached to the developing seed near the

A. Testa

B. Hilum

C. Micropyle

D. Chalaza

**Answer: D**

Watch Video Solution

**55.** Which of the following plants is used to extract the blue dye ?

A. Trifolium

B. Indigofera

C. Lupin

D. Cassia

**Answer: B**

Watch Video Solution

**56.** Match the followings and choose correct option.

| Group A | Group B |
|---|---|
| $A$. Aleurone layer | $(ii)$Without fertilisation |
| $B$. Parthenocarpic fruit | $(ii)$Nutrition |
| $C$. Ovule | $(iii)$Double fertilisation |
| $D$. Endosperm | $(iv)$Seed |

A. A-i, B-ii, C-iii, D-iv

B. A-ii, B-i, C-iv, D-iii

C. A-iv, B-ii, C-i, D-iii

D. A-ii, B-iv, C-i, D-iii

**Answer: B**

Watch Video Solution

**57.** The scutellum observed in a grain of wheat or maize is comparable to which part of the seed in other monocotyledons

A. Plumule

B. Cotyledons

C. Endosperm

D. Aleurone layer

**Answer: B**

Watch Video Solution

**58.** Aestivation of petals in the flower of cotton is correctly shown in

A.

B.

C.

D.

**Answer: D**

Watch Video Solution

**59.** Find the correct match

A. mustard plant-leaves opposite

B. mustard plant-leaves alternate

C. guava plant-leaves alternate

D. guava plant-leaves whorled

**Answer: B**

Watch Video Solution

**60.** Which of the following is not correctly paired

A. Fabaceae-legume family

B. Solanaceae-Potato family

C. Liliaceae-Sunflower family

D. Brassicaceae-Mustard family

**Answer: C**

Watch Video Solution

**61.** Find the incorrect statement.

A. root hairs are univellular elongation

B. root hairs absord water and minerals

C. trichomes are unicellular elongations

D. trichomes are multicellular elongations

**Answer: C**

Watch Video Solution

**62.** Find the incorrect match.

A. tap root-correct

B. adventitious root-sweet potato

C. poop root-banyantree

D. stilt root-turnip

**Answer: D**

Watch Video Solution

**63.** Example of corm is

A. Ginger

B. Colocasia

C. Onion

D. Potato

**Answer: B**

Watch Video Solution

**64.** The 'Eyes' of the potato tuber are :-

A. root buds

B. flower buds

C. shoot buds

D. axillary buds

**Answer: D**

Watch Video Solution

**65.** Which one of the following statements is correct?

A. in tomato, fruit is a capsule

B. seeds of orchids have oil-rich endosperm

C. placentation in primose is basal

D. flower of tulip is a modified shoot

**Answer: D**

Watch Video Solution

**66.** The floral formula of solanaceae (Chilli) is

A.

B.

C.

D.

**Answer: B**

Watch Video Solution

**67.** The ovary is half inferior in flowers of

A. peach

B. cucumber

C. cotton

D. guava

**Answer: A**

Watch Video Solution

**68.** A drupe develops in

A. mango

B. wheat

C. pea

D. tomato

**Answer: A**

Watch Video Solution

**69.** Which one the following pairs is wrongly matched while the remaining three and correct

A. Penicillium - conidia

B. Water hyacinth - runner

C. Bryophyllum - leaf buds

D. Agave - bulbils

**Answer: B**

Watch Video Solution

**70.** Which one of the following diagrams represents the placentation in Dianthus

A.

B.

C.

D.

**Answer: B**

Watch Video Solution

**71.** Whorled, simple leaves with reticulate venation are present in

A. Calotropis

B. neem

C. China rose

D. Alstonia

**Answer: D**

Watch Video Solution

**72.** Flowers are Zygomorphic in

A. mustard

B. gulmohar

C. tomato

D. datura

**Answer: B**

Watch Video Solution

**73.** The mesocarp in coconut is

A. fleshy

B. stony

C. fibrous

D. milky

**Answer: C**

Watch Video Solution

**74.** Placentation in tomato and lemon is

A. Parietal

B. Free central

C. Marginal

D. Axile

**Answer: D**

Watch Video Solution

**75.** Cymose inflorescence is present in

A. Solanum

B. Sesbania

C. Trifolium

D. Brassica

**Answer: A**

**76.** Phyllode is present in :-

A. Asparagus

B. Euphorbia

C. Australian Acacia

D. Opuntia

**Answer: C**

Watch Video Solution

**77.** The gynoecium consists of many free pistils in flowers of

A. Aloe

B. Tomato

C. Papaver

D. Michelia

**Answer: D**

Watch Video Solution

**78.** Which one of the following is correctly matched

A. Onion - Bulb

B. Ginger - Sucker

C. Chlamydomonas - Conidia

D. Yeast-Zoospores

**Answer: A**

Watch Video Solution

**79.** How many plants in the list given below have composite fruits that develop from an inflorescence.

Walnut, poppy, radish, fig, pineapple, apple, totato, mulberry.

A. Four

B. Five

C. Two

D. Three

**Answer: D**

Watch Video Solution

**80.** The coconut water and the edible part of coconut are equivalent to or the morphological nature of the edible part of coconut is

A. Endosperm

B. Endocarp

C. Mesocarp

D. Embryo

**Answer: A**

Watch Video Solution

**81.** Vexillary aestivation is charactetic of the family

A. Fabaceae

B. Asteraceae

C. Solanaceae

D. Brassicaceae

**Answer: A**

Watch Video Solution

**82.** In china rose the flowers are

A. actinomorphie, epigynous with valvate aestivation

B. zygomorphic, hypogynous with imbricate aestivation

C. zygomorphic, epigynous with twisted aestivation

D. actinomorphic, hypogynous with twisted aestivation

**Answer: D**

Watch Video Solution

**83.** Age of tree can be estimated by

A. biomass

B. number of annual rings

C. diameter of its heartwood

D. its height and girth

**Answer: B**

Watch Video Solution

**84.** The essential chemical components of many coenzymes are

A. nucleic acids

B. carbohydrates

C. vitamins

D. proteins

**Answer: C**

Watch Video Solution

**85.** Perisperm differs from endosperm in

A. having no reserve food

B. being a diploid tissue

C. its formation by fusion of secondary nucleus with several sperms

D. being a haploid tissue

**Answer: B**

Watch Video Solution

**86.** Seed coat is not thin, membranous in

A. Coconut

B. Groundnut

C. Gram

D. Maize

**Answer: A**

Watch Video Solution

**87.** Among bitter gourd. Mustard, brinjal, pumpkin, chinarose, lupin, cucumber, sunnehemp, gram, guava, bean, chilli, plum,petunia, tomato, rose,withania, potato, onion, aloe and tulip how many plants havehypogynous flower

A. Ten

B. Fifteen

C. Eighteen

D. Six

**Answer: B**

Watch Video Solution

**88.** An example of edible underground stem is

A. Carrot

B. Groundnut

C. Sweet potato

D. Potato

**Answer: D**

Watch Video Solution

**89.** Which one of the following stateme its is correct ?

A. The seed in grasses is not endospermic.

B. Mango is a parthenocarpic fruit.

C. A proteinaceous aleurone layer is present in maize grain.

D. A sterile pistil is called a staminode.

**Answer: C**

Watch Video Solution

**90.** Placenta and pericarp are both edible portions in

A. Apple

B. Banana

C. Tomato

D. Potato

**Answer: C**

Watch Video Solution

**91.** When the margins of sepals or petals overlap one another without any particular direction, the condition in termed as

A. Vexillary

B. lmbricate

C. Twisted

D. Valvate

**Answer: B**

Watch Video Solution

**92.** An aggregate fruit is the one which develops from

A. Multicarpellary syncarpous gynoecium

B. Multicarpellary apocarpus gynoecium

C. Complete inflorescence

D. Multicarpellary superior ovary

**Answer: B**

Watch Video Solution

**93.** Non-albuminous seed is produced in

A. Maize

B. Castor

C. Wheat

D. Pea

**Answer: D**

Watch Video Solution

**94.** Cladode' is a characteristic morphological feature of

A. Asparagus and Ruscus

B. Casuarina and Opuntia

C. Cladophora and Cactus

D. Citrus and Euphorbia

**Answer: A**

**95.** Censer mechanism of seed dispersal is found in

A. Papaveraceae

B. Liliaceae

C. Leguminoceae

D. Rosaceae

**Answer: A**

Watch Video Solution

**96.** Elater mechanism for seed dispersal is exhibited by

A. Riccia

B. Marchantia

C. Dryopteris

D. Funaria

**Answer: B**

Watch Video Solution

**97.** Among china rose, mustard, Brinjal, potato, guava,cucumber onion and tulip, how many plants have superior ovary

A. Four

B. Five

C. Six

D. Three

**Answer: C**

Watch Video Solution

**98.** Flowers are unisexual in

A. Onion

B. Pea

C. Cucumber

D. China rose

**Answer: C**

Watch Video Solution

**99.** Roots play insignificant role in absorption of water in

A. Wheat

B. Sunflower

C. Pistia

D. Pea

**Answer: C**

Watch Video Solution

**100.** The small projections present on the surface of pineapple fruit represent

A. persistent styles

B. persistent stamens

C. persistent calyx

D. persistent bracts

**Answer: D**

Watch Video Solution

**101.** Stems modlfted into flat green organs performing the functions of leaves are known as

A. Phyllodes

B. Phylloclades

C. Scales

D. Cladodes

**Answer: B**

Watch Video Solution

**102.** The standard petal of a papilionaceous corola is also called

A. Pappus

B. Vexillum

C. Corona

D. Carina

**Answer: B**

Watch Video Solution

**103.** Tricarpellary syncarpous gynoecium is found in flowers of

A. Solanaceae

B. Fabaceae

C. Poaceae

D. Liliaceae

**Answer: D**

Watch Video Solution

**104.** Cotyledon of maize grain is called

A. Coleorhiza

B. Coleoptile

C. Scutellum

D. Plumule

**Answer: C**

Watch Video Solution

**105.** Which of the following is not a stem modification

A. Thorns of citrus

B. Tendrils of cucumber

C. Flattened structures of Opuntia

D. Pitcher of Nepenthes

**Answer: D**

Watch Video Solution

**106.** Proximal end of the filament of stamen is attached to the

A. Connective

B. Placenta

C. Thalamus or petal

D. Anther

**Answer: C**

Watch Video Solution

**107.** The term 'polyadelphous' is related to

A. gynoecium

B. androecium

C. corolla

D. calyx

**Answer: B**

Watch Video Solution

**108.** Many plants among Indigofera, Sesbania, Salvia, Allium, Aloe, mustard, groundant,radish,gram and turnip have stamens with different in their flowers

A. Three

B. Four

C. Five

D. Six

**Answer: B**

Watch Video Solution

**109.** Radial symmetry is found in the flowers of

A. Brassica

B. Trifolium

C. Pisum

D. Cassia

**Answer: A**

Watch Video Solution

**110.** Free-central placentation is found in

A. Dianthus

B. Argemone

C. Brassica

D. Citrus

**Answer: A**

Watch Video Solution

**111.** Match Column-I with Column-II and select the correct option using the codes given below

| | Column-I | | Column-II |
|---|---|---|---|
| (A) | Pistils fused together | (i) | Gametogenesis |
| (B) | Formation of gametes | (ii) | Pistillate |
| (C) | Hyphae of higher Ascomycetes | (iii) | Syncarpous |
| (D) | Unisexual female flower | (iv) | Dikaryotic |

A. $\begin{matrix}(1) & (2) & (3) & (4)\\ (iv) & (iii) & (i) & (ii)\end{matrix}$

B. $\begin{matrix}(1) & (2) & (3) & (4)\\ (ii) & (i) & (iv) & (iii)\end{matrix}$

C. $\begin{matrix}(1) & (2) & (3) & (4)\\ (i) & (ii) & (iv) & (iii)\end{matrix}$

D. $\begin{matrix}(1) & (2) & (3) & (4)\\ (iii) & (i) & (iv) & (ii)\end{matrix}$

**Answer: D**

Watch Video Solution

**112.** Coconut fruit is a

A. berry

B. nut

C. capsule

D. drupe

**Answer: D**

Watch Video Solution

**113.** In Bougainvillea, thorns are the modifications of

A. adventitious root

B. stem

C. leaf

D. stipules

**Answer: B**

Watch Video Solution

**114.** The morphological nature of the edible part of coconut is

A. cotyledon

B. endosperm

C. pericarp

D. perisperm

**Answer: B**

Watch Video Solution

**115.** Sweet potato is a modified

A. stem

B. adventitious root

C. tap root

D. rhizome

**Answer: B**

Watch Video Solution

## Assertion Reason Type Questions

**1.** Assertion. An exposed potato tuber develops chlorophyll and turns green.

Reason. Potato tuber is a modified stem and its cells have ability to develop chlorophyll if exposed to light.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: A**

Watch Video Solution

**2.** Assertion. The leaflet of a compound leaf is not a true leaf.

Reason. There is no bud in the axil of a leaflet.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: A**

Watch Video Solution

**3.** Assertion. The fruits of groundnuts are not nuts but underground pods.

Reason. The pods will not develop until the fertilized ovary is pushed under the soil.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: B**

Watch Video Solution

**4.** Assertion. A flower is a modified shoot .

Reason . All the floral parts are borne on the receptacle ( or thalamus).

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: B**

Watch Video Solution

**5.** Assertion. Dispersal of fruits in Xanthium takes place by wind and water.

Reason. Fruits are light and floating.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: D**

Watch Video Solution

**6.** Assertion. Fruits are formed only in angiosperm .

Reason . Fruits develop from ovaries which are found only in angiosperm .

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: A**

Watch Video Solution

**7.** Assertion. Seeded plants are highly evolved.

Reason. Seeds are found in gymnosperms and angiosperms.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: B**

Watch Video Solution

**8.** Assertion : An ovule is transformed into a fruit after fertilization.

Reason : The fruits are formed only from ovules.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: D**

Watch Video Solution

**9.** Assertion. Flowers are structures of sexual reproduction

.

Reason. Various embryological processess of plants occur in the flower.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: A**

Watch Video Solution

**10.** Assertion. Epigynous ovary is always inferior.

Reason. All the floral parts lie above the level of ovary in epigynous condition and, therefore, ovary becomes inferior.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: A**

Watch Video Solution

**11.** Assertion. The placenta is swollen in the ovary of Solanaceae.

Reason. The ovary is obliquely placed on the thalamus in Solanaceae.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: B**

Watch Video Solution

**12.** Assertion. The flowers of the family-Liliaceae are pentamerous.

Reason. Flowers of Liliaceae are pentamerous because it is a dicot family.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: D**

Watch Video Solution

**13.** Assertion. The corolla of the family Fabaceae is papilionaceous.

Reason: The petals of Lamiaceae are fused in such a manner that the corolla looks bilipped.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: B**

Watch Video Solution

**14.** Assertion. Leaves of Smilax show parallel venation.

Reason. Leaves of all the plants belonging to family Liliaceae show parallel venation.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: D**

**15.** Assertion : Ginger has a prostrate-growing rhizome.

Reason : Shoot growth is not affected by gravity

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: C**

Watch Video Solution

**16.** Assertion : Coconut tree is distributed in coastal areas over a large part of the world.

Reason : Coconut fruit can float and get dispersed over thousands of Kilometers before losing viability.

A. If both Assertion and Reason are true and the Reason is a correct explanation of the Assertion.

B. If both Assertion and Reason are true but Reason is not a correct explanation of the Assertion.

C. If Assertion is true but the Reason is false.

D. If both Assertion and Reason are false.

**Answer: B**

Watch Video Solution