



# PHYSICS

## BOOKS - NCERT FINGERTIPS PHYSICS (HINGLISH)

### KINETIC THEORY

#### Molecular Nature Of Matter

1. Molecular motion shows itself as

A. temperature

B. internal energy

C. friction

D. viscosity

**Answer: A**



**Watch Video Solution**

2. Cooking gas container are kept in a lorry moving with uniform speed. The temperature of the gas molecules inside will

A. increase

B. decrease

C. remains the same

D. decrease for some and increase for others

**Answer: C**



**Watch Video Solution**

**Behaviour Of Gases**

1. A real gas behaves like an ideal gas if its





- A. both pressure and temperature are high
- B. both pressure and temperature are low
- C. pressure is high and temperature is low
- D. pressure is low and temperature is high

**Answer: D**



**Watch Video Solution**

2. Which of the following graphs represent the behaviour of an ideal gas ?



**Answer: A**



Watch Video Solution

3. If the pressure and the volume of certain quantity of ideal gas are halved, then its temperature

- A. is doubled
- B. becomes one-fourth
- C. remains constant
- D. become four times

**Answer: B**



Watch Video Solution

4. An air bubble of volume  $1.0\text{cm}^3$  rises from the bottom of a lake 40 m deep at a temperature of  $12^\circ\text{C}$ . To what volume does it grow when it reaches the surface, which is at a temperature of  $35^\circ\text{C}$ . ? Given  $1\text{atm} = 1.01 \times 10^5\text{Pa}$ .

A.  $10.6 \times 10^{-6}\text{m}^3$

B.  $5.3 \times 10^{-6}\text{m}^3$

C.  $2.8 \times 10^{-6}\text{m}^3$

D.  $15.6 \times 10^{-6} m^3$

**Answer: B**



**Watch Video Solution**

5. The diameter of an oxygen molecule is  $3 \text{ \AA}$ . The ratio of molecular volume to the actual volume occupied by the oxygen gas at STP is

A.  $2 \times 10^{-4}$



B.  $1 \times 10^{-4}$

C.  $1.5 \times 10^{-4}$

D.  $4 \times 10^{-4}$

**Answer: D**



**Watch Video Solution**

6. A balloon contains  $1500m^3$  of helium at  $27^\circ C$  and 4 atmospheric pressure. The volume of helium at  $-3^\circ C$  temperature and 2 atmospheric pressure will be

A.  $1500m^3$

B.  $1700m^3$

C.  $1900m^3$

D.  $2700m^3$


**Answer: D**



**Watch Video Solution**

7. Pressure versus temperature graph of an ideal gas of equal number of moles of different volumes is plotted as shown in Fig.

Choose the correct alternative.



- A.  $V_1 = V_2 = V_3 = V_4$
- B.  $V_4 > V_3 > V_2 > V_1$
- C.  $V_1 = V_2, V_3 = V_4$  and  $V_2 > V_3$
- D.  $V_1 = V_2, V_3 = V_4$  and  $V_2 < V_3$


**Answer: C**



Watch Video Solution

8. Given is the graph between  $\frac{PV}{T}$  and P for 1 gm of oxygen gas at two different temperatures  $T_1$  and  $T_2$  Fig. Given, density of oxygen =  $1.427 \text{kgm}^{-3}$ . The value of  $(PV)/(T)$  at the point A and the relation

between  $T_1$  and  $T_2$  are respectively :



A.  $0.259 \text{ JK}^{-1}$  and  $T_1 < T_2$

B.  $8.314 \text{ gJmol}^{-1} \text{ K}^{-1}$  and  $T_1 > T_2$

C.  $0.259 \text{ JK}^{-1}$  and  $T_1 > T_2$

D.  $4.28 \text{ gJK}^{-1}$  and  $T_1 < T_2$

**Answer: C**



Watch Video Solution

9. A vessel has 6 g of oxygen at pressure  $P$  and temperature 400 K. A small hole is made in it so that oxygen leaks out. How much oxygen leaks out if the final pressure is  $\frac{P}{2}$  and temperature 300 K?

A. 5 g

B. 4 g

C. 2 g

D. 3 g

**Answer: C**



**Watch Video Solution**

**10.** A vessel has 6g of hydrogen at pressure  $P$  and temperature 500K. A small hole is made in it so that hydrogen leaks out. How much hydrogen leaks out if the final pressure is  $P/2$  and temperature falls to 300 K ?

A. 2 g

B. 3 g

C. 4 g

D. 1 g

**Answer: D**



**Watch Video Solution**

**11.** A vessel contains two non-reactive gases neon (monoatomic) and oxygen (diatomic). The ratio of their partial pressures is 3:2. Estimate the ratio of



(i) number of molecules, and

(ii) mass density of neon and oxygen in the vessel.

Atomic mass of neon = 20.2 u, and molecular mass of oxygen = 32.0 u.

A.  $\frac{3}{2}$

B.  $\frac{2}{3}$

C.  $\frac{1}{3}$

D.  $\frac{1}{2}$

**Answer: A**



12. A cylinder contained  $10\text{kg}$  of gas at pressure  $10^7 \frac{N}{m^2}$ . The quantity of gas taken out of cylinder if final pressure is  $2.5 \times 10^6 N/m^2$  is (Assume temperature of gas is constant)

- A. 9.5 kg
- B. 7.5 kg
- C. 14.2 kg
- D. zero

**Answer: B**



**Watch Video Solution**

**13.** A gas at 300 K has pressure  $4 \times 10^{-10} N/m^2$ . IF  $k = 1.38 \times 10^{-23} J/K$ , the number of molecule/ $cm^3$  is of the order of

A.  $10^3$

B.  $10^5$

C.  $10^6$

D.  $10^9$

**Answer: B**



**Watch Video Solution**

**14.** The volume of vessel A is twice the volume of another vessel B and both of them are filled with the same gas. If the gas in A is at twice the temperature and twice the pressure in comparison to the gas in B, what is the ratio of number of gas molecule in A and B ?

A.  $\frac{1}{2}$

B.  $\frac{2}{1}$

C.  $\frac{3}{2}$

D.  $\frac{2}{3}$

**Answer: B**



**Watch Video Solution**

**15.** When the temperature of a gas filled in a closed vessel is increased by  $1^{\circ}C$ , its pressure

increases by 0.4 percent. The initial temperature of gas was

A.  $250^{\circ}C$

B.  $25^{\circ}C$

C. 250 K

D. 25 K

**Answer: C**



**Watch Video Solution**

**16.** The equation of state for 5 g of oxygen at a pressure  $P$  and temperature  $T$ , when occupying a volume  $V$ , will be

A.  $PV = (5/32) RT$

B.  $PV = 5RT$

C.  $PV = (5/2) RT$

D.  $PV = (5/16) RT$

**Answer: A**



**Watch Video Solution**

17. A vessel contains 1 mole of  $O_2$  gas (relative molar mass 32) at a temperature  $T$ . The pressure of the gas is  $P$ . An identical vessel containing one mole of He gas (relative molar mass 4) at temperature  $2T$  has a pressure of

A.  $\frac{P}{8}$

B.  $P$

C.  $2P$

D.  $8P$

**Answer: C**





Watch Video Solution

**18.** In a certain region of space there are only 5 gaseous there is 3 K. The pressure of this gas is

$$\left(k_B = 1.38 \times 10^{-23} \text{Jmol}^{-1} \text{K}^{-1}\right)$$

A.  $20.7 \times 10^{-16} \text{Nm}^{-2}$

B.  $20.4 \times 10^{-17} \text{Nm}^{-2}$

C.  $10.7 \times 10^{-16} \text{Nm}^{-2}$

D.  $10.7 \times 10^{-17} \text{Nm}^{-2}$

**Answer: B**



**Watch Video Solution**

**19.** A sample of an ideal gas occupies a volume  $V$  at pressure  $P$  and absolute temperature  $T$ . The mass of each molecule is  $m$ , then the density of the gas is

A.  $mKT$

B.  $\frac{Pm}{KT}$

C.  $\frac{P}{km}$

D.  $\frac{P}{KT}$

**Answer: B**



**Watch Video Solution**

**20.** One half mole each of nitrogen, oxygen and carbon dioxide are mixed in enclosure of volume 5 litres and temperature  $27^{\circ}C$ .

Calculate the pressure exerted by the mixture.

Given  $R = 8.31 J mol^{-1} K^{-1}$ .

A.  $7.48 \times 10^5 Nm^{-2}$

B.  $5 \times 10^5 Nm^2$

C.  $6 \times 10^5 Nm^2$

D.  $3 \times 10^5 Nm^{-2}$

**Answer: A**



**Watch Video Solution**

**21.** From a certain apparatus, the diffusion rate of hydrogen has an average value of  $28.7cm^3s^{-1}$ . The diffusion of another gas under the same condition is measured to have

an average rate of  $7.2\text{cm}^3\text{s}^{-1}$ . Identify the gas.

A. Nitrogen

B. Helium

C. Argon

D. Oxygen

**Answer: D**



**Watch Video Solution**

22. The volume of water molecule is

Take, density of water is  $10^3 \text{ kg m}^{-3}$  and  
avogadro's number =  $6.023 \times 10^{23} \text{ mole}^{-1}$ )

A.  $3 \times 10^{-28} \text{ m}^3$

B.  $3 \times 10^{-29} \text{ m}^3$

C.  $1.5 \times 10^{-28} \text{ m}^3$

D.  $1.5 \times 10^{-29} \text{ m}^3$

**Answer: B**



**Watch Video Solution**

# Kinetic Theory Of An Ideal Gas

1. Which one of the following is not an assumption in the kinetic theory of gases?

A. The volume occupied by the molecules of the gas is negligible.

B. The force of attraction between the molecules is negligible.

C. The collision between the molecules are elastic.

D. All molecules have same speed.

**Answer: D**



**Watch Video Solution**

2. Pressure of a gas at constant volume is proportional to

A. total internal energy of the gas

B. average kinetic energy of the molecules



C. average potential energy of the molecules

D. total energy of the gas

**Answer: A**



**Watch Video Solution**

3. The kinetic theory of gases gives the formula  $PV = \frac{1}{3}Nm\bar{v}^2$  for the pressure P exerted by a gas enclosed in a volume V. The term Nm represents

A. the mass of a mole of the gas

B. the mass of the present in the volume  $V$

C. the averatge mass of one molecule of  
the gas

D. the total number of molecules present  
in volume  $V$

**Answer: B**



**View Text Solution**

4. When an ideal gas is compressed adiabatically, its temperature rises the molecules on the average have more energy than before. The kinetic energy increases,

A. because of collision with moving parts of the wall only.

B. because of collision with the entire wall.

C. because the molecules get accelerated in their motion inside the volume.

D. because the redistribution of energy amongst the molecules.

**Answer: A**



**Watch Video Solution**

5. A gas is filled in a container at pressure  $P_0$ . If the mass of molecules is halved and their rms speed is doubled, then the resultant pressure would be

A.  $2P_0$

B.  $4P_0$

C.  $\frac{P_0}{4}$

D.  $\frac{P_0}{2}$

**Answer: A**



**Watch Video Solution**

6. A vessel is filled with a gas at a pressure of 76 cm of mercury at a certain temperature. The mass of the gas is increased by 50% by

introducing more gas in the vessel at the same temperature. Find the resultant pressure of the gas.

- A. 76 cm of mercury
- B. 108 cm of mercury
- C. 112 cm mercury
- D. 114 cm of mercury

**Answer: D**



**Watch Video Solution**

7. 0.014 kg of nitrogen is enclosed in a vessel at a temperature of  $24^{\circ}C$ . At which temperature the rms velocity of nitrogen gas is twice of its the rms velocity at  $27^{\circ}C$ ?

A. 1200 K

B. 600 K

C. 300 K

D. 150 K

**Answer: A**



**Watch Video Solution**

8. At what temperature is the rms velocity of a hydrogen molecule equal to that of an oxygen molecule at  $47^{\circ}C$ ?

A. 10 K

B. 20 K

C. 30 K

D. 40 K

**Answer: B**



**Watch Video Solution**



9. If three molecules have velocities  $0.5\text{km s}^{-1}$ ,  $1\text{km s}^{-1}$  and  $2\text{km s}^{-1}$ , the ratio of the rms speed and average speed is

A. 2.15

B. 1.13

C. 0.53

D. 3.96

**Answer: B**



10. The kinetic energy of 1 g molecule of a gas, at normal temperature and pressure, is

A.  $0.56 \times 10^{-4} J$

B.  $2.4 \times 10^{-2} J$

C.  $1.3 \times 10^2 J$

D.  $3.4 \times 10^3 J$

**Answer: D**



11. An insulated container containing monoatomic gas of molar mass  $m$  is moving with a velocity  $V_0$ . If the container is suddenly stopped, find the change in temperature.

A.  $\frac{mv_0^2}{2R}$

B.  $\frac{mv_0^2}{3R}$

C.  $\frac{R}{mv_0^2}$

D.  $\frac{3mv_0^2}{2R}$

**Answer: B**



Watch Video Solution

12. Two moles of gas A at  $27^{\circ} C$  mixed with a 3 moles of gas at  $37^{\circ} C$ . If both are monatomic ideal gases, what will be the temperature of the mixture ?

A.  $66^{\circ} C$

B.  $11^{\circ} C$

C.  $22^{\circ} C$

D.  $33^{\circ} C$

**Answer: D**



**Watch Video Solution**

**13.** The average kinetic energy of  $O_2$  at a particular temperature is 0.768 eV. The average kinetic energy of  $N_2$  molecules in eV at the same temperature is

A. 0.0015

B. 0.0030

C. 0.048

D. 0.768

**Answer: D**



**Watch Video Solution**

**14.** Calculate the rms speed of smoke particles of mass  $5 \times 10^{-17} \text{ kg}$  in their Brownian motion in air at NTP. Given

$$k_B = 1.38 \times 10^{-23} \text{ J/K}$$

A.  $3 \times 10^{-2} \text{ ms}^{-1}$

B.  $1.5 \times 10^{-2} \text{ms}^{-1}$

C.  $3 \times 10^{-3} \text{ms}^{-1}$

D.  $1.5 \times 10^{-3} \text{ms}^{-1}$

**Answer: B**



**Watch Video Solution**

**15.** The molecules of a given mass of a gas have root mean square speeds of  $100 \text{ms}^{-1}$  at  $27^\circ \text{C}$  and 1.00 atmospheric pressure. What will be the root mean square

speeds of the molecules of the gas at  $127^{\circ}C$   
and 2.0 atmospheric pressure?

A.  $\frac{200}{\sqrt{3}}$

B.  $\frac{100}{\sqrt{3}}$

C.  $\frac{400}{3}$

D.  $\frac{200}{3}$

**Answer: A**



**Watch Video Solution**



16. The temperature of an ideal gas is increased from  $27^{\circ}C \rightarrow 127^{\circ}C$ , the percentage increase in  $V_{rms}$  is

A. 37 %

B. 11 %

C. 33 %

D. 15.5 %

**Answer: D**



**Watch Video Solution**

17. At what temperature is the root mean square speed of an atom in an argon gas cylinder equal to the r.m.s. speed of a helium gas atom at  $-20^{\circ}C$  ? (Atomic mass of Ar = 39.9 u, of He = 4.0 u).

A.  $2.52 \times 10^3 K$

B.  $2.52 \times 10^2 K$

C.  $4.03 \times 10^3 K$

D.  $4.03 \times 10^2 K$

**Answer: A**



Watch Video Solution

18. The temperature of an ideal gas is increased from 120 K to 480 K. If at 120 K the root mean square velocity of the gas molecules is  $v$ , at 480 K it becomes

A.  $4v_{rms}$

B.  $2v_{rms}$

C.  $\frac{2v_{rms}}{2}$

D.  $\frac{v_{rms}}{4}$

**Answer: B**



**Watch Video Solution**

## Law Of Equipartition Of Energy

1. According to the law of equipartition of energy, the energy associated with each degree of freedom is :

A.  $E = k_B T$

B.  $E = \frac{1}{2} k_B T$

C.  $E = 3k_B T$

D.  $E = \frac{3}{2}k_B T$

**Answer: D**



**Watch Video Solution**

2. The internal energy of one gram of helium at 100 K and one atmospheric pressure is

A. 100 J

B. 1200 J

C. 300 J

D. 500 J

**Answer: C**



**Watch Video Solution**

**3.** One kg of a diatomic gas is at pressure of  $8 \times 10^4 \text{ N/m}^2$ . The density of the gas is  $4 \text{ kg/m}^3$ . What is the energy of the gas due to its thermal motion?

A.  $3 \times 10^4 J$

B.  $5 \times 10^4 J$

C.  $6 \times 10^4 J$

D.  $4 \times 10^4 J$

**Answer: B**



**Watch Video Solution**

4. A gas mixture consists of 2 moles of oxygen and 4 of Argon at temperature T. Neglecting

all vibrational modes, the total internal energy of the system is

A. 4 RT

B. 9 RT

C. 11 RT

D. 15 RT

**Answer: C**



**Watch Video Solution**



5. The average energy per molecule of a triatomic gas at room temperature  $T$  is

A.  $3kT$

B.  $\frac{1}{2}kT$

C.  $\frac{3}{2}kT$

D.  $\frac{5}{2}kT$

**Answer: A**



**Watch Video Solution**

## Specific Heat Capacity

1. If for a gas,  $\frac{R}{C_V} = 0.67$ , the gas is

A. monatomic

B. diatomic

C. polyatomic

D. mixture of diatomic and polyatomic molecules

**Answer: A**



**Watch Video Solution**

2. If  $C_p$  and  $C_v$  denoted the specific heats of unit mass of nitrogen at constant pressure and volume respectively, then

A.  $C_p = C_v = \frac{R}{28}$

B.  $C_p - C_v = \frac{R}{7}$

C.  $C_p - C_v = \frac{R}{14}$

D.  $C_p - C_v = R$

**Answer: A**



Watch Video Solution

3. The ratio of the molar heat capacities of a diatomic gas at constant pressure to that at constant volume is

A.  $\frac{7}{5}$

B.  $\frac{3}{2}$

C.  $\frac{3}{5}$

D.  $\frac{5}{2}$

**Answer: A**

 [Watch Video Solution](#)

4. The heat capacity per mole of water is ( $R$  is universal gas constant)

A.  $9R$

B.  $\frac{9}{2}R$

C.  $6R$

D.  $5R$

**Answer: A**

 [Watch Video Solution](#)

5. Three moles of oxygen are mixed with two moles of helium. What will be the ratio of specific heats at constant pressure and constant volume for the mixture ?

A. 2.5

B. 3.5

C. 1.5

D. 1

**Answer: C**



**Watch Video Solution**

6. A gaseous mixture enclosed in a vessel consists of one gram mole of a gas A with

$\gamma = \left(\frac{5}{3}\right)$  and some amount of gas B with

$\gamma = \frac{7}{5}$  at a temperature T.

The gases A and B do not react with each other and are assumed to be ideal. Find the number of gram moles of the gas B if  $\gamma$  for the gaseous mixture is

$\left(\frac{19}{13}\right)$ .

A. 2

B. 12

C. 16

D. 8

**Answer: A**



**Watch Video Solution**

7. A cylinder of fixed capacity 44.8 litres contains helium gas at standard temperature and pressure. What is the amount of heat



needed to raise the temperature of the gas in the cylinder by  $15^{\circ}C$  ? Given  $R = 8.31 \text{ J mole}^{-1} \text{ K}^{-1}$  . (For monoatomic gas,  $C_v = 3R/2$ )

A. 265 J

B. 310.10 J

C. 373.95 J

D. 387.97 J

**Answer: C**



**Watch Video Solution**

8.  $(1/2)$  mole of helium is contained in a container at STP how much heat energy is needed to double the pressure of the gas, keeping the volume constant? Heat capacity of gas is  $3Jg^{-1}K^{-1}$ .

A. 1436 J

B. 736 J

C. 1638 J

D. 5698 J

**Answer: C**



**Watch Video Solution**

9. One mole of an ideal monoatomic gas at temperature  $T_0$  expands slowly according to the law  $\frac{p}{V} = \text{constant}$ . If the final temperature is  $2T_0$ , heat supplied to the gas is

A.  $2RT_0$

B.  $RT_0$

C.  $\frac{3}{2}RT_0$

D.  $\frac{1}{2}RT_0$

**Answer: A**



**Watch Video Solution**

**10.** If a gas has  $n$  degrees of freedom ratio of specific heats of gas is

A.  $\frac{1 + n}{2}$

B.  $1 + \frac{1}{n}$

C.  $1 + \frac{n}{2}$

D.  $1 + \frac{2}{n}$

**Answer: D**



**Watch Video Solution**

**11.** 1 mole of a gas with  $\gamma = 7/5$  is mixed with 1 mole of a gas with  $\gamma = 5/3$ , then the value of  $\gamma$  for the resulting mixture is

A.  $\frac{7}{5}$

B.  $\frac{2}{5}$

C.  $\frac{3}{2}$

D.  $\frac{12}{7}$

**Answer: C**



**Watch Video Solution**

**12.** A molecule of a gas has six degrees of freedom. Then the molar specific heat of the gas at constant volume is

A.  $\frac{R}{2}$

B.  $R$

C.  $\frac{3R}{2}$

D.  $3R$

**Answer: D**



**Watch Video Solution**

**13.** Two moles of oxygen are mixed with eight moles of helium. The effective specific heat of the mixture at constant volume is

A.  $1.3R$

B.  $1.4R$

C.  $1.7R$

D.  $1.9R$

**Answer: C**



**Watch Video Solution**

**Higher Order Thinking Skills**



1.  $N$  molecules each of mass  $m$  of gas  $A$  and  $2N$  molecules each of mass  $2m$  of gas  $B$  are contained in the same vessel which is maintained at a temperature  $T$ . The mean square velocity of the molecules of  $B$  type is denoted by  $v^2$  and the mean square of the x-component of the velocity of  $A$  type is denoted by  $\omega^2$ . What is the ratio of  $\omega^2 / v^2 = ?$

A. 3:2

B. 1:3

C. 2:3

D. 1:1

**Answer: C**



**Watch Video Solution**

2. Two cylinder having  $m_1g$  and  $m_2g$  of a gas at pressure  $P_1$  and  $P_2$  respectively are put in communication with each other, temperature remaining constant. The common pressure reached will be

A.  $\frac{P_1 P_2 (m_1 + m_2)}{P_2 m_1 + P_1 m_2}$

B.  $\frac{P_1 P_2 m_1}{P_2 m_1 + P_1 m_2}$

C.  $\frac{m_1 m_2 (P_1 + P_2)}{P_2 m_1 (P_1 + P_2)}$

D.  $\frac{m_1 m_2 P_2}{P_2 m_1 + P_1 m_2}$

**Answer: A**



**Watch Video Solution**

3. A gas in a cylinder. Its temperature is increased by 20% on kelvin scale and volume is reduced to 90% how much percentage of

the gas has to leak for pressure to remain constant?

A. 20 %

B. 25 %

C. 30 %

D. 40 %

**Answer: B**



**Watch Video Solution**

4. A thin tube of uniform cross-section is sealed at both ends. It lies horizontally, the middle 5 cm containing mercury and the two equal end containing air at the same pressure  $P$ . When the tube is held at an angle of  $60^\circ$  with the vertical direction, the length of the air column above and below the mercury column are 46 cm and 44.5 cm respectively. Calculate the pressure  $P$  in centimeters of mercury. (The temperature of the system is kept at  $30^\circ C$ ).

A. 75.4

B. 45.8

C. 67.5

D. 89.3

**Answer: A**



**Watch Video Solution**

5. An insulated container containing monoatomic gas of molar mass  $m$  is moving

with a velocity  $V_0$ . If the container is suddenly stopped, find the change in temperature.

A.  $\frac{mv_0^2}{2R}$

B.  $\frac{mv_0^2}{3R}$

C.  $\frac{3mv_0^2}{2R}$

D.  $\frac{1}{2} \frac{mv_0^2}{R}$

**Answer: B**



**Watch Video Solution**

6. A gas has molar heat capacity  $C = 37.55 \text{ J mole}^{-1} \text{ K}^{-1}$ , in the process  $PT = \text{constant}$ , find the number of degree of freedom of the molecules of the gas.

A. 6

B. 3

C. 1

D. 5

**Answer: D**



Watch Video Solution



7. Calculate the mean free path of nitrogen at  $27^\circ C$  when pressure is 1.0 atm. Given, diameter of nitrogen molecule =  $1.5\text{\AA}$ ,  $k = 1.38 \times 10^{-23} JK^{-1}$ . If the average speed of nitrogen molecules is  $675ms^{-1}$ , find the time taken by the molecule between two successive collisions and the frequency of collisions.

A. 0.6 ns

B. 0.4 ns

C. 0.8 ns

D. 0.3 ns

**Answer: A**



**Watch Video Solution**

8. Ten small planes are flying at a speed of  $150\text{km}/\text{h}$  in total darkness in an air space that is  $20 \times 20 \times 1.5\text{km}^3$  in volume. You are in one of the planes, flying at random within this space with no way of knowing where the other

planes are, On the average about how long a time will elapse between near collision with your plane. Assume for this rough computation that a safety region around the plane can be approximately by a sphere of radius 10 m.

A. 125 h

B. 220 h

C. 432 h

D. 225 h

**Answer: D**



Watch Video Solution

9. Two identical containers A and B with frictionless pistons contain the same ideal gas at the same temperature and the same velocity  $V$ . The mass of the gas in A is  $m_A$ , and that in B is  $m_B$ . The gas in each cylinder is now allowed to expand isothermally to the same final volume  $2V$ . The changes in the pressure in A and B are found to be  $\Delta P$  and  $1.5\Delta P$  respectively. Then

A.  $4m_A = 9m_B$

B.  $2m_A = 3m_B$

C.  $3m_A = 2m_B$

D.  $9m_A = 4m_B$

**Answer: C**



**Watch Video Solution**

**10.** Consider a rectangular block of wood moving with a velocity  $v_0$  in a gas at temperature  $T$  and mass density  $p$ . Assume the

velocity is along x-axis and the area of cross-section of the block perpendicular to  $v_0$  is  $A$ .

show that the drag force on the block is

$4rAv_0\sqrt{\frac{kT}{m}}$  where,  $m$  is the mass of the gas molecule.

A.  $4pAv_0\sqrt{\frac{kT}{m}}$

B.  $2pAv_0\sqrt{\frac{kT}{3m}}$

C.  $\frac{pA}{2v_0}\sqrt{\frac{kT}{m}}$

D.  $\frac{v_0}{pA}\sqrt{\frac{kT}{2m}}$

**Answer: A**





Watch Video Solution

## Ncert Exemplar

1. A cubic vessel (with face horizontal + vertical ) contains an ideal gas at NTP. The vessel is being carried by a rocket which is moving at a speed of  $500\text{ms}^{-1}$  in vertical direction. The pressure of the gas inside the vessel as observed by us on the ground.

A. remains the same because 500 m/s is very much smaller than  $v_{rms}$  of the gas

B. remains the same because motion of the vessel as a whole does not affect the relative motion of the gas molecules and the walls

C. will increase by a factor equal to

$$\left[ v_{rms}^2 + (500)^2 \right] / v_{rms}^2, \text{ where } v_{rms} \text{ was}$$

the original mean square velocity of the gas



D. will be different on the top wall and bottom wall of the vessel.


**Answer: B**



**Watch Video Solution**

2. Mole of an ideal gas is contained in a cubical volume  $V$ , ABCDEFGH at 300 K (figure). One face of the cube (EFGH) is made up of a material which totally absorbs any gas

molecule incident on it .At any given time.



A. the pressure on EFGH would be zero

B. the pressure on all the faced will be equal

C. the pressure of EFGH would be double the pressure on ABCD

D. the pressure on EFGH would be half that  
on ABCD.

**Answer: D**



**Watch Video Solution**

**3. Boyle's law is applicable for an**

A. adiabatic process

B. isothermal process

C. isobaric process

D. isochoric process


**Answer: B**



**Watch Video Solution**

4. A cylinder containing an ideal gas is in vertical position and has a piston of mass  $M$  that is able to move up or down without friction (figure). If the temperature is

increased



- A. both  $P$  and  $V$  of the gas will change
- B. only  $P$  will increase according to Charle's law.
- C.  $V$  will change but not  $P$
- D.  $P$  will change but not  $V$


**Answer: C**



**Watch Video Solution**

5. Volume versus temperature graphs for a given mass of an ideal gas are shown in figure. At two different values of constant pressure. What can be inferred about relation between

$P_1$  and  $P_2$  ?



A.  $P_1 > P_2$

B.  $P_1 = P_2$

C.  $P_1 < P_2$

D. data is insufficient

**Answer: A**



**Watch Video Solution**

6. 1 mole of  $H_2$  gas is contained in box of volume  $V = 1.00m^3$  at  $T = 300K$ . The gas is heated to a temperature of  $T = 3000 K$  and the gas gets converted to a gas of hydrogen atoms. The final pressure would be (considering all gases to be ideal)

A. same as the pressure initially



B. 2 times the pressure initially

C. 10 times the pressure initially

D. 20 times the pressure initially

**Answer: D**



**Watch Video Solution**

7. A vessel of volume  $V$  contains a mixture of 1 mole of hydrogen and 1 mole oxygen (both considered as ideal). Let  $f_1(v)dv$ , denote the fraction of molecules with speed between  $v$

and  $(v+ dv)$  with  $f_2(v)dv$ , similarly for oxygen .

Then ,

A.  $f_1 (v) + f_2 (v) =$  obeys the Maxwell's  
distribution law

B.  $f_1(v), f_2(v)$  will obey the Maxwell's  
distribution law separately

C. Neither  $f_1 (v)$  nor  $f_2 (v)$  will obey the  
Maxwell's distribution law

D.  $f_2 (v)$  and  $f_1 (v)$  will be the same

**Answer: B**



Watch Video Solution

8. An inflated rubber balloon contains one mole of an ideal gas has a pressure  $p$ , volume  $V$  and temperature  $T$ . if the temperature rises to  $1.1 T$ , and the volume is increased to  $1.05 V$ , the final pressure will be

A.  $1.1 P$

B.  $P$

C. less than  $P$

D. between P and 1.1 P.

**Answer: D**



**Watch Video Solution**

## Assertion And Reason

**1. Assertion:** All molecules of an ideal gas move with the same speed.

**Reason:** There is no attraction between the molecules in an ideal gas.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not be correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

**Answer: A**



**Watch Video Solution**

2. Assertion: In a mixture of gases at a fixed temperature, the heavier molecule has the lower average speed.

Reason: Temperature of a gas is a measure of the average kinetic energy of a molecule.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not be correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

**Answer: B**



**View Text Solution**

**3. Assertion:** In case of collision of gas molecules in a given amount of gas, total kinetic energy is conserved.

**Reason:** All collisions of the gas molecules in a given amount of gas are elastic.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not be correct explanation of



assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

**Answer: A**



**View Text Solution**

**4.** Assertion : The rms velocity and most probable speeds of the molecules in a gas are same.

The Maxwell distribution curve for the speed of the molecules in a gas is symmetrical.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not be correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

**Answer: D**



**Watch Video Solution**

5. Assertion: The ratio of rms speed and average speed of a gas molecules at a given temperature is  $\sqrt{3} : \sqrt{8/\pi}$

Reason:  $c_{rms} > c_{av}$ .

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

**Answer: B**



**Watch Video Solution**

6. Assertion: Average kinetic energy per molecule of any ideal monoatomic gas is

$$\frac{3}{2}k_B T$$

Reason: Average kinetic energy depends only on temperature and is independent of the nature of the gas.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not be correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

**Answer: B**



**View Text Solution**

7. Assertion: For a mixture of non reactive ideal gases, the total pressure gets contribution from each gas in the mixture.

Reason: In equilibrium, the average kinetic energy of the molecules of different gases will be equal.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not be correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

**Answer: B**



[View Text Solution](#)



**8. Assertion:** Each vibrational mode gives two degrees of freedom.

**Reason:** By law of equipartition of energy, the energy for each degree of freedom in thermal equilibrium is  $2k_B T$ .

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not be correct explanation of

assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

**Answer: C**



**View Text Solution**

**9. Assertion :** Specific heat of a gas at constant pressure is greater than its specific heat at constant volume.

This is because at constant pressure, some heat is spent in expansion of the gas.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not be correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

**Answer: A**



**Watch Video Solution**

**10. Assertion :** The ratio  $C_P / C_v$  for a diatomic gas is more than that for a monoatomic gas.

**Reason :** The molecules of a monoatomic gas have more degrees of freedom than those of a diatomic gas.

A. If both assertion and reason are true and reason is the correct explanation of

assertion.

B. If both assertion and reason are true but reason is not be correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

**Answer: D**



**Watch Video Solution**