



PHYSICS

BOOKS - NCERT FINGERTIPS PHYSICS (HINGLISH)

THERMAL PROPERTIES OF MATTER

Temperature And Heat

1. In order that the heat flows from one part of a solid to another part, what is required

- A. uniform density
- B. temperature gradient
- C. density gradient
- D. uniform temperature

Answer: B



Watch Video Solution

2. How much work can be done by 100 calories of heat?

A. 100 J

B. 420 J

C. 42 J

D. 4200 J

Answer: B



Watch Video Solution

Measurement Of Temperature

1. At what temperature do the Celsius and Fahrenheit readings have the same numerical value ?

A. 273

B. -273

C. -40

D. 40

Answer: C



Watch Video Solution

2. When a thermometer is taken from the melting ice to warm liquid , the mercury level rises to $\frac{2^{th}}{5}$ of the distance between the lower and the upper fixed points . The temperature of liquid in K is

A. 217.15

B. 313.15

C. 220

D. 330

Answer: B



Watch Video Solution

Thermal Expansion

1. The volume of a block of a metal changes by 0.12% when it is heated through $20^\circ C$. The coefficient of linear expansion of the metal is

A. $4 \times 10^{-5} \text{ } ^\circ C^{-1}$

B. $2 \times 10^{-5} \text{ } ^\circ C^{-1}$

C. $0.5 \times 10^{-5} \text{ } ^\circ C^{-1}$

D. $4 \times 10^{-4} \text{ } ^\circ C^{-1}$

Answer: B



Watch Video Solution

2. There is a hole in the middle of a copper plate . When heating the plate , diameter of hole would

- A. always increase
- B. always decrease
- C. remains the same
- D. none of these

Answer: A



Watch Video Solution

3. Length of wire at room temperature is 4.55 m , when the temperature increases up to $100^{\circ}C$ then its length becomes 4.57 m . The coefficient of linear expansion (α) of the given wire is

A. $5.021 \times 10^{-5} K^{-1}$

B. $6.021 \times 10^{-5} K^{-1}$

C. $7.021 \times 10^{-5} K^{-1}$

$$D. 8.021 \times 10^{-5} K^{-1}$$

Answer: D



Watch Video Solution

4. To increase the length of brass rod by 2% its temperature should increase by
($\alpha = 0.00002^\circ C^{-1}$)

A. $800^\circ C$

B. $900^\circ C$

C. $1000^\circ C$

D. $1100^{\circ}C$

Answer: C



Watch Video Solution

5. At $50^{\circ}C$, a brass rod has a length 50 cm and a diameter 2 mm. It is joined to a steel rod of the same length and diameter at the same temperature. The change in the length of the composite rod when it is heated to $250^{\circ}C$ is (Coefficient of linear expansion of brass =

$2.0 \times 10^{-5} \text{ } ^\circ\text{C}^{-1}$, coefficient of linear expansion of steel = $1.2 \times 10^{-5} \text{ } ^\circ\text{C}^{-1}$)

A. 0.28 cm

B. 0.30 cm

C. 0.32 cm

D. 0.34 cm

Answer: C



Watch Video Solution

6. The moment of inertia of a rod about its perpendicular bisector is I . When the temperature of the rod is increased by ΔT , the increase in the moment of inertia of the rod about the same axis is (Here , α is the coefficient of linear expansion of the rod)

A. $\alpha I \Delta T$

B. $2\alpha I \Delta T$

C. $\frac{\alpha I \Delta T}{2}$

D. $\frac{2I \Delta T}{\alpha}$

Answer: B



Watch Video Solution

7. A brass wire 1.8 m long at $27^{\circ}C$ is held taut with negligible tension between two rigid supports. Diameter of the wire is 2 mm, its coefficient of linear expansion, $\alpha_{\text{Brass}} = 2 \times 10^{-5} \text{ } ^{\circ}C^{-1}$ and its young's modulus, $Y_{\text{Brass}} = 9 \times 10^{10} \text{ Nm}^{-2}$. If the wire is cooled to a temperature $-39^{\circ}C$, tension developed in the wire is

A. $2.7 \times 10^2 N$

B. $3.7 \times 10^2 N$

C. $4.7 \times 10^2 N$

D. $5.7 \times 10^2 N$

Answer: B



Watch Video Solution

8. The coefficient of volume expansion of liquid is γ . The fractional change in its density for ΔT rise in temperature is

A. $\gamma\Delta T$

B. $\frac{\Delta T}{\gamma}$

C. $1 + \gamma\Delta t$

D. $1 - \gamma\Delta T$

Answer: A



Watch Video Solution

9. A rectangular block is heated from $0^\circ C$ to $100^\circ C$. The percentage increase in its length is 0.2% . The percentage increase in its volume is

A. 0.6 %

B. 0.10 %

C. 0.2 %

D. 0.4 %





Answer: A



Watch Video Solution

10. Which of the following graphs correctly shows variation of coefficient of volume

expansion of copper as a function of temperature ?



Answer: C



Watch Video Solution

11. If α , β and γ coefficient of linear, superficial and volume expansion respectively, then

A. $\frac{\beta}{\alpha} = \frac{1}{2}$

B. $\frac{\beta}{\gamma} = \frac{2}{3}$

C. $\frac{\gamma}{\alpha} = \frac{3}{2}$

D. $\frac{\beta}{\alpha} = \frac{\gamma}{\beta}$

Answer: B



Watch Video Solution

12. Two spheres A and B are made of the same material and have the same radius. Sphere A is hollow and sphere B is solid. Both the spheres are heated to the same temperature. Which of the following is correct ?

- A. A expands more than B.
- B. A expands less than B.
- C. Both the spheres expand equally.
- D. Data is insufficient

Answer: C



13. The volume of a metal sphere increases by 0.24 % when its temperature is raised by $40^{\circ} C$. The coefficient of linear expansion (*per $^{\circ} C$*) of the metal is

A. $2 \times 10^5 .^{\circ} C^{-1}$

B. $6 \times 10^{-5} .^{\circ} C^{-1}$

C. $18 \times 10^{-5} .^{\circ} C^{-1}$

D. $1.2 \times 10^{-5} .^{\circ} C^{-1}$

Answer: A



Watch Video Solution

14. Calculate the stress developed inside a tooth cavity filled with copper when hot tea at temperature of $57^{\circ}C$ is drunk. You can take body (tooth) temperature to be $37^{\circ}C$ and $\alpha_{Cu} = 1.7 \times 10^{-5} / ^{\circ}C$ bulk modulus for copper $B_{Cu} = 140 \times 10^9 N/m^2$.

A. $1.43 \times 10^8 Nm^{-2}$

B. $4.13 \times 10^8 Nm^{-2}$

C. $2.12 \times 10^4 Nm^{-2}$

D. $3.12 \times 10^4 Nm^{-2}$


Answer: A




Watch Video Solution

15. Which of the following graph shows the variation of density of water increase in temperature ?


A.




B.



C.



D.



Answer: A



Watch Video Solution

Specific Heat Capacity

1. Which one of the following substances has highest specific heat capacity at room temperature and atmospheric pressure ?

A. Water

B. Ice

C. Aluminium

D. Mercury

Answer: A



Watch Video Solution

2. If specific heat of a substance is infinite, it means

A. heat is given out

B. heat is taken in

C. no change in temperature whether heat is taken in or given out

D. all of these

Answer: C



Watch Video Solution

3. Water is used as a coolant because

- A. it has lower density
- B. it has low specific heat.
- C. it has high specific heat.
- D. It is easily available.

Answer: C



Watch Video Solution

4. A 10 kW drilling machine is used to drill a bore in a small aluminium block of mass 8 kg. Find the rise in temperature of the block in 2.5 minutes, assuming 50% power is used up in heating the machine itself or lost to the surroundings.

(Specific heat of aluminium
 $= 0.91 J g^{-1} \cdot ^\circ C^{-1}$)

A. $100.^\circ C$

B. $103.^\circ C$

C. $150.^\circ C$

D. $155.^\circ C$

Answer: B



Watch Video Solution

5. A person weighing 50 kg takes in 1500 kcal diet per day. If this energy were to be used in heating the body of person without any losses, then the rise in his temperature is (specific heat of human body $= 0.83 \text{ cal g}^{-1} \text{ C}^{-1}$)

A. $30.^\circ C$

B. $48.^\circ C$

C. $40.16.^\circ C$

D. $36.14.^\circ C$

Answer: D



Watch Video Solution

Calorimetry

1. 10 g of ice of $0^{\circ}C$ is mixed with 100g of water at $50^{\circ}C$ in a calorimeter. The final temperature of the mixture is [Specific heat of water $= 1\text{calg}^{-1}\cdot^{\circ}C^{-1}$, latent of fusion of ice $= 80\text{calg}^{-1}$]

A. $31.2^{\circ}C$

B. $32.8^{\circ}C$

C. $36.7^{\circ}C$

D. $38.2^{\circ}C$

Answer: D



2. When 1.5 kg of ice at 0°C mixed with 2 kg of water at 70°C in a container, the resulting temperature is 5°C the heat of fusion of ice is (

$$s_{\text{water}} = 4186\text{ jkg}^{-1}\text{K}^{-1})$$

A. $1.42 \times 10^5\text{ jkg}^{-1}$

B. $2.42 \times 10^5\text{ jkg}^{-1}$

C. $3.42 \times 10^5\text{ jkg}^{-1}$

D. $4.42 \times 10^5\text{ jkg}^{-1}$

Answer: C



Watch Video Solution

3. The temperature of equal masses of three different liquids A, B and C are $12^{\circ}C$, $19^{\circ}C$ and $28^{\circ}C$ respectively. The temperature when A and B are mixed is $16^{\circ}C$ and when B and C are mixed it is $23^{\circ}C$. What should be the temperature when A and C are mixed?

A. $18.2^{\circ}C$

B. $22.^\circ C$

C. $20.3.^\circ C$

D. $24.2.^\circ C$

Answer: C



Watch Video Solution

4. An ice cube of mass 0.1 kg at $0^\circ C$ is placed in an isolated container which is at $227^\circ C$. The specific heat s of the container varies with temperature T according to the empirical

relation

$$s = A + BT,$$

where

$$A = 100 \text{ cal / kg. K} \text{ and } B = 2 \times 10^{-2} \text{ cal / kg. K}^2$$

. If the final temperature of the container is 27°C , determine the mass of the container.

(Latent heat of fusion for water = $8 \times 10^4 \text{ cal / kg}$, specific heat of water = 10^3 cal / kg. K).

A. 0.495 kg

B. 0.595 kg

C. 0.695 kg

D. 0.795 kg

Answer: A



Watch Video Solution

Change Of State

1. The change from solid state to vapour state without passing through the liquid state is called.....

A. Fusion

B. Regulation

C. vaporation

D. sublimation

Answer: D



Watch Video Solution

2. Match the following.

Column I		Column II	
(A)	Conversion of a liquid into solid is	(p)	Regelation
(B)	Conversion of a liquid into vapour is	(q)	Sublimation
(C)	Conversion of solid into vapour directly	(r)	Fusion
(D)	Melting of ice caused by pressure is	(s)	Vaporisation

A. $A - r, B - q, C - p, D - s$

B. $A - r, B - s, C - q, D - p$

C. $A - q, B - p, C - s, D - r$

D. $A - p, B - q, C - r, D - s$

Answer: B



Watch Video Solution

3. Which of the following statement is correct ?

A. The triple point of water is $253.16K$.

B. Burns from steam are less severe than those from boiling water.

C. Ethyl alcohol expands less than mercury for the same rise in temperature.

D. When fully inflated ballon is immersed in cold water, it will contract.


Answer: D



View Text Solution

4. In the phase diagram shows, the point Q corresponds to the triple point of water. The region I, II and III respectively correspond to

phases




- A. liquid, solid, vapour
- B. solid, liquid, vapour
- C. liquid, vapour, solid
- D. solid, vapour, liquid

Answer: A



View Text Solution

5. Refer to the plot of temperature versus time showing the changes in the state of ice on heating (not to scale).



Which of the following is correct ?

A. AB represents ice and water are not in the thermal equilibrium

B. At B water starts boiling.

C. At C all the water gets converted into steam.




D. CD represents water and steam in equilibrium at boiling point.


Answer: D



Watch Video Solution

6. A block of ice at -10°C is slowly heated and converted to steam at 100°C . Which of the following curves represents the phenomenon qualitatively?





Answer: A



Watch Video Solution

7. Two absolute scales A and B have triple points of water defined to be $200A$ and $350B$.

What is the relation between T_A and T_B ?

A. $T_A = 4/7T_B$

B. $T_B = 4/7T_A$

$$C. T_A = 2/7T_B$$

$$D. T_B = 2/7T_A$$

Answer: A



Watch Video Solution

8. The triple point of carbon dioxide is 216.55 K
the corresponding temperature on the celsius
and Fahrenheit scale respectively are

A. $56.45^{\circ} C, - 69.61^{\circ} F$

B. $- 56.45^{\circ} C, 69.61^{\circ} F$

C. $54.45^{\circ} C$, $69.61^{\circ} F$

D. -5445° , $-6961^{\circ} F$

Answer: D



Watch Video Solution

9. The latent heat of vaporization of a substance is always

A. greater than its latent heat of fusion

B. greater than its latent heat of sublimation

C. equal to its latent heat of sublimation

D. less than its latent heat of fusion

Answer: A



Watch Video Solution

10. The sprinkling of water slightly reduces the temperature of a closed room because

A. temperature of water of less than that of the room

B. specific heat of water is high

C. water has large latent heat of vapourisation

D. water is a bad conductor of heat.

Answer: C



Watch Video Solution

11. If 10 g of ice is added to 40 g of water at $15^{\circ}C$, then the temperature of the mixture is (specific heat of water = $4.2 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$,

Latent heat of fusion of ice = $3.36 \times 10^5 \text{ jkg}^{-1}$

)

A. 15°C

B. 12°C

C. 10^2C

D. 0°C

Answer: D



Watch Video Solution

12. If a ball of 80 kg mass hits an ice cube and temperature of ball is $100.^{\circ} C$, then how much ice converted into water ? (Specific heat of ball is $0.2calg^{-1}$, Latent heat of ice = $80calg^{-1}$)

A. 20 g

B. 200g

C. $2 \times 10^3 kg$

D. $2 \times 10^4 g$

Answer: D



Watch Video Solution

13. Rays from the sun are focussed by a lens of diameter 5 cm on to a block of ice and 10 g of ice is melted in 20 min. Therefore the heat from the sun reaching the earth per min per square centimetre is

(Latent heat of ice $L = 80 \text{ cal g}^{-1}$)

A. 2.04 cal

B. 0.51 cal

C. 4.08 cal

D. 3.02 cal

Answer: A



Watch Video Solution

Heat Transfer

1. For transmission of heat from one place to the other, medium is required in

A. conduction

B. convection

C. radiation

D. both (a) and (b)

Answer: D



Watch Video Solution

2. One end of a 0.25 m long metal bar is in steam and the other is in contact with ice . If 12 g of ice melts per minute, what is the thermal conductivity of the metal? Given cross-section of the bar $= 5 \times 10^{-4} m^2$ and latent heat of ice is $80 cal g^{-1}$

A. $20 \text{ cal s}^{-1} \text{ m}^{-1} \cdot ^\circ \text{ C}^{-1}$

B. $10 \text{ cal s}^{-1} \text{ m}^{-1} \cdot ^\circ \text{ C}^{-1}$

C. $40 \text{ cal s}^{-1} \text{ m}^{-1} \cdot ^\circ \text{ C}^{-1}$

D. $80 \text{ cal s}^{-1} \text{ m}^{-1} \cdot ^\circ \text{ C}^{-1}$

Answer: D



Watch Video Solution

3. A pan filled with hot food cools from 94° C to 86° C in 2 minutes when the room temperature is at 20° C . How long will it take to cool from

$71^{\circ}C$ to $69^{\circ}C$? Here cooling takes place according to Newton's law of cooling.

A. 12 s

B. 22 s

C. 32 s


D. 40 s

Answer: D



Watch Video Solution

4. Two bars of same length and same cross-sectional area but of different thermal conductivities K_1 and K_2 are joined end to end as shown in the figure. One end of the compound bar is at temperature T_1 and the opposite end at temperature T_2 (where $T_1 > T_2$). The temperature of the junction is



A.
$$\frac{K_1 T_1 + K_2 T_2}{K_1 + K_2}$$

B.
$$\frac{K_1 T_2 + K_2 T_1}{K_1 + K_2}$$

C. $\frac{K_1(T_1 + T_2)}{K_1 + K_2}$

D. $K_2 \frac{T_1 + T_2}{K_1}$

Answer: A



Watch Video Solution

5. In the question number of 45, the equivalent thermal conductivity of the compound bar is

A. $\frac{K_1 K_2}{K_1 + K_2}$

B. $\frac{2K_1 K_2}{K_1 + K_2}$

C. $\frac{K_1}{K_1 + K_2}$

D. $\frac{K_2}{K_1 + K_2}$

Answer: B



View Text Solution

6. Consider a compound slab consisting of two different material having equal thickness and thermal conductivities K and $2K$ respectively. The equivalent thermal conductivity of the slab is

A. $\frac{2}{3}K$

B. $\sqrt{2}K$

C. $3K$

D. $\frac{4}{3}K$

Answer: D



Watch Video Solution

7. Two rods of equal length and diameter have thermal conductivity 3 and 4 units respectively.

If they are joined in series, the thermal

conductivity of the combination in the given units would be

A. 3.43 units

B. 4.43 units

C. 5.43 units


D. 2.43 units

Answer: A



Watch Video Solution

8. Three metal rods of the same material and identical in all respect are joined as shown in the figure. The temperatures at the ends are maintained as indicated. Assuming no loss of heat from the curved surface of the rods, the temperature at the junction X would be



A. 45°C

B. $60^{\circ} C$

C. $30^{\circ} C$

D. $20^{\circ} C$

Answer: B



Watch Video Solution

9. Three very large plates of same area are kept parallel and close to each other. They are considered as ideal black surfaces and have high thermal conductivity. The first and third plates

are maintained at temperatures $2T$ and $3T$ respectively. The temperature of the middle (i.e., second) plate under steady state condition is

A. $\left(\frac{65}{2}\right)^{1/4} T$

B. $\left(\frac{97}{4}\right)^{1/4} T$

C. $\left(\frac{97}{2}\right)^{1/4} T$

D. $(97)^{1/4} T$

Answer: C



Watch Video Solution

10. A cubical ice box of side 50 cm has a thickness of 5.0 cm. If 5 kg of ice is put in the box, estimate the amount of ice remaining after 4 hours. The outside temperature is 40°C and coefficient of thermal conductivity of the material of the box = $0.01\text{Js}^{-1}\text{m}^{-1}\text{.}^{\circ}\text{C}^{-1}$.
Heat of fusion of ice = 335Jg^{-1} .

A. 3.7kg

B. 3.9kg

C. 4.7kg

D. 4.9kg

Answer: C



Watch Video Solution

11. A wall has two layers A and B, each made of different material. Both the layers have the same thickness. The thermal conductivity of the material of A is twice that of B . Under thermal equilibrium, the temperature difference across the wall is $36^{\circ}C$. The temperature difference across the layer A is

A. $12^{\circ}C$

B. $18^{\circ}C$

C. $6^{\circ}C$

D. $24^{\circ}C$

Answer: A



Watch Video Solution

12. Mud houses are cooler in summer and warmer in winter because

A. mud is a good conductor of heat

B. mud is a superconductor of heat

C. mud is a bad conductor of heat

D. none of these


Answer: C



Watch Video Solution

13. Which of the following is the $\nu_m = T$ graph for a perfectly black body (ν_m = maximum frequency

of radiation)



A. A

B. B

C. C

D. D

Answer: C



Watch Video Solution

14. The equatorial and polar regions of the earth receive unequal solar heat. The convection current arising due to this is called

A. land breeze

B. sea breeze

C. trade wind

D. tornado

Answer: C



View Text Solution

15. In which of the following process, convection does not take place primarily

A. Sea and land breeze

B. Trade wind

C. Boiling of water

D. Warming of glass of bulb due to filament

Answer: D



Watch Video Solution

16. Wien's displacement law expresses relation between

A. colour of light and temperature

B. wavelength and temperature

C. radiation energy and wavelength

D. wavelength corresponding to maximum energy and temperature

Answer: D



Watch Video Solution

17. If λ_m denotes

A. $\lambda_m \propto T$

B. $\lambda_m \propto T^{-1}$

C. $\lambda_m \propto T$

D. λ_m is independent on T

Answer: B



Watch Video Solution

18. A black body has maximum wavelength λ_m at temperature $2000K$. Its corresponding wavelength at temperature 3000 will be

A. $\frac{3}{2}\lambda_m$

B. $\frac{2}{3}\lambda_m$

C. $\frac{16}{81}\lambda_m$

D. $\frac{81}{16}\lambda_m$

Answer: B



Watch Video Solution

19. The thermal radiation from a hot body travels with a velocity of

A. $330ms^{-1}$

B. $2 \times 10^8ms^{-1}$

C. $1200ms^{-1}$

D. $3 \times 10^8ms^{-1}$

Answer: D



Watch Video Solution

20. Experimental investigations show that the intensity of solar radiation is maximum for a wavelength $480nm$ in the visible region. Estimate the surface temperature of sun. (Given Wien's constant $b = 2.88 \times 10^{-3}mK$).

A. $4000K$

B. $6000K$

C. $8000K$

D. 10^6K

Answer: B



21. The wavelength of maximum intensity of radiation emitted by a star is 289.8 nm . The radiation intensity for the star is : (Stefan's constant $5.67 \times 10^{-8} W m^{-2} K^{-4}$, constant $b = 2898 \mu m K$)-

A. $5.67 \times 10^8 W m^{-2}$

B. $5.67 \times 10^{12} W m^{-2}$

C. $10.67 \times 10^7 W m^{-2}$

D. $10.67 \times 10^{14} W m^{-2}$

Answer: A



Watch Video Solution

22. The temperature of a radiation body increases by 30%. Then the increase in the amount of radiation is?

A. 185 %

B. 285 %

C. 325 %

D. 130 %

Answer: A



Watch Video Solution

23. If the temperature of hot black body is raised by 5%, rate of heat energy radiated would be increased by how much percentage ?

A. 12 %

B. 22 %

C. 32 %

D. 42 %

Answer: B



Watch Video Solution

24. Two spheres of same material have radius 1 m and 4 m and temperature 4000 K and 2000 K respectively. The energy radiated per second by the first sphere is

- A. greater than that by the second
- B. less than that by the second
- C. equal in both cases

D. the information is incomplete to draw any conclusion

Answer: C



Watch Video Solution

25. If the temperature of the Sun were to increase from T to $2T$ and its radius from R to $2R$. The ratio of power radiated by it would become

A. 64 times

B. 16 times

C. 32 times

D. 4 times

Answer: A



Watch Video Solution

26. The rate of cooling at $600K$, If surrounding temperature is $300K$, is H . The rate of cooling at $900K$ is

A. $\frac{16}{3}H$

B. $2H$

C. $3H$

D. $\frac{2}{3}H$

Answer: A




Watch Video Solution

Higher Order Thinking Skills

1. The temperature of the two outer surfaces of a composite slab, consisting of two materials

having coefficients of thermal conductivity K and $2K$ and thickness x and $4x$ respectively. Temperatures on the opposite faces of composite slab are T_1 and T_2 where $T_2 > T_1$, as shown in fig. what is the rate of flow of heat through the slab in a steady state?



A. 1

B. $\frac{1}{2}$

C. $\frac{2}{3}$

D. $\frac{1}{3}$

Answer: D



Watch Video Solution

2. A wall of dimensions 2.00 m by 3.50m has a single-pane window of dimensions 0.75m by 1.20m . If the inside temperature is $20^{\circ}C$ and the outside temperature is $-10^{\circ}C$, effective thermal resistance of the opaque wall and

window are $2.10m^2 KW^{-1}$ and $0.21m^2 KW^{-1}$ respectively. The heat flow through the entire wall will be .

A. $215W$

B. $205W$

C. $175W$


D. $110W$

Answer: A



View Text Solution

3. The figure show a cross-section of a double glass unit of a window on a vertical wall. A graph of the temperature at different points within the unit is show next to it. The temperature difference across the unit is $13K$. It has a cross-sectional area of $1.3m^2$ and the rate of heat flow through it is $65W$. Then the correct statement is (Glass has a thermal conductivity of $1Wm^{-1}K^{-1}$)



A. The unit is in steady state and in thermal equilibrium

B. The unit is in steady state but not in thermal equilibrium.

C. The unit is not in steady state but is in thermal

D. The unit neither in steady state nor in thermal equilibrium

Answer: B



View Text Solution

4. In question number 3, the thermal conductivity (in $Wm^{-1}K^{-1}$) if air is

A. $\frac{1}{10}$

B. $\frac{1}{12}$

C. $\frac{1}{14}$

D. $\frac{9}{130}$

Answer: C



View Text Solution

5. A lake surface is exposed to an atmosphere where the temperature is $< 0^\circ C$. If the thickness of the ice layer formed on the surface grows from 2cm to 4cm in 1 hour. The atmospheric temperature, T_a will be- (Thermal conductivity of ice $K = 4 \times 10^{-3} \text{cal/cm/s/}^\circ C$, density of ice $= 0.9\text{gm/ice}$. Latent heat of fusion of ice $= 80\text{cal/m}$. Neglect the change of density during the state change. Assume that the water below the ice has 0° temperature every where)

A. $-20.^\circ C$

B. $0.^\circ C$

C. $-30.^\circ C$

D. $15.^\circ C$

Answer: C



Watch Video Solution

6. Two chunks of metal with heat capacities C_1 and C_2 , are interconnected by a rod length l and cross-sectional area S and fairly low heat conductivity K . The whole system is thermally

insulated from the environment. At a moment $t = 0$ the temperature difference between the two chunks of metal equals $(\Delta T)_0$. Assuming the heat capacity of the rod to be negligible, find the temperature difference between the chunks as a function of time.

A. $T_0 e^{\frac{-KA(C_1+C_2)t}{C_1C_2}}$

B. $T_0 e^{\frac{-KA(C_1+C_2)}{C_1C_2}}$

C. $T_0 e^{\frac{KA(C_1+C_2)t}{C_1C_2}}$

D. $T_0 e^{\frac{KA(C_1+C_2)t^2}{C_1C_2}}$

Answer: A



Watch Video Solution

7. A double pan window used for insulating a room thermally from outside consists of two glass sheets each of area 1 m^2 and thickness 0.01 m separated by 0.05 m thick stagnant air space. In the steady state, the room-glass interface and the glass-outdoor interface are at constant temperatures of 27°C and 0°C respectively. The thermal conductivity of glass is $0.8 \text{ W m}^{-1} \text{ K}^{-1}$ and of air $0.08 \text{ W m}^{-1} \text{ K}^{-1}$. Answer the following questions.

(a) Calculate the temperature of the inner glass-air interface.

(b) Calculate the temperature of the outer glass-air interface.

(c) Calculate the rate of flow of heat through the window pane.

A. $41.5W$

B. $31.5W$

C. $21.5W$

D. $11.5W$

Answer: A



[Watch Video Solution](#)

8. In question number 7, the temperature of other interfaces (in $^{\circ}C$) is

A. 26.5, 0.5

B. 27.5, 1

C. 28.5, 2

D. 29.5, 3

Answer: A



[View Text Solution](#)

Ncert Exemplar

1. A bimetallic strip made of aluminium and steel ($\alpha_{Al} > \alpha_{steel}$) on heating the strip will

A. remain straight

B. get twisted

C. will bend with aluminium on concave side

D. will bend with steel on concave side.

Answer: D



Watch Video Solution

2. A uniform metallic rod rotates about its perpendicular bisector with constant angular speed. If it is heated uniformly to raise its temperature slightly, then

- A. its speed of rotation increases
- B. its speed of rotation decreases
- C. its speed of rotation remains same
- D. its speed increases because its moment of inertia increases


Answer: B



Watch Video Solution

3. The graph between two temperature scales A and B is shown in Fig. Between upper fixed point and lower fixed point there are 150 equal divisions on scales A and 100 on scale B . The relation between the temperature in two scales

is given by_



A. $\frac{T_A - 180}{100} = \frac{T_B}{150}$

B. $\frac{T_A - 30}{150} = \frac{T_B}{100}$

C. $\frac{T_B - 180}{150} = \frac{T_A}{100}$

D. $\frac{T_B - 40}{100} = \frac{T_A}{180}$

Answer: B



Watch Video Solution

4. An aluminium sphere is dipped into water.

Which of the following is true ?

A. Buoyancy will be less in water at $0^{\circ}C$ than that in water at $4^{\circ}C$.

B. Buoyancy will be more in water at $0^{\circ}C$ than that in water at $4^{\circ}C$.

C. Buoyancy in water at $0^{\circ}C$ will be same as that in water at $4^{\circ}C$.

D. Buoyancy may be more or less in water at $4^{\circ}C$ depending on the radius of the sphere.

Answer: A



Watch Video Solution

5. As the temperature is increased, the period of a pendulum

A. Increases as its effective length increases
even through its centre of mass still
remains at the centre of the bob.

B. decreases as its effective length increases
even though its centre of mass still
remains at the centre of the bob.

C. Increases as its effective length increases
due to shifting of centre of mass below
the centre of the bob.

D. decreases as its effective length remains same but the centre of mass shifts above the centre of the bob.

Answer: A



Watch Video Solution

6. Heat is associated with

A. kinetic energy of random motion of molecules

B. kinetic energy of orderly motion of molecules

C. total kinetic energy of random and orderly motion of molecules

D. kinetic energy of random motion in some cases and kinetic energy orderly motion in other.

Answer: A



Watch Video Solution

7. The radius of metal sphere at room temperature T is R and the coefficient of linear expansion of the metal is α . The sphere is heated a little by a temperature T , so that new temperature is $T + \Delta T$. The increase in volume of sphere is approximately

A. $2\pi R\alpha\Delta T$

B. $\pi R^2\alpha\Delta T$

C. $4\pi R^3\alpha\Delta T / 3$

D. $4\pi R^3\alpha\Delta T$

Answer: D



Watch Video Solution

8. A sphere, a cube and a thin circular plate, all of same material and same mass are initially heated to same high temperature.

A. Plate will cool fastest and cube the slowest.

B. Sphere will cool fastest and cube the slowest.

C. Plate will cool fastest and sphere the slowest.

D. Cube will cool fastest and plate the slowest.

Answer: C



Watch Video Solution

Assertion And Reason

1. Assertion : A change in the temperature of a body cause change in dimentions.

Reason : The dimensions of a body decrease due to the increase in its temperature.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true and reason is not the correct explanation of assertion.

C. If assertion is true and reason is false.

D. If both assertion and reason are false.

Answer: C



Watch Video Solution

2. The Kelvin scale temperature is 0 K. What is the corresponding Celsius scale temperature ?

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true and reason is not the correct explanation of

assertion.

C. If assertion is true and reason is false.

D. If both assertion and reason are false.

Answer: B



Watch Video Solution

3. Assertion : The density of water remains constant as it is cooled from room temperature until its temperature reached $4^{\circ}C$.

Reason : Below $4^{\circ}C$, the density increases.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true and reason is not the correct explanation of assertion.

C. If assertion is true and reason is false.

D. If both assertion and reason are false.

Answer: D



Watch Video Solution

4. Water is used in car radiators as coolant because

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true and reason is not the correct explanation of assertion.

C. If assertion is true and reason is false.

D. If both assertion and reason are false.

Answer: A



Watch Video Solution

5. Assertion : The specific heat capacity of a given solid can be determined by using the principle of calorimetry

Reason : Heat gained is equal to the heat lost.

A. If both assertion and reason are true and reason is the correct explanation of

assertion.

B. If both assertion and reason are true and reason is not the correct explanation of assertion.

C. If assertion is true and reason is false.

D. If both assertion and reason are false.

Answer: A



Watch Video Solution

6. Assertion : In change of state from solid to liquid the temperature decreases until the entire amount of the solid substance melts.

Reason : The phenomenon of refreezing is called melting

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true and reason is not the correct explanation of

assertion.

C. If assertion is true and reason is false.

D. If both assertion and reason are false.

Answer: D



Watch Video Solution

7. Assertion : Cooking food is difficult on hills.

Reason : The boiling point decreases with increase in pressure

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true and reason is not the correct explanation of assertion.

C. If assertion is true and reason is false.

D. If both assertion and reason are false.

Answer: C



Watch Video Solution

8. Assertion : In the human body the heart acts as the pump that circulates blood through differences parts of the body, transferring heat by force convection.

Reason : In forced convection , material is force to move by a pump or by some other physics means.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true and reason is not the correct explanation of assertion.

C. If assertion is true and reason is false.

D. If both assertion and reason are false.

Answer: a



Watch Video Solution

9. Assertion : All bodies emit radiant energy whether they are solid, liquid or gases.

Reason : Black bodies absorb and emit radiant energy better than bodies of lighter colours.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true and reason is not the correct explanation of assertion.

C. If assertion is true and reason is false.

D. If both assertion and reason are false.

Answer: B



Watch Video Solution

10. Assertion: According to Newton's law of cooling, the rate of loss of heat, $-dQ/dt$ of the body is directly proportional to the difference of temperature.

Reason :This law holds for all type of temperature differences.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true and reason is not the correct explanation of assertion.

C. If assertion is true and reason is false.

D. If both assertion and reason are false.

Answer: c



Watch Video Solution

