



PHYSICS

BOOKS - NCERT FINGERTIPS PHYSICS (HINGLISH)

ATOMS

Alpha Particle Scattering And Rutherford Model

1. The first method of atom in 1998 was proposed by

A. Ernst Rutherford

B. Albert Einstein

C. J.J. Thomson

D. Niels Bohr

Answer: C



Watch Video Solution

2. In Geiger-Marsden scattering experiment, the trajectory traced by α -particle depends on

A. number of collision

B. number of scattered α -particles

C. impact parameter

D. none of these

Answer: C



Watch Video Solution

3. In a geiger - marsden experiment. Find the distance of closest approach to the nucleus of a $7.7 \text{ me v } \alpha$ - particle before it comes

momentarily to rest and reverses its direction.

(z for gold nucleus = 79) .

A. 30 fm

B. 20 fm

C. 40 fm

D. 10 fm

Answer: A



Watch Video Solution

4. In the Geiger-Marsden scattering experiment, the number of scattered particles detected are maximum and minimum at the scattering angles respectively at

A. 0° and 180°

B. 180° and 0°

C. 90° and 180°

D. 45° and 90°

Answer: A



Watch Video Solution

5. In the geiger -marsden scattering experiment, in case of head- on collision the impact parameter should be

A. maximum

B. minimum


C. infinite

D. zero


Answer: B




6. The graph of the total number of α -particles scattered at different angles in a given interval of time for α - particles scattering in the geiger- marsden experiment is given by




A.



B.



C.



D.

Answer: A



View Text Solution

7. Rutherford experiment suggested that the size of the nucleus is about

A. $10^{-14}m$ to $10^{-12}m$

B. 10^{-15m} to $10^{-13}m$

C. 10^{-15m} to $10^{-14}m$

D. 10^{-15m} to $10^{-12}m$

Answer: C



Watch Video Solution

8. In an atom the ratio of radius of orbit of electron to the radius of nucleus is

A. 10^3

B. 10^4

C. 10^5

D. 10^6

Answer: C



Watch Video Solution

9. The relation between the orbit radius and the electron velocity for a dynamically stable orbit in a hydrogen atom is (where, all notations have their usual meanings)

$$\text{A. } v = \sqrt{\frac{4\pi\epsilon_0}{me^2r}}$$

$$\text{B. } r = \sqrt{\frac{e^2}{4\pi\epsilon_0v}}$$

$$\text{C. } v = \sqrt{\frac{e^2}{4\pi\epsilon_0mr}}$$

$$\text{D. } r = \sqrt{\frac{ve^2}{4\pi\epsilon_0m}}$$

Answer: C



Watch Video Solution

10. The relationship between kinetic energy (K) and potential energy (U) of electron moving in

a orbit around the nucleus is

A. $U = -K$

B. $U = -2K$

C. $U = -3K$

D. $U = -\frac{1}{2}K$

Answer: B



Watch Video Solution

11. The volume occupied by an atom is greater than the volume of the nucleus by factor of about

A. 10^1

B. 10^5

C. 10^{10}

D. 10^{15}

Answer: D



Watch Video Solution

12. Consider aiming a beam of free electrons towards free atoms. When they scatter, an electron and a proton cannot combine to produce a H-atom,

A. energy conservation

B. simultaneously releasing energy in the form of radiation

C. momentum conservation

D. angular momentum conservation

Answer: A



Watch Video Solution

13. In an experiment on α -particle scattering, α -particles are directed towards a gold foil and detectors are placed in position P, Q and R . What is the distribution of α -particles as recorded at P, Q and R?



A.	P	Q	R
	all	none	none

- | | | | |
|----|--------------|-------------|--------------|
| B. | P
none | Q
none | R
all |
| C. | P
a few | Q
some | R
most |
| D. | P
most | Q
some | R
a few |

Answer: C



View Text Solution

Atomic Spectra

1. The first spectral series was discovered by

A. balmer

B. lyman

C. paschen

D. Pfund

Answer: A



Watch Video Solution

2. which of the following spectral series falls within the visible range of electromagnetic radiation ?

A. lyman series

B. balmer series

C. paschen series

D. pfund series

Answer: B



Watch Video Solution

3. What is the shortest wavelength present in the Paschen series of spectral lines?

A. 720 nm

B. 790 nm

C. 800 nm

D. 820 nm

Answer: D



Watch Video Solution

4. The shortest wavelength in the balmer series is ($R = 1.097 \times 10^7 m^{-1}$)

A. 200 nm

B. 256.8 nm

C. 300 nm

D. 364.6 nm

Answer: D



Watch Video Solution

5. the wavelength limit present in the pfund

series is $(R = 1 - 097 \times 10^7 m^{-1})$

A. 1572 nm

B. 1898 nm

C. 2278 nm

D. 2535 nm

Answer: C



Watch Video Solution

6. when an electron jumps from the fourth orbit to the second orbit, one gets the

A. second line of paschen series

B. second line of balmer serie

C. first line pfund series

D. second line of lyman series

Answer: B



Watch Video Solution

7. The balmer series for the H-atom can be observed

- A. if we measure the frequencies of light emitted when an excited atom falls to the ground state
- B. if we measure the frequencies of light emitted due to transitions between excited states and the first excited state
- C. in any transition in a H-atom
- D. none of these

Answer: B



Watch Video Solution

8. when an atomic gas or vapour is excited at low pressure, by passing an electric current through it then

- A. emission spectrum is observed
- B. absorption spectrum is observed
- C. band spectrum is observed
- D. both a and c

Answer: A



View Text Solution

9. In balmer series of emission spectrum of hydrogen, first four lines with different wavelength H_α , H_β , H_γ and H_δ are obtained. Which line has maximum frequency out of these ?

A. 1. H_α

B. 2. H_β

C. 3. H_γ

D. 4. H_δ

Answer: D



Watch Video Solution

10. hydrogen atom emits light when it changes from $n = 5$ energy level to $n = 2$ energy level.

Which colour of light would the atom emit ?

A. red

B. yellow

C. green

D. violet.

Answer: D



Watch Video Solution

11. the wavelength of the first line of lyman series is 1215\AA , the wavelength of first line of balmer series will be

A. 4545\AA

B. 5295\AA

C. 6561\AA

D. 6750\AA

Answer: C



Watch Video Solution

12. the wavelength of radiation emitted is λ_0 when an electron jumps. From the third to second orbit of hydrogen atom. For the electron jumping from the fourth to the second orbit of the hydrogen atom, the wavelength of radiation emitted will be

A. $(16/25)\lambda_0$

B. $(20 / 27)\lambda_0$

C. $(27 / 20)\lambda_0$

D. $(25 / 16)\lambda_0$

Answer: B



Watch Video Solution

13. the wavelength of the first line of lyman series for hydrogen atom is equal to that of the second line of balmer series for a

hydrogen like ion. The atomic number Z of hydrogen like ion is

A. 3

B. 4

C. 1

D. 2

Answer: D



Watch Video Solution

14. According to bohr's theory, the wave number of last line of balmer series is

$$(R = 1.1 \times 10^7 m^{-1})$$

A. $5.5 \times 10^5 m^{-1}$

B. $4.4 \times 10^7 m^{-1}$

C. $2.75 \times 10^6 m^{-1}$

D. $2.75 \times 10^8 m^{-1}$

Answer: C



Watch Video Solution

15. The first line of the Lyman series in a hydrogen spectrum has a wavelength of 1210 \AA . The corresponding line of a hydrogen like atom of $Z = 11$ is equal to

A. 4000 \AA

B. 100 \AA

C. 40 \AA

D. 10 \AA

Answer: D



Watch Video Solution

16. What is the ratio of the shortest wavelength of the balmer to the shortest of the lyman series ?

A. 4 : 1

B. 4 : 3

C. 4 : 9

D. 5 : 9

Answer: A



17. If the wavelength of the first line of the Balmer series of hydrogen is 6561\AA , the wavelength of the second line of the series should be

A. 13122\AA

B. 3280\AA

C. 4860\AA

D. 2187\AA

Answer: C



Watch Video Solution

Emission series		Make transitions from higher levels to following levels	
A	Lyman series	P	$n = 1$
B	Paschen series	Q	$n = 2$
C	Balmer series	R	$n = 3$
D	Brackett series	S	$n = 4$
		T	$n = 5$

18.

A. A-P,B-R,C-Q,D-S

B. A-P,B-Q,C-R,D-T

C. A-Q,B-R,C-S,D-T

D. A-T,B-S,C-R,D-Q

Answer: A



View Text Solution

19. If ν_1 is the frequency of the series limit of lyman series, ν_2 is the frequency of the first line of lyman series and ν_3 is the frequency of the series limit of the balmer series, then

A. $\nu_1 - \nu_2 = \nu_3$

B. $v_1 = v_2 - v_3$

C. $\frac{1}{v_2} = \frac{1}{v_1} + \frac{1}{v_3}$

D. $\frac{1}{v_1} = \frac{1}{v_2} + \frac{1}{v_3}$

Answer: A



Watch Video Solution

Bohr Model Of Hydrogen Atom

1. which of the following postulates of the Bohr model led to the quantization of energy

of the hydrogen atom ?

A. The electron goes around the nucleus in circular orbits.

B. The angular momentum of the electron can only be an integral multiple of $h / 2\pi$.

C. The magnitude of the linear momentum of the electron is quantized.

D. Quantization of energy is itself a postulate of the Bohr model.

Answer: B



Watch Video Solution

2. The Bohr model of atoms

A. assumes that the angular momentum of electrons is quantized .

B. uses Einstein's photoelectric equation.

C. predicts continuous emission spectra for atoms.

D. predicts the same emission spectra for all types of atoms.

Answer: A



Watch Video Solution

3. The angular speed of the electron in the n^{th} Bohr orbit of the hydrogen atom is proportional to

A. directly proportional to n

B. inversely proportional to \sqrt{n}

C. inversely proportional to n^2

D. inversely proportional to n^3

Answer: D



Watch Video Solution

4. The electric current I created by the electron in the ground state of H atom bohr model in terms of bohr radius (a_0) and velocity of electron in first orbit v_0 is

A. $\frac{ev_0}{2\pi a_0}$

B. $\frac{2\pi a}{ev_0}$

C. $\frac{2\pi a}{v_0}$

D. $\frac{v_0}{2\pi a}$

Answer: A



Watch Video Solution

5. Total energy of electron in n th stationary orbit of hydrogen atom is

A. $\frac{e^2}{4\pi\epsilon_0 r}$

B. $\frac{-e^2}{4\pi\epsilon_0 r}$

C. $\frac{-e^2}{8\pi\epsilon_0 r}$

D. $\frac{e^2}{8\pi\epsilon_0 r}$

Answer: C



Watch Video Solution

6. According to Bohr's theory the moment of momentum of an electron revolving in second orbit of hydrogen atom will be

A. $\frac{h}{\pi}$

B. $2\pi h$

C. $\frac{2h}{\pi}$

D. $\frac{\pi}{h}$

Answer: A



Watch Video Solution

7. According to second postulate of bohr model, the angular momentum (L_n) of n^{th} possible orbit of hydrogen atom is given by

A. $\frac{h}{2\pi n}$

B. $\frac{nh}{2\pi}$

C. $\frac{2\pi n}{h}$

D. $\frac{2\pi}{nh}$

Answer: B



Watch Video Solution

8. Which of the following statements is true of hydrogen atom ?

A. Angular moment $\propto \frac{1}{n}$

B. Linear moment $\propto \frac{1}{n}$

C. Radius $\propto \frac{1}{n}$

D. Energy $\propto \frac{1}{n}$

Answer: B



Watch Video Solution

9. Total energy of electron in n th stationary orbit of hydrogen atom is

A. $\frac{-13.6}{n} eV$

B. $\frac{-13.6}{n^2} eV$

C. $\frac{-136}{n} eV$

D. $\frac{-136}{n^2} eV$

Answer: B



Watch Video Solution

10. The radius of n^{th} orbit r_n in the terms of Bohr radius (a_0) for a hydrogen atom is given by the relation

A. na_0

B. $\sqrt{na_0}$

C. n^2a_0

D. n^3a_0

Answer: C



Watch Video Solution

11. In Bohr model of the hydrogen atom, the lowest orbit corresponds to

A. infinite energy

B. maximum energy

C. minimum energy

D. zero energy

Answer: C



Watch Video Solution

12. Define ionisation energy. What is the value for a hydrogen atom?

A. $3.4E\text{v}$

B. $10.4E\text{v}$

C. $12.09E\text{v}$

D. $13.6E\text{v}$

Answer: D



Watch Video Solution

13. energy is absorbed in the hydrogen atom giving absorption spectra when transition takes place from

A. $n = 1 \rightarrow n'$ where ' n' ' > 1

B. $n = 2 \rightarrow 1$

C. $n' \rightarrow n$

D. $n \rightarrow n' = \infty$

Answer: A



View Text Solution

14. If n is the orbit number of the electron in a hydrogen atom, the correct statement among the following is

A. electron energy increases as n increases.

B. hydrogen emits infrared rays for the

electron transition from $n = \infty$ to

$n = 1$.

C. electron energy is zero for $n = 1$.

D. electron energy varies as n^2 .

Answer: A



View Text Solution

15. Which of the following is not correct about bohr model of the hydrogen atom ?

A. 1. An electron in an atom could revolve in certain stable orbits without the emission of radiant energy.

B. 2. Electron revolves around the nucleus only in those orbits for which angular momentum $L_n = \frac{nh}{2\pi}$.

C. 3. when electron make a transition from one of its stable orbit to lower orbit then a photon emitted with energy

$$h\nu = E_f - E_i.$$

D. 4. Bohr model is applicable to all atoms.

Answer: D



Watch Video Solution

16. In which of the following systems will the radius of the first orbit ($n = 1$) be minimum ?

A. doubly ionized lithium

B. singly ionized helium

C. deuterium atom

D. hydrogen atom

Answer: A



Watch Video Solution

17. when an electron falls from a higher energy to a lower energy level the difference in the energies appears in the form of

A. electromagnetic radiation only

B. thermal radiation only

C. both electromagnetic and thermal radiations

D. none of these

Answer: A



Watch Video Solution

18. Let $E = \frac{-1me^4}{8\varepsilon_0^2 n^2 h^2}$ be the energy of the n^{th} level of H-atom state and radiation of frequency $(E_2 - E_1) / h$ falls on it ,

A. it will not be absorbed at all

B. some of atoms will move to the first excited state

C. all atoms will be excited to the $n = 2$ state

D. all atoms will make a transition to the

$$n = 3 \text{ state}$$

Answer: B



Watch Video Solution

19. An ionized H-molecules consists of an electron and two protons. The protons are separated by a small distance of the order of angstrom. In the ground state,

A. the electron would not move in circular orbits

B. the energy would be $(2)^4$ times that of a H-atom

C. the molecule will soon decay in to a proton and a H-atom

D. none of these

Answer: A



Watch Video Solution

20. From quantisation of angular momentum one gets for hydrogen atom, the radius of the

$$n^{\text{th}} \text{ orbit as } r_n = \left(\frac{n^2}{m_e} \right) \left(\frac{h}{2\pi} \right)^2 \left(\frac{4\pi^2 \epsilon_0}{e^2} \right)$$

For a hydrogen like atom of atomic number Z,

- A. the radius of the first orbit will be the same
- B. r_n will be greater for larger Z values
- C. r_n will be smaller for larger Z values
- D. none of these

Answer: C



View Text Solution

21. Bohr's basic idea of discrete energy levels in atoms and the process of emission of photons from the higher levels to lower levels was experimentally confirmed by experiments performed by

A. Michelson- morley

B. millikan

C. joule

D. franck and hertz

Answer: D



View Text Solution

22. Suppose an electron is attracted toward the origin by a force $\frac{k}{r}$ where k is a constant and r is the distance of the electron from the origin. By applying Bohr model to this system the radius of the n^{th} orbital of the electron is

found to be r_n and the kinetic energy of the electron to be T_n , Then which of the following is true ?

A. $T_n \propto \frac{1}{n^2}$

B. T_n is independent of n , $r_n \propto n$

C. $T_n \propto \frac{1}{n}$ and r_n

D. $T_n \propto \frac{1}{n}$ and $r_n \propto n^2$

Answer: B



Watch Video Solution

23. In Bohr model of the hydrogen atom, let R , v and E represent the radius of the orbit, speed of the electron and the total energy respectively. Which of the following quantities are directly proportional to the quantum number n ?

A. VR

B. RE

C. R/E

D. none of these

Answer: A



Watch Video Solution

24. suppose an electron is attracted towards the origin by a force k/r , where k is a constant and r is the distance of the electron from the origin. By applying bohr model to this system, the radius of n^{th} orbit of the electron is found to be r_n and

A. E

B. 2E

C. 3E

D. 4E

Answer: D



View Text Solution

25. Which of the transitions in hydrogen atom emits a photon of lowest frequency ($n =$ quantum number)?

A. $n = 2$ to $n = 1$

B. $n = 4$ to $n = 2$

C. $n = 4$ to $n = 1$

D. $n = 4$ to $n = 3$

Answer: D



Watch Video Solution

26. The transition from the state $n = 3$ to $n = 1$ in a hydrogen-like atom results in

ultraviolet radiation. Infrared radiation will be obtained in the transition from

A. $2 \rightarrow 1$

B. $3 \rightarrow 2$

C. $4 \rightarrow 2$

D. $4 \rightarrow 3$

Answer: D



Watch Video Solution

27. If 13.6eV energy is required to separate a hydrogen atom into a proton and an electron, then the orbital radius of electron in a hydrogen atom is

A. $5.3 \times 10^{-11}\text{m}$

B. $4.3 \times 10^{-11}\text{m}$

C. $6.3 \times 10^{-11}\text{m}$

D. $7.3 \times 10^{-11}\text{m}$

Answer: A



Watch Video Solution

28. In the question number 59, the value of velocity of the revolving electron is

A. $1.2 \times 10^6 \text{ms}^{-1}$

B. $2.2 \times 10^6 \text{ms}^{-1}$

C. $3.2 \times 10^6 \text{ms}^{-1}$

D. $4.2 \times 10^6 \text{ms}^{-1}$

Answer: B



View Text Solution

29. A 10kg satellite circles earth once every 2hr in an orbit having a radius of 8000km . Assuming that Bohr's angular momentum postulate applies to satellites just as it does to an electron in the hydrogen atom, find the quantum number of the orbit of the satellite.

A. 5.3×10^{40}

B. 5.3×10^{45}

C. 7.8×10^{48}

D. 7.8×10^{50}

Answer: B



Watch Video Solution

30. if an electron is revolving in its bohr orbit having bohr radius of 0.529\AA , then the radius of third orbit is

A. 4234\AA

B. 4496\AA

C. 4.761\AA

D. $5125nm$

Answer: C



Watch Video Solution

31. the energy required to excite an electron in hydrogen atom to its first excited state is

A. $8.5eV$

B. $10.2eV$

C. $12.7eV$

D. $13.6eV$

Answer: B



Watch Video Solution

32. in the question number 63, the frequency of emitted photon due to the given transition is

$$(h = 6.64 \times 10^{-34} Js, 1eV = 1.6 \times 10^{-19} J)$$

A. $2.46 \times 10^{10} Hz$

B. $2.46 \times 10^{12} Hz$

C. $2.46 \times 10^{15} Hz$

D. $2.46 \times 10^{18} \text{ Hz}$

Answer: C



View Text Solution

33. Which state of the triply ionized Beryllium (Be^{3+}) has the same orbit radius as that of the ground state of hydrogen atom?

A. $n = 1$

B. $n = 2$

C. $n = 3$

D. $n = 4$

Answer: B



Watch Video Solution

34. A difference of 2.3 eV separates two energy levels in an atom. What is the frequency of radiation emitted when the atom transits from the upper level to the lower level.

A. $2.6 \times 10^{13} \text{ Hz}$

B. $2.6 \times 10^{14} \text{ Hz}$

C. $2.6 \times 10^{18} \text{ Hz}$

D. $2.6 \times 10^{18} \text{ Hz}$

Answer: B



Watch Video Solution

35. The ground state energy of hydrogen atom is -13.6eV . What are P.E. and K.E. of electron in this state?

A. $2.18 \times 10^{-14} J$

B. $2.18 \times 10^{-16} J$

C. $2.18 \times 10^{-18} J$

D. $2.18 \times 10^{-19} J$

Answer: C



Watch Video Solution

36. In the question number 67, find the potential energy of electron (in joule) in the given state.

A. $-4.36 \times 10^{-14} J$

B. $-4.36 \times 10^{-16} J$

C. $-4.36 \times 10^{-17} J$

D. $-4.36 \times 10^{-18} J$

Answer: D



View Text Solution

37. If the radius of inner most electronic orbit of a hydrogen atom is 5.3×10^{-11} m, then the radii of $n = 2$ orbit is

A. 1.12\AA

B. 2.12\AA

C. 3.22\AA

D. 4.54\AA

Answer: B



Watch Video Solution

38. A hydrogen atom initially in the ground level absorbs a photon and is excited to $n = 4$ level then the wavelength of photon is

A. 790\AA

B. 870\AA

C. 970\AA

D. 1070\AA

Answer: C



Watch Video Solution

39. In question number 70, what is frequency of photon ?

A. $3.1 \times 10^{15} \text{ Hz}$

B. $3.1 \times 10^{18} \text{ Hz}$

C. $9.1 \times 10^{15} \text{ Hz}$

D. $9.1 \times 10^{18} \text{ Hz}$

Answer: A



View Text Solution

40. The radius of electron orbit and the speed of electron in the ground state of hydrogen atom is $5.30 \times 10^{-11} \text{ m}$ and $22 \times 10^6 \text{ ms}^{-1}$

respectively, then the orbital period of this electron in second excited state will be

A. $1.21 \times 10^{-14} \text{ s}$

B. $1.21 \times 10^{-12} \text{ s}$

C. $1.21 \times 10^{-10} \text{ s}$

D. $1.21 \times 10^{-15} \text{ s}$

Answer: D



Watch Video Solution

41. In accordance with the Bohr's model, find the quantum number that characterizes the earth's revolution around the sun in an orbit of radius $1.5 \times 10^{11}m$ with orbital speed $3 \times 10^4m / s$. (Mass of earth= $6.0 \times 10^{24}kg$)

A. 5.98×10^{86}

B. 2.57×10^{38}

C. 8.57×10^{64}

D. 2.57×10^{74}

Answer: D



Watch Video Solution

42. if speed of electron is ground state energy level is $2.2 \times 10^6 \text{ms}^{-1}$, then its speed in fourth excited state will be

A. $6.8 \times 10^6 \text{ms}^{-1}$

B. $8.8 \times 10^5 \text{ms}^{-1}$

C. $5.5 \times 10^5 \text{ms}^{-1}$

D. $5.5 \times 10^6 \text{ms}^{-1}$

Answer: C



[View Text Solution](#)

43. if muonic hydrogen atom is an atom in which a negatively charged muon (μ) of mass about $207m_e$ revolves around a proton, then first bohr radius of this atom is ($r_e = 0.53 \times 10^{-10}m$)

A. $2.56 \times 10^{-10}m$

B. $2.56 \times 10^{-11}m$

C. $2.56 \times 10^{-12}m$

D. $2.56 \times 10^{-13} m$

Answer: D



Watch Video Solution

44. In the question number 75, if ground state energy of electron is $-13.6 eV$ then what is the ground state energy of muonic hydrogen atom?

A. $1.8 keV$

B. -2.8keV

C. -3.8keV

D. 4.8keV

Answer: B



Watch Video Solution

45. An electron is revolving in the n^{th} orbit of radius 4.2\AA , then the value of n is
($r - 1 = 0.529\text{\AA}$)

A. 4

B. 5

C. 6

D. 3

Answer: D



Watch Video Solution

46. Positronium is just like a H-atom with the proton replaced by the positively charged anti-particle of the electron (called the positron

which is as massive as the electron). What would be the ground state energy of positronium?

A. $3.4eV$

B. $-5.2eV$

C. $-6.8eV$

D. $-10.2eV$

Answer: C



Watch Video Solution

47. The minimum energy that must be given to a H atom in ground state so that it can emit an H γ line in balmer series is

A. $12.4eV$

B. $10.2eV$

C. $13.06eV$

D. $13.6eV$

Answer: C



Watch Video Solution

48. In the question number 79, what would be the angular momentum of H_γ photon if the angular momentum of the system is conserved

A. \hbar

B. $2\hbar$

C. $3\hbar$

D. $4\hbar$

Answer: C



View Text Solution

49. Energy E of a hydrogen atom with principle quantum number n is given by

$$E = \frac{-13.6}{n^2} eV. \text{ The energy of a photon}$$

ejected when the electron jumps from $n = 3$ state to $n = 2$ state of hydrogen is approximately

A. $1.5eV$

B. $0.85eV$

C. $3.4eV$

D. $1.9eV$

Answer: D



Watch Video Solution

50. Out of the following which one is not a possible energy for a photon to be emitted by hydrogen atom according to Bohr's atomic model?

A. $0.65eV$

B. 1.9eV

C. 11.1eV

D. 13.6eV

Answer: C



Watch Video Solution

51. The ratio of the speed of the electrons in the ground state of hydrogen to the speed of light in vacuum is

A. $\frac{1}{2}$

B. $\frac{2}{237}$

C. $\frac{1}{137}$

D. $\frac{1}{237}$

Answer: C



Watch Video Solution

52. In an excited state of hydrogen like atom an electron has total energy of $-3.4eV$. If the

kinetic energy of the electron is E and its de-Broglie wavelength is λ , then

A. $E = 6.8eV, \lambda = 6.6 \times 10^{-10}m$

B. $E = 3.4eV, \lambda = 6.6 \times 10^{-10}m$

C. $E = 3.4eV, \lambda = 6.6 \times 10^{-11}m$

D. $E = 6.8eV, \lambda = 6.6 \times 10^{-11}m$

Answer: B



Watch Video Solution

53. The binding energy of the electron in the ground state of He atom is equal to $E_0 = 24.6eV$. Find the energy required to remove both the electrons from the atom.


- A. 49.2eV
- B. 54.4eV
- C. 79eV
- D. 108.8eV

Answer: C



Watch Video Solution

54. The diagram shows the energy levels for an electron in a certain atom. Which transition shown represents the emission of photon with the most energy ?



A. I

B. II

C. III





D. IV

Answer: C



Watch Video Solution

55. In a hydrogen atom, the radius of n^{th} bohr orbit is r_n . The graph between $\log (r_n / r_1)$ and $\log n$ will be



Answer: A

 [View Text Solution](#)

56. An electron in the ground state of hydrogen atom is revolving in anticlockwise direction in circular orbit of radius R . The orbital magnetic dipole moment of the electron will be

A. $\frac{eh}{4\pi m}$

B. $\frac{eh}{2\pi m}$

C. $\frac{eh^2}{4\pi m}$

D. $\frac{e^2h}{4\pi m}$

Answer: A



Watch Video Solution

57. The electron in hydrogen atom makes a transition $n_1 \rightarrow n_2$ where n_1 and n_2 are the principal quantum number of two states. Assuming the Bohr model to be valid, the time period of the electron in the initial state is eight times that in the final state. The possible value of n_1 and n_2 are:

A. $n_1 = 4, n_2 = 2$

B. $n_1 = 8, n_2 = 2$

C. $n_1 = 8, n_2 = 1$

D. $n_1 = 6, n_2 = 2$

Answer: A



Watch Video Solution

58. the ionization energy of Li^{++} is equal to

A. $9hcR$

B. $6 hcR$

C. $2hcR$

D. hcR

Answer: A



Watch Video Solution

59. A hydrogen atom and a Li^{2+} ion are both in the second excited state. If l_H and l_{Li} are their respective electronic angular momenta, and E_H and E_{Li} their respective energies, then

(a) $l_H > l_{Li}$ and $|E_H| > |E_{Li}|$

(b) $l_H = l_{Li}$ and $|E_H| < |E_{Li}|$

(C) $l_H = l_{Li}$ and $|E_H| > |E_{Li}|$

(d) $l_H < l_{Li}$ and $|E_H| < |E_{Li}|$

A. $l_H > l_{Li}$ and $|E_H| > |E_{Li}|$

B. $l_H = l_{Li}$ and $|E_H| > |E_{Li}|$

C. $l_H = l_{Li}$ and $|E_H| < |E_{Li}|$

D. $l_H < l_{Li}$ and $|E_H| < |E_{Li}|$

Answer: C



Watch Video Solution

The Line Spectra Of Hydrogen Atom

1. The de-Broglie wavelength of an electron in the first Bohr orbit is

A. equal to one-fourth the circumference of the first orbit

B. equal to half the circumference of first orbit

C. equal to twice the circumference of first orbit.

D. equal to the circumference of the first orbit.

Answer: D



Watch Video Solution

2. the wavelength of spectral line in the lyman series of a H-atom is 1028\AA . If instead of hydrogen, we consider deuterium then shift in the wavelength of this line be ($m_p = 1860m_e$)

A. 1027.7\AA

B. 1036\AA

C. 1028\AA

D. 1021\AA

Answer: A



Watch Video Solution

3. The rydberg formula, for the spectrum of the hydrogen atom where all terms have their usual meaning is

$$\text{A. } h\nu_{\text{if}} = \frac{me^4}{8\varepsilon_0^2 h^2} \left(\frac{1}{n_f} - \frac{1}{n_i} \right)$$

$$\text{B. } h\nu_{\text{if}} = \frac{me^4}{8\varepsilon_0^2 h^2} \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$$

$$\text{C. } h\nu_{\text{if}} = \frac{8\varepsilon_0^2 h^2}{me^4} \left(\frac{1}{n_f} - \frac{1}{n_i} \right)$$

$$\text{D. } h\nu_{\text{if}} = \frac{8\varepsilon_0^2 h^2}{me^4} \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$$

Answer: B



Watch Video Solution

4. Hydrogen atom from excited state comes to the ground state by emitting a photon of

wavelength λ . If R is the Rydberg constant, then the principal quantum number n of the excited state is

A. $\sqrt{\frac{\lambda R}{\lambda R - 1}}$

B. $\sqrt{\frac{\lambda}{\lambda R - 1}}$

C. $\sqrt{\frac{\lambda R^2}{\lambda R - 1}}$

D. $\sqrt{\frac{\lambda R}{\lambda - 1}}$

Answer: A



Watch Video Solution

5. Ionization potential of hydrogen atom is $13.6V$. Hydrogen atoms in the ground state are excited by monochromatic radiation of photon energy $12.1eV$. The spectral lines emitted by hydrogen atoms according to Bohr's theory will be

A. one

B. two

C. three

D. four

Answer: C



Watch Video Solution

6. the excitation energy of lyman last lines is

A. the same as ionisation energy

B. the same as the last absorption line in

lyman series

C. both (a) and (b)

D. different from (a) and (b)

Answer: C



Watch Video Solution

De Broglie Explanation Of Bohr Model

1. the number of de broglie wavelength contained in the second bohr orbit of hydrogen atom is

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

2. if there are N atoms in a source of laser light and each atom is emitting light with intensity I , then the total intensity produced by it is

A. NI

B. $N^2 I$

C. $N^3 I$

D. $N^4 I$

Answer: B



View Text Solution

3. What does the word LASER stand for ?

A. Light amplification by stimulated emission of radiation

B. light amplitude by stimulated emission of radiation

C. light amplification by strong emission of radiation

D. light amplification by stimulated emission of radiowave

Answer: A



Watch Video Solution

Higher Order Thinking Skills

1. Deuterium was discovered in 1932 by Harold Urey by measuring the small change in wavelength for a particular transition in ${}^1\text{H}$ and ${}^2\text{H}$. This is because, the wavelength of transition depend to a certain extent on the nuclear mass. If nuclear motion is taken into account, then the electrons and nucleus revolve around their common centre of mass. Such a system is equivalent to a single particle

with a reduced mass μ , revolving around the nucleus at a distance equal to the electron - nucleus separation. Here

$\mu = m_e M / (m_e + M)$, where M is the nuclear mass and m_e is the electronic mass.

Estimate the percentage difference in wavelength for the 1st line of the Lyman series

in ${}^1_1\text{H}$ and ${}^2_1\text{H}$. (mass of ${}^1_1\text{H}$ nucleus is

1.6725×10^{-27} kg, mass of ${}^2_1\text{H}$ nucleus is

3.3374×10^{-27} kg, Mass of electron

$= 9.109 \times 10^{-31}$ kg.)

A. $2.7 \times 10^{-1} \%$.

B. $2.7 \times 10^{-2} \%$.

C. $3.5 \times 10^{-2} \%$.

D. $3.5 \times 10^{-1} \%$.

Answer: B



Watch Video Solution

2. The inverse square law in electrostatic is

$$|F| = \frac{e^2}{(4\pi\epsilon_0)r^2}$$

for the force between an

electron and a proton. The $\left(\frac{1}{r}\right)$ dependence

of $|F|$ can be understood in quantum theory

as being due to the fact that the particle of light (photon) is massless. If photons had a mass m_p , force would be modified to

$$|F| = \frac{e^2}{(4\pi\epsilon_0)\pi^2} \left[\frac{1}{r^2} + \frac{\lambda}{r} \right] \cdot \exp(-\lambda r)$$

where $\lambda = \frac{m_p c}{h}$ and $h = \frac{h}{2\pi}$. Estimate the

change in the ground state energy of a H-atom if m_p were 10^{-6} times the mass of the electron.

A. $18.6\lambda r_B$

B. -27.2

C. $27.2\lambda r_B$

$$D. -\lambda r_b$$

Answer: C



Watch Video Solution

3. The Bohr model for the H-atom relies on the Coulomb's law of electrostatics . Coulomb's law has not directly been varified for very short distances of the order of angstroms. Suppos-ing Coulomb's law between two oppsite charge $+q_1, -q_2$ is modified to

$$\left| \vec{F} \right| = \frac{q_1 q_2}{(4\pi\epsilon_0)r^2} \frac{1}{r^2}, r \geq R_0$$
$$= \frac{q_1 q_2}{(4\pi\epsilon_0)r^2} \frac{1}{R_0^2} \left(\frac{R_0}{r} \right)^\epsilon, r \leq R_0$$

Calculate in such a case , the ground state energy of H-atom , if $\epsilon = 0.1$, $R_0 = 1\text{\AA}$

A. -11.4

B. -17.3

C. 5.9

D. -23.2

Answer: A



Watch Video Solution

4. In the Auger process as atom makes a transition to a lower state without emitting a photon. The excess energy is transferred to an outer electron which may be ejected by the atom. (this is called an Auger electrons) . Assuming the nucleus to be massive , calculate the kinetic energy of an $n = 4$ Auger electron emitted by chromium by absorbing the energy from a $n = 2$ to $n = 1$ transition .

A. 4.6

B. 7.5

C. 5.38

D. 3.36

Answer: C



Watch Video Solution

5. If a proton had a radius R and the charge was uniformly distributed, the ground state energy (in eV) of a H-atom for $R = 0.1\text{\AA}$ is

A. -13.6

B. -27.2

C. -3.4

D. -30.8

Answer: A



Watch Video Solution

6. The ground state energy of an atom is $-13.6eV$. The photon emitted during the transition of electron from $n = 3$ to $n = 1$

state, is incident on a photosensitive material of unknown work function. The photoelectrons are emitted from the materials with a maximum kinetic energy of 9 eV. the threshold wavelength of the material used is

A. $0.9 \times 10^{-7} m$

B. $4 \times 10^{-7} m$

C. $0.47 \times 10^{-7} m$

D. $9 \times 10^{-7} m$

Answer: A



7. For scattering by an inverse-square field (such as that produced by a charged nucleus in Rutherford's model) the relation between impact parameter b and the scattering angle θ is given by, the scattering angle for $b = 0$ is

A. 180°

B. 90°

C. 45°

D. 120°

Answer: A



Watch Video Solution

8. The impact parameter at which the scattering angle is 90^0 , $z = 79$ and initial energy $10MeV$ is



Watch Video Solution

Ncert Exemplar

1. Taking the Bohr radius $a_0 = 53$ pm, the radius of Li^{++} ion in its ground state, on the basis of Bohr's model, will be about.

A. 53pm

B. 27pm

C. 18pm

D. 13pm

Answer: C



Watch Video Solution

2. The binding energy of a H-atom considering an electron moving around a fixed nuclei (proton), is

$$B = - \frac{me^4}{8n^2\epsilon_0^2h^2} \quad (m = \text{electron mass})$$

If one decides to work in a frame of reference where the electron is at rest, the proton would be moving around it. By similar arguments, the binding energy would be :

$$B = - \frac{me^4}{8n^2\epsilon_0^2h^2} \quad (M = \text{proton mass})$$

This last expression is not correct, because

A. n would not be integral

B. bohr- quantisation applies only to
electron

C. the frame in which the electron is at rest
is not inertial

D. the motion of the proton would not be
in circular orbits, even approximately

Answer: C



Watch Video Solution

3. The simple Bohr model cannot be directly applied to calculate the energy level of an atom with many electrons . This is because.

A. of the electrons not being subject to a central force

B. of the electrons colliding with each other

C. of screening effects

D. the force between the nucleus and an electron will no longer be given by

coulomb's law.

Answer: A



Watch Video Solution

4. For the ground state, the electron in the H-atom has an angular momentum $= h$, according to the simple Bohr model. Angular momentum is a vector and hence there will be infinitely many orbits with the vector pointing

in all possible direction . In actuality , this is not true,

A. because bohr model gives incorrect values of angular momentum

B. because only one of these would have a minimum energy

C. angular momentum must be in the direction of spin of electron

D. because electrons go around only in horizontal orbits

Answer: A



Watch Video Solution

5. O_2 molecules consists of two oxygen atoms.

In the molecules , nuclear force between the nuclei of the two atoms

A. is not important because nuclear forces are short-ranged.

B. is as important as electrostatic force for binding the two atoms

C. cancels the repulsive electrostatic force
between the nuclei

D. is not important because oxygen
nucleus has equal number of neutrons
and protons.

Answer: A



Watch Video Solution

6. Two H atoms in the ground state collide in elastically. The maximum amount by which their combined kinetic energy is reduced is

A. 10.2eV

B. 20.4eV

C. 13.6eV

D. 27.2eV

Answer: A



Watch Video Solution

7. A set of atoms in an excited state decays

A. in general to any of the states with lower energy

B. into a lower state only when excited by an external electric field

C. all together simultaneously into a lower state

D. to emit photons only when they collide

Answer: A



Watch Video Solution

Assertion And Reason

1. (A) atoms of each element are stable and emit characteristic spectrum.

(R) the spectrum provides useful information about the atomic structure.

A. if both assertion and reason are true
and reason is the correct explanation of

assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: B



Watch Video Solution

2. (A) atom as a whole is electrically neutral.

(R)atom contains equal amount of positive and negative charges.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: A



Watch Video Solution

3. (A) according to electromagnetic theory an accelerated particle continuously emits radiation.

(R) according to classical theory, the proposed path of an electron in rutherford atom model will be parabolic.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: C



Watch Video Solution

4. (A) In alpha particle scattering, the number of alpha particles undergoing head-on collision is small.

(R) A small fraction of the number of incident particles rebound back.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: B



Watch Video Solution

5. (A) most of the mass of the atom is concentrated in its nucleus.

(R) all alpha particles striking a gold sheet are scattered in different directions.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: C



Watch Video Solution

6. (A) the trajectory traced by an incident particle depends on the impact parameter of collision.

(R) the impact parameter is the perpendicular distance of the initial velocity vector of the

incident particle from the centre of the target nucleus.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: B



View Text Solution

7. (A) in the experiment of alpha particle scattering, extremely thin gold foils are preferred over other metals.

(R)gold is a ductile material.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: B



Watch Video Solution

8. (A) the total energy of an electron revolving in any stationary orbit is negative.

(R) energy can have positive or negative values.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: B



Watch Video Solution

9. Statement -1 : Large angle scattering of alpha particles led to the discovery of atomic nucleus.

Statement -2 : Entire positive charge of atom is concentrated in the central core.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: A



Watch Video Solution

10. Assertion: For the scattering of α -particles at a large angles, only the nucleus of the atom is responsible.

Reason: Nucleus is very heavy in comparison to electrons.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: A



Watch Video Solution

11. Assertion: Hydrogen atom consists of only one electron but its emission spectrum has many lines.

Reason: Only Lyman series is found in the absorption spectrum of hydrogen atom whereas in the emission spectrum, all the series are found.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: B



Watch Video Solution

12. (A) bohr model can not be extended to two or more electron atoms.

(R) each electron in the atom interacts not only with the positively charged nucleus but also with all other electrons.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of

assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: A



Watch Video Solution

13. Assertion: Bohr had to postulate that the electrons in stationary orbits around the nucleus do not radiate.

Reason: According to classical physical all moving electrons radiate.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: C



Watch Video Solution

14. (A) bohr's third postulaate states that the stationary orbits are those for which the angular momentum is some integral multiple of $\frac{h}{2\pi}$.

(R) linear momentum of the electron in the atom is quantised.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: D



Watch Video Solution

15. Assertion: Electrons in the atom are held due to coulomb forces.

Reason: The atom is stable only because the centripetal force due to Coulomb's law is balanced by the centrifugal force.

A. if both assertion and reason are true and reason is the correct explanation of assertion.

B. if both assertion and reason are true but reason is not the correct explanation of assertion.

C. if assertion is true but reason is false.

D. if both assertion and reason are false.

Answer: C



Watch Video Solution