



PHYSICS

BOOKS - NCERT FINGERTIPS PHYSICS


(HINGLISH)

NUCLEI


Atomic Masses

1. The set which represents the isotope , isobar and isotone respectively is


A.




B.



C.



D.



Answer: D



View Text Solution

2. The natural boron of atomic weight 10.81 is found to have two isotopes ^{10}B and ^{11}B . The ratio of abundance of isotopes of natural boron should be

A. 11 : 10

B. 81 : 19

C. 10 : 11

D. 19 : 81

Answer: d



Watch Video Solution

3. Two stable isotopes of lithium ${}^6_3\text{Li}$ and ${}^7_3\text{Li}$ have respective abundances of 7.5% and 92.5% . These isotopes have masses 6.0152 u and 7.016004 u respectively. Find the atomic weight of lithium

A. 6.941 u

B. 3.321 u

C. 2.561 u

D. 0.621 u

Answer: a



Watch Video Solution

Size Of The Nucleus

1. The most common kind of iron nucleus has a mass number of 56. Find the radius, approximate mass, and approximate density of the nucleus.



[Watch Video Solution](#)

2. Order of magnitude of density of uranium nucleus is , [m = 1.67×10^{-27} kg]

A. $10^{20} \text{ kg m}^{-3}$

B. $10^{17} \text{ kg m}^{-3}$

C. 10^{14}kg m^{-3}

D. 10^{11}kg m^{-3}

Answer: B



Watch Video Solution

3. Two nuclei have their mass numbers in the ratio of 1:3. The ratio of their nuclear densities would be

A. $(3)^{1/3} : 1$

B. 1 : 1

C. 1 : 3

D. 3:1

Answer: B



Watch Video Solution

4. The radius of a spherical nucleus as measured by electron scattering is 3.6 fm. What is the mass number of the nucleus most likely to be ?

A. 27

B. 40

C. 56

D. 120

Answer: A



Watch Video Solution

5. The ratio of the nuclear radii of the gold isotope

${}_{79}^{197}Au$ and silver isotope ${}_{47}^{107}Ag$ is

A. 1.23

B. 0.216

C. 2.13

D. 3.46

Answer: A



Watch Video Solution

6. If the nucleus of ${}_{13}\text{Al}^{27}$ has a nuclear radius of about 3.6 fm, then ${}_{52}\text{Te}^{125}$ would have its radius approximately as

A. 9.6 fm

B. 12 fm

C. 4.8 fm

D. 6 fm

Answer: D



Mass Energy And Nuclear Binding Energy

1. How much mass has to be converted into energy to produce electric power of 500 MW for one hour ?

A. 2×10^{-5} kg

B. 1×10^{-5} kg

C. 3×10^{-5} kg

D. 4×10^{-5} kg

Answer: A



Watch Video Solution

2. if in a nuclear fusion reaction, mass defect to 0.3% ,
then energy released in fusion of 1 kg mass

A. 27×10^{10} J

B. 27×10^{11} J

C. 27×10^{12} J

D. 27×10^{13} J

Answer: D



Watch Video Solution

3. The equivalent energy of 1 g of substance is

A. $9 \times 10^{13} \text{ J}$

B. $6 \times 10^{12} \text{ J}$

C. $3 \times 10^{13} \text{ J}$

D. $6 \times 10^{13} \text{ J}$

Answer: A



Watch Video Solution

4. Fission of nuclei is possible because the binding energy per nucleon in them

A. increases with mass number at low mass numbers

B. decreases with mass number at low mass numbers

C. increases with mass number at high mass numbers

D. decreases with mass number at high mass numbers

Answer: d



Watch Video Solution

5. The mass of ${}^7_3\text{Li}$ is 0.042 amu less than the sum of masses of its constituents. The binding energy per nucleon is

A. 5.586 MeV

B. 10.522 MeV

C. 2.433 MeV

D. 3.739 MeV

Answer: a



Watch Video Solution

6. The binding energy per nucleon of deuterium and helium nuclei are 1.1 MeV and 7.0 MeV respectively. When two deuterium nuclei fuse to form a helium nucleus the energy released in the fusion is

- A. 23.6 MeV
- B. 2.2 MeV
- C. 28.0 MeV
- D. 30.2 MeV

Answer: A



Watch Video Solution

7. Let m_p be the mass of a proton, m_n the mass of a neutron, M_1 the mass of a ${}^{20}_{10}\text{Ne}$ nucleus and M_2 the mass of a ${}^{40}_{20}\text{Ca}$ nucleus. Then

A. $M_2 = M_1$

B. $M_2 > M_1$

C. $M_2 < M_1$

D. $M_1 < 10(m_n + m_p)$

Answer: D



Watch Video Solution

8. The mass of proton is 1.0073 u and that of neutron is 1.0087 u (u=atomic mass unit). The binding energy of ${}^4_2\text{He}$, if mass of ${}^4_2\text{He}$ is $4.0015u$, is

A. 0.0305 erg

B. 0.0305 J

C. 28.4 MeV

D. 0.061 u

Answer: C



Watch Video Solution

9. If the binding energy per nucleon of deuterium is 1.115 MeV, its mass defect in atomic mass unit is

A. 0.0048

B. 0.0024

C. 0.0012

D. 0.0006

Answer: b



Watch Video Solution

1. The half - life period of a radioactive element x is same as the mean life time of another radioactive element y Initially both of them have the same number of atoms. Then

- A. X and Y decay at same rate always
- B. X will decay faster than Y
- C. Y will decay faster than X
- D. X and Y have same decay rate initially

Answer: C



Watch Video Solution

2. Sample of two radioactive nuclides A and B are taken. λ_A and λ_B are the disintegration constants of A and B respectively. In which of the following cases, the two sample can simultaneously have the same decay rate at any time ?

A. Initial rate of decay of A is twice the initial rate of decay of B and $\lambda_A = \lambda_B$.

B. Initial rate of decay of A is twice the initial rate of decay of B and $\lambda_A > \lambda_B$.

C. Initial rate of decay of B is twice the initial rate of decay of A and $\lambda_A > \lambda_B$.

D. Initial rate of decay of B is twice the initial rate of decay of A at $t=2h$ and $\lambda_B = \lambda_A$

Answer: B

 **Watch Video Solution**

3. The half life of ${}_{92}^{238}U$ undergoing α -decay is 4.5×10^9 years. The activity of 1 g sample of ${}_{92}^{238}U$ is

A. 1.23×10^4 Bq

B. 1.23×10^5 Bq

C. 1.23×10^3 Bq

D. 1.23×10^6 Bq

Answer: A



Watch Video Solution

4. The half life of a radioactive substance is 20s, the time taken for the sample to decay by $7/8^{th}$ of its initial value is

A. 20 s

B. 40 s

C. 60 s

D. 80 s

Answer: C



Watch Video Solution

5. 1 mg radium has 2.68×10^{18} atoms. Its half life is 1620 years. How many radium atoms will disintegrate from 1 mg of pure radium in 3240 years ?

- A. 2.01×10^9
- B. 2.01×10^{18}
- C. 1.01×10^9
- D. 1.01×10^{18}

Answer: B



Watch Video Solution

6. Radon has 3.8 days as its half-life . How much radon will be left out of 15 mg mass after 38 days ?

A. 1.05 mg

B. 0.015 mg

C. 0.231 mg

D. 0.50 mg

Answer: B



Watch Video Solution

7. The half life of a radioactive substance is 30 days.

What is the time taken to disintegrate to $3/4^{th}$ of its original mass ?

A. 30 days

B. 15 days

C. 60 days

D. 90 days

Answer: C



Watch Video Solution

8. The half life of polonium is 140 days. In what time will 15 g of polonium be disintegrated out of its initial mass of 16 g?

A. 230 days

B. 560 days

C. 730 days

D. 160 days

Answer: B



Watch Video Solution

9. In a sample of radioactive material , what fraction of the initial number of active nuclei will remain undisintegrated after half of the half life of the sample ?

A. $\frac{1}{4}$

B. $\frac{1}{2\sqrt{2}}$

C. $\frac{1}{\sqrt{2}}$

D. $\sqrt{2} - 1$

Answer: c



Watch Video Solution

10. At a given instant, there are 25% undecayed radioactive nuclei in a sample. After 10 seconds the number of undecayed nuclei reduces to 12.5%, the mean life of the nuclei is

A. 10.21 s

B. 14.43 s

C. 5.31 s

D. 7.43 s

Answer: B



Watch Video Solution

11. Two radioactive nuclei A and B have disintegration constants λ_A and λ_B and initially N_A and N_B number of nuclei of them are taken, then the time after which their undisintegrated nuclei are same is

A. $\frac{\lambda_A \lambda_B}{(\lambda_A - \lambda_B)} \ln \left(\frac{N_B}{N_A} \right)$

B. $\frac{1}{(\lambda_A + \lambda_B)} \ln \left(\frac{N_B}{N_A} \right)$

C. $\frac{1}{(\lambda_B - \lambda_A)} \ln \left(\frac{N_B}{N_A} \right)$

D. $\frac{1}{(\lambda_A - \lambda_B)} \ln \left(\frac{N_B}{N_A} \right)$

Answer: C



Watch Video Solution

12. A freshly prepared radioactive source of half-life $2h$ emits radiation of intensity which is 64 times the permissible safe level. The minimum time after which it would be possible to work safely with this source is

A. 128 h

B. 24 h

C. 6 h

D. 12 h

Answer: D



Watch Video Solution

13. Two samples X and Y contain equal amount of radioactive substances. If $\frac{1^{th}}{16}$ of the sample Y , remain after 8 hours, then the ratio of half life periods of X and Y is

A. 2:1

B. 1:2

C. 1:4

D. 4:1

Answer: A



View Text Solution

14. The decay constant of a radioactive isotope is λ . If A_1 and A_2 are its activities at time t_1 and t_2 respectively, then the number of nuclei which have decayed the time $(t_1 - t_2)$

A. $A_1 t_1 - A_2 t_2$

B. $A_1 - A_2$

C. $(A_1 - A_2) / \lambda$

D. $\lambda(A_1 - A_2)$

Answer: C



Watch Video Solution

15. A 280 day old radioactive substances shows an activity of 6000 dps, 140 days later its activity becomes 3000 dps. What was its initial activity ?

A. 20000 dps

B. 24000 dps

C. 12000 dps

D. 6000 dps

Answer: b



Watch Video Solution

16. Tritium has a half life of 12.5 years against beta decay. What fraction of a sample of pure tritium will remain undecayed after 25 years?

- A. one half
- B. one fourth
- C. one third
- D. can't say

Answer: B



Watch Video Solution

17. The decay constant , for a given radioactive sample, is 0.3465 day^{-1} .What percentage of this sample will get decayed in a period of 4 days ?

A. 100 %

B. 50 %

C. 75 %

D. 10 %

Answer: C



Watch Video Solution

18. The count rate of a radioactive sample falls from $4.0 \times 10^6 \text{ s}^{-1}$ to $1.0 \times 10^6 \text{ s}^{-1}$ in 20 hours. What will be the count rate after 100 hours from beginning ?

A. $3.91 \times 10^3 \text{ s}^{-1}$

B. $3.91 \times 10^2 \text{ s}^{-1}$

C. $3.91 \times 10^4 \text{ s}^{-1}$

D. $3.91 \times 10^6 \text{ s}^{-1}$

Answer: A



Watch Video Solution

19. Carbon dating is best suited for determining the age of fossils of their age in years is of the order of

A. 10^3

B. 10^4

C. 10^5

D. 10^6

Answer: b



Watch Video Solution

20. The half life of radioactive Radon is 3.8days . The time at the end of which $\frac{1}{20}$ th of the radon sample will remain undecayed is (*given* $\log e = 0.4343$)

A. 3.8 days

B. 16.5 days

C. 33 days

D. 76 days

Answer: B



Watch Video Solution

21. A radioactive element X with half life 2 h decays giving a stable element Y. After a time t, ratio of X and Y atoms is 1:16 .Time t is

A. 6 h

B. 4 h

C. 8 h

D. 16 h

Answer: C



Watch Video Solution

22. The half life of ${}_{38}^{90}\text{Sr}$ is 28 years. The disintegration rate of 15 mg of this isotope is of the order of

A. 10^{11} Bq

B. 10^{10} Bq

C. 10^7 Bq

D. 10^9 Bq

Answer: b



Watch Video Solution

23. A sample of a radioactive element has a mass of 10 g at an instant $t=0$. The approximate mass of this element in the sample left after two mean lives is

A. 1.35 g

B. 2.50 g

C. 3.70 g

D. 6.30 g

Answer: A



Watch Video Solution

24. The count rate from 100cm^3 of a radioactive liquid is c . Some of this liquid is now discarded. The count rate of the remaining liquid is found to be $c/10$ after three half-lives. The volume of the remaining liquid, in cm^3 , is

A. 20

B. 40

C. 60


D. 80

Answer: D



Watch Video Solution

25. The variation of decay rate of two radioactive samples A and B with time is shown in fig.



Which of the following statements are true?

- A. Decay constant of A is greater than that of B,
hence A always decays faster than B.

- B. Decay constant of A is greater than that of B,
but it does not always decays faster than B.
- C. Decay constant of B is smaller than that of A
but still its decay rate becomes equal to that of
A at a later instant.
- D. Both (b) and (c)

Answer: D



Watch Video Solution

26. A sample of radioactive material has mass m ,
decay constant λ , and molecular weight M . Avogadro

constant = N_A . The initial activity of the sample is:

A. λm

B. $\frac{\lambda m}{M}$

C. $\frac{\lambda m N_A}{M}$

D. $m N_A \lambda$

Answer: c



Watch Video Solution

27. Two radioactive substance A and B have decay constants 5λ and λ respectively. At $t = 0$ they have the same number of nuclei. The ratio of number of

nuclei of nuclei of A to those of B will be $\left(\frac{1}{e}\right)^2$
after a time interval

A. 4λ

B. 2λ

C. $1/2\lambda$

D. $1/4\lambda$

Answer: C



Watch Video Solution

28. Plutonium decays with half life of 24000 years. If plutonium is stored for 72000 years, the fraction of it

that remains is

A. $1/8$

B. $1/3$

C. $1/4$

D. $1/2$

Answer: a



Watch Video Solution

29. A fraction f_1 of a radioactive sample decays in one mean life, and a fraction f_2 decays in one half life.

Then

A. $f_1 > f_2$

B. $f_1 < f_2$

C. $f_1 = f_2$

D. either of (a),(b) or (c) depending on the values of the mean life and half life.

Answer: a



Watch Video Solution

30. The activity of a radioactive sample is measured as N_0 counts per minute at $t = 0$ and N_0/e counts per

minute at $t = 5 \text{ min}$. The time (in minute) at which the activity reduces to half its value is.

A. $\frac{\log_e 2}{5}$

B. $\frac{5}{\log_e 2}$

C. $5 \log_{10} 2$

D. $5 \log_e 2$

Answer: D



Watch Video Solution

31. The number of beta particles emitter by radioactive substance is twice the number of alpha

particles emitter by it. The resulting daughter is an

- A. isomer of parent
- B. isotone of parent
- C. isotope of parent
- D. isobar of parent

Answer: C



Watch Video Solution

32. During negative β decay, an anti- neutrino is also emitted along with the ejected electron. Then

A. only linear momentum will be conserved

B. total linear momentum and total angular momentum but not total energy will be conserved

C. total linear momentum and total energy but not total angular momentum will be conserved

D. total linear momentum , total angular momentum and total energy will be conserved.

Answer: d



Watch Video Solution

33. Which of the following cannot be emitted by radioactive substances during their decay ?

A. Neutrinos

B. Protons

C. Electrons

D. Helium nuclei

Answer: b



Watch Video Solution

34. A radioactive decay can form an isotope of the original nucleus with the emission of particles

A. one α and four β

B. one α and two β

C. one α and one β

D. four α and one β

Answer: b



Watch Video Solution

35. Pick out the incorrect statement from the following .

- A. β^- emission from the nucleus is always accompanied with a neutrino
- B. The energy of the α -particle emitted from a given nucleus is always constant.
- C. γ -ray emission makes the nucleus more stable.
- D. Nuclear force is charge-independent

Answer: A



Watch Video Solution

36. Consider α - , β^- particles and γ - rays, each having an energy fo 0.5Mev . In increasing order of

penetration power, the radiations are:

A. α , β , γ

B. α , γ , β

C. β , γ , α

D. γ , β , α

Answer: a



Watch Video Solution

37. The electron emitted in beta radiation originates from
from

- A. inner orbits of atom
- B. free electrons existing in the nuclei
- C. decay of a neutron in a nuclei
- D. photon escaping from the nucleus

Answer: c

 [Watch Video Solution](#)

38. Radioactive ${}_{27}^{60}\text{Co}$ is transformed into stable ${}_{28}^{60}\text{Ni}$ by emitting two γ -rays of energies

- A. 1.33 MeV and 1.17 MeV in succession

B. 1.17 MeV and 1.33 MeV in succession

C. 1.37 MeV and 1.13 MeV in succession

D. 1.13 MeV and 1.37 MeV in succession

Answer: b



Watch Video Solution

39. Complete the series ${}^6\text{He} \rightarrow e^{-} + {}^6\text{Li} +$

A. neutrino

B. antineutrino

C. proton

D. neutron

Answer: B



Watch Video Solution

40. A nucleus of Ux_1 has a half life of 24.1 days. How long a sample of Ux_1 will take to change to 90% of Ux_1 .

A. 80 days

B. 40 days

C. 20 days


D. 10 days

Answer: A



Watch Video Solution

41. An element A decays into element C by a two-step process :



Then.

A. A and C are isotopes

B. A and C are isobars

C. B and C are isotopes


D. A and B are isobars

Answer: A



Watch Video Solution

42. The equation



A. β -decay

B. γ -decay

C. fusion

D. fission

Answer: c



Watch Video Solution

Nuclear Energy

1. Light energy emitted by star is due to

- A. breaking of nuclei
- B. joining of nuclei
- C. burning of nuclei
- D. reflection of solar light

Answer: b



Watch Video Solution

2. ENERGY CONSERVATION IN NUCLEAR REACTION

A. mass only

B. energy only

C. momentum only

D. mass, energy and momentum

Answer: D



Watch Video Solution

3. which of the following are used as control rods in a nuclear reaction ?

A. cadmium

B. graphite

C. krypton

D. plutonium

Answer: A



Watch Video Solution

4. Fast neutrons can easily be slowed down by





- A. the use of lead shielding
- B. passing them through water
- C. elastic collisions with heavy nuclei
- D. applying a strong electric field

Answer: b



Watch Video Solution

5. Which of the following equations pick out the possible nuclear fusion reactions?



Answer: A



Watch Video Solution

6. If 200MeV energy is released in the fission of a single U^{235} nucleus, the number of fissions required

per second to produce 1 kilowatt power shall be

(Given $1eV = 1.6 \times 10^{-19} J$).

A. 3.125×10^{13}

B. 1.52×10^6

C. 3.125×10^{12}

D. 3.125×10^{14}

Answer: A



Watch Video Solution

7. In a nuclear fusion reaction, two nuclei, A & B , fuse to produce a nucleus C , releasing an amount of

energy ΔE in the process. If the mass defects of the three nuclei are ΔM_A , ΔM_B & ΔM_C respectively, then which of the following relations holds? Here, c is the speed of light.

A. $\Delta M_A + \Delta M_B = \Delta M_C - \Delta E / c^2$

B. $\Delta M_A + \Delta M_B = \Delta M_C + \Delta E / c^2$

C. $\Delta M_A - \Delta M_B = \Delta M_C - \Delta E / c^2$

D. $\Delta M_A - \Delta M_B = \Delta M_C + \Delta E / c^2$

Answer: A



Watch Video Solution

8. Mass numbers of the elements A,B , C and D are 30,60,90 and 120 respectively. The specific binding energy of them are 5 MeV, 8.5 MeV , 6 MeV and 7 MeV respectively. Then, in which of the following reaction/s energy is released ?

(1) $D \rightarrow 2B$, (2) $C \rightarrow B + A$, (3) $B \rightarrow 2A$

A. only in (1)

B. in (2),(3)

C. in (1) , (3)

D. in (1), (2) and (3)

Answer: A



9. The fission properties of ${}_{94}^{239}\text{Pu}$ are very similar to those of ${}_{92}^{235}\text{U}$. The average energy released per fission is 180 MeV. If all the atoms in 1 kg of pure ${}_{94}^{239}\text{Pu}$ undergo fission, then the total energy released in MeV is

A. 4.53×10^{26} MeV

B. 2.21×10^{14} MeV

C. 1×10^{13} MeV

D. 6.33×10^{24} MeV

Answer: A



Watch Video Solution

10. A nucleus with mass number 220 initially at rest emits an α -particle. If the Q-value of the reaction is 5.5MeV , calculate the kinetic energy of the α -particle.


- A. 4.4 MeV
- B. 5.4 MeV
- C. 5.6 MeV
- D. 6.5 MeV

Answer: B



Higher Order Thinking Skills

1. Sometimes a radioactive nucleus decays into a nucleus which itself is radioactive . An example is :



Assume that we start with 1000 ${}^{38}\text{S}$ nuclei at time $t=0$

. The number of ${}^{38}\text{Cl}$ is of count zero at $t=0$ and will again be zero at $t = \infty$. At what value of t , would the number of counts be a maximum ?

A. 1.65 h

B. 2.62 h

C. 3.24 h

D. 3.95 h

Answer: a



Watch Video Solution

2. The deuteron is bound by nuclear forces just as H-atom is made up of p and e bound by electrostatic forces. If we consider the forces between neutron and proton in deuteron as given in the form of a Coulomb

force but with an effective charge e' :
$$F = \frac{1}{4\pi\epsilon_0} \frac{e'^2}{r^2}$$

estimate the value of (e'/e) given that the following binding energy of a deuteron is 2.2MeV.

A. 1.89

B. 9.24

C. 3.64


D. 7.62

Answer: c



Watch Video Solution

3. A fission reaction is given by



x and y are two particles. Considering ${}_{92}^{236}\text{U}$ to be at

rest, the kinetic energies of the products are denoted

by $K_{xe}, K_{Sr}, K_x(2MeV)$ and $K_y(2MeV)$, respectively. Let the binding energies per nucleon of ${}_{92}^{236}U$, ${}_{54}^{140}Xe$ and ${}_{38}^{94}Sr$ be $7.5MeV$, $8.5MeV$ and $8.5MeV$, respectively. Considering different conservation laws, the correct options (*s*) is (are)


- A. $x=n, y=n, K_{Sr}=129\text{ MeV}, K_{Xe}=86\text{ MeV}$
- B. $x=p, y=e^{-}, K_{Sr}=129\text{ MeV}, K_{Xe}=86\text{ MeV}$
- C. $x=p, y=n, K_{Sr}=129\text{ MeV}, K_{Xe}=86\text{ MeV}$
- D. $x=n, y=n, K_{Sr}=86\text{ MeV}, K_{Xe}=129\text{ MeV}$

Answer: A



Watch Video Solution

4. How long can an electric lamp of 100W be kept glowing by fusion of 2.0 kg of deuterium? The fusion reaction can be taken as



A. 2.4×10^6 years

B. 7.4×10^4 years

C. 1.6×10^6 years

D. 4.9×10^4 years

Answer: d



Watch Video Solution

5. Nuclei of a radioactive element A are being produced at a constant rate α . The element has a decay constant λ . At time $t=0$, there are N_0 nuclei of the element. The number N of nuclei of A at time t is

A. $\frac{1}{\lambda} [\alpha + (\alpha - N_0\lambda)e^{-\lambda t}]$

B. $\frac{1}{\lambda} [\alpha - (\alpha - N_0\lambda)e^{-\lambda t}]$

C. $\lambda [\alpha - (\alpha - N_0\lambda)e^{-\lambda t}]$

D. $[\alpha - (N_0\lambda - \alpha)e^{-\lambda t}]$

Answer: B



Watch Video Solution

6. If $\alpha = 2N_0\lambda$, calculate the number of nuclei of A after one half-life of A, and also the limiting value of N as

A. $2N_0, \frac{5}{2}N_0$

B. $3N_0, 2N_0$

C. $4N_0, 2N_0$

D. $\frac{3}{2}N_0, 2N_0$

Answer: d



View Text Solution

7. The element curium ${}_{96}^{248}\text{Cm}$ has a mean life of 10^{13}s . Its primary decay modes are spontaneous fission and α -decay, the former with a probability of 8% and the latter with a probability of 92%, each fission releases 200MeV of energy. The masses involved in decay are as follows

$${}_{96}^{248}\text{Cm} = 248.072220u,$$

$${}_{94}^{244}\text{Pu} = 244.064100u \quad \text{and} \quad {}_2^4\text{He} = 4.002603u.$$

Calculate the power output from a sample of 10^{20} Cm atoms. ($1u = 931\text{MeV}/c^2$)

A. $4.42 \times 10^{-3} \text{ W}$

B. $3.32 \times 10^{-5} \text{ W}$

$$\text{C. } 4.42 \times 10^{-5} \text{ W}$$



$$\text{D. } 3.32 \times 10^{-3} \text{ W}$$

Answer: b



Watch Video Solution

8. The alongside is a plot of binding energy per nucleon E_b , against the nuclear mass M , A, B, C, D, E, F correspond to different nuclei. Consider four reactions.



energy released. In which reactions , is ε positive?



A. (i) and (iii)

B. (ii) and (iv)

C. (ii) and (iii)

D. (i) and (iv)

Answer: D



Watch Video Solution

Ncert Exemplar

1. Suppose we consider a large number of containers each containing initially 10000 atoms of a radioactive material with a half life of 1 year. After 1 year.

A. all the containers will have 5000 atoms of the material.

B. all the containers will contain the same number of atoms of the material but that number will only be approximately 5000

C. the containers will in general have different number of the atoms of the material but their average will be close to 5000

D. none of containers can have more than 5000 atoms.

Answer: c

 **Watch Video Solution**

2. The gravitational force between a H-atom and another particle of mass m will be given by Newton's

law: $F = G \frac{M \cdot m}{r^2}$, where r is in km and

A. $M = m_{\text{proton}} + m_{\text{electron}}$

B. $M = m_{\text{proton}} + m_{\text{electron}} - \frac{B}{c^2}$ (B=13.6 eV)

C. M is not related to the mass of hydrogen atom

D. $M = m_{\text{proton}} + m_{\text{electron}} - \frac{|V|}{c^2}$ ($|V|$ =magnitude of the potential energy of electron in the H-atom)

Answer: b



Watch Video Solution

3. When a nucleus in an atom undergoes a radioactive decay, the electronic energy levels of the atom

A. do not change for any type of radioactivity

B. change for α and β radioactivity but not for γ -radioactivity

C. change for α -radioactivity but not for others

D. change for β -radioactivity but not for others

Answer: b



View Text Solution

4. M_x and M_y denote the atomic masses of the parent and the daughter nuclei respectively in a radioactive decay. The Q - value for a β^- decay is Q_1 and that for a β^+ decay is Q_2 . If m_e denotes the

mass of an electrons, then which of the following statements is correct?

A. $Q_1 = (M_x - M_y)c^2$ and

$$Q_2 = (M_x - M_y - 2m_e)c^2$$

B. $Q_1 = (M_x - M_y)c^2$ and $Q_2 = (M_x - M_y)c^2$

C. $Q_1 = (M_x - M_y - 2m_e)c^2$ and

$$Q_2 = (M_x - M_y + 2m_e)c^2$$

D. $Q_1 = (M_x - M_y + 2m_e)c^2$ and

$$Q_2 = (M_x - M_y + 2m_e)c^2$$

Answer: A



Watch Video Solution

5. Tritium is an isotope of hydrogen whose nucleus triton decays , it would transform into He^3 nucleus . This does not happen .This is because

- A. Triton energy is less than that of a He^3 nucleus
- B. The electron created in the beta decay process cannot remain in the nucleus
- C. both the neutrons in triton have a decay simultaneously resulting in a nucleus with 3 protons which is not a He^3 nucleus .
- D. because free neutrons decay due to external perturbations which is absent in a triton

nucleus .

Answer: A



View Text Solution

6. Heavy stable nuclei have more neutrons than protons. This is because of the fact that

A. neutrons are heavier than protons .

B. electrostatic force between protons are repulsive

C. neutrons decay into protons through beta decay

D. nuclear forces between neutrons are weaker than that between protons.

Answer: B

 [Watch Video Solution](#)

7. In a nuclear reactor, moderators slow down the neutrons which come out in a fission process. The moderator used have light nuclei. Heavy nuclei will not serve the purpose because

A. they will break up

B. elastic collision of neutrons with heavy nuclei

will not slow them down

C. the net weight of the reactor would be

unbearably high

D. substances with heavy nuclei do not occur in

liquid or gaseous state at room temperature .

Answer: b



Watch Video Solution

1. Assertion: The whole mass of the atom is concentrated in the nucleus.

Reason: The mass of a nucleus can be either less than or more than the sum of the masses of nucleons present in it.

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: c



Watch Video Solution

2. Assertion : The radius of a nucleus determined by electron scattering is found to be slightly different from that determined by alpha particle scattering .

Reason : Electron scattering senses the charge distribution of the nucleus whereas alpha and similar particles sense the nuclear matter.

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: A



View Text Solution

3. Assertion: Isotopes of an element can be separated by using a mass spectrometer.

Reason: Separation of isotopes is possible because of difference in electron number of isotopes .

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: c



View Text Solution

4. Assertion: When a nucleus is in an excited state, it can make a transition to a lower energy state by the emission of gamma rays .

Reason: These are energy levels for a nucleus just like there are energy levels in atoms .

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: B



Watch Video Solution

5. Assertion: Binding energy per nucleon is nearly constant for element in the range $A=30$ to $A=170$.

Reason : The nuclear force between two nucleons

falls rapidly to zero as their distance is more than a few femtometres.

- A. If both assertion and reason are true and reason is the correct explanation of assertion .
- B. If both assertion and reason are true but reason is not the correct explanation of assertion .
- C. If assertion is true but reason is false .
- D. If both assertion and reason are false

Answer: A



View Text Solution

6. Assertion: Nuclear force between neutron-neutron, proton-neutron and proton-proton is approximately the same

Reason : The nuclear force does not depend on the electric charge

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: A



Watch Video Solution

7. Assertion: A free neutron is unstable

Reason : Free neutron disintegrates into proton, electron and an antineutrino i.e. $n \rightarrow p + e^{-} + \bar{\nu}$

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of

assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: A



Watch Video Solution

8. Assertion:The detection of neutrinos is extremely difficult .

Reason : Neutrinos interact only very weakly with matter.

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: a



Watch Video Solution

9. Assertion :An α -particle is emitted when uranium 238 decays into thorium

Reason : The decay of uranium 238 to thorium is represented by ${}_{92}^{238}U \rightarrow {}_{90}^{234}Th + {}_2^4He$. The helium nuclei is called an alpha particle.

- A. If both assertion and reason are true and reason is the correct explanation of assertion .
- B. If both assertion and reason are true but reason is not the correct explanation of assertion .
- C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: A

 [Watch Video Solution](#)

10. Assertion: The mass of β -particles when they are emitted is higher than the mass of electrons obtained by other means.

Reason: β -particle and electron , both are similar particles.

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: b



View Text Solution

11. Assertion: Neutrons penetrate matter more readily as compared to protons.

Reason: A neutron has no charge .

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: a



Watch Video Solution

12. Assertion: There occurs a chain reaction when uranium is bombarded with slow neutrons.

Reason: When uranium is bombarded with slow neutrons more neutrons are produced

- A. If both assertion and reason are true and reason is the correct explanation of assertion .
- B. If both assertion and reason are true but reason is not the correct explanation of assertion .
- C. If assertion is true but reason is false .
- D. If both assertion and reason are false

Answer: a



Watch Video Solution

13. Assertion: Fusion of hydrogen nuclei into helium nuclei is the source of energy of all stars.

Reason: In fusion heavier nuclei split to form lighter nuclei.

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of

assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: C



Watch Video Solution

14. Assertion: Nuclear sources will give a million times larger energy than conventional sources.

Reason: Nuclear energy sources are massive than conventional energy sources.

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: C



Watch Video Solution

15. Assertion: Naturally , thermonuclear fusion reaction is not possible on earth.

Reason : For thermonuclear fusion to take place, extreme condition of temperature and pressure are required .

A. If both assertion and reason are true and reason is the correct explanation of assertion .

B. If both assertion and reason are true but reason is not the correct explanation of assertion .

C. If assertion is true but reason is false .

D. If both assertion and reason are false

Answer: A



Watch Video Solution