



CHEMISTRY

BOOKS - DISHA CHEMISTRY (HINGLISH)

SOLUTIONS

Mcqs

1. How many grams of concentrated nitric acid solution should be used to prepare 250 mL of $2.0M HNO_3$? The concentrated acid is 70 % HNO_3

A. 90.0 g conc. HNO_3

B. 70.0 g conc. HNO_3

C. 54.0 g conc. HNO_3

D. 45.0 g conc. HNO_3

Answer: D



[View Text Solution](#)

2. For a solution of two liquids A and B it was proved that

$P_S = x_A(p_A^\circ - p_A^\circ + P^{\text{B}})$. The resulting solution will be

- A. Non-ideal
- B. ideal
- C. semi-ideal
- D. None of these

Answer: B



[View Text Solution](#)

3. If the elevation in boiling point of a solution of 10 gm of solute (mol wt. = 100) in 100 gm of water is ΔT_b , the ebullioscopic constant of water is

A. 10

B. $10\Delta T_b$

C. ΔT_b

D. $\frac{\Delta T_b}{10}$

Answer: C



[View Text Solution](#)

4. Which of the following aqueous solution will have highest depression in freezing point?

A. 0.1 M urea

B. 0.1 M Sucrose

C. 0.1M $AlCl_3$

D. 0.1M $K_4[Fe(CN)_6]$

Answer: D



[View Text Solution](#)

5. Two liquids X and Y form an ideal solution. At 300 K, vapour pressure of the solution containing 1 mol of X and 3 mol of Y is 550 mmHg. At the same temperature, if 1 mol of Y is further added to this solution, vapour pressure of the solution increases by 10 mmHg. Vapour pressure (in mmHg) of X and Y in their pure states will be, respectively:

A. 300 and 400

B. 400 and 600

C. 500 and 600

D. 200 and 300

Answer: B



[View Text Solution](#)

6. 6.02×10^{20} molecules of urea are present in 100 ml of its solution.

The concentration of urea solution is

A. $0.02M$

B. $0.01M$

C. $0.001M$

D. $0.1M$

Answer: B



[View Text Solution](#)

7. To neutralise completely 20 mL of 0.1 M aqueous solution of phosphorous acid (H_3PO_3), the value of 0.1 M aqueous KOH solution required is

A. 40 mL

B. 20 mL

C. 10 mL

D. 60 mL

Answer: A



[View Text Solution](#)

8. Two 1-litre flask A and B are connected to each other by a valve which is closed. Flask A has benzene in equilibrium with its vapours at $30^\circ C$. The flask B, is evacuated, and the valve is opened. Which of the following is true. If temperature is kept constant.'

A. Some of the benzene molecules would move to flask B from flask

A.

B. Vapour pressure will be half the initial value.

C. The vapour pressure remains unchanged

D. Some more of the liquid benzene in flask A would evaporate.

Answer: C



[View Text Solution](#)

9. Two Aqueous solutions S_1 and S_2 are separated by a semipermeable membrane. Solution S_1 has got a greater vapour pressure than solution S_2 . Water will be flowing

A. from S_1 to S_2

B. from S_2 to S_1

C. in both the directions

D. in either direction depending upon the nature of the solute

Answer: A

 [View Text Solution](#)

10. Henry's law constant of oxygen is $1.4 \times 10^{-3} \text{ mol lit}^{-1} \cdot \text{Atm}^{-1}$ at 298 K. How much of oxygen is dissolved in 100 ml at 298 K when the partial pressure of oxygen is 0.5 atm ?

A. 1.4 g

B. 3.2g

C. 22.4 g

D. 2.24 g

Answer: D

 [View Text Solution](#)

11. Which of the following liquid pairs shows a positive deviation from Raoult's law?

- A. Water- nitric acid
- B. Benzene-methanol
- C. Water - hydrochloric acid
- D. Acetone- chloroform

Answer: B



[View Text Solution](#)

12. Dissolving 120 g of urca (mol . W_i 60) in 1000 g of water gave a solution of density 1.15 g/mL the molarity of the solution is

- A. 1.78 M
- B. 2.00 M

C. 2.05 M

D. 2.22 M

Answer: C

 [View Text Solution](#)

13. The vapour pressure of a solution of the liquids A ($p^\circ = 80\text{mmHg}$ and $X_A = 0.4$) and ($p^\circ = 120\text{mmHg}$ and $X_B = 0.6$) is found to be 100 mm Hg. It shows that the solution exhibits

A. positive deviation from ideal behaviour

B. negative deviation from ideal behaviour

C. ideal behaviour

D. positive deviation for lower cone. and negative for higher cone.

Answer: B

 [View Text Solution](#)

14. The vapour pressure of two liquids X and Y are 80 and 60 torr respectively. The total vapour pressure of the ideal solution obtained by mixing 3 moles of X and 2 moles of Y would be

- A. 68 Torr
- B. 140 Torr
- C. 48 Torr
- D. 72 Torr

Answer: D



[View Text Solution](#)

15. Iodine and sulphur dissolve in

- A. water

B. ben/cnc

C. carbon disulphide

D. cthanol

Answer: C



[View Text Solution](#)

16. A 5% solution of cane sugar (molar mass 342) is isotonic with 1% of a solution of an unknown solute. The molar mass of unknown solute in g/mol is :

A. 171.2

B. 68.4

C. 34.2

D. 136.2

Answer: D



[View Text Solution](#)

17. Coolant used in car radiator is aqueous solution of ethylene glycol. In order to prevent the solution from freezing at $-0.3^{\circ}C$. How much ethylene glycol must be added to 5 kg of water ?

$$(K_f = 1.86 \text{ kg mol}^{-1})$$

A. 50 kg

B. 55 g

C. 45 kg

D. 40 g

Answer: B



[View Text Solution](#)

18. A solution of urea (mol. Mass 56 g mol^{-1}) boils at 100.18°C at the atmospheric pressure. If K_f and K_b for water are 1.86 and $0.512 \text{ K kg mol}^{-1}$ respectively, the above solution will freeze at

A. 0.654°C

B. -0.654°C

C. 6.54°C

D. -6.54°C

Answer: B

 [View Text Solution](#)

19. A solution is prepared by mixing 8.5 g of CH_2Cl_2 and 11.95 g of CHCl_3 . If vapour pressure of CH_2Cl_2 and CHCl_3 at 298 K are 415 and 200 mmHg respectively. The mole fraction of CHCl_3

in vapour form is:

(Molar mass of $Cl = 35.5 \text{ g mol}^{-1}$)

A. 0.162

B. 0.675

C. 0.325

D. 0.486

Answer: C



[View Text Solution](#)

20. If α is the degree of dissociation of Na_2SO_4 the Van't Hoff's factor

(i) used for calculating the molecular mass is

A. $1 + \alpha$

B. $1 - \alpha$

C. $1 + 2\alpha$

$$D. 1 - 2\alpha$$

Answer: D



View Text Solution

21. The molecular mass of a solute cannot be calculated by which of the following?

$$A. M_B = \frac{W_B \times RT}{\pi V}$$

$$B. M_B = \frac{p^\circ W_B M_A}{(p^\circ - p) W_A}$$

$$C. M_B = \frac{\Delta T_b W_B \times 1000}{K_b W_A}$$

$$D. M_B = \frac{K_b W_B \times 1000}{\Delta T_b \times W_A}$$

Answer: C



View Text Solution

22. We have three aqueous solutions of NaCl labelled as 'A', 'B' and 'C' with concentrations 0.1M, 0.01M and 0.001M, respectively. The value of van't Hoff factor for these solutions will be in the order _____

A. $i_A < i_B < i_C$

B. $i_A > i_B > i_C$

C. $i_A = i_B = i_C$

D. $i_A < i_B > i_C$

Answer: C



[View Text Solution](#)

23. The value of Henry's constant k_{H_2} _____.

A. greater for gases with higher solubility.

B. greater for gases with lower solubility.

C. constant for all gases.

D. not related to the solubility of gases.

Answer: B



[View Text Solution](#)

24. Which one of the following gases has the lowest value of Henry's law constant?

A. N_2

B. He

C. H_2

D. CO_2

Answer: D



[View Text Solution](#)

25. A binary liquid solution is prepared by mixing n-heptane and ethanol. Which one of the following statements is correct regarding the behaviour of the solution?

A. The solution is non-ideal, showing -ve deviation from Raoult's Law.

B. The solution is non-ideal, showing + ve deviation from Raoult's Law.

C. n-heptane shows + ve deviation while ethanol shows -ve deviation from Raoult's Law.




D. The solution formed is an ideal solution.

Answer: B



[View Text Solution](#)

26. Which one of the following salts will have the same value of van't Hoff factor (i) as that of $K_4[Fe(CN)_6]$.



Answer: A



[View Text Solution](#)

27. Relation between partial pressure and mole fraction is stated by

A. Graham's law

B. Raoult's law

C. Le-Chatelier

D. Avogadro law

Answer: B



[View Text Solution](#)

28. Which is an application of Henry's law?

A. Spray paint

B. Bottled water

C. Filling up a tire

D. Soft drinks (soda)

Answer: D



[View Text Solution](#)

29. For which of the following parameters the structural isomers C_2H_5OH and CH_3OCH_3 would be expected to have the same values ?

(Assume ideal behaviour)

- A. Boiling points
- B. Vapour pressure at the same temperature
- C. Heat of vapourization
- D. Gaseous densities at the same temperature and pressure

Answer: D



[View Text Solution](#)

30. 5 g of Na_2SO_4 was dissolved in x g of H_2O . The change in freezing point was found to be $3.82^\circ C$. If Na_2SO_4 is 81.5 % ionised, the value of x

(K_f for water = $1.86^\circ Ckgmol^{-1}$) is approximately:

(molar mass of $S = 32gmol^{-1}$) and that of $Na = 23gmol^{-1}$)

A. 15 g

B. 25 g

C. 45 g

D. 65 g

Answer: C



[View Text Solution](#)

31. The vapour pressure of acetone at $20^\circ C$ is 185 torr. When 1.2 g of a non-volatile substance was dissolved in 100 g of acetone at $20^\circ C$, its vapour pressure was 183 torr. The molar mass ($g\ mol^{-1}$) of the substance is :

A. 128

B. 488

C. 32

D. 64

Answer: D



[View Text Solution](#)

32. In mixture A and 8 components show -ve deviation as

A. $\Delta V_{mix} > 0$

B. $\Delta H_{mix} < 0$

C. A-B interaction is weaker than A-A and B-B

D. A-B interaction is stronger than A-A and B-B interaction.

Answer: B



[View Text Solution](#)

33. Which among the following will show maximum osmotic pressure?

A. $1MNaCl$

B. $1MMgCl_2$

C. $1M(NH_4)_3PO_4$

D. $1MNa_2SO_4$

Answer: C



[View Text Solution](#)

34. At $80^\circ C$, the vapour pressure of pure liquid 'A' is 520 mm Hg and that of pure liquid 'B' is 1000 mm Hg. If a mixture of solution of 'A' and 'B' boils at $80^\circ C$ and 1 atm pressure the amount of 'A' in the mixture is (1 atm = 760 mm Hg)

A. 52 mol percent

B. 34 mol percent

C. 48 mol percent

D. 50 mol percent

Answer: D

 [View Text Solution](#)

35. The observed osmotic pressure for a 0.10 M solution of $Fe(NH_4)_2(SO_4)_2$ at $25^\circ C$ is 10.8 atm. The expected and experimental (observed) values of van't Hoff factor (i) will be respectively .

A. 5 and 4.42

B. 4 and 4.00

C. 5 and 3.42





D. 3 and 5.42

Answer: A



[View Text Solution](#)

36. The freezing point of equimolal aqueous solution will be highest for



Answer: D



[View Text Solution](#)

37. If the solution boils at a temperature T_1 and the solvent at a temperature T_2 the elevation of boiling point is given by

A. $T_1 + T_2$

B. $T_1 - T_2$

C. $T_2 - T_1$

D. $T_1 + T_2$

Answer: B



[View Text Solution](#)

38. The freezing point of a 1.00 m aqueous solution of HF is found to be $-1.91^\circ C$. The freezing point constant of water, K_f is $1.86 K kg mol^{-1}$

The percentage dissociation of HF at this concentration is

A. 30 %

B. 10 %

C. 5.2 %

D. 2.7 %

Answer: D



[View Text Solution](#)

39. A solution containing 0.85 g of $ZnCl_2$ in 125.0 g of water freezes at $-0.23^\circ C$. The apparent degree of dissociation of the salt is (K_f for water = 1.86 kkgmol^{-1} , atomic mas: $Zn = 65.3$ and $Cl = 35.5$)

A. 1.36 %

B. 73.5 %

C. 7.35 %

D. 2.47 %

Answer: B



[View Text Solution](#)

40. During depression of freezing point in a solution the following are in equilibrium

- A. liquid solvent, solid solvent
- B. liquid solvent, solid solute
- C. liquid solute, solid solute
- D. liquid solute, solid solvent

Answer: A



[View Text Solution](#)

41. The molecular weight of benzoic acid in benzene as determined by depression in freezing point method corresponds to

- A. ionization of benzoic acid
- B. dimerization of benzoic acid
- C. trimerization of benzoic acid
- D. solvation of benzoic acid

Answer: B



[View Text Solution](#)

42. How many grams of methyl alcohol should be added to 10 litre tank of water to prevent its freezing at 268 K ? (K_f for water is $1.86 \text{ K kg mol}^{-1}$)

- A. 880.07g
- B. 899.04g
- C. 886.02g
- D. 868.06g

Answer: D



[View Text Solution](#)

43. The solubility of N_2 in water at 300 K and 500 torr partial pressure is $0.01gL^{-1}$. The solubility (in gL^{-1}) at 750 torr partial pressure is :

A. 0.0075

B. 0.005

C. 0.02

D. 0.015

Answer: D



[View Text Solution](#)

44. When mercuric iodide is added to the aqueous solution of potassium iodide then

- A. freezing point is raised.
- B. freezing point does not change.
- C. freezing point does not change.
- D. boiling point does not change.

Answer: A



[View Text Solution](#)

45. Azeotropic mixture of HCl and H_2 has

- A. 48% HCl
- B. 22.2 % HCl
- C. 36 % HCl

D. 20.2 % HCl

Answer: D



View Text Solution